

AD-A239 993



(2)

OFFICE OF NAVAL RESEARCH

Grant N00014-90-J-1193

TECHNICAL REPORT No. 60

Ground-State Properties of a ^3He Impurity in Liquid ^4He Monolayers

by

C. I. Um, S. K. Yoo and Thomas F. George

Prepared for publication

in

Journal of Low Temperature Physics

Departments of Chemistry and Physics
Washington State University
Pullman, WA 99164-1046



August 1991

Reproduction in whole or in part is permitted for any purpose of the United States Government.

This document has been approved for public release and sale; its distribution is unlimited.

91-09522



UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

1a. REPORT SECURITY CLASSIFICATION Unclassified				1b. RESTRICTIVE MARKINGS			
2a. SECURITY CLASSIFICATION AUTHORITY				3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited			
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE							
4. PERFORMING ORGANIZATION REPORT NUMBER(S) WSU/DC/91/TR-60				5. MONITORING ORGANIZATION REPORT NUMBER(S)			
6a. NAME OF PERFORMING ORGANIZATION Depts. Chemistry & Physics Washington State University			6b. OFFICE SYMBOL (If applicable)		7a. NAME OF MONITORING ORGANIZATION		
6c. ADDRESS (City, State, and ZIP Code) 428 French Administration Building Pullman, WA 99164-1046				7b. ADDRESS (City, State, and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217			
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Office of Naval Research			8b. OFFICE SYMBOL (If applicable)		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER Grant N00014-90-J-1193		
8c. ADDRESS (City, State, and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217				10. SOURCE OF FUNDING NUMBERS			
				PROGRAM ELEMENT NO.		PROJECT NO.	
				TASK NO.		WORK UNIT ACCESSION NO.	
11. TITLE (Include Security Classification) Ground-State Properties of a ³ He Impurity in Liquid ⁴ He Monolayers							
12. PERSONAL AUTHOR(S) C. I. Um, S. K. Yoo and Thomas F. George							
13a. TYPE OF REPORT		13b. TIME COVERED FROM _____ TO _____		14. DATE OF REPORT (Year, Month, Day) August, 1991		15. PAGE COUNT 26	
16. SUPPLEMENTARY NOTATION Prepared for publication in <u>Journal of Low Temperature Physics</u>							
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)				
FIELD	GROUP	SUB-GROUP	³ He IMPURITY LIQUID ⁴ He MONOLAYER GROUND-STATE PROPERTIES RADIAL DISTRIBUTION FUNCTION HYPERNETTED-CHAIN APPROXIMATION JASTROW WAVEFUNCTION				
19. ABSTRACT (Continue on reverse if necessary and identify by block number) We present the results of ground-state energies, radial distribution functions, liquid structure functions and effective interactions for a ³ He impurity in a ⁴ He background in two dimensions. The hypernetted-chain scheme for the system described by a Jastrow-type wavefunction is used, taking into account the triplet correlations and elementary diagrams up to fifth order. Solving the Euler-Lagrange equations for the two-body distribution functions, which contain triplet correlation and elementary diagrams, improves the results considerably. Furthermore, as a ³ He impurity is inserted into the ⁴ He background, the ground-state energy increases, but the equilibrium density decreases from 0.0350 Å ⁻² to 0.0336 Å ⁻² . The radial distribution function is broadened, while its maximum is lowered and shifted to the right (the direction of increasing radial distance) due to its larger zero-point energy, with therefore less localization of the ³ He particle. The results are compared with Monte Carlo results and other studies.							
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT ☒ UNCLASSIFIED/UNLIMITED ☒ SAME AS RPT. ☐ DTIC USERS				21. ABSTRACT SECURITY CLASSIFICATION Unclassified			
22a. NAME OF RESPONSIBLE INDIVIDUAL Dr. David L. Nelson				22b. TELEPHONE (Include Area Code) (202) 696-4410		22c. OFFICE SYMBOL	

Ground-State Properties of a ^3He Impurity in Liquid ^4He Monolayers

Chung-In Um and Sahng-Kyoon Yoo
Department of Physics
College of Science
Korea University
Seoul 136-701, Korea

Thomas F. George
Departments of Chemistry and Physics
Washington State University
Pullman, Washington 99164-1046

We present the results of ground-state energies, radial distribution functions, liquid structure functions and effective interactions for a ^3He impurity in a ^4He background in two dimensions. The hypernetted-chain scheme for the system described by a Jastrow-type wavefunction is used, taking into account the triplet correlations and elementary diagrams up to fifth order. Solving the Euler-Lagrange equations for the two-body distribution functions, which contain triplet correlation and elementary diagrams, improves the results considerably. Furthermore, as a ^3He impurity is inserted into the ^4He background, the ground-state energy increases, but the equilibrium density decreases from $0.0350 \text{ \AA}^{-2}$ to $0.0336 \text{ \AA}^{-2}$. The radial distribution function is broadened, while its maximum is lowered and shifted to the right (the direction of increasing radial distance) due to its larger zero-point energy, with therefore less localization of the ^3He particle. The results are compared with Monte Carlo results and other studies.

PACS No.: 67.60.-g, 67.60.Fp, 67.40.Db



Accession For	
NTIS GRA&I J	
DTIC TAB	
Unannounced	
Justification	
By	
Dist. Statement	
Availability	
Date	Availability
A-1	Unannounced

I. Introduction

In the past three decades there has been considerable interest in single-component Bose-fluid¹ and binary boson mixtures² at very low temperatures, such as stable bulk spin-polarized hydrogen atom and its isotope.³ Among the Bose systems, liquid ^4He and ^3He - ^4He mixtures have been analyzed successfully by introducing a Jastrow-type wavefunction. In the investigation of ground states, the Green's function Monte Carlo (GFMC) technique,⁴⁻⁵ paired-phonon analysis (PPA),⁶⁻⁷ average correlation approximation (ACA)⁸ and hypernetted-chain (HNC) approximation scheme⁹⁻¹⁰ provide powerful tools in both two and three dimensions. First, these calculations give the exact ground- and excited-state energies, but less information about the optimized correlation function and other quantities which depend on the long-range behavior, although much experimental data about them have been reported. However, in spite of its weakness in the calculation of the ground-state energy, the HNC approximation can be improved by including the triplet correlation function and some elementary diagrams¹¹ in the energy and Euler-Lagrange equations.¹²⁻¹⁷ Woo and Coldwell¹²⁻¹³ have considered the wavefunction consisting of the Jastrow function multiplied by a three-particle function and obtained the improved energy and liquid structure function. Recently, Fabrocini and Polls¹⁶ have extended this idea to the ideal boson ^3He - ^4He mixture in the zero ^3He concentration limit.

In the two-dimensional case, Miller and Woo¹⁸ have studied the ground state of a ^4He monolayer assuming no significant modifications by the existence of a substrate in the mobile limit. Chang⁷ has evaluated the ground-state energy and liquid structure function using the optimal Jastrow function determined from a self-consistent PPA, and Hatzikostantinou¹⁰ has also investigated ground-state properties by using a Jastrow-type wavefunction and solving a Euler-Lagrange equation within the HNC approximation without a three-particle factor.

The main purpose of the present paper is to evaluate the ground-state properties of two-dimensional ^4He and ^3He - ^4He mixtures in the zero ^3He concentration limit by optimizing a Jastrow-type wavefunction and solving the Euler-Lagrange equations within the HNC approximation, including a triplet correlation function and elementary diagrams up to fifth order.

In Sec. II, we summarize the construction of the Euler-Lagrange equations for zero ^3He concentration obtained from the energy minimization condition. The Jastrow wavefunction consisting of the product of the two- and three-body correlation factors and the Jackson and Feenberg energy form are adopted. We apply our results to the ^4He monolayer and the ^3He - ^4He mixture in Sec. III to obtain the ground-state energies, radial distribution functions, liquid structure functions and the effective potential, assuming a Lennard-Jones 6-12 potential. The improvements in the ground-state energies and other quantities are also discussed. In Sec. IV, we make conclusions.

II. Euler-Lagrange Equation in the Zero-Concentration Limit

In this section we consider a homogeneous binary boson mixture, which can be described the the Hamiltonian

$$\begin{aligned}
 H = & -\frac{\hbar^2}{2m_\alpha} \sum_{i=1}^{N_\alpha} \nabla_i^2 - \frac{\hbar^2}{2m_\beta} \sum_{i=1}^{N_\beta} \nabla_i^2 \\
 & + \sum_{i<j}^{N_\alpha} v_{\alpha\alpha}(r_{ij}) + \sum_{i<j}^{N_\beta} v_{\beta\beta}(r_{ij}) + \sum_{i=1}^{N_\alpha} \sum_{j=1}^{N_\beta} v_{\alpha\beta}(r_{ij}) .
 \end{aligned} \tag{2.1}$$

Here, the first two terms are kinetic energies of components α and β , respectively, and $V_{\alpha\beta}(r)$ is the pairwise interaction potential for the three different pairs of particles. We assume that the total number of particles in an area A is $N = N_\alpha + N_\beta$. We choose the Jastrow-type wavefunction in terms of two- and three-particle correlation functions of the form

$$\begin{aligned}
 \Psi(\vec{r}_{\alpha,1}, \dots, \vec{r}_{\alpha,N_\alpha}; \vec{r}_{\beta,1}, \dots, \vec{r}_{\beta,N_\beta}) = & \prod_{i<j} f_{\alpha\alpha}^{(2)}(\vec{r}_{ij}) \prod_{i<j} f_{\beta\beta}^{(2)}(\vec{r}_{ij}) \prod_{i=1}^{N_\alpha} \prod_{j=1}^{N_\beta} f_{\alpha\beta}^{(2)}(\vec{r}_{ij}) \\
 & \times \prod_{i<j<k} f_{\alpha\alpha\alpha}^{(3)}(\vec{r}_i, \vec{r}_j, \vec{r}_k) \prod_{i<j<k} f_{\beta\beta\beta}^{(3)}(\vec{r}_i, \vec{r}_j, \vec{r}_k) \prod_{i<j<k} f_{\alpha\alpha\beta}^{(3)}(\vec{r}_i, \vec{r}_j, \vec{r}_k) \prod_{i<j<k} f_{\beta\beta\alpha}^{(3)}(\vec{r}_i, \vec{r}_j, \vec{r}_k) \\
 = & \exp\left(\frac{1}{2} \left[\sum_{i<j} u_{\alpha\alpha}(|\vec{r}_{\alpha,i} - \vec{r}_{\alpha,j}|) + \sum_{i<j} u_{\beta\beta}(|\vec{r}_{\beta,i} - \vec{r}_{\beta,j}|) \right. \right. \\
 & + \sum_{i<j} u_{\alpha\beta}(|\vec{r}_{\alpha,i} - \vec{r}_{\beta,j}|) \left. \right] \exp\left(\frac{1}{6} \left[\sum_{i<j<k} u_{\alpha\alpha\alpha}(\vec{r}_{\alpha,i}, \vec{r}_{\alpha,j}, \vec{r}_{\alpha,k}) \right. \right. \\
 & + \sum_{i<j<k} u_{\beta\beta\beta}(\vec{r}_{\beta,i}, \vec{r}_{\beta,j}, \vec{r}_{\beta,k}) + \sum_{i<j<k} u_{\alpha\alpha\beta}(\vec{r}_{\alpha,i}, \vec{r}_{\alpha,j}, \vec{r}_{\beta,k}) \\
 & \left. \left. + \sum_{i<j<k} u_{\beta\beta\alpha}(\vec{r}_{\beta,i}, \vec{r}_{\beta,j}, \vec{r}_{\alpha,k}) \right] \right) , \tag{2.2}
 \end{aligned}$$

where $u_{\alpha\beta}(r)$ is the two-body correlation function which satisfies the boundary conditions

$$\lim_{r \rightarrow 0} u_{\alpha\beta}(r) \rightarrow \infty$$

$$\lim_{r \rightarrow \infty} u_{\alpha\beta}(r) = 0, \quad (2.3)$$

and $u_{\alpha\beta\gamma}(r)$ is the three-body correlation factor. The two-body distribution function can be defined as

$$g_{\alpha\beta}(r) = \frac{N_\alpha(N_\beta - \delta_{\alpha\beta})}{\rho_\alpha \rho_\beta} \frac{\int d\vec{r}(i_\alpha, i_\beta) \Psi^* \Psi}{\int d\vec{r}_1 \dots d\vec{r}_N \Psi^* \Psi}, \quad (2.4)$$

where $d\vec{r}(i_\alpha, i_\beta)$ denotes $d\vec{r}_1 \dots d\vec{r}_N$ with the exclusion of $d\vec{r}_{i_\alpha}$ and $d\vec{r}_{i_\beta}$, and $\rho_\alpha = N_\alpha/A$ is the partial density of component α . The radial distribution function $g_{\alpha\beta}^{(2)}(r)$ must also satisfy the conditions

$$\lim_{r \rightarrow \infty} [g_{\alpha\beta}(r) - 1] = 0$$

$$\rho \int d^2\vec{r} [g_{\alpha\beta}(r) - 1] + \delta_{\alpha\beta} = 0. \quad (2.5)$$

Defining $x_\alpha = \rho_\alpha/\rho$ as the concentration of component α , where $\rho = (N_\alpha + N_\beta)/A$ is the total density, we can express the energy per particle of the system as

$$E = E^{(2)} + E^{(3)}, \quad (2.6)$$

with

$$E^{(2)} = x_\alpha^2 E_{\alpha\alpha} + 2x_\alpha x_\beta E_{\alpha\beta} + x_\beta^2 E_{\beta\beta}. \quad (2.7)$$

and

$$E^{(3)} = x_\alpha^3 E_{\alpha\alpha\alpha} + 3[x_\alpha^2 x_\beta E_{\alpha\alpha\beta} + x_\alpha x_\beta^2 E_{\beta\beta\alpha}] + x_\beta^3 E_{\beta\beta\beta} \quad (2.8)$$

The two- and three-body energies are

$$E_{\alpha\beta}^{(2)} = \frac{1}{2\rho} \int d^2\vec{r} g_{\alpha\beta}^{(2)}(\vec{r}) \tilde{v}_{\alpha\beta}(\vec{r}) \quad (2.9)$$

with the effective interaction

$$\tilde{v}_{\alpha\beta}(\vec{r}) = v_{\alpha\beta}(\vec{r}) - \frac{\hbar^2}{4m_{\alpha\beta}} \nabla^2 u_{\alpha\beta}(\vec{r})$$

$$m_{\alpha\beta}^{-1} = \frac{1}{2}(m_\alpha^{-1} + m_\beta^{-1}) \quad (2.10)$$

and

$$E_{\alpha\beta\gamma}^{(3)} = \frac{1}{6\rho} \int d\vec{r} g_{\alpha\beta\gamma}^{(3)}(\vec{r}_\alpha, \vec{r}_\beta, \vec{r}_\gamma) v_{\alpha\beta\gamma}(\vec{r}) \quad (2.11)$$

with

$$v_{\alpha\beta\gamma}(\vec{r}) = \frac{\hbar^2}{2} \left[\frac{\nabla_\alpha^2}{m_\alpha} + \frac{\nabla_\beta^2}{m_\beta} + \frac{\nabla_\gamma^2}{m_\gamma} \right] u_{\alpha\beta\gamma}^{(3)}(\vec{r}_\alpha, \vec{r}_\beta, \vec{r}_\gamma) \quad (2.12)$$

where $g_{\alpha\beta\gamma}^{(3)}(\vec{r}_\alpha, \vec{r}_\beta, \vec{r}_\gamma)$ is the three-body distribution function. This is written as

$$g_{\alpha\beta\gamma}(\vec{r}_\alpha, \vec{r}_\beta, \vec{r}_\gamma) = g_{\alpha\beta}^{(2)}(r_{\alpha\beta}) g_{\gamma\alpha}^{(2)}(r_{\gamma\alpha}) g_{\gamma\beta}^{(2)}(r_{\gamma\beta}) \\ \times [f^{(3)}(\vec{r}_\alpha, \vec{r}_\beta, \vec{r}_\gamma)]^2 e^{A(\vec{r}_\alpha, \vec{r}_\beta, \vec{r}_\gamma)} \quad (2.13)$$

where $A(\vec{r}_\alpha, \vec{r}_\beta, \vec{r}_\gamma)$ is the sum of the contributions of all the Abe terms.

In order to establish a relation between $g_{\alpha\beta}^{(2)}(r)$ and $f_{\alpha\beta}^{(2)}(r)$, we introduce the HNC approximation, which consists of the equation¹

$$g_{\alpha\beta}^{(2)}(r) = [f_{\alpha\beta}^{(2)}(r)]^2 \exp[C_{\alpha\beta}(r) + T_{\alpha\beta}(r) + E_{\alpha\beta}(r)] \quad , \quad (2.14)$$

where

$$C_{\alpha\beta}(r) = \sum_i \rho_i \int d\vec{r}_i [g_{\alpha\beta}^{(2)}(r) - 1 - C_{\alpha\beta}(r)] [g_{\alpha\beta}^{(2)}(r) - 1] \quad , \quad (2.15)$$

$T_{\alpha\beta}(r)$ is the integral of the triplet function

$$T_{\alpha\beta}(r) = \sum_i \rho_i \int d\vec{r}_i [g_{\alpha\beta}^{(2)}(r_{ij}) g_{\alpha\beta}^{(2)}(r_{ik}) \{ [f_{\alpha\beta\gamma}^{(3)}(\vec{r}_i, \vec{r}_j, \vec{r}_k)]^2 - 1 \}] \quad , \quad (2.16)$$

and $E_{\alpha\beta}(r)$ is the sum of all the elementary diagrams. Rewriting Eq. (2.9) and taking $C_{\alpha\beta}(r)$ in momentum space, we get

$$\begin{aligned} E_{\alpha\beta}^{(2)} = & \frac{1}{2} \rho \int d^2\vec{r} g_{\alpha\beta}^{(2)}(r) v_{\alpha\beta}(r) - \frac{\hbar^2 \rho}{8m_{\alpha\beta}} \int d^2\vec{r} g_{\alpha\beta}^{(2)}(r) \nabla^2 [T_{\alpha\beta}(r) + E_{\alpha\beta}(r)] \\ & - \frac{\hbar^2 \rho}{8m_{\alpha\beta}} \frac{\delta_{\alpha\beta}}{(2\pi)^2 \rho_\alpha \rho_\beta} \int d^2\vec{k} k^2 [S_{\alpha\beta}(k) \{S_{\alpha\beta}(k) - 3\} - 2 \\ & + \frac{S_{\gamma\beta}(k) [1 - S_{\alpha\beta}(k)]}{D(k)}] - \frac{\hbar^2 \rho}{8m_{\alpha\beta}} \int d^2\vec{r} g_{\alpha\beta}^{(2)}(r) \nabla^2 [T_{\alpha\beta}(r) + E_{\alpha\beta}(r)] \quad , \quad (2.17) \end{aligned}$$

where $S_{\alpha\beta}(k)$ is the liquid structure function of the form

$$S_{\alpha\beta}(k) = \delta_{\alpha\beta} + 2\pi\rho \int dr r [g_{\alpha\beta}^{(2)}(r) - 1] J_0(kr) , \quad (2.18)$$

$D(k) = S_{11}S_{22} - S_{12}^2$ and $\alpha \neq \gamma$. Here $J_0(kr)$ is the zeroth-order Bessel function. To minimize the energy, we must calculate the variational equations $\delta E/\delta f^{(2)}(r) = 0$ and $\delta E/\delta f^{(3)}(r) = 0$ and solve the resultant Euler-Lagrange equations. In the presence of triplet correlations this is very complicated. To solve this problem we take the limit of zero concentration of component α and assume that $f^{(3)}(r)$ is fixed for all the triplets. Minimizing the energy, we finally get the Euler-Lagrange equations

$$\left[-\frac{\hbar^2}{m_{\alpha 2}} \nabla^2 + v_{\alpha 2}(r) + w_{\alpha 2}^0(r) + w_{\alpha 2}^T(r) + w_{\alpha 2}^E(r) \right] [g_{\alpha 2}^{(2)}(r)]^L = 0 , \quad (2.19)$$

where $\alpha = 1, 2$. Here $w_{\alpha 2}^0(r)$, $w_{\alpha 2}^T(r)$ and $w_{\alpha 2}^E(r)$ are the induced potentials due to two-body correlations, triplet correlations and elementary diagrams, respectively. $w_{\alpha 2}^0(r)$ is expressed in momentum space as

$$w_{12}^0(k) = -\frac{\hbar^2 k^2}{4m_2 \rho_2} \frac{S_{12}(S_{22} - 1)}{S_{22}^2} (2m_{12}^{-1} m_2 S_{22} + 1)$$

$$w_{22}^0(k) = -\frac{\hbar^2 k^2}{4m_2 \rho_2} \frac{(S_{22} - 1)^2 (2S_{22} + 1)}{S_{22}^2} , \quad (2.20)$$

and

$$w_{\alpha 2}^T(r) = \frac{\hbar^2}{8m_{\alpha 2}} \nabla^2 T_{\alpha 2}(r) . \quad (2.21)$$

$$w_{\alpha 2}^E(r) = \frac{\hbar^2}{8m_{\alpha 2}} \nabla^2 E_{\alpha 2}(r) . \quad (2.22)$$

Here we assume $E_{12}(r) = E_{21}(r)$. We may rewrite Eq. (2.6) for slight changes in x_1 as

$$E^{(2)} = E_{22}^{(2)} + 2x_1[E_{12}^{(2)} - E_{22}^{(2)}] + x_1^2[E_{11}^{(2)} - 2E_{12}^{(2)} + E_{22}^{(2)}] \quad (2.23)$$

Assuming that there is an impurity in the mixture, then x_1 would be $1/N$, and approximately to order $1/N$ we can minimize $E^{(2)}$ through minimization of $E_{22}(r)$, that is $g_{\alpha\beta}^{(2)}(r)$ can be estimated consecutively.

From the Euler-Lagrange equations we can construct the n -th step iteration scheme and find $g_{\alpha 2, n+1}^{(2)}(r) = g_{\alpha 2, n}^{(2)}(r) + \delta g_{\alpha 2, n}^{(2)}(r)$ for a given $g_{\alpha 2, n}^{(2)}(r)$. Then by means of the convolution theorem, the Euler-Lagrange equations can be converted to

$$\begin{aligned} & [\nabla^2 - \frac{m_{\alpha 2}}{\hbar^2}(v_{\alpha 2} + W_{\alpha 2, n}^0 + W_{\alpha 2, n}^T + W_{\alpha 2, n}^E)] \delta g_{\alpha 2, n}^{(2)}(r) \\ & + [\nabla^2 - \frac{m_{\alpha 2}}{\hbar^2}(v_{\alpha 2} - W_{\alpha 2, n}^0 - W_{\alpha 2, n}^T - W_{\alpha 2, n}^E)] g_{\alpha 2, n}^{(2)}(r) \\ & + g_{\alpha 2, n}^{(2)}(r) \int d^3r' W_{\alpha 2, n}^{0'}(r - r') g_{\alpha 2, n}^{(2)}(r') \delta g_{\alpha 2, n}^{(2)}(r') = 0 \quad (2.24) \end{aligned}$$

where $W_{\alpha 2}'$ is given by

$$\begin{aligned} W_{12}^{0'} &= \frac{k^2 [2m_{12}^{-1} m_2 S_{22}^2 + (1 - 2m_{12}^{-1} m_2) S_{22} - 1]}{S_{22}^2} \\ W_{22}^{0'} &= \frac{k^2 (S_{22}^3 - 1)}{2S_{22}^3} \quad (2.25) \end{aligned}$$

This variational calculation consists of two parts: (a) we consider a system of a background component and a trial wavefunction of the form of Eq. (2.2) and determine $g_{22}^{(2)}(r)$ variationally, and (b) we replace the background system with an impurity and introduce the trial wavefunction

$$\begin{aligned} \Psi(\vec{r}_1, \dots, \vec{r}_{n-1}, \vec{r}_n) = & \exp\left(\frac{1}{2}\left[\sum_{i<j}^{n-1} u_{22}(\vec{r}_{ij}) + \sum_{k=1}^{n-1} u_{12}(\vec{r}_{kn})\right]\right) \\ & \times \exp\left(\frac{1}{6}\left[\sum_{i<j<k}^{n-1} u_{222}(\vec{r}_i, \vec{r}_j, \vec{r}_k) + \sum_{i<j=k}^{n-1} u_{221}(\vec{r}_i, \vec{r}_j, \vec{r}_n)\right]\right) \end{aligned} \quad (2.26)$$

Through step (b), $g_{12}^{(2)}(r)$ can be determined variationally.

Equation (2.24) represents a set of integro-differential equations. Under the assumption that $\delta g_{\alpha 2}^{(2)}(r)$ vanishes outside some radius R , Eq. (2.24) can be replaced by a finite sum over a set of equally-spaced points in coordinate space, and the Laplacian term for $\delta g_{\alpha 2}^{(2)}(r)$ can also be expressed by finite differences. Then we get a set of inhomogeneous linear equations, which can be written in matrix form and diagonalized by the standard Gauss elimination technique. By making a good choice for the initial $g_{\alpha 2}^{(2)}(r)$, the rate of convergence will be rapid, so that the optimized distribution function can be readily obtained.

III. Numerical Results and Discussion

In this section we apply the results obtained in previous sections to two-dimensional ^3He and (Bose) ^3He - ^4He mixtures in the zero ^3He concentration limit. To be realistic in the two-dimensional mixture, it is necessary to consider the ^3He coverage of a few atomic layers on the ^4He background.¹⁹ At $T = 0$ K a ^3He particle behaves nearly like ^4He particle, and thus we assume that the potential $V_{\alpha\beta}(r)$

appearing in the Hamiltonian [Eq. (2.1)] are all of the form of a Lennard-Jones potential given as

$$V_{\alpha\beta}(r) = 4\epsilon[(\frac{\sigma}{r})^{12} - (\frac{\sigma}{r})^6] \quad (3.1)$$

with $\epsilon = 10.22$ K and $\sigma = 2.556$ Å. As an initial trial guess for $g_{22}^{(2)}(r)$, we adopt the results for $g_{22}^{(2)}(r)$ and $S_{22}(k)$ obtained from the Jastrow wavefunction, which are only known for $r \leq 5$ Å and $k \leq 5$ Å⁻¹, such that the short-range behavior of $g(r)$ at a density of 0.03 Å⁻² coincides with the results obtained from the PPA.⁷ The initial form of $g_{22}^{(2)}(r)$ is evaluated using an iterative scheme,¹⁰ since the long-range behavior of $g(r)$ depends on the value of $S(k)$ for small k , and the exact value of $S(k)$ for $k \geq 5$ Å⁻¹ depends on the value of $g(r)$ for small r . The initial value of $g_{12}^{(2)}(r)$ is determined by optimizing $g_{22}^{(2)}(r)$.

For the triplet correlation functions we assume that $f^{(3)}(r)$ has the same form for all the triplets and choose the parametrized form given as

$$f^{(3)}(\vec{r}_1, \vec{r}_2, \vec{r}_3) = \exp\left[\sum_{\text{cyclic}} \lambda \xi(r_{ij}) \zeta(r_{ik}) r_{ij}^2 \hat{r}_{ij} \cdot \hat{r}_{ik}\right] \quad (3.2)$$

with

$$\zeta(r) = F(r) \exp\left[-\left(\frac{r - r_c}{w}\right)^2\right] \quad (3.3)$$

$$F(r \leq b) = [(r - r_b)/r_b]^3, \quad F(r > r_b) = 0$$

Here $\hat{r}_{ij}$ and $\hat{r}_{ik}$ are unit vectors in the direction of radial vectors $\vec{r}_{ij}$ and $\vec{r}_{ik}$, respectively. We choose the parameters r_b , r_c , λ and w as those which are adopted in

the three-dimensional case, and note that the energies and radial distribution functions are not significantly affected by the choice of the above parameters. Taking the terms of elementary diagrams up to fifth order with the triplet correlation factor and approximating $E_{12}(r)$ by $E_{22}(r)$, the calculation of the integro-differential equation [Eq. (2.24)] is carried out by using the Gauss elimination technique until $r = 20 \text{ \AA}$.

The ground-state energies per particle for pure liquid ^4He and ^3He - ^4He mixtures are listed in Table 1. By considering the triplet correlation factor and elementary diagrams, the energies are lowered by about 21-96% compared with those considering the two-body factor only, and the equilibrium density increases for the pure case but decreases for the mixture case. The changes by the triplet correlation and elementary diagrams are shown in Figure 1. For the pure case (solid line), the ground-state energy is -0.646 K at the equilibrium density of $\rho = 0.0350 \text{ \AA}^{-2}$ and similar to the results of the BBGKY,¹⁸ the molecular dynamics²⁰ and the Monte Carlo studies.⁴ For the mixture case (dashed line), the equilibrium density is $0.0336 \text{ \AA}^{-2}$ with a ground-state energy -0.542 K . The rate of drifts in energies are similar to that in the three-dimensional system. Moreover, we can view the chemical potential of ^3He at $T = 0 \text{ K}$ as independent of the density of the ^4He background due to complete phase separation.⁸

We note that one of us (CIU) has evaluated the ground-state energies of one- and two-dimensional Bose liquids²¹ and charged boson systems using ring diagrams²² and the self-consistent field approximation.²³ In these results the ground-state energies of one- and two-dimensional Bose fluids consist of repulsive and attractive parts with the adoption of a soft potential with a Lennard-Jones tail. For our current two-dimensional boson system, the ground-state energy is proportional to $R_a^{-2/3}$ with an additive positive constant, where R_a is the ratio of the mean

separation between particles to the Bohr radius within the self-consistent approximation. This result is different from that of the ring diagram approximation, which does not include an additive positive constant.

Figure 2 illustrates the optimal radial distributions of $g_{22}^{(2)}(r)$ (dashed line) and $g_{12}^{(2)}(r)$ (solid line) at the respective equilibrium density, while their numerical values at various density are listed in Table 2. As the density increases, the maximum values of $g_{22}^{(2)}(r)$ and $g_{12}^{(2)}(r)$ increase. At the equilibrium density the maximum position of $g_{12}^{(2)}(r)$ shifts to the right (direction of increasing radial distance) in comparison with the fixed $g_{22}^{(2)}(r)$. This behavior is due to the fact that the volume swept by ^3He is larger than that by ^4He . The maximum value of $g_{12}^{(2)}(r)$ is smaller than that of $g_{22}^{(2)}(r)$ and broadened around the maximum point because of less localization of the ^3He impurity with a larger zero-point energy. In this sense the behavior of $g_{12}^{(2)}(r)$ agrees with the three-dimensional case.²³ With the consideration of the triplet correlation function, the maxima of $g_{12}^{(2)}(r)$ and $g_{22}^{(2)}(r)$ increase but are hardly affected by elementary diagrams. However, the ground-state energies are lowered not only by triplet factors but by elementary diagrams. We note that the optimized radial distribution functions^{21,22,25} of one-, two- and three-dimensional hard-sphere Bose systems decrease as r^{-2} , r^{-3} and r^{-4} , respectively, at short distances, while at long distances they oscillate and tend to go to unity as $r \rightarrow \infty$, which is due to the contribution from the core part of the soft potential. These three proportionalities represent the main long-distance behaviors of phonons in each dimensional Bose liquid.

Figure 3 shows the optimized liquid structure functions at various densities. The initial slope decreases, which implies a decrease of sound velocity, but the maximum increases and shifts to the right with increasing density. A comparison between $S_{12}(k)$ (solid line) and $S_{22}(k) - 1$ (dashed line) at equilibrium density is shown in Figure 4. The addition of the impurity ^3He leads to a decrease of the

maximum and shift of the maximum position to the right in k -space. This behavior obtained is as expected in a scattering problem and agrees with that of the three-dimensional case²⁵ except in the region of $k \rightarrow 0$.

In Figure 5 we plot the bare Lennard-Jones potential V_{LJ} , the induced potential $W_{12}(r)$ (V_{ind}) and the effective potential V_{eff} . According to Table 3, we observe for the effective potential that as the density increases, its hard core radius ($V_{eff} = 0$) decreases and potential becomes deeper, while the repulsive barrier height grows at about the same position.

IV. Conclusions

We have investigated the ground-state properties and behaviors of the ^3He - ^4He system. The introduction of the triplet correlation and elementary diagrams up to fifth order in the HNC approximation leads to successful improvement in the ground-state energy and optimized two-body distribution functions. Furthermore, the triplet correlation factor contributes to the change of both energy and the distribution function, while elementary diagrams do not change the distribution function. By addition of an impurity ^3He into a ^4He background, the energy increases, the distribution function moves to the right, and its maximum decreases and is broadened around the maximum position, due to larger zero-point energy and less localization of the ^3He particle.

Acknowledgments

This research was supported by a Grant to Korea University from Korea Science and Engineering Foundation under Contract No. 891-0203-013-2, and by the U.S. Office of Naval Research.

References

1. K. Hiroike, *Prog. Theor. Phys.* 27, 342 (1962); L. J. Lantto, A. D. Jackson and P. J. Siemens, *Phys. Lett. B* 68, 311 (1977); *Nucl. Phys. A* 317, 70 (1979).
2. T. Chakraborty, *J. Low Temp. Phys.* 48, 151, (1982).
3. I. F. Silvera and J. T. M. Walraven, *Phys. Rev. Lett.* 44, 164 (1980); J. T. M. Walraven, I. F. Silvera and A. P. M. Matthey, *Phys. Rev. Lett.* 45, 449 (1980); *J. Phys.* 41, C7-137 (1980).
4. D. S. Hyman, Report No. 1273, Material Science Center, Cornell University (1970), unpublished; K. S. Liu, M. H. Karlos and G. V. Chester, *Phys. Rev. B* 13, 1971 (1976).
5. P. A. Whitlock and G. V. Chester, *Phys. Rev. B* 38, 2418 (1988).
6. C. E. Campbell, *Ann. Phys. (N.Y.)* 74, 43 (1980).
7. C. C. Chang, *J. Low Temp. Phys.* 38, 691 (1980).
8. M. D. Miller, *Ann. Phys. (N.Y.)* 127, 367 (1980); *Phys. Rev. B* 17, 1139 (1978); *Phys. Rev. B* 18, 4730 (1979).
9. T. Chakraborty, *Phys. Rev. B* 25, 3177 (1982).
10. P. Hatzikonstantinou, *J. Phys. C* 18, 2393 (1985), and private communication.
11. R. A. Smith, *Phys. Lett.* 85 B, 183 (1979).
12. C. W. Woo, *Phys. Rev. Lett.* 28, 1442 (1972).
13. C. W. Woo and R. L. Coldwell, *Phys. Rev. Lett.* 29, 1062 (1972).
14. P. Berdahl, *Phys. Rev. A* 10, 2378 (1974).
15. V. R. Pandharipande, *Phys. Rev. B* 18, 218 (1978).
16. A. Fabrocini and A. Polls, *Phys. Rev. B* 30, 1200 (1984).
17. E. Krotscheck, *Phys. Rev. B* 33, 3158 (1986).
18. M. D. Miller and C. W. Woo, *Phys. Rev. A* 6, 1942 (1972).
19. C. E. Campbell and M. Schick, *Phys. Rev. A* 3, 691 (1971).

20. A. Isihara and C. I. Um, *Phys. Rev. B* 19, 5725 (1979); C. I. Um, W. H. Kahng and E. S. Yim, *J. Korean Phys. Soc.* 16, 120 (1983); *ibid.* 17, 75 (1984); C. I. Um., W. H. Kahng, E. S. Yim and K. T. Cheong, *Phys. Rev. B* 29, 6626 (1984).
21. C. I. Um, A. Isihara and S. T. Choh, *J. Korean Phys. Soc.* 14, 39 (1981).
22. C. I. Um, W. H. Kahng, S. M. Kahng, S. M. Hong and E. S. Yim, *J. Korean Phys. Soc.* 22, 32 (1989); *Phys. Rev. B*, in press.
23. W. E. Massey, C. W. Woo and H. T. Tan, *Phys. Rev. A* 1, 519 (1970); M. L. Ristig, S. Fantoni and K. E. Kurten, *Z. Phys. B* 51, 1 (1983).
24. A. Isihara and D. W. Jepsen, *Phys. Rev.* 158, 112 (1967).
25. M. B. Yim and W. E. Massey, *Phys. Rev. B* 19, 3529 (1979); A. Fabrocini and A. Polls, *Phys. Rev. B* 26, 1438 (1982).

Table 1. Ground-state energies E_0 due to the two-body factor only (2), two- and three-body factors (2+3) and two- and three-body factors plus elementary diagrams (tot) for the pure and mixture cases at various densities.

ρ ($\text{\AA}^{-2}$)	$E_{0,\text{mix}}^{(2)}$ (K)	$E_{0,\text{mix}}^{(2+3)}$ (K)	$E_{0,\text{mix}}^{(\text{tot})}$ (K)	$E_{0,\text{pure}}^{(2)}$ (K)	$E_{0,\text{pure}}^{(2+3)}$ (K)	$E_{0,\text{pure}}^{(\text{tot})}$ (K)
0.0250	-0.28	-0.38	-0.45	-0.39	-0.45	-0.50
0.0275	-0.30	-0.40	-0.48	-0.44	-0.48	-0.55
0.0300	-0.32	-0.43	-0.50	-0.48	-0.53	-0.58
0.0325	-0.35	-0.45	-0.53	-0.46	-0.56	-0.62
0.0350	-0.38	-0.47	-0.54	-0.46	-0.59	-0.65
0.0375	-0.35	-0.45	-0.52	-0.43	-0.58	-0.63
0.0400	-0.33	-0.44	-0.51	-0.40	-0.57	-0.61

Table 2. Radial distribution functions at various densities.

	$\rho = 0.025 \text{ \AA}^{-2}$		$\rho = 0.030 \text{ \AA}^{-2}$		$\rho = 0.035 \text{ \AA}^{-2}$	
$r(\text{\AA})$	$g_{22}(r)$	$g_{12}(r)$	$g_{22}(r)$	$g_{12}(r)$	$g_{22}(r)$	$g_{12}(r)$
2.0	0.000	0.000	0.000	0.000	0.000	0.000
2.2	0.010	0.010	0.010	0.011	0.010	0.011
2.4	0.067	0.065	0.070	0.066	0.071	0.069
2.6	0.208	0.207	0.211	0.209	0.215	0.209
2.8	0.409	0.404	0.418	0.408	0.425	0.413
3.0	0.628	0.608	0.640	0.621	0.656	0.630
3.2	0.835	0.801	0.839	0.815	0.841	0.828
3.4	0.988	0.960	0.992	0.963	0.997	0.970
3.6	1.086	1.055	1.095	1.066	1.101	1.075
3.8	1.140	1.109	1.159	1.117	1.160	1.126
4.0	1.188	1.140	1.189	1.155	1.198	1.160
4.2	1.181	1.145	1.182	1.158	1.190	1.162
4.4	1.169	1.131	1.168	1.135	1.173	1.140
4.6	1.121	1.095	1.121	1.099	1.123	1.103
4.8	1.084	1.077	1.084	1.078	1.085	1.075
5.0	1.050	1.048	1.043	1.043	1.043	1.040
5.2	1.015	1.019	1.015	1.018	1.010	1.011
5.4	0.990	0.996	0.987	0.992	0.986	0.990
5.6	0.972	0.978	0.969	0.974	0.967	0.972
5.8	0.957	0.963	0.956	0.961	0.955	0.959
6.0	0.951	0.956	0.948	0.953	0.947	0.952

Table 3. Induced potentials V_{ind} and effective potentials V_{eff} at various densities.

	$\rho = 0.025 \text{ \AA}^{-2}$		$\rho = 0.030 \text{ \AA}^{-2}$		$\rho = 0.35 \text{ \AA}^{-2}$	
$r(\text{\AA})$	$V_{\text{ind}}(\text{K})$	$V_{\text{eff}}(\text{K})$	V_{ind}	V_{eff}	V_{ind}	V_{eff}
2.4	-1.056	26.332	-1.625	25.763	-2.401	24.987
2.6	-0.756	-4.347	-1.162	-4.754	-1.752	-5.344
2.8	-0.480	-10.447	-0.942	-10.709	-1.159	-11.126
3.0	-0.226	-9.882	-0.367	-10.023	-0.623	-10.279
3.2	0.009	-7.850	-0.027	-7.887	-0.141	-8.000
3.4	0.231	-5.816	0.290	-5.758	0.302	-5.744
3.6	0.437	-4.129	0.579	-3.987	0.702	-3.863
3.8	0.607	-2.828	0.812	-2.624	1.024	-2.412
4.0	0.757	-1.828	1.026	-1.568	1.315	-1.279
4.2	0.950	-1.021	1.279	-0.692	1.643	-0.329
4.4	1.216	-0.294	1.637	0.126	2.086	0.575
4.6	1.546	0.378	2.052	0.884	2.588	1.421
4.8	1.828	0.917	2.364	1.453	2.964	2.053
5.0	1.885	1.168	2.359	1.642	2.960	2.244
5.2	1.625	1.057	1.968	1.490	2.501	1.932
5.4	1.123	0.669	1.328	0.874	1.745	1.291
5.6	0.576	0.210	0.694	0.328	0.995	0.629
5.8	0.154	-0.143	0.249	-0.048	0.466	0.168
6.0	-0.088	-0.331	-0.022	-0.221	0.191	-0.052

Figure Captions

1. Ground-state energies of the pure ^4He system (solid line) and (Bose) ^3He - ^4He mixture in the impurity ^3He limit (dashed line) as a function of density. $E(2)$, $E(2+3)$ and $E(\text{tot})$ represent the energy containing the two-body correlation factor only, energy by two- and three-body factors, and total energy by the above factors plus elementary diagrams, respectively.
2. Optimized radial distribution functions $g_{12}^{(2)}(r)$ (solid line) and $g_{22}^{(2)}(r)$ (dashed line) at equilibrium density.
3. Optimized liquid structure function $S_{12}(k)$ at various densities.
4. Optimized liquid structure function $S_{12}(k)$ (solid line) and $S_{22}(k)-1$ (dashed line) at the equilibrium density.
5. Bare Lennard-Jones potential (V_{LJ}), induced potential (V_{ind}) and effective potential (V_{eff}) at the equilibrium density.

Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5

