

AD-A241 993



PAGE

Form Approved
OMB No. 0704-0188Public reporting burden
gathering and
collection of information
about this document
through this form
is estimated to average
1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, reviewing and completing the form, and reviewing the instructions. Send comments regarding this burden estimate or any other aspect of this form, including suggestions for reducing the burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Road, Suite 1204, Washington, DC 20540.

per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, reviewing and completing the form, and reviewing the instructions. Send comments regarding this burden estimate or any other aspect of this form, including suggestions for reducing the burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Road, Suite 1204, Washington, DC 20540.

1. AGENCY USE ONLY (Leave blank)

3. REPORT TYPE AND DATES COVERED

Final Report 01Mar86/28Feb91

4. TITLE AND SUBTITLE

A Study of Gamma-Ray generation from channeled
electrons and positions

5. FUNDING NUMBERS

F49620-86-K-0015

6. AUTHOR(S)

Dr Pantell

7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)

Sanford University
Stanford California 943058. PERFORMING ORGANIZATION
REPORT NUMBER

AFOSR-TR-91-0843

9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)

AFOSR/NP
Bldg 410
Bolling AFB DC 20332-6448
Dr Barker10. SPONSORING/MONITORING
AGENCY REPORT NUMBER

2301/A8

11. SUPPLEMENTARY NOTES

DTIC
ELECTE
OCT 22 1991
S B D

12a. DISTRIBUTION / AVAILABILITY STATEMENT

UNLIMITED

DISTRIBUTION STATEMENT A

Approved for public release
Distribution Unlimited

12b. DISTRIBUTION CODE

13. ABSTRACT (Maximum 200 words)

Operating at peak current levels 10(13) times greater than previously attempted, channeling radiation photon fluxes of 10(19) photons/sr-keV-sec over a picosecond duration at a wavelength of 0.42 angstrom were observed. The experimental efforts were hampered by the shutdown of John Madey's storage ring accelerator at Stanford. However the experiments located at the Naval Postgraduate School, the Saskatchewan Accelerator Laboratory and the Institute for Atomic Physics at Darmstadt (Germany). Thusfar, four publications and one PhD thesis have resulted directly from this work.

91-13770



91 10 22 051

14. SUBJECT TERMS

15. NUMBER OF PAGES

16. PRICE CODE

17. SECURITY CLASSIFICATION
OF REPORT

UNCLASS

18. SECURITY CLASSIFICATION
OF THIS PAGE

UNCLASS

19. SECURITY CLASSIFICATION
OF ABSTRACT

UNCLASS

20. LIMITATION OF ABSTRACT

UL

Final Report

Contract No. AF-F49620-86-K-0015

Object: To investigate the generation of radiation from channeled particles.

This project was intended to coordinate with the construction of John Madey's linear accelerator and positron storage ring. On the Mark III linac we assembled a channeling radiation beamline which included a goniometer, focussing and steering magnets, vacuum pumps, TV cameras, and retractable screens for e-beam viewing.

The primary purpose of the experiments on the linac were to demonstrate that channeling radiation can be an inexpensive source of bright, hard x-rays with picosecond duration. Channeled particle trajectories are similar to the trajectories in a magnetic wiggler, but the equivalent magnetic field would have to be about ten megagauss. Indeed, we measured⁽¹⁾ a photon flux of 10^{19} photons/sr-keV-sec over a picosecond duration at a wavelength of $0.42 \text{ \AA}$. Our peak current levels were $\approx 10^{13}$ times greater than the currents used in previous channeling experiments and average currents were $\approx 10^8$ times greater.

To perform these measurements we had to develop a spectrometer capable of operating at high photon fluxes with several percent energy resolution at x-ray wavelengths, and in a bremsstrahlung background. This was accomplished using a graphite crystal Bragg reflector, a photomultiplier detector, and specially designed apertures⁽²⁾.

During the course of this project the Madey accelerator was shut down, but we continued the experimental work on accelerators located at the Naval Postgraduate School in Monterey, at the Saskatchewan Accelerator Laboratory in Saskatoon, Canada, and the accelerator at the Institut für Keruphysik in Darmstadt, Germany.

Thusfar, four publications have resulted from these efforts ⁽¹⁻⁴⁾, along with one Ph.D thesis⁽⁵⁾. It is anticipated that there will be one more publication and one more Ph.D. thesis resulting from this project.

In addition to the high power channeling radiation research, we also studied channeling radiation in superlattices, both theoretically, and experimentally at Darmstadt.

Superlattices offer a means for increasing the radiation, utilizing the periodicity of the layers.

The Madey storage ring was to provide a bright positron current source for seeking x-ray laser action by means of channeling radiation. A current density of $10^7 - 10^8 \text{ A/cm}^2$ could be obtained, which would provide significant stimulated gain over a picosecond time interval. (The time duration is determined by the interval over which the crystal remains intact.) Unfortunately, the storage ring was never constructed so that this portion of the program could not be carried out.

References:

1. C.K. Gary, A.S. Fisher, R.H. Pantell, J. Harris, and M.A. Piestrup, "Channeling of Electrons in Si Produces Intense Quasimonochromatic, Tunable, Picosecond x-ray Bursts", Phys. Rev. B. No1, **42**, 7 (1990).
2. C.K. Gary, A.S. Fisher, R.H. Pantell, J. Harris, M. Piestrup, "The measurement of channeling radiation from a linear-accelerator electron beam at high currents" Nuclear Instruments and Methods in Physics Research B51, 458-465 (1990).
3. J.O. Kephart, B.L. Berman, R.H. Pantell, S. Datz, R.K. Klein, H. Park, "Thermal vibration amplitudes of silicon determined by channeling-radiation measurements," Physical Review B, **44**, No. 6, 1992-2002, 1990.
4. C.K. Gary, R.H. Pantell, M. Ozcan, M.A. Piestrup, D.G. Boyers, "Optimization of the Channelling Radiation Source Crystal to Produce Intense Quasimonochromatic X-Rays", Journal of Applied Physics, Sept. 15, 1991.
5. C.K. Gary, "Intense Channeling Radiation as an X-ray Source," & Stanford University, Ph.D. Thesis in the Department of Electrical Engineering, September, 1990.



Accession For	
NTIS GRA&I	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	