

AD-A251 110



ENTATION PAGE

Form Approved
OMB No. 0704-0188

2

estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering the collection of information, Send comments regarding this burden estimate or any other aspect of this estimate, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Avenue, Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE May 1992	3. REPORT TYPE AND DATES COVERED Technical Rep. (Interim) 7/1/91-6/30/92	
4. TITLE AND SUBTITLE Ultra-Fast Solvation Dynamics in a Stockmayer Fluid			5. FUNDING NUMBERS Grant N00014-89-J-1169 RT 4131036	
6. AUTHOR(S) L. Perera and M. Berkowitz				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of North Carolina Department of Chemistry CB# 3290 Chapel Hill, NC 27599			8. PERFORMING ORGANIZATION REPORT NUMBER Technical Report #10	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Office of Naval Research 800 N. Quincy Street Arlington, VA 22217-5000 (Code 1113PS, P.P. Schmidt)			10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES Submitted to the Journal of Chemical Physics				
12a. DISTRIBUTION/AVAILABILITY STATEMENT This document has been approved for public release and sale; its distribution is unlimited.			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) SDTIC ELECTE JUN 03 1992 A D				
14. SUBJECT TERMS Stockmayer fluid, ultra-fast solvation			15. NUMBER OF PAGES	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT	

OFFICE OF NAVAL RESEARCH

Grant N00014-89-J-1169

R&T Code 4131036

Technical Report # 10

" Ultra-Fast Solvation Dynamics in a Stockmayer Fluid."

by

L. Perera and Max L. Berkowitz

Submitted to the
Journal of Chemical Physics

The University of North Carolina at Chapel Hill
Department of Chemistry
Chapel Hill, NC 27599-3290

MAY 1992



Accession For	
NTIS	CRA&I ☒
DTIC	TAB ☐
Unannounced	☐
Justification	
By	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
A-1	

Reproduction in whole or in part is permitted for any purpose of the United States Government.

This document has been approved for public release and sale; its distribution is unlimited.

92-14400

92 6 01 069

May 1992
J. Chem. Phys. (Submitted)

ULTRA-FAST SOLVATION DYNAMICS IN A STOCKMAYER FLUID

Lalith Perera and Max L. Berkowitz

Department of Chemistry, University of North Carolina, Chapel Hill, North Carolina, 27599

Abstract: Computer simulations of charge solvation dynamics in Stockmayer solvent display the existence of two time regimes. In this communication we show that the short time regime is due to inertial independent motion of solvent particles.

Dynamics of charge (or charge distribution) solvation is a subject of considerable interest [1]. Recent molecular dynamics computer simulation studies revealed that solvent relaxation occurs in two time regimes [2-7]. During the initial regime, which can be described by a Gaussian function, the relaxation occurs very fast (25-150 fs). The fast regime contributes about 60-80 % towards the total solvent relaxation and the motion of the solvent molecules in this regime has an inertial character. The short time regime is followed by a long time regime, which can be described by an exponential function. The motion of solvent molecules in this regime has a diffusional character. The two phase behavior of the solvent relaxation was observed in MD simulations performed in solvents such as water[2], acetonitrile[3], methanol[4] and methyl chloride [5]. Very recently the two regime behavior of solvent relaxation was also observed in MD simulations performed in simple polar fluids described by a Stockmayer potential [6-7]. The presence of two time regimes in the solvation dynamics was also confirmed by the experiment [8].

While the existence of the two regimes in the relaxation is firmly established, the nature of the solvent dynamics in these regimes is not completely clear. Thus for example, using the results of his simulations in acetonitrile, Maroncelli has assigned the fast regime to a small amplitude independent, inertial (mostly rotational) motion of molecules in the first solvation shell [3]. Contrary to this assignment Bagchi and Chandra recently proposed that the short time decay of the relaxation in simple Stockmayer solvents can be described in terms of collective excitation [9].

To separate between these two different proposed mechanisms we performed one extra set of non-equilibrium computer simulations in addition to our previous simulations [6] where we studied the dynamics of the Stockmayer solvent following a sudden change in the solute charge. In our previous study we performed three Non-Equilibrium Simulation sets NESI, NESII and NESIII in which we followed the time development of the normalized response function $S(t)$ defined as

$$S(t) = \frac{\delta E(t) - \delta E(\infty)}{\delta E(0) - \delta E(\infty)} \quad (1)$$

In equation (1) $\delta E(t)$ is the fluctuation in the solvent-solute interaction energy at a given time t .

To study the effect of the rotational and translational degrees of freedom of the solvent we froze the translational motion in the first simulation (NESI). In NESII both rotational and translational motion participated in the relaxation. The Stockmayer solvent in our simulations was characterized (in reduced units) by a dipole moment $(\mu^*)^2 = 3.0$, a moment of inertia $I^* = 0.025$, a density $\rho^* = 0.822$ and a temperature $T^* = 1.15$. More details on the simulations were given in ref. [6]. In NESIII we changed the moment of inertia of the solvent in order to study the relative importance of the translational and rotational modes.

The additional simulation set performed now (NESIV) consisted of 50 trajectories which had the same starting configurations as the trajectories from the set NESI. The difference between the trajectories in NESIV set and NESI set is that the molecular dynamics in NESIV set was performed with no solvent-solvent interactions. Figure 1 displays the behavior of the relaxation function $S(t)$ obtained from the sets NESI, NESII and NESIV. As we can see from this figure, up to ~ 0.09 ps (at this time the energy relaxed $\sim 70\%$) the decay functions obtained from NESI and NESIV are practically indistinguishable. We also know from our previous work and from the other simulations that the main contribution into relaxation comes from the rotational motion of molecules in the first solvation shell (the difference in $S(t)$ for the NESI and NESII is very small). Therefore from figure 1 we conclude that the initial relaxation in our system is due to the independent inertial rotational motion of solvent molecules.

To find out how large in amplitude is this rotational motion we present the average value of the rotational angle as a function of time in figure 2. As we can see from this figure, after 0.09 ps the average displacement angle for the dipoles in the first solvation shell is $\sim 45^\circ$, which is twice as large as the average displacement angle for the Stockmayer dipole in a sample of pure solvent under the same conditions. Also note that the average displacement angle for the dipoles in secondary shells around the ion is $\sim 25^\circ$, the same value as in pure solvent. Due to the acceleration the average angular displacement of the first shell dipoles grows quickly and reaches its maximum value of $\sim 120^\circ$ overshooting the equilibrium value of $\sim 85^\circ$. While the angular motion of the first shell dipoles is underdamped, the secondary shell dipoles display an overdamped motion.

The picture of the dynamic solvation of the ion in Stockmayer fluid that emerges from

our simulation is rather simple. Once the charge on the solute is turned on, the solvent dipoles of the first shell rush towards a favorable orientation along the field emanating from the ion. The motion of the solvent particles has an inertial independent character. The solvent particles practically do not see each other during the short time regime (~ 0.1 ps) and only after its completion the interaction between particles starts influencing their motion. At this time the motion becomes correlated. At long time scale the motion has diffusional character and this is reflected in a slow exponential decay of the relaxation function. Finally we want to mention that this picture of the solvent dynamics of the Stockmayer solvent is probably correct when a large perturbation ($\Delta q = 1.0$) is applied to the solute.

We thank Prof. B. Bagchi for providing us with the manuscript of ref. [9] prior to its publication. This work was supported by the Office of Naval Research.

References

- [1] For recent reviews and references therein see, for example: P.F. Barbara and W. Jarzeba, *Adv. Photochem*, **15**, 1 (1990); M. Maroncelli, J. MacInnis, and G.R. Fleming, *Science*, **243**, 1674 (1989); B. Bagchi, *Annu. Rev. Phys. Chem.* **40**, 115 (1989); B. Bagchi, and A. Chandra, *Adv. Chem. Phys.* **LXXX**, 1 (1991); J.D. Simon, *Acc. Chem. Res.* **21**, 128 (1988).
- [2] M. Maroncelli, and G.R. Fleming, *J. Chem. Phys.* **89**, 5044 (1988).
- [3] M. Maroncelli, *J. Chem. Phys.* **94**, 2084 (1991).
- [4] T. Fonseca, and B.M. Ladanyi, *J. Phys. Chem.* **95**, 2116 (1991).
- [5] E.A. Carter, and J.T. Hynes, *J. Chem. Phys.* **94**, 5961 (1991).
- [6] L. Perera and M. Berkowitz, *J. Chem. Phys.* **96**, 3092 (1992).
- [7] E. Neria and A. Nitzan, *J. Chem. Phys.* **96**, 5433 (1992).
- [8] S.J. Rosenthal, X. Xie, M. Du, and G.R. Fleming, *J. Chem. Phys.* **95**, 4715 (1991).
- [9] B. Bagchi and A. Chandra "Ultra-fast Solvation Dynamics: Molecular Explanation of Computer Simulation Results in a Simple Dipolar Solvent" (preprint)

Figure captions.

Fig 1. Response function $S(t)$ obtained from: NESII (solid line), NESI (dashed line) and NESIV (dotted line).

Fig 2. Average orientational angle of the dipole in the: first solvation shell (solid line), second solvation shell (dashed line) and third solvation shell (dotted line).




Fig 1




Fig. 2

TECHNICAL REPORT DISTRIBUTION LIST - GENERAL

Office of Naval Research (2)*
Chemistry Division, Code 1113
800 North Quincy Street
Arlington, Virginia 22217-5000

Dr. James S. Murday (1)
Chemistry Division, Code 6100
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. Robert Green, Director (1)
Chemistry Division, Code 385
Naval Air Weapons Center
Weapons Division
China Lake, CA 93555-6001

Dr. Elek Lindner (1)
Naval Command, Control and Ocean
Surveillance Center
RDT&E Division
San Diego, CA 92152-5000

Dr. Bernard E. Douda (1)
Crane Division
Naval Surface Warfare Center
Crane, Indiana 47522-5000

Dr. Richard W. Drisko (1)
Naval Civil Engineering
Laboratory
Code L52
Port Hueneme, CA 93043

Dr. Harold H. Singerman (1)
Naval Surface Warfare Center
Carderock Division Detachment
Annapolis, MD 21402-1198

Dr. Eugene C. Fischer (1)
Code 2840
Naval Surface Warfare Center
Carderock Division Detachment
Annapolis, MD 21402-1198

Defense Technical Information
Center (2)
Building 5, Cameron Station
Alexandria, VA 22314

* Number of copies to forward