

AD-A254 978

TION PAGE 1

Form Approved
GAS No. 0704-0100

Notice: How are you using this form for reporting information? Reporting covering data received, the collection of information, and comments regarding the current status of any other aspect of the Washington Headquarters System, Directorate for Information Operations and Reports, 1215 Jefferson Management and Budget, Performance Assessment Project (704-0100), Washington, DC 20503.

1. REPORT DATE		2. REPORT TYPE AND DATES COVERED	
		ANNUAL 15 Jan 91 TO 14 Jan 92	
3. TITLE AND SUBTITLE		5. FUNDING NUMBERS	
GPS Measurements at Vandenberg AFB		61102F AFOSR-90-0339 2309 AS	
4. AUTHOR(S)			
Dr Robert King			
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)		8. PERFORMING ORGANIZATION REPORT NUMBER	
MASSACHUSETTS INSTITUTE OF TECHNOLOGY Dept of Earth, Atmospheric, and Planetary Sciences Cambridge, MA 02139-4307		AFOSR-TR-92 0822	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)		10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
Dr Dickinson AFOSR/NL Building 410 Bolling AFB DC 20332-6448			
11. SUPPLEMENTARY NOTES			
DTIC ELECTE S A D AUG 27 1992			
12a. DISTRIBUTION/AVAILABILITY STATEMENT		12b. DISTRIBUTION CODE	
Approved for public release; distribution unlimited			
13. ABSTRACT (Maximum 200 words)			
The objective was to measure directly the tectonic deformation in the Santa Maria Fold and Thrust Belt (SMFTB), northwest of Santa Barbara California. The principal measurement technique was space geodesy; using microwave signals transmitted by Block II satellites of the Global Positioning System (GPS), from March to August 1990. Geodetic observations were analyzed to resolve tectonic deformation across the SMFTB. The geodetic network forms a braced quadrilateral with 40 km sides whose southwest corner is the Vandenberg very long baseline interferometry (VLBI) station. Three different types of data were combined to estimate two-dimensional station position and strain rate parameters simultaneously. Significant strain rates were discovered using a model which constrains the relative velocity field to be linear in space and constant in time. The maximum compressive strain is oriented N17E±5E, and the compressive strain rate in that direction is 0.13±0.03 strain/yr. Under the assumption that the unresolved rotational component of the velocity field is zero, the integrated rate of deformation across the basin was estimated at 7±1 mm/yr oriented at N3E±13. This vector can be decomposed into 6±2 mm/yr of crustal shortening on the general structural trend of N30E and 3±1 mm/yr of right-lateral shear across the axis. On the basis of these values and earthquake			
14. SUBJECT TERMS		15. NUMBER OF PAGES	
16. SECURITY CLASSIFICATION OF THIS PAGE		17. SECURITY CLASSIFICATION OF ABSTRACT	
(U)		(U)	
18. SECURITY CLASSIFICATION OF REPORT		19. LIMITATION OF ABSTRACT	
(U)		(U)	

focal mechanisms in the area, deformation was inferred on northwest trending folds and thrusts within the belt. These results are consistent with the rate and direction of deformation across the central California Coast Ranges inferred by balancing Pacific-North America plate motion against San Andreas slip and Basin and Range extension. They imply that the SMFTB is the primary active element in transforming motion from the Coast Ranges to the western Transverse Ranges and the Santa Barbara Channel. Five years of GPS measurements were used to measure the relative velocities of five stations in the SMFTB. When considered with respect to the GPS mark nearest the Vandenberg VLBI station, the horizontal velocities of three stations are significantly different from zero at the 95% confidence level. The velocities indicate the importance of N-S compressional deformation in the area, as indicated by geologic cross sections, seismicity, earthquake focal mechanisms, and borehole breakout data. The compressional components of the three most significant velocities are grossly consistent with the deformation predicted by a simple dislocation model involving two thrust faults in the SMFTB, in addition to deep slip on the San Andreas fault.

**Investigation of Crustal Deformation in the Vicinity of
Vandenberg Air Force Base**

**Grant AFOSR-90-0339
(MIT OSP No. 74324)**

Annual Technical Report

for the period

22 August 1990 - 31 August 1991

Submitted to

Air Force Office of Scientific Research

**James F. Lewkowicz
Program Manager
PL/LWH Hanscom AFB MA 01731-5000**

**Capt Helen R. Tyson
Contracting Officer
AFOSR/PKZA
Bolling AFB, DC 20332-6448**

by

**Robert W. King
Principal Investigator
Department of Earth, Atmospheric, and Planetary Sciences
Massachusetts Institute of Technology
Cambridge, MA 02139**

30 September 1991

Accession For	
NTIS CRA&I	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
A-1	

DTIC QUALITY INSPECTED 3

92-23768
5pgs
404784

1 82 8 26 078

11 AUG 1992

INTRODUCTION

Recent geological and geodetic studies have suggested that the region surrounding Vandenberg AFB is undergoing active crustal deformation, with important implications for both the geodetic stability and the seismogenic potential of the Western Test Range (WTR) [Feigl *et al.*, 1990]. Part of the evidence for significant deformation was obtained from GPS measurements over a broad area of central and southern California which we carried out in cooperation with other university and government scientists from 1987 through 1991. Although useful in defining the regional tectonic setting, these measurements are of insufficient spatial and temporal density to answer many important questions about the seismogenic potential of Vandenberg.

In 1989 we received funding from AFOSR under the Commander's Reserve Fund (Grant AFOSR-89-0400), with matching funds from MIT, to purchase five GPS receivers and to begin a series of measurements designed to determine the magnitude and spatial distribution of deformation in a region encompassing the major faults and folds within 50 km of Vandenberg. The present grant (AFSOF 90-0339) supports a follow-on, 3-yr study including four major elements:

- Remeasurement and extension of the GPS network
- Analysis of the GPS data to determine horizontal strains with a precision of 5×10^{-8} /yr within the network
- Compilation of the historical geodetic data available for the area
- Synthesis of the geodetic, geological, and seismological data to assess the tectonic setting and seismogenic potential of Vandenberg

The pre-grant studies of Vandenberg tectonics and our progress during the first year of this grant have been completely described in the doctoral thesis of Kurt L. Feigl, completed at M. I. T. in September 1991 and attached to this report. We give here only an overview of our work, along with reference to the appropriate chapters of the thesis [Feigl, 1991].

GPS MEASUREMENTS

Since February, 1990, we have carried out four GPS experiments involving the stations of the Vandenberg network. Two of these experiments (February and September 1990) included most or all of the current network; the other two (March 1990 and February 1991) included two or three Vandenberg stations as part of a remeasurement of the regional central and southern California networks. The details of these experiments are given in Chapter 4 of Feigl [1991], especially Figures 4.1–4.3, Tables 4.1–4.6, and the text on pp. 124–125 and 137.

DATA ANALYSIS

In order to determine the most accurate (vector) motions of the stations in the Vandenberg network, it is necessary to estimate the geocentric positions of the sites and the motions of the GPS satellites using a continental- or global-scale GPS network. Thus, our analysis of the Vandenberg data has been carried out simultaneously with data from the central and southern California network and the stations of the (global) Cooperative International GPS Network (CIGNET). This analysis is described in Chapter 4 of *Feigl* [1991], especially pp. 138–145, and in *Murray* [1991]. A discussion of GPS strain estimates from our preliminary analysis is also given in Chapter 4 of *Feigl*, on pp. 162–185.

Coincident with our March, 1990, experiment in central and southern California the DoD activated Selected Availability (SA) in the Block II GPS satellites. This presented a particular problem in our analysis since for this experiment we used receivers which sampled the GPS phase at times separated by up to .08 seconds. Such a mismatch of sample times can cause errors of several centimeters in station-position estimates unless the variations in transmitted frequency caused by SA are properly modeled. Our work in developing a model led to the publication [*Feigl et al.*, 1991] of a simple algorithm that should be of use to other investigators. This model is described in greater detail in Chapter 2 of *Feigl* [1991].

SYNTHESIS OF GEODETIC, GEOLOGIC, AND SEISMIC DATA

In *Feigl et al.* [1990], we summarized the geologic observations, as well as the historical geodetic and seismic data available for the Vandenberg; this paper is reproduced as Chapter 1 of *Feigl* [1991]. In Chapter 4 (pp. 191–201) *Feigl* discusses some inferences that can be made from our GPS measurements and some additional geologic studies completed within the past two years. The greater spatial resolution provided by the recent studies allow him to suggest likely locations for the measured deformation, guiding the future course of our measurements.

CITED PUBLICATIONS

- Feigl, K. L., R. W. King, and T. H. Jordan, Geodetic measurement of tectonic deformation in the Santa Maria Fold and Thrust Belt, California, *J. Geophys. Res.*, 95, 2679–2699, 1990.
- Feigl, K. L., *Geodetic measurement of tectonic deformation in Central California*, Ph.D. thesis, 222 pp., MIT, Cambridge, MA, 1991.
- Feigl, K. L., R. W. King, T. A. Herring, and M. Rothacher, A scheme for reducing the effect of Selective Availability on precise GPS carrier phase measurements, *Geophys. Res. Lett.*, 18, 1289–1292, 1991.
- Murray, M. H., *Global Positioning System measurements of crustal deformation in central California*, Ph. D. thesis, 273 pp., MIT, Cambridge, MA, 1991.