


AD-A256 281



2



DTIC
ELECTE
OCT 5 1992
S C D

PUBLICATION REPORT

1693

8/92

ISOLATION AND CHARACTERIZATION OF LEISHMANIA MAJOR
FROM PHLEBOTOMUS PAPATASI AND MILITARY PERSONNEL
IN NORTH SINAI, EGYPT

BY

N.S. Mansour, D.J. Fryauff, G.B. Modi,
E.M. Mikhail and F.G. Youssef

U.S. NAVAL MEDICAL RESEARCH UNIT NO. 3
(CAIRO, ARAB REPUBLIC OF EGYPT)
PSC 452, BOX 5000
FPO AE 09835-0007

92 10 2 038

92-26392



249977

Short Report

Isolation and characterization of *Leishmania major* from *Phlebotomus papatasi* and military personnel in north Sinai, Egypt*

N. S. Mansour, D. J. Fryauff, G. B. Modi, E. M. Mikhail and F. G. Youssef US Naval Medical Research Unit No. 3, Cairo, Egypt

Since October 1982, cutaneous leishmaniasis has been repeatedly diagnosed among military members of the multinational force and observers (MFO), an international peace keeping force based in the north Sinai desert of Egypt (DUNN & SMERZ, 1983; MANSOUR *et al.*, 1987). In an effort to assist in the recognition, prevention, and treatment of this disease among MFO personnel, the US Naval Medical Research Unit no. 3 initiated epidemiological studies in July 1989 at locations in north-eastern Sinai.

As part of these epidemiological investigations, an entomological survey sought to determine the seasonal dynamics of local sandflies and the prevalence of *Leishmania* infection among these potential vectors. Sandflies were obtained from human landing and resting collections during the months of July, August and September, 1989. Flies were cryopreserved following a modification of the procedure developed by YOUNG *et al.* (1987). Sandfly pools in cryopreservant were slowly frozen on dry ice and transferred to liquid nitrogen for storage until examined. Cryopreserved sandflies were thawed and individually dissected on autoclaved microscope slides in a drop of sterile phosphate-buffered saline (PBS) containing 200 iu penicillin and 200 µg streptomycin per ml to determine the species, reproductive status, and presence and location of promastigote infections. The intact midgut and surrounding media from those specimens containing promastigotes were removed from the slide, triturated in approximately 1.0 ml of sterile PBS with antibiotics (as above), and inoculated into NNN medium and into the hind footpads of BALB/c mice. Tanabe's medium (TANABE, 1923) was used for initial isolation and maintenance of *Leishmania* obtained by aspiration and biopsy of suspected cutaneous leishmaniasis lesions from patients presenting with typical lesions. Both sandfly and human isolates were subsequently passaged into El On's medium (EL-ON, 1969) and harvested at log growth

phase. The promastigotes were centrifuged, washed, resuspended in PBS, and stored at -80°C until isoenzyme electrophoresis was performed. Promastigotes were serotyped according to the method of SCHNUR & ZUCKERMAN (1977), using specific parasite excretory factors (EF) present in the culture medium. Comparative cellulose acetate electrophoresis against World Health Organization reference strains of *L. major* MHOM/IL/67/Jericho-11 (=LRC-L137), *L. tropica* MHOM/SU/58/OD (=LRC-L39), and *L. donovani* MHOM/IN/80/DD8 was used to identify 2 sandfly isolates, IPAP/EG/89/Si-177 and IPAP/EG/90/Si-1614, and 3 human isolates, MHOM/EG/89/Si-15, MHOM/EG/89/Si-17 and MHOM/EG/90/Si-18, from MFO personnel, all from the Sinai study area. The electrophoretic methods of KREUTZER & CHRISTENSEN (1980) were used for the separation of glucose-6-phosphate dehydrogenase (G6PD; EC.1.1.1.49), glucose phosphate isomerase (GPI; EC.5.3.1.9), mannose phosphate isomerase (=phosphomannose isomerase: MPI; EC.5.3.1.8), phosphoglucomutase (PGM; EC.2.7.5.1) and 6-phosphogluconic dehydrogenase (6PGD; EC.1.1.1.44), and that of HARRIS & HOPKINSON (1976) for the visualization of malate dehydrogenase (MDH; EC.1.1.1.37).

Phlebotomus papatasi was the only man-biting sandfly species identified in the 1594 specimens dissected. The promastigote infection rate for this sample, representing the period July–September 1989, was approximately 0.7% (11/1594). Promastigotes from 2 sandfly infections were successfully cultured in NNN medium, and produced lesions after inoculation into the footpads of BALB/c mice. Serotyping of both the sandfly isolates revealed that they were of the same serotype as the 2 recent MFO isolates (EF serotype A₁B₂) and also several other human *Leishmania* isolates from the same area in the Sinai (MANSOUR *et al.*, 1987). In addition, each isolate yielded isoenzyme profiles identical to those of the *L. major* reference strain used for comparison (Figure).




Figure. Diagrammatic representation of 6 isoenzyme electrophoretic patterns on cellulose acetate for *Leishmania* isolated from two *Phlebotomus papatasi* and 3 human cases with cutaneous lesions (acquired in Sinai) compared to World Health Organization (WHO) reference strains. Numbers correspond to isolates and WHO strains, as follows: (1) *L. major* MHOM/IL/67/Jericho-11; (2) MHOM/EG/89/Si-15; (3) MHOM/EG/89/Si-17; (4) MHOM/EG/90/Si-18; (5) IPAP/EG/89/Si-177; (6) IPAP/EG/90/Si-1614; (7) *L. tropica* MHOM/SU/58/OD; (8) *L. donovani* MHOM/IN/80/DD8. See text for explanation of abbreviated names of enzymes.

*This work was supported by the US Naval Medical Research and Development Command, Bethesda, Maryland, Work Unit no. 3M162770A870.AR.322. The opinions and assertions contained herein are the private ones of the authors and are not to be construed as official or reflecting the views of the Navy Department, Department of Defense, or the US Government.

Requests for offprints: Research Publications Branch, US Naval Medical Research Unit no. 3 (NAMRU-3), FPO, AB 09835-1600, USA.

It is reasonable to consider the 10 unidentified promastigote infections in the wild-caught sandflies as *L. major*. This consideration is based upon their position in the sandfly midgut and the tendency of *P. papatasi* to feed preferentially upon warm-blooded hosts. Their position in the midgut, and not in the hind-gut, excludes the possibility of their being trypanosomatids of reptiles, upon which *P. papatasi* only occasionally feeds (KILLICK-KENDRICK, 1979). Because *Leishmania* isolated from the naturally-infected sandflies in this study was indistinguishable from the parasite isolated from cutaneous lesions in man, *P. papatasi* is considered to be a grade 3 vector (KILLICK-KENDRICK & WARD, 1981) in Egypt. WAHBA *et al.* (1990) had previously characterized a single isolate of *L. major* from Egyptian collections of *P. papatasi*. However, this is the first report of repeated, concurrent isolations of this parasite from both sandflies and humans in Egypt.

Acknowledgements

We acknowledge, with appreciation, confirmatory isoenzyme electrophoresis of human and sandfly isolates by Dr Richard Kreutzer of the Youngstown State University, Youngstown, Ohio, USA. Grateful thanks are extended to members of the multinational force and observers for their participation.

References

- Dunn, M. A. & Smerz, R. W. (1983). Medical support of the Sinai multinational force and observers, 1982. *Military Medicine*, **148**, 773-778.
- El-On, J. (1969). [A comparison of the disc electrophoretic profiles of in vivo (*trypan. astigote*) and in vitro (*epimastigote*) forms of trypanosomes]. MSc thesis, The Hebrew University, Jerusalem. [In Hebrew.]
- Harris, M. & Hopkinson, D. A. (1976). The enzymes. In: *Handbook of Enzyme Electrophoresis in Human Genetics*. Amsterdam: North Holland, chapter 4.
- Killick-Kendrick, R. (1979). Biology of *Leishmania* in phlebotomine sandflies. In: *Biology of the Kinetoplastida*, vol. 2, Lumsden, W. H. R. & Evans, D. A. (Editors). London and New York: Academic Press, pp. 395-460.
- Killick-Kendrick, R. & Ward, R. D. (1981). Ecology of *Leishmania*. *Parasitology*, **82**, 143-152.
- Kreutzer, R. D. & Christensen, H. A. (1980). Characterization of *Leishmania* sp. by isoenzyme electrophoresis. *American Journal of Tropical Medicine and Hygiene*, **29**, 199-208.
- Mansour, N. S., Youssef, F. G., Mohareb, E. W., Dees, W. H. & Karuru, E. R. (1987). Cutaneous leishmaniasis in north Sinai. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, **81**, 747.
- Schnur, L. F. & Zuckerman, A. (1977). Leishmanial excreted factor (EF) serotypes in Sudan, Kenya and Ethiopia. *Annals of Tropical Medicine and Parasitology*, **71**, 273-294.
- Tanabe, M. (1923). [On the condition necessary for the development of *Leishmania donovani*]. *Saikingaku, Zasshi*, **333**, 425-436 [In Japanese: English summary in *Japan Medical World* (1924), **4**, 46.]
- Wahba, M. M., Schnur, L. F., Morsy, T. A. & Merdan, A. (1990). The characterization of *Leishmania major* from *Phlebotomus papatasi* (Scopoli) caught in northern Sinai, Egypt. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, **84**, 785-786.
- Young D. G., Morales, A., Kreutzer, R. D., Alexander, J. B., Corredor, A. & Tesh, R. B. (1987). Isolation of *Leishmania braziliensis* (Kinetoplastida: Trypanosomatidae) from cryopreserved Colombian sandflies (Diptera: Psychodidae). *Journal of Medical Entomology*, **24**, 587-589.

Received 20 September 1990; revised 7 February 1991; accepted for publication 8 February 1991

Announcement

Preliminary Announcement

The second residential meeting of the Royal Society of Tropical Medicine and Hygiene (to include the Annual General Meeting) and other European Societies of Tropical Medicine will be held at the Royal College of Physicians, Edinburgh, Scotland from Monday 5th to Wednesday 7th July 1993. Further details available shortly from the Administrator, Royal Society of Tropical Medicine and Hygiene, Manson House, 26 Portland Place, London, W1N 4EY, UK (Tel: 071 580 2127; Fax: 071 436 1389).

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE 20 SEP 90	3. REPORT TYPE AND DATES COVERED		
4. TITLE AND SUBTITLE Isolation and Characterization of <u>Leishmania major</u> from <u>Phlebotomus papatasi</u> and Military Personnel in North Sinai, Egypt.		5. FUNDING NUMBERS PE- 62770A WU- 3M162770A870.AR.322		
6. AUTHOR(S) Mansour, N.S., Fryauff, D.J., Modi, G.B., Mikhail, E.M. and Youssef, F.G.		8. PERFORMING ORGANIZATION REPORT NUMBER 8/92		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) U.S. Naval Medical Research Unit No. 3 PSC 452, Box 5000 FPO AE 09835-0007				
9. SPONSORING, MONITORING AGENCY NAME(S) AND ADDRESS(ES) Naval Medical Research and Development Command, National Naval Medical Center Building 1, Tower 12 Bethesda, MD 20889-5044		10. SPONSORING/MONITORING AGENCY REPORT NUMBER Accession For NTIS 02161 DTIC TAB Unannounced Justification By Distribution 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100		
11. SUPPLEMENTARY NOTES Published in: Trans. R. Soc. Trop. Med. Hyg., 85:590-591, 1991; Acc. No. 1693.		12. DISTRIBUTION CODE Availability Codes Dist. Avail. or Special A-1 20		
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; Distribution is unlimited.		13. ABSTRACT (Maximum 200 words) Cutaneous leishmaniasis (CL) was repeatedly diagnosed among members of the Multinational Forces and Observers (MFO) in the north Sinai. <u>Leishmania</u> parasites were isolated from both sandflies and suspected human cases of (CL). This study revealed that <u>Phlebotomus papatasi</u> is the only man-biting species in this area. <u>Leishmania</u> parasites produced lesions in the footpads of BALB/c mice, their excretory factor (EF) gave the serotype A ₁ B ₂ , and each isolate yielded isoenzyme profiles identical to those of the <u>L. major</u> reference strain. This is the first report of repeated, concurrent isolation of <u>L. major</u> from both sandflies and humans in Egypt.		
14. SUBJECT TERMS Cutaneous leishmaniasis; Diagnosis; <u>Leishmania major</u>; <u>Phlebotomus papatasi</u>; MFO personnel; North Sinai, Egypt.		15. NUMBER OF PAGES 2		
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED		16. PRICE CODE		
17. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED	18. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED	20. LIMITATION OF ABSTRACT		