

AD-A256 756

2

WL-TR-92-7019



Improved Synthesis of 3,5-Diamino-2,4,6-Trinitrotoluene

Dr. Alan P. Marchand
Dr. G. Madhusudhan Reddy

Department of Chemistry
University of North Texas
NT Station, Box 5068
Denton TX 76203-0068

SEPTEMBER 1992



FINAL REPORT FOR PERIOD AUGUST 1990 - FEBRUARY 1992

Approved for public release; distribution is unlimited.

92

020

409358

92-27442



141p8

WRIGHT LABORATORY, ARMAMENT DIRECTORATE

Air Force Material Command ■ United States Air Force ■ Eglin Air Force Base

NOTICE

When Government drawings, specifications, or other data are used for any purpose other than in connection with a definitely Government-related procurement, the United States Government incurs no responsibility or any obligation whatsoever. The fact that the Government may have formulated or in any way supplied the said drawings, specifications, or other data, is not to be regarded by implication, or otherwise as in any manner construed, as licensing the holder, or any other person or corporation; or as conveying any rights or permission to manufacture, use, or sell any patented invention that may in any way be related thereto.

This technical report has been reviewed and is approved for publication.

FOR THE COMMANDER


STEPHEN C. KORN
Chief, Munitions Division

Even though this report may contain special release rights held by the controlling office, please do not request copies from the Wright Laboratory, Armament Directorate. If you qualify as a recipient, release approval will be obtained from the originating activity by DTIC. Address your request for additional copies to:

Defense Technical Information Center
Cameron Station
Alexandria VA 22304-6145

If your address has changed, if you wish to be removed from our mailing list, or if your organization no longer employs the addressee, please notify WL/MNME, Eglin AFB FL 32542-5000, to help us maintain a current mailing list.

Do not return copies of this report unless contractual obligations or notice on a specific document requires that it be returned.

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE September 1992	3. REPORT TYPE AND DATES COVERED Final, August 1990 - February 1992	
4. TITLE AND SUBTITLE Improved Synthesis of 3,5-Diamino-2,4,6-Trinitrotoluene			5. FUNDING NUMBERS C: F08635-90-K-0203 PE: 62602F PR: 2502 TA: 10 WU: 29	
6. AUTHOR(S) Dr. Alan P. Marchand Dr. G. Madhusudhan Reddy				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Department of Chemistry University of North Texas NT Station, Box 5068 Denton TX 76203-0068			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) Wright Laboratory, Armament Directorate Munitions Division Energetic Materials Branch (WL/MNME) Eglin AFB FL 32542-5000			10. SPONSORING / MONITORING AGENCY REPORT NUMBER WL-TR-92-7019	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution is unlimited.			12b. DISTRIBUTION CODE A	
13. ABSTRACT (Maximum 200 words) A three-step efficient synthesis is described that provides the title compound in 58 percent overall yield from orcinol monohydrate.				
14. SUBJECT TERMS Synthesis Polynitroaromatic Compounds Diaminotrinitrotoluene (DATNT)			15. NUMBER OF PAGES 14	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED	18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED	19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED	20. LIMITATION OF ABSTRACT SAR	

PREFACE

This report describes the results of work performed by personnel at the Department of Chemistry, University of North Texas, NT Station, Box 5068, Denton, TX 76203-0068, during the period August 1990 through February 1992. The contract program manager was Dr. Alan P. Marchand, and the WL/MNME program manager was Lt Maria E. Boado. Funding was provided by WL/MNME, Eglin AFB FL 32542-5000, under Contract F08635-90-K-0203.

TABLE OF CONTENTS

Section	Title	Page
I	INTRODUCTION	1
II	DISCUSSION	2
III	EXPERIMENTAL	5
IV	SUMMARY AND CONCLUSIONS	7
	REFERENCES	8

Accession For	
NTIS CRA&I	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
A-1	

SECTION 1

INTRODUCTION

Polynitroaromatic compounds constitute an important class of high density energetic materials that are utilized as explosives. Polynitroaromatic compounds that contain an α -C-H group (e.g., 2,4,6-trinitrotoluene, TNT) consistently show greater impact sensitivity than similar compounds that contain the same oxygen balance but do not possess the α -C-H functionality (References 1 and 2). It has been demonstrated that the impact sensitivity of polynitrotoluenes can be reduced significantly by the introduction of one or more amino groups (Reference 3). 3,5-Diamino-2,4,6-trinitrotoluene (DATNT, 1) is of particular interest in this connection, as this compound is proposed to be a more powerful explosive than TNT but is less impact-sensitive (Reference 4). Herein, a three-step, efficient synthesis is described that provides DATNT in 58 percent overall yield from orcinol monohydrate.

SECTION II

DISCUSSION

DATNT has been synthesized previously in two steps by starting with 3,5-dichloro- (Reference 4) or 3,5-dibromotoluene (References 5 and 6). See Figure 1, **2a** or **2b**, respectively. These syntheses suffer from the following limitations:

- o One of the starting materials, **2a**, is not available commercially; the other is available and can be purchased but is very expensive.
- o Due to the cumulative ring-deactivating influence of the halogen substituents, harsh conditions (e.g., fuming nitric acid-concentrated sulfuric acid) are required to effect trinitration of **2a** and **2b**.
- o The reaction of **3b** with ethanolic ammonia has been reported (Reference 6) to require elevated pressure; thus, this reaction must be performed in a sealed reaction vessel.

More recently, Atkins and coworkers (Reference 7) have reported that DATNT can be synthesized by reacting either pentanitrotoluene or 3-amino-2,4,5,6-tetranitrotoluene with ammonia in dioxane at ambient temperature and atmospheric pressure. However, these procedures employ starting materials that are powerful explosives not readily available and must be prepared via multistep syntheses.

In an effort to overcome the limitations described above, a simple and convenient three-step procedure has been developed that affords DATNT in 58 percent overall yield. The procedure employs as starting material 3,5-dihydroxytoluene (orcinol, **2c**), which is both commercially available (as the monohydrate) and relatively inexpensive.




Figure 1. Approaches to the Rational Synthesis of DATNT

The hydroxy groups in **2c** activate the aromatic ring toward electrophilic nitration. Thus, trinitration of this material is performed conveniently and in high yield (77 percent) simply by using concentrated nitric acid-concentrated sulfuric acid (Reference 8). *O*-methylation of the hydroxy groups in the trinitration product, **3c**, is carried out in high yield (94 percent) by using ethereal diazomethane (Reference 9). Finally, nucleophilic aromatic substitution of the methoxy groups in **4** is executed conveniently and in high yield (80 percent) by reacting a methanol solution of this compound with ammonia gas under ambient conditions.

Alternative methods to replace the use of diazomethane for *O*-methylation of **3c** were sought, as diazomethane was regarded as being unsuitable for use in large-scale reactions. The following methods were attempted: (a) dimethyl sulphate, acetone, solid K_2CO_3 , reflux 4 hours; (b) CH_3I , HMPA, aqueous NaOH, 25 °C, stir 6 hours; and (c) CH_3I , DMSO, powdered KOH, 25 °C, stir 30 minutes. In each case, decomposition of the starting material resulted; no **4** could be isolated from any of these alternative procedures for performing *O*-methylation of **3c**.

SECTION III

EXPERIMENTAL

Melting points are uncorrected. *WARNING!* Polynitroaromatics are powerful explosives and should be handled with extreme caution.

1. 3,5-DIHYDROXY-2,4,6-TRINITROTOLUENE (3c)

A solution of orcinol monohydrate (2c, 284 mg, 2.0 mmol) in concentrated H_2SO_4 (15 mL) is cooled externally (ice-water bath) to 0 °C. A precooled (0 °C) mixture of concentrated HNO_3 (0.8 mL) and concentrated H_2SO_4 (1.6 mL) is added dropwise with stirring to the reaction vessel. This is followed by dropwise addition of cold, concentrated HNO_3 (1.5 mL) to the stirred reaction mixture. The external cold bath is removed, and the reaction mixture is allowed to warm gradually to room temperature during 1 hour. The reaction mixture then is poured into ice water (25 mL). The product, which precipitates immediately, is isolated via suction filtration and air-dried. Pure 3c (398 mg, 77 percent) is thereby obtained as a yellow-orange microcrystalline solid: mp 166-168 °C (literature mp 171-172 °C [Reference 8]; 162 °C [Reference 10]); IR (nujol) 3160 (s), 1628 (m), 1592 (s), 1541 (s), 1469 (s), 1375 (s), 1188 cm^{-1} (s); ^1H NMR (acetone- d_6) δ 2.36 (s, 3 H), 10.06 (br s, 2 H); ^{13}C NMR (acetone- d_6) δ 14.02 (q), 125.77 (s), 133.64 (s), 134.99 (s), 149.75 (s).

2. 3,5-DIMETHOXY-2,4,6-TRINITROTOLUENE (4)

A solution of diazomethane in diethyl ether is added dropwise with stirring to a solution of 3c (130 mg, 0.50 mmol) in ethyl acetate (10 mL) (Reference 9). The addition is continued until the yellow color of diazomethane persists, thereby indicating the presence of a slight excess of this reagent. The resulting solution is stirred overnight (15 hours) at room temperature and then is concentrated in vacuo. Pure 4 (135 mg, 94 percent) is thereby obtained as a yellow

microcrystalline solid: mp 66-67 °C (literature [Reference 9] mp 65-66 °C); IR (nujol) 1592 (s), 1538 (s), 1466 (s), 1365 (s), 1136 cm⁻¹ (s); ¹H NMR (CDCl₃) δ 2.26 (s, 3 H), 3.99 (s, 6 H); ¹³C NMR (CDCl₃) δ 13.25 (1), 64.56 (q), 127.63 (s), 138.55 (s), 141.54 (s), 146.19 (s).

3. 3,5-DIAMINO-2,4,6-TRINITROTOLUENE (1)

A solution of **4** (100 mg, 0.35 mmol) in methanol (10 mL) under argon is cooled externally (ice-water bath) to 0 °C and is stirred vigorously. Ammonia gas is bubbled continuously through this cooled solution for 1.5 hours. The external cold bath is removed, and slow passage of ammonia gas through the reaction mixture is continued with vigorous stirring for an additional 1.5 hours. The gas bubbling apparatus is removed, the flask is stoppered, and the reaction mixture is stirred at room temperature for 1.5 hours. The stopper is removed, and excess ammonia is allowed to evaporate slowly from the reaction mixture. During this time, DATNT slowly crystallizes from the reaction mixture. The product is collected by suction filtration, and the residue is washed with methanol and then is air-dried. Pure DATNT (72 mg, 80.4 percent) is thereby obtained as yellow needles: mp 223-224 °C (literature mp 222 °C [Reference 4], 222.5-224 °C [Reference 7]); IR (nujol) 3439 (m), 3320 (m), 1585 (m), 1469 (s), 1340 cm⁻¹ (m); ¹H NMR (CDCl₃) δ 2.41 (s, 3 H), 8.0-8.3 (br s, 4 H).

SECTION IV

SUMMARY AND CONCLUSIONS

In summary, our three-step synthesis of DATNT, described within this report, offers the following advantages:

- o Each synthetic step is easy to set up, to perform, and to work up.
- o Only inexpensive and readily available reagents are required.
- o Mild conditions are utilized for the trinitration step.
- o Handling of polynitroaromatics is minimized, and the procedure does not subject them to harsh environmental conditions.
- o The use of elevated pressure in the amination step is avoided.
- o The desired material is produced in high overall yield.

It is anticipated that this procedure will lend itself readily to scaleup and thus may find application for large-scale synthesis of DATNT.

REFERENCES

1. M. J. Kamlet and H. J. Adolph, *Propellants and Explosives*, Vol. 4, Page 30, 1979.
2. W. S. Wilson, D. Bliss, S. L. Christian, and D. J. Knight, *Explosive Properties of Polynitroaromatics*, Naval Weapons Center Technical Publication 7073, Naval Weapons Center, China Lake CA 93555-5000, April 1990.
3. S. Iyer, *Propellants, Explosives, and Pyrotechnics*, Vol. 7, Page 37, 1982.
4. S. Iyer, *Journal of Energetic Materials*, Vol. 2, Page 151, 1984.
5. S. S. Joshi and S. P. Gupta, *Journal of the Indian Chemical Society*, Vol. 33, Page 329, 1959.
6. A. W. Palmer, *Chemisches Berichte*, Vol. 21, Page 3501, 1889.
7. R. L. Atkins, R. A. Hollins, and W. S. Wilson, *Journal of Organic Chemistry*, Vol. 51, Page 3261, 1986.
8. A. J. Birch, R. A. Massy-Westropp, R. W. Rickards, and H. Smith, *Journal of the Chemical Society*, Page 360, 1958.
9. K. V. Rao and H. S. Kuo, *Journal of Heterocyclic Chemistry*, Vol. 16, Page 1241, 1979.
10. J. Stenhouse, *Journal of the Chemical Society*, Page 357, 1871.

DISTRIBUTION LIST
(WL-TR-92-7019)

DTIC/DDAC		Eglin AFB offices:	
Cameron Station	2	WL/CA-N	1
Alexandria VA 22304-6145		WL/MNOI (Scientific and Tech. Info. Facility)	1
AFSAA/SAI		WL/FIES/SURVIAC	1
Washington DC 20330-5420	1	Wright-Patterson AFB OH 45433-6553	
AUL/LSE			
Maxwell AFB AL 36112-5564	1		
HQ USAFE/INATW			
APO NY 09012-5001	1		
Eglin AFB offices:		Department of Chemistry	
		University of North Texas	1
WL/MNME	5	NT Station, Box 5068	
WL/MNM	1	Denton TX 76203-0068	
Director		Naval Weapons Center	
U.S. Army Ballistic Research Lab	1	Technical Library	1
Attn: SLC-BR-DD-T		China Lake CA 93555	
Aberdeen Proving Ground MD 21005-5066			
Naval Air Systems Command			
Tech. Library (AIR-723)	1		
Washington DC 20361-9320			
Naval Coastal Systems Center			
Attn: Tech. Library, Code 71121	1		
Panama City Beach FL 32407			
Director			
U.S. Army TRADOC Systems Analysis			
Activity	1		
Attn: ATRC-WSS-R			
White Sands Missile Range NM 88002-5502			