

A CLASS OF PLANAR WELL-COVERED GRAPHS WITH GIRTH FOUR

by

Michael R. Pinter*
Department of Mathematics
Belmont University
Nashville, Tennessee, USA

DTIC
ELECTE
MAR 18 1993
S c I

AD-A262 424


20000920252

DISTRIBUTION STATEMENT A

Approved for public release
Distribution Unlimited

Reproduced From
Best Available Copy

412 272
93-05545



* work partially supported by ONR Contracts #N00014-85-K-0488 and #N00014-91-J-1142

98 3 17 011

~~98 2 18 307~~

DTIC

17 2

Abstract

A well-covered graph is a graph in which every maximal independent set is a maximum independent set; Plummer introduced the concept in a 1970 paper. The notion of a 1-well-covered graph was introduced by Staples in her 1975 dissertation: a well-covered graph G is 1-well-covered if and only if $G-v$ is also well-covered for every point v in G . Except for K_2 and C_5 , every 1-well-covered graph contains triangles or 4-cycles. Thus, triangle-free 1-well-covered graphs necessarily have girth 4. We show that all planar 1-well-covered graphs of girth 4 belong to a specific infinite family, and we give a characterization of this family.

A CLASS OF PLANAR WELL-COVERED GRAPHS WITH GIRTH FOUR

INTRODUCTION

A set of points in a graph is independent if no two points in the graph are joined by a line. The maximum size possible for a set of independent points in a graph G is called the independence number of G and is denoted by $\alpha(G)$. A set of independent points which attains the maximum size is referred to as a maximum independent set. A set S of independent points in a graph is maximal (with respect to set inclusion) if the addition to S of any other point in the graph destroys the independence. In general, a maximal independent set in a graph is not necessarily maximum.

In a 1970 paper, Plummer [14] introduced the notion of considering graphs in which every maximal independent set is also maximum; he called a graph having this property a well-covered graph. The work on well-covered graphs that has appeared in the literature has focused on certain subclasses of well-covered graphs. Campbell [2] characterized all cubic well-covered graphs with connectivity at most two, and Campbell and Plummer [3] proved that there are only four 3-connected cubic planar well-covered graphs. Royle and Ellingham [16] have recently completed the picture for cubic well-covered graphs by determining all 3-connected cubic well-covered graphs.

For a well-covered graph with no isolated points, the independence number is at most one-half the size of the graph. Well-covered graphs whose independence number is exactly one-half the size of the graph are called very well-covered graphs. The subclass of very well-covered graphs was characterized by Staples [17] and includes all well-covered trees and all well-covered bipartite graphs. Independently, Ravindra [15] characterized bipartite well-covered graphs and Favaron [6] characterized the very well-covered graphs. Recently, Dean and Zito [4] characterized the very well-covered graphs as a subset of a more general (than well-covered) class of graphs.

CHAS
TAB
UNCLAS
CONF

by <u>Rec 1</u>	
Distribution /	
Availabil	
Dist	Avail Sp
<u>A-1</u>	

Finbow and Hartnell [7] and Finbow, Hartnell, and Nowakowski [8] studied well-covered graphs relative to the concept of dominating sets. Finbow, Hartnell, and Nowakowski have also obtained a characterization of well-covered graphs with girth at least five [9].

A well-covered graph is 1-well-covered if and only if the deletion of any point from the graph leaves a graph which is also well-covered. A well-covered graph is in the class W_2 if and only if any two disjoint independent sets in the graph can be extended to disjoint maximum independent sets. Staples [18] showed that a well-covered graph is 1-well-covered if and only if it is in W_2 . Since we will appeal mostly to the notion of extending two disjoint independent sets to disjoint maximum independent sets, henceforth we use the W_2 nomenclature instead of referring to 1-well-covered graphs.

The class of well-covered graphs contains all complete graphs and all complete bipartite graphs of the form $K_{n,n}$. The only cycles which are well-covered are C_3 , C_4 , C_5 , and C_7 . We note that all complete graphs are also in W_2 , but no complete bipartite graphs (except $K_{1,1}$) are in W_2 . The cycles C_3 and C_5 are the only cycles in W_2 .

PRELIMINARY RESULTS

We assume that all graphs are connected, unless otherwise stated. The reader is referred to [1] for terminology and notation not defined here. Note that a disconnected graph is in W_2 if and only if each of its components is in W_2 . Suppose G is well-covered, $G \neq K_1$. Let v be a point in G and consider the graph $G-v$. Since $G \neq K_1$, there exists a point $u \sim v$. Since G is well-covered, the point u is contained in a maximum independent set I in G . Clearly, v is not in I . Thus, I is also independent in $G-v$. Consequently, $\alpha(G-v) = \alpha(G)$ for any point v . Hence, from a result of Erdős and Gallai [5] it follows that $\alpha(G) \leq IV(G)/2$. Thus, W_2 graphs inherit this bound on independence number.

Staples [18] proved that a W_2 graph cannot have an endpoint.

Theorem 1. If $G \in W_2$ and G is not complete, then $\delta \geq 2$.

In the next theorem, we prove that a point on a 4-cycle in a W_2 graph must have at least three neighbors.

Theorem 2. If $G \in W_2$ and v is a point on a 4-cycle in G , then $\deg(v) \geq 3$.

Proof. Suppose v is on the 4-cycle $vabc$ in G . Also suppose that $\deg(v) = 2$. Then $\{v\}$ and $\{b\}$ cannot be extended to disjoint maximum independent sets in G , a contradiction since $G \in W_2$. Thus, $\deg(v) \geq 3$. []

In Theorem 3, we show that any point in a W_2 graph that is not on a triangle must be on a 5-cycle.

Theorem 3. If $G \in W_2$ and v is a point in G , then v is on either a triangle or a 5-cycle.

Proof. Suppose v is not on a triangle. Suppose also that v is not on a 5-cycle. Let $N_1 = N(v)$ and $N_2 = \{x \in V(G) : d(x, v) = 2\}$. Since v is not on a triangle, then N_1 is independent. Since $\delta \geq 2$ by Theorem 1, each point in N_1 has a neighbor in N_2 . For each $x \in N_1$, let $N'(x) = N(x) \cap N_2$. Pick $a_x \in N'(x)$ and let $A = \{a_x : x \in N_1\}$. If $a_x \neq a_y$, then a_x is not adjacent to a_y ; otherwise, $vx a_x a_y v$ is a 5-cycle in G containing v . Thus, A is independent. Since A dominates N_1 , then it follows that A and $\{v\}$ don't extend to disjoint maximum independent sets in G . This is the desired contradiction. []

A W_2 graph can have a point of degree two or possibly two adjacent points of degree two. However, we show in the next theorem that a W_2 graph cannot have a point of degree two with each of its neighbors of degree two.

Theorem 4. If G is well-covered (and not a cycle) with a path of three consecutive points of degree two, then G is not in W_2 .

Proof. Suppose a, b and c are points in G such that $a \sim b, b \sim c$ and $\deg(a) = \deg(b) = \deg(c) = 2$. Since G is not a cycle, then a is not adjacent to c . Assume to the contrary that G is in W_2 . Then by Theorem 3, the point b must lie on a 5-cycle. Suppose a 5-cycle containing b is $C = xabcy$. Since G is not a cycle, then either $\deg(x) > 2$ or $\deg(y) > 2$. Without loss of generality, assume $\deg(x) > 2$. Let $u \sim x$ such that $u \notin C$. Then $\{u, c\}$ is independent. So $\{u, c\}$ and $\{a\}$ don't extend to disjoint maximum independent sets in G , a contradiction.

Therefore, G is not in W_2 . []

Consider a graph G which is not complete and point v in G . By deleting v and its neighbors, we obtain a subgraph of G . Specifically, we define the subgraph $G_v = G - N[v]$.

In Theorem 5, we state a *necessary* condition for a well-covered graph to be in W_2 , which is proved in [13]. We will reference Theorem 5 on several occasions in this paper.

Theorem 5. If a graph G is in W_2 and G is not complete, then G_v is in W_2 for all v in G .

The girth of a graph is the size of a smallest cycle in the graph. We say a graph with no cycles has infinite girth. In [13], we prove the following theorem.

Theorem 6. If $G \in W_2$ ($G \neq K_2$ or C_5), then $\text{girth } G \leq 4$.

Hence, a W_2 graph (other than K_2 and C_5) must contain a triangle or a 4-cycle. Thus, a triangle-free W_2 graph (other than K_2 and C_5) has girth 4. In [13], we construct infinite families of W_2 graphs with girth 4. We study *planar* W_2 graphs of girth four for the remainder of this paper.

In general, a W_2 graph can have a cutpoint. However, we prove in the next theorem that a W_2 graph of girth four cannot have a cutpoint.

Theorem 7. If G is a W_2 graph of girth 4, then G is 2-connected.

Proof. Assume to the contrary that G has a cutpoint v . Let $G_1, G_2, \dots, G_n$ be the components of $G-v$. By Theorem 1.20, graphs $G_1, \dots, G_n$ are W_2 graphs. Let $N_i = N(v) \cap G_i$, for $i = 1, \dots, n$. Since G has girth 4, then N_i is independent for all i . Since $G_i \in W_2$, there exists maximum independent sets J_i in G_i such that $J_i \cap N_i = \emptyset$, for all i . Clearly, $J = J_1 \cup \dots \cup J_n$ is an independent set in G . Consequently, J and $\{v\}$ are disjoint independent sets in G which do not extend to disjoint maximum independent sets in G . This is a contradiction since $G \in W_2$. Hence, G is 2-connected. $\square$

A line in a graph G is a critical line if its removal increases the independence number. A line-critical graph is a graph with only critical lines. Staples proved in [17] that a triangle-free W_2 graph is line-critical. Hence, all graphs given subsequently in this paper are line-critical.

PLANAR W_2 GRAPHS OF GIRTH FOUR

In this section, we will characterize all *planar* W_2 graphs of girth 4. For graphs drawn in the plane, we say two faces are adjacent if they share a line. If a face F contains point v , we say F is incident to v . The size of a face is the number of points it contains. We refer to the order and sizes of the faces incident to a point v as the face configuration at v .

Lebesgue [10] developed the theory of Euler contributions for planar graphs and Ore [11] and Ore and Plummer [12] used the theory to study plane graph colorings. The Euler contribution of a point v , $\phi(v)$, is defined as the quantity $\phi(v) = 1 - (1/2)\deg(v) +$

$\sum (1/x_i)$, where the sum is taken over all faces F_i incident to v and x_i is the size of F_i . If $|F(G)|$ denotes the number of faces in the plane graph G , then it follows that $\sum_v \phi(v) = |V(G)| - |E(G)| + |F(G)|$. Here the sum is taken over all points v in G . Since Euler's formula for plane graphs says $|V(G)| - |E(G)| + |F(G)| = 2$, then we have $\sum_v \phi(v) = 2$. Thus, $\phi(v)$ must be positive for some v in G . If $\phi(v) > 0$, we say v is a point with positive Euler contribution.

An infinite family.

The following construction allows us to build larger planar W_2 graphs of girth 4 from a given such graph. It can be verified directly from the definition of a W_2 graph that the construction indeed yields a W_2 graph.

Construction 1. Suppose G is a W_2 graph with adjacent degree two points x and y which are not on a triangle. Let $N(x) = \{u, y\}$, $N(y) = \{x, v\}$, and let a, b and c be new points. Form a new graph H with

$$V(H) = V(G) \cup \{a, b, c\}, \text{ and}$$

$$E(H) = E(G) \cup \{xa, ab, bc, cy, cu\}. \text{ See Figure 1.}$$

Then H is a W_2 graph with $\alpha(H) = \alpha(G) + 1$.




Figure 1

In Construction 1, if G is planar and has girth 4, then clearly H is also a planar W_2 graph of girth 4. In the following theorem, we recursively construct an infinite family of planar W_2 graphs of girth 4.

Theorem 8. Let $n \geq 3$ be a positive integer. Then there exists a planar W_2 graph of girth 4, denoted G_n , such that $\alpha(G_n) = n$ and $|V(G_n)| = 3n - 1$.

Proof. (By induction on n .) For $n = 3$, let G_3 be the graph on eight points given in Figure 2. Then $\alpha(G_3) = 3$ and $|V(G_3)| = 3(3) - 1$. For $k \geq 3$, let G_{k+1} be a graph obtained from G_k by the construction given in Construction 1. Assume $\alpha(G_k) = k$ and $|V(G_k)| = 3k - 1$. From the observation preceding Theorem 8, graph G_{k+1} is a planar W_2 graph of girth 4. Also, $|V(G_{k+1})| = |V(G_k)| + 3 = 3k - 1 + 3 = 3(k+1) - 1$, and $\alpha(G_{k+1}) = \alpha(G_k) + 1 = k + 1$.

Therefore, G_{k+1} satisfies the statement of the theorem. The result follows by induction. □




Figure 2

Now that we have an infinite family of planar W_2 graphs of girth 4, we work toward showing that all planar W_2 graphs of girth 4 are in the family in Theorem 8.

A characterization.

Since the smallest cycle in a W_2 graph of girth 4 is a 4-cycle, it is of interest to learn what we can about 4-cycles in these graphs. The next lemma will help us to determine those W_2 graphs of girth 4 which have exactly one 4-cycle.

Lemma 9. Suppose G is a W_2 graph of girth 4. Let C be a 4-cycle in G . If $\deg(v) = 3$ for all points v in C , then G is isomorphic to the graph given in Figure 2.

Proof. Let $C = v_1v_2v_3v_4$. Assume $\deg(v_i) = 3$, for all i . Since G has girth 4, then v_1 is not adjacent to v_3 and v_2 is not adjacent to v_4 . Let $u_i \sim v_i$ such that u_i is not in C , for all i . Since G has girth 4, then $u_i \neq u_{i+1}$ for all i (addition mod 4).

Suppose $u_1 = u_3$. Then $N(v_1) = N(v_3)$. So $\{v_1\}$ and $\{v_3\}$ don't extend to disjoint maximum independent sets in G , contradicting $G \in W_2$. Thus, $u_1 \neq u_3$ and, similarly, $u_2 \neq u_4$. So we can assume that $i \neq j$ implies $u_i \neq u_j$.

Suppose $u_1 \sim u_4$. Since $\deg(v_1) = 3$, then $\{u_4, v_3\}$ is independent and dominates $N(v_1)$. Thus, $\{u_4, v_3\}$ and $\{v_1\}$ don't extend to disjoint maximum independent sets in G , a contradiction. So u_1 is not adjacent to u_4 . Similarly, u_1 is not adjacent to u_2 , u_2 is not adjacent to u_3 , and u_3 is not adjacent to u_4 .

Since $G \in W_2$, $\deg(u_1) \geq 2$. Since $u_1 \neq u_i$, $i \neq 1$, then u_1 is not adjacent to v_i , $i \neq 1$. Thus, there exists $y \sim u_1$ such that $y \notin C$. If y is not adjacent to v_3 , then $\{y, v_3\}$ and $\{v_1\}$ don't extend to disjoint maximum independent sets in G . So we assume $y \sim v_3$; that is, $y = u_3$ and $u_1 \sim u_3$. Moreover, we have shown that $\deg(u_1) = 2$. By symmetry, $\deg(u_3) = 2$.

By a symmetrical argument, $u_2 \sim u_4$ and $\deg(u_2) = \deg(u_4) = 2$. So $\deg(v_i) = 3$ for all i and $\deg(u_i) = 2$ for all i . Therefore G can be drawn in the plane as the graph given in Figure 2. □

Now we show in Theorem 10 that there is only one W_2 graph of girth 4 with exactly one 4-cycle.

Theorem 10. If G is a W_2 graph of girth 4 with exactly one 4-cycle, then G is isomorphic to the graph in Figure 2.

Proof. Suppose G is a W_2 graph of girth 4 with exactly one 4-cycle. Let $C = abcd$ be the 4-cycle in G . By Theorem 5, $\text{graph } G_v \in W_2$ for all points v in G . By Theorem 2, $\deg(x) \geq 3$ for all x in C .

Suppose $\deg(x) \geq 4$ for some x in C . Without loss of generality, assume $\deg(b) \geq 4$. Let w and y be neighbors of b such that $\{w, y\} \cap \{a, c\} = \emptyset$. Since G has only one 4-cycle, then d is adjacent to neither w nor y . Consider the W_2 graph G_d . Note that b, y and w are in the same component of G_d . Since G has only one 4-cycle, then G_d has no 4-cycles. Thus, G_d is a W_2 graph with girth > 4 .

By Theorem 6, each component of G_d is a line or a 5-cycle. So the component H of G_d containing b, y and w must be a 5-cycle, say $H = stybw$. See Figure 3.




Figure 3

Since G has only one 4-cycle, then s is adjacent to neither a nor c . Thus, the points a, b, c and y are all in the same component of the W_2 graph G_s . But then the component of

G , containing a, b, c and y is neither a line nor a 5-cycle. Since G is a W_2 graph with girth > 4 , we obtain a contradiction via Theorem 6.

Hence, $\deg(b) = 3$. It follows that $\deg(x) = 3$ for all x in C . By Lemma 9, the result follows. []

In the following theorem, we prove that if a W_2 graph has a point of degree two which does not have a neighbor of degree two, then the graph is not planar. As a consequence, we prove in Corollary 12 that if a planar W_2 graph of girth 4 has points of degree two, then those points of degree two must occur in adjacent pairs.

Theorem 11. Suppose G is in W_2 and contains a point v of degree two which is not on a triangle and whose neighbors have degree ≥ 3 . Then G is not planar.

Proof. Let $N(v) = \{a, b\}$. Since v is not on a triangle, then a is not adjacent to b . Let $N_1 = N(a) - v$ and $N_2 = N(b) - v$. By Theorem 2, a 4-cycle in a W_2 graph cannot have a point of degree two. Thus, $N_1 \cap N_2 = \emptyset$. Suppose there exist points x and y such that $x \in N_1$, $y \in N_2$ and x is not adjacent to y . Then $\{x, y\}$ and $\{v\}$ don't extend to disjoint maximum independent sets in G , contradicting $G \in W_2$. Hence, $x \in N_1$ and $y \in N_2$ implies $x \sim y$. Since $\deg(a) \geq 3$ and $\deg(b) \geq 3$, then there exist points u_1 and v_1 in N_1 and points u_2 and v_2 in N_2 . Since $x \sim y$ for all $x \in N_1$, for all $y \in N_2$, it follows that $u_1 \sim u_2$, $u_1 \sim v_2$, $u_2 \sim v_1$ and $v_1 \sim v_2$. Thus, G is not planar. []

Corollary 12. If G is a planar W_2 graph of girth 4 with $\delta = 2$, then the points of degree two occur as adjacent pairs.

Proof. Suppose v is a point of degree two in G . Since G has girth 4, by Theorem 11 it follows that v has a neighbor of degree two. By Theorem 4, the point v cannot have two neighbors of degree two. Thus, v has exactly one neighbor with degree two. Hence, the points of degree two occur as adjacent pairs in G . []

We note that it is possible for a W_2 graph of girth 4 to have a point of degree two whose neighbors have degree greater than two. The graph in Figure 4 is one such example. Moreover, using the graph in Figure 4 as a starting graph, an infinite family of W_2 graphs of girth 4 can be recursively constructed via Construction 1. Each graph in this family has a point of degree two whose neighbors have degree greater than two.




Figure 4

We return to consideration of planar W_2 graphs of girth 4 with points of degree two. Since we know degree two points occur in pairs, we consider the structure around adjacent points of degree two.

Lemma 13. Suppose G is a planar W_2 graph of girth 4 with adjacent degree two points x and y . Let $N(x) = \{u, y\}$ and $N(y) = \{v, x\}$. Then $\deg(u) = \deg(v) = 3$. Moreover, u and v have two common neighbors.

Proof. By Theorem 7, graph G is 2-connected. By Theorem 3, the points x and y are on a 5-cycle C . Then $C = xyvwu$. By Theorem 4, $\deg(u) \geq 3$ and $\deg(v) \geq 3$. Thus, $\deg(w) > 2$ by Theorem 11. Let $N'(w) = N(w) - \{u, v\}$.

Let $U_x = N(u) - x$. Suppose there exists some $p \in U_x$ such that p is not adjacent to v . Then $\{p, v\}$ and $\{x\}$ don't extend to disjoint maximum independent sets in G , contradicting $G \in W_2$. Thus, $p \in U_x$ implies $p \sim v$.

Assume that $\deg(u) > 3$.

Suppose U_x has at least two points outside C and no points inside C . Let $a \in U_x$ such that no point of U_x is in the interior of cycle $uwva$, and let $b \in U_x$ such that a is the only member of U_x in the interior of cycle $uwvb$. Since G has girth 4, then $\{a, b, w\}$ is independent (see Figure 5).




Figure 5

Suppose $z \in N'(w)$ implies $z \sim a$. Then $\{a\}$ and $\{w\}$ don't extend to disjoint maximum independent sets in G , contradicting $G \in W_2$. Thus, there exists some $z \in N'(w)$ such that z is not adjacent to a . Consider the graph G_z . By Theorem 5, graph $G_z \in W_2$.

Let $A_1 = \{c \sim a: c \text{ is inside the cycle } uwva\}$ and $A_2 = \{d \sim a: d \text{ is inside cycle } uavb\}$. If $A_1 = \emptyset$, then w is a cutpoint for G , contradicting the 2-connectedness of G . Thus $A_1 \neq \emptyset$. If there exists $c \in A_1$ such that c is not adjacent to z , then a is a cutpoint for G_z . Since G_z is 2-connected by Theorem 7, we obtain a contradiction. Thus, $c \in A_1$ implies $c \sim z$. Now, if $A_2 = \emptyset$, then $\{b, z\}$ and $\{a\}$ don't extend to disjoint maximum independent sets in G . So $A_2 \neq \emptyset$.

Let $B = \{f \sim b: f \text{ is inside cycle } uavb\}$. Suppose there exists some $d \in A_2$ such that $d \notin B$. Then a is a cutpoint in the graph G_b , a contradiction by Theorem 7. Thus, $d \in A_2$ implies $d \in B$; that is, A_2 is contained in B . But then $\{b, z\}$ and $\{a\}$ don't extend to disjoint maximum independent sets in G .

Hence, it is not possible that U_x has at least two points outside C and no points inside C . By symmetry, we cannot have at least two points of U_x inside C and no points of U_x outside C .

If U_x has at least one point inside C , then rename w as the innermost point of U_x inside C ; that is, choose w so that no point of U_x is inside the 5-cycle $xyvwu$. Now we can proceed as above to obtain a contradiction.

Hence, $\deg(u) = 3$. By symmetry, $\deg(v) = 3$. Let $N(u) = \{x, w, t\}$. From above, $p \in U_x$ implies $p \sim v$. Thus, $N(v) = \{y, w, t\}$. []

Hence, if x is a point of degree two in a planar W_2 graph of girth 4, then x must have one neighbor, say y , with degree two and one neighbor, say u , with degree three. In addition, if v is the second neighbor of y , then $\deg(v) = 3$ and u and v have two common neighbors.

We show in the next theorem that a planar W_2 graph of girth 4 with points of degree two must be in the family constructed in Theorem 8.

Theorem 14. Suppose G is a planar W_2 graph of girth 4 with $\delta = 2$. Then G is a member of the family of graphs given in Theorem 8.

Proof. (By induction on the number of 4-cycles.) Suppose G is a planar W_2 graph of girth 4 with $\delta = 2$. Suppose G has exactly one 4-cycle. Then by Theorem 10, it follows that $G = G_3$ given in Theorem 8. Assume the inductive hypothesis: if G is a planar W_2 graph of girth 4 with $\delta = 2$ and the number of 4-cycles in G is exactly $k-1$ ($k \geq 2$), then G is a member of the family of graphs given in Theorem 8.

Suppose G is a planar W_2 graph of girth 4 with $\delta = 2$ and the number of 4-cycles in G is exactly k ($k \geq 2$). By Corollary 12, graph G has adjacent degree two points, say x and y . Let $N(x) = \{u, y\}$ and $N(y) = \{v, x\}$. It follows from Lemma 13 that $\deg(u) = \deg(v) = 3$ and u and v have two common neighbors. So let $N(u) = \{x, w, a\}$ and $N(v) =$

$\{y, w, a\}$. Without loss of generality, assume a is exterior to cycle $xyvwu$ (see Figure 6).

By Corollary 12, $\deg(w) > 2$ and $\deg(a) > 2$.




Figure 6

If there exists some point $s, s \notin \{u, v\}$, such that $s \sim w$ and $s \sim a$, then $\{s, x\}$ and $\{v\}$ don't extend to disjoint maximum independent sets in G , contradicting $G \in W_2$. Thus, $N'(w) \cap N'(a) = \emptyset$, where $N'(w) = N(w) - \{u, v\}$ and $N'(a) = N(a) - \{u, v\}$.

Case 1. Suppose $\deg(w) \geq 4$.

Case 1.1. Also suppose $\deg(a) \geq 4$. Since we are assuming $\deg(w) \geq 4$ and $\deg(a) \geq 4$, then it follows that there exist distinct points z_1, z_2, a_1, a_2 such that $z_i \in N'(w)$ and $a_i \in N'(a)$, $i = 1, 2$.

Suppose there exist i and j such that a_i is not adjacent to z_j . Then $\{a_i, z_j, y\}$ is independent and so $\{a_i, z_j, y\}$ and $\{u\}$ don't extend to disjoint maximum independent sets in G , contradicting $G \in W_2$. Thus $a_i \sim z_j$ for all i and all j . But this contradicts the planarity of G .

Case 1.2. Thus we must have $\deg(a) = 3$. Consider the graph G_y . By Theorem 5, graph G_y is in W_2 . In G_y , the points a and u are adjacent points of degree two. Since $N'(w) \cap N'(a) = \emptyset$, it follows that G_y has exactly one less 4-cycle than G ($uwva$ is the only 4-cycle in G containing v). Since the number of 4-cycles in G is exactly k , then the number of 4-cycles in G_y is exactly $k-1$. So G_y is a planar W_2 graph of girth 4 with $\delta = 2$ and the number of 4-cycles in G_y is exactly $k-1$. By the inductive assumption, the graph

G_y is in the family of graphs given in Theorem 8. Then G can be obtained from G_y via the construction in Construction 1, with x , y and v playing the roles of a , b and c , respectively (see Figure 7). Thus, G is a member of the family of graphs given in Theorem 8.




Figure 7

Case 2. Suppose $\deg(w) = 3$. Then in the graph G_y , points w and u are adjacent points of degree two. As in Case 1.2, the number of 4-cycles in G_y is exactly $k-1$ and, hence, G_y is in the family of graphs given in Theorem 8. Once again, G can be obtained from G_y via the construction in Construction 1, with x , y and v playing the roles of a , b and c . Thus, G is a member of the family of graphs given in Theorem 8. []

In order to complete the discussion on planar W_2 graphs of girth 4, we will show that a planar W_2 graph of girth 4 must have points of degree two. The theory of Euler contributions will be used for this.

We need to consider possible face configurations at a point of degree three in a planar W_2 graph of girth 4. In a 2-connected planar graph, the faces incident at a point can intersect in various ways. However, for a planar W_2 graph of girth 4, the following lemma

shows that adjacent faces which are incident to a point of degree three always have a line as their intersection. We omit the proof of the lemma.

Lemma 15. Suppose G is planar W_2 graph of girth 4 with $\delta \geq 3$. Suppose v is a point of degree three with $N(v) = \{u_1, u_2, u_3\}$, incident faces F_1, F_2 and F_3 (where face F_1 contains lines vu_1 and vu_2 , face F_2 contains lines vu_2 and vu_3 , and face F_3 contains lines vu_3 and vu_1) and positive Euler contribution, $\phi(v)$. Then $F_1 \cap F_2 = vu_2$, $F_2 \cap F_3 = vu_3$ and $F_3 \cap F_1 = vu_1$.

In Theorem 16, by considering all possible face configurations at a point v with $\deg(v) = 3$ and $\phi(v) > 0$, we conclude that a planar W_2 graph of girth 4 must have points of degree two.

Theorem 16. If G is a planar W_2 graph of girth 4, then $\delta = 2$.

Proof. Assume to the contrary that $\delta \geq 3$. Let $\phi(v)$ be the Euler contribution of point v in G . If $\deg(v) = 4$, then $\phi(v) = -1 + \sum(1/x_i)$, where the sum is taken over the four faces incident to v . Since G has girth 4, the largest possible value for $\sum(1/x_i)$ is 1, when v has face configuration $(4,4,4,4)$. Hence, $\phi(v) \leq 0$ whenever $\deg(v) = 4$. If $\deg(v) = 5$, then $\phi(v) = -3/2 + \sum(1/x_i)$, where the sum is taken over the five faces incident to v . The largest possible value for $\sum(1/x_i)$ here is $5/4$, when v has face configuration $(4,4,4,4,4)$. Hence, $\phi(v) < 0$ whenever $\deg(v) = 5$. In fact, $\phi(v) < 0$ whenever $\deg(v) \geq 5$. Since G must have a point v with $\phi(v) > 0$ and we are assuming $\delta \geq 3$, then we must have $\delta = 3$.

So assume v is a point in G with $\deg(v) = 3$ and $\phi(v) > 0$. Then $\phi(v) = -1/2 + \sum(1/x_i)$, where the sum is taken over the three faces incident to v ; $\phi(v) > 0$ implies that $\sum(1/x_i) > 1/2$. Since G has girth 4, the only possible face configurations at v are the following solutions to the Diophantine inequality $\sum(1/x_i) > 1/2$:

1. $(4,4,n)$, for $n \geq 4$;
2. $(4,5,n)$, for $5 \leq n \leq 19$;
3. $(4,6,n)$, for $6 \leq n \leq 11$;
4. $(4,7,n)$, for $7 \leq n \leq 9$;
5. $(5,5,n)$, for $5 \leq n \leq 9$;
6. $(5,6,n)$, for $6 \leq n \leq 7$.

Let $N(v) = \{u_1, u_2, u_3\}$. Let F_1, F_2 and F_3 be the faces incident to v (where face F_1 contains lines vu_1 and vu_2 , face F_2 contains lines vu_2 and vu_3 , and face F_3 contains lines vu_3 and vu_1). It follows from Lemma 15 that $F_1 \cap F_2 = vu_2$, $F_1 \cap F_3 = vu_1$ and $F_2 \cap F_3 = vu_3$.

Case 1. Suppose v has face configuration $(4,4,n)$, $n \geq 4$. Let $F_1 = vu_1au_2$ and $F_2 = vu_2bu_3$. Since G has girth 4, then a is not adjacent to b . Then $\{a, b\}$ and $\{v\}$ don't extend to disjoint maximum independent sets in G , contradicting $G \in W_2$. Therefore, $(4,4,n)$, $n \geq 4$, cannot occur as a face configuration for v .

Case 2. Suppose v has face configuration $(4,5,n)$, $5 \leq n \leq 19$. Let $F_1 = vu_1au_2$, $F_2 = vu_2bcu_3$ and $F_3 = vu_3df...eu_1$ ($e = f$ when $n = 5$). If a is not adjacent to c , then $\{a, c\}$ and $\{v\}$ don't extend to disjoint maximum independent sets in G . So $a \sim c$. Similarly, $a \sim d$. Since G is planar, point b is adjacent to none of points d, e or u_1 , point c is not adjacent to e , and d is not adjacent to u_2 . See Figure 8.




Figure 8

Suppose $\deg(u_3) \neq 3$. Then there exists $w \sim u_3$ such that $w \notin \{c, d, v\}$ and $\{w, b, e\}$ is independent. Then $\{w, b, e\}$ and $\{v\}$ don't extend to disjoint maximum independent sets in G , a contradiction. Thus, $\deg(u_3) = 3$.

Let $N = N(c) - \{a, b, u_3\}$. Assume $t \in N$ implies $t \sim u_2$. Then $\{d, u_2\}$ and $\{c\}$ don't extend to disjoint maximum independent sets in G . Thus, there exists some $t \in N$ such that t is not adjacent to u_2 . But then $\{t, u_2, f\}$ is independent and so $\{t, u_2, f\}$ and $\{u_3\}$ don't extend to disjoint maximum independent sets in G .

Thus, v cannot have face configuration $(4, 5, n)$, $5 \leq n \leq 19$.

Case 3. Suppose v has face configuration $(4, 6, n)$, for $6 \leq n \leq 11$. Let $F_1 = vu_1au_2$, $F_2 = vu_2bwcu_3$ and $F_3 = vu_3d...eu_1$. As in Case 2, we have $a \sim c$ and $a \sim d$. Thus, since G is planar, e is adjacent to neither b nor c . Since G has girth 4, then b is not adjacent to c . Hence, $\{b, c, e\}$ is independent and so $\{b, c, e\}$ and $\{v\}$ don't extend to disjoint maximum independent sets in G , contradicting $G \in W_2$.

Thus, v cannot have face configuration $(4, 6, n)$, $6 \leq n \leq 11$.

Case 4. Suppose v has face configuration $(4, 7, n)$, $7 \leq n \leq 9$. Let $F_1 = vu_1au_2$, $F_2 = vu_2bxycu_3$ and $F_3 = vu_3dz...eu_1$. As in Case 2, we have $a \sim c$ and $a \sim d$. Also as in Case 2, $\deg(u_3) = 3$. If b is not adjacent to c , then $\{b, c, e\}$ is independent (since G is planar). Thus, $\{b, c, e\}$ and $\{v\}$ don't extend to disjoint maximum independent sets in G , a contradiction. So $b \sim c$. It follows that y is not adjacent to u_2 . But then $\{y, u_2, z\}$ is independent and so $\{y, u_2, z\}$ and $\{u_3\}$ don't extend to disjoint maximum independent sets in G .

Thus, v cannot have face configuration $(4, 7, n)$, $7 \leq n \leq 9$.

Case 5. Suppose v has face configuration $(5, 5, n)$, $5 \leq n \leq 9$. Let $F_1 = vu_1abu_2$, $F_2 = vu_2cdu_3$ and $F_3 = vu_3ex...fu_1$ ($x = f$ when $n = 5$).

Case 5.1. Suppose a is not adjacent to c . If neither a nor c is adjacent to e , then $\{a, c, e\}$ and $\{v\}$ don't extend to disjoint maximum independent sets in G . Thus, either $a \sim e$ or $c \sim e$.

Case 5.1.1. Suppose $a \sim e$. Then neither b nor d is adjacent to f . See Figure 9.




Figure 9

If b is not adjacent to d , then $\{b, d, f\}$ and $\{v\}$ don't extend to disjoint maximum independent sets in G . Thus, $b \sim d$. As in Case 2, $\deg(u_3) = 3$. Now we can apply the argument given in Case 2 to obtain a contradiction.

Case 5.1.2. So $c \sim e$. If b is not adjacent to f , then $\{b, f, d\}$ and $\{v\}$ don't extend to disjoint maximum independent sets in G . So $b \sim f$. See Figure 10.




Figure 10

As in Case 2, with u_2 playing the role of u_3 , we have $\deg(u_2) = 3$. Now we can apply the argument given in Case 2, with $N = N(c) - \{d, e, u_2\}$, to show that this configuration cannot occur.

Case 5.2. Hence, $a \sim c$. If f is not adjacent to d , then $\{f, d, b\}$ and $\{v\}$ don't extend to disjoint maximum independent sets in G . So assume $f \sim d$. See Figure 11.




Figure 11

As in Case 2, with u_1 playing the role of u_3 , we have $\deg(u_1) = 3$. Now we can apply the argument given in Case 2 to obtain a contradiction.

Thus, v cannot have face configuration $(5, 5, n)$, $5 \leq n \leq 9$.

Case 6. Suppose v has face configuration $(5, 6, n)$, $n = 6$ or 7 . Let $F_1 = vu_1abu_2$, $F_2 = vu_2cxdu_3$ and $F_3 = vu_3ewyfu_1$ ($w = y$ when $n = 6$). Since G has girth 4, then c is not adjacent to d .

Case 6.1. Suppose $a \sim c$. If f is not adjacent to d , then $\{b, d, f\}$ is independent and so $\{b, d, f\}$ and $\{v\}$ don't extend to disjoint maximum independent sets in G . So assume $f \sim d$. See Figure 12.




Figure 12

As in Case 2, with u_1 playing the role of u_3 , we have $\deg(u_1) = 3$. Now we can apply the argument given in Case 2 to obtain a contradiction.

Case 6.2. So a is not adjacent to c . If a is not adjacent to d , then $\{a, c, d\}$ and $\{v\}$ don't extend to disjoint maximum independent sets in G , a contradiction. So assume $a \sim d$.

Case 6.2.1. Suppose $n = 6$. Since G has girth 4, then f is not adjacent to e . Then $\{b, f, e\}$ is independent and so $\{b, f, e\}$ and $\{v\}$ don't extend to disjoint maximum independent sets in G . So $n = 6$ cannot occur.

Case 6.2.2. Suppose $n = 7$. If f is not adjacent to d , then $\{c, d, f\}$ and $\{v\}$ don't extend to disjoint maximum independent sets in G . So assume $f \sim d$. If f is not adjacent to e , then $\{f, b, e\}$ and $\{v\}$ don't extend to disjoint maximum independent sets in G . So assume $f \sim e$. See Figure 13.




Figure 13

As in Case 2, with u_1 playing the role of u_3 , we have $\deg(u_1) = 3$. Then $\{b, y, u_3\}$ is independent and so $\{b, y, u_3\}$ and $\{u_1\}$ don't extend to disjoint maximum independent sets in G . Thus, $n = 7$ cannot occur.

Hence, v cannot have face configuration $(5, 6, n)$, for $n = 6$ or 7 . Thus, $\deg(v) = 3$ with $\phi(v) > 0$ leads to a contradiction in all possible cases. Therefore, $\delta \leq 2$. Since $G \in W_2$ and $G \neq K_2$, then by Theorem 1 it follows that $\delta \geq 2$. We conclude that $\delta = 2$. $\square$

Hence, we are able to completely characterize the planar W_2 graphs of girth 4. In particular, the next corollary shows that the family of graphs in Theorem 8 is identical to the family of planar W_2 graphs of girth 4.

Corollary 17. If G is a planar W_2 graph of girth 4, then G is a member of the family of graphs given in Theorem 8.

Proof. This follows immediately from Theorem 16 and Theorem 14. $\square$

REFERENCES

1. J. A. Bondy and U. S. R. Murty, *Graph Theory With Applications*, American Elsevier, New York (1976).
2. S. R. Campbell, Some results on planar well-covered graphs, *Ph.D. Dissertation*, Vanderbilt University, Nashville, TN (1987).
3. S. R. Campbell and M. D. Plummer, On well-covered 3-polytopes, *Ars Combin.* 25-A (1988) 215-242.
4. N. Dean and J. Zito, Well-covered graphs and extendability, preprint, Bellcore, Morristown, NJ, and The Johns Hopkins University, Baltimore, MD (1990).
5. P. Erdős and T. Gallai, On the minimal number of vertices representing the edges of a graph, *Publ. Math. Inst. Hung. Acad. Sci.* 6 (1961) 181-203.
6. O. Favaron, Very well covered graphs, *Discrete Math.* 42 (1982) 177-187.
7. A. Finbow and B. Hartnell, On locating dominating sets and well-covered graphs, *Congr. Numer.* 65 (1988) 191-200.

8. A. Finbow, B. Hartnell, and R. Nowakowski, Well-dominated graphs: a collection of well-covered ones, *Ars Combin.* 25-A (1988) 5-10.
9. A. Finbow, B. Hartnell, and R. Nowakowski, A characterization of well-covered graphs of girth 5 or greater, to appear.
10. H. Lebesgue, Quelques conséquences simples de la formule d'Euler, *Jour. de Math.* 9 (1940) 27-43.
11. O. Ore, *The Four-color Problem*, Academic Press, New York (1967) 54-61.
12. O. Ore and M. D. Plummer, Cyclic coloration of plane graphs, *Recent Progress in Combinatorics*, (ed. W. T. Tutte) Academic Press, New York (1969) 287-293.
13. M. R. Pinter, A class of well-covered graphs with girth 4, submitted, 1991.
14. M. D. Plummer, Some covering concepts in graphs, *J. Combinatorial Theory* 8 (1970) 91-98.
15. G. Ravindra, Well-covered graphs, *J. Combin. Inform. System. Sci.* 2(1) (1977) 20-21.
16. G. F. Royle and M. N. Ellingham, A characterization of well-covered cubic graphs, preprint, Vanderbilt University, Nashville, TN (1991).
17. J. W. Staples, On some subclasses of well-covered graphs, *Ph.D. Dissertation*, Vanderbilt University, Nashville, TN (1975).
18. J. W. Staples, On some subclasses of well-covered graphs, *J. Graph Theory* 3 (1979) 197-204.

**END
FILMED**

DATE:

4-93

DTIC