

# AD-A264 779 REPORT DOCUMENTATION PAGE



2

|                                                                                                                                                                                   |                              |                                                                                                     |                                        |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------|-----------------------------------------------------------------------------------------------------|----------------------------------------|
| 2b. DECLASSIFICATION/DOWNGRADING SCHEDULE                                                                                                                                         |                              | 1b. RESTRICTIVE MARKINGS                                                                            |                                        |
| 4. PERFORMING ORGANIZATION REPORT NUMBER(S)<br>NMRI 92-128                                                                                                                        |                              | 3. DISTRIBUTION/AVAILABILITY OF REPORT<br>Approved for public release;<br>distribution is unlimited |                                        |
| 6a. NAME OF PERFORMING ORGANIZATION<br>Naval Medical Research<br>Institute                                                                                                        |                              | 6b. OFFICE SYMBOL<br>(If applicable)                                                                |                                        |
| 7a. NAME OF MONITORING ORGANIZATION<br>Naval Medical Command                                                                                                                      |                              | 7b. ADDRESS (City, State, and ZIP Code)<br>Department of the Navy<br>Washington, DC 20372-5120      |                                        |
| 8a. NAME OF FUNDING/SPONSORING<br>ORGANIZATION Naval Medical<br>Research & Development Command                                                                                    |                              | 8b. OFFICE SYMBOL<br>(If applicable)                                                                |                                        |
| 9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER                                                                                                                                   |                              | 10. SOURCE OF FUNDING NUMBERS                                                                       |                                        |
| 8c. ADDRESS (City, State, and ZIP Code)<br>8901 Wisconsin Avenue<br>Bethesda, MD 20889-5044                                                                                       |                              | PROGRAM<br>ELEMENT NO.<br>63706N                                                                    | PROJECT<br>NO.<br>M0095.003            |
|                                                                                                                                                                                   |                              | TASK<br>NO.<br>1007                                                                                 | WORK UNIT<br>ACCESSION NO.<br>DN677130 |
| 11. TITLE (Include Security Classification)<br>T cell recognition of MIs-like superantigens: analysis of TCR requirements, superantigenic ligands, and signal<br>transduction     |                              |                                                                                                     |                                        |
| 12. PERSONAL AUTHOR(S) Hodes RJ, Abe R                                                                                                                                            |                              |                                                                                                     |                                        |
| 13a. TYPE OF REPORT<br>journal article                                                                                                                                            | 13b. TIME COVERED<br>FROM TO | 14. DATE OF REPORT (Year, Month, Day)<br>1992                                                       | 15. PAGE COUNT<br>9                    |
| 16. SUPPLEMENTARY NOTATION<br>Reprinted from: Seminars in Immunology 1992 vol.4 pp.319-327                                                                                        |                              |                                                                                                     |                                        |
| 17. COSATI CODES                                                                                                                                                                  |                              | 18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)                   |                                        |
| FIELD                                                                                                                                                                             | GROUP                        | SUB-GROUP                                                                                           |                                        |
|                                                                                                                                                                                   |                              | T cells, Superantigenic, Calcium, Murine mammary tumor virus.                                       |                                        |
| 19. ABSTRACT (Continue on reverse if necessary and identify by block number)                                                                                                      |                              |                                                                                                     |                                        |
| <div style="text-align: center;"> <p><b>DTIC</b><br/><b>ELECTE</b><br/><b>S MAY 24 1993 D</b><br/><b>B</b></p> </div>                                                             |                              |                                                                                                     |                                        |
| 20. DISTRIBUTION/AVAILABILITY OF ABSTRACT<br><input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS |                              | 21. ABSTRACT SECURITY CLASSIFICATION<br>Unclassified                                                |                                        |
| 22a. NAME OF RESPONSIBLE INDIVIDUAL<br>Phyllis Blum, Librarian.                                                                                                                   |                              | 22b. TELEPHONE (Include Area Code)<br>(301) 295-2188                                                | 22c. OFFICE SYMBOL<br>MRL/NMRI         |

Reproduced From  
Best Available Copy

## T cell recognition of Mls-like superantigens: analysis of TCR requirements, superantigenic ligands, and signal transduction

Richard J. Hodes and Ryo Abe\*

*Mls-like endogenous superantigens have been reported to be unique, differing from conventional antigens in the T cell receptor (TCR) requirements for their recognition, in the signal transduction events which they elicit in T cells, and in the identity of the stimulatory ligand. This review describes recent findings indicating that TCR V $\alpha$  as well as V $\beta$  products play a role in Mls recognition. Data are also summarized which indicate that T cell recognition of Mls leads to increased intracellular [Ca<sup>2+</sup>] and phosphatidyl inositol hydrolysis. Finally, genetic linkage between mouse mammary tumor virus (MMTV) proviruses and endogenous superantigens is analyzed, and the possible role of non-MMTV products is examined.*

**Key words:** T cell / T cell receptor / superantigen / MMTV

MLS (MINOR LYMPHOCYTE STIMULATING) antigens were first characterized by Hilliard Festenstein and his colleagues as antigens expressed by certain mouse strains which were capable of stimulating vigorous primary proliferative responses by T cells from major histocompatibility complex (MHC)-identical strains which did not express these determinants.<sup>1</sup> This unique ability to stimulate large proliferative T cell responses was the source of considerable interest for those interested in antigen recognition by T cells. Initially, Mls antigens Mls<sup>a</sup>, Mls<sup>c</sup> and Mls<sup>d</sup> were regarded as the allelic products of a single genetic locus, for which the non-stimulatory allele was Mls<sup>b</sup>. However, subsequent findings from our own laboratory as well as others revealed several surprising aspects of the Mls antigens. First, it was demonstrated by conventional genetic analysis that the stimulatory

Mls determinants were not expressed on allelic products of a single locus. Rather, Mls<sup>a</sup> and Mls<sup>c</sup> determinants were the products of distinct and non-allelic genes, and the Mls<sup>d</sup> phenotype represented the co-expression of the independently encoded Mls<sup>a</sup> and Mls<sup>c</sup> products.<sup>2-7</sup> Thus, there was no evidence for polymorphic allelism at any Mls locus, but rather the expression or non-expression of a T cell stimulatory product of each locus. Further analysis revealed another unique characteristic of Mls determinants, first demonstrated for Mls<sup>c</sup>, namely a 'genetic redundancy' such that two or more unlinked genes existed either one of which was sufficient to encode expression of an indistinguishable Mls determinant.<sup>8,9</sup>

These unique characteristics of the Mls system, the absence of allelic polymorphism and the presence of genetic redundancy, were explained by the recent discovery by several laboratories of the relationship between Mls-like 'superantigens' and endogenous mouse mammary tumor virus (MMTV) proviruses.<sup>10-12</sup> Different proviruses, with distinct sequences in regions including the open reading frames of their LTRs, have different strain distributions.<sup>10-13</sup> Each provirus is integrated at one genomic site and is either present (giving an Mls stimulatory phenotype) or absent (the non-stimulatory phenotype), without structurally allelic forms of its product, consistent with the behavior of Mls determinants. In addition, distinct proviruses integrated in unlinked loci can share sufficient sequence homology to encode highly cross-reactive or indistinguishable Mls-like determinants,<sup>13</sup> accounting for the 'genetic redundancy' observed in the encoding of Mls.

The other recent development which has provided critical insights into the functional behavior of Mls antigens is the characterization of antigen recognition via the antigen-specific T cell receptor (TCR). Antigen recognition by most T cells is mediated by an antigen-specific receptor which is an  $\alpha\beta$  heterodimer, each chain of which is the product of multiple variable gene segments (V, D, and J for the  $\beta$  chain, and V and J for the  $\alpha$  chain) as well as non-germline

*From the Experimental Immunology Branch, National Cancer Institute, Bethesda, MD 20892, USA and \*Immune Cell Biology Program, Naval Medical Research Institute, Bethesda, MD 20889-5055, USA*

©1992 Academic Press Ltd

1044-5323/92/050319 + 09\$8.00/0



encoded junctional residues.<sup>14</sup> T cell recognition of conventional protein antigens involves recognition of peptides presented in association with self MHC molecules, and is dependent upon the precise combinatorial specificity of the TCR, requiring the appropriate co-expression of multiple variable  $\alpha$  and  $\beta$  chain segments.<sup>15</sup> In contrast, a class of 'superantigens' has been described, the recognition of which appears to depend primarily upon expression of an appropriate V $\beta$  product, with little or no influence of other TCR components. Thus, all or nearly all T cells expressing an appropriate TCR V $\beta$  product are responsive to a superantigen. In addition to exogenous superantigens, which are mycoplasma, bacterial, or viral products, the Mls-like products expressed in the mouse represent a unique class of endogenous superantigens.<sup>9,16-20</sup> Each of the endogenous superantigens which has been characterized to date appears to be encoded by a distinct MMTV provirus, each of which has a unique distribution among mouse strains.<sup>10-13</sup> As noted above, many of the endogenous superantigens are capable of stimulating primary proliferative responses by T cells from mice that do not express the corresponding superantigen. In addition, these endogenous superantigens mediate the clonal deletion during T cell development of cells which express a V $\beta$  product capable of reacting with that 'self' superantigen.<sup>9,17,19-21</sup>

This review summarizes our recent analysis of the ligands which serve as Mls-like superantigens, of TCR expression by Mls<sup>a</sup>-reactive T cells, and of the nature of signal transduction events which are associated with T cell recognition of Mls<sup>a</sup>.

### Association of MMTV proviruses with deletion of T cells expressing V $\beta$ 5, 11, and 12

A group of V $\beta$  families, which includes V $\beta$  5, 11, 12, and 16, was initially characterized as 'I-E' related in that most strains which express an E $\alpha$ E $\beta$  class II MHC product delete T cells expressing the products of these V $\beta$  genes.<sup>22,23</sup> More recently, it was shown that the self ligands which mediate deletion of V $\beta$ 5, 11, and 12-expressing cells are determined by multiple non-MHC-linked genes<sup>19,20,24</sup> and it was subsequently reported that the ligand(s) for V $\beta$ 5 and V $\beta$ 11 map to MMTV proviruses Mtv-8, Mtv-9, and Mtv-11.<sup>11,12</sup> At least some of these V $\beta$ -deleting MMTV proviral products also serve as superantigens in the Mls-like stimulation of primary T cell

proliferative responses.<sup>19,20</sup> The relationship between specific TCR V $\beta$  deletions and expression of particular MMTV proviruses was analyzed in more detail for this newly characterized set of superantigens. In order to relate V $\beta$  specificity to MMTV LTR sequences, a series of backcross animals was analyzed in which individual Mtv-8, Mtv-9, and Mtv-11 proviruses detected by Southern blotting segregated independently, and in which MMTV expression could be correlated with deletion of V $\beta$ 5, 11, and 12. The linkage of specific MMTVs to these deletions was found to be complex. Expression of Mtv-9 was sufficient to delete essentially all mature T cells expressing V $\beta$ s 5, 11 and 12; Mtv-8 caused complete deletion of V $\beta$ 12, but only partial deletion of V $\beta$ 11, and little or no deletion of V $\beta$ 5; and Mtv-11 caused complete deletion of V $\beta$ s 11 and 12, but no deletion of V $\beta$ 5 (Figure 1). If these differences in Mtv-8, 9, and 11 are related to sequence differences in the C-terminus of the LTR open reading frame, as suggested by previous studies,<sup>13</sup> the limited sequence differences among these three MMTVs either produce different determinants for V $\beta$ -specific recognition or lead to affinity differences in recognition of the same determinant by different TCR V $\beta$  products.<sup>25</sup>

### T cell receptor V $\beta$ repertoire in the absence of MMTV proviruses

The effect of endogenous proviruses on the developing TCR repertoire is considerable, potentially resulting in the deletion of large portions of the potential repertoire. This has stimulated considerable speculation upon the potential advantages or disadvantages of such effects for the host.<sup>25,26</sup> To date, however, the existence of endogenous superantigens has been established only in the mouse. In addition, each of the superantigens which has been identified in the mouse has been associated with a MMTV provirus. An important question, which remains to be clarified, is therefore whether endogenous superantigens are limited to the MMTV proviruses expressed in mice, or whether they represent a more generalized biological phenomenon. We have addressed this issue by two approaches, the first of which involved the study of feral mice that do not express detectable MMTV proviruses, and the second of which examined the ability of another species, the rat, to contribute to the ligands for V $\beta$ -specific negative selection in mice.

Availability Codes

and/or

Dist Special


A-1 20

|        | V $\beta$ 5       | V $\beta$ 11      | V $\beta$ 12      |
|--------|-------------------|-------------------|-------------------|
| Mtv-8  | No deletion       | Partial deletion  | Complete deletion |
| Mtv-9  | Complete deletion | Complete deletion | Complete deletion |
| Mtv-11 | No deletion       | Complete deletion | Complete deletion |

**Figure 1.** Relationship between TCR V $\beta$  expression and MMTV provirus. Segregation analyses were carried out in multiple strain combinations in which the presence or absence of multiple MMTV genomic proviruses segregated independently. Individual mice were then analyzed for genomic MMTV and for TCR V $\beta$  expression. The presence of proviral Mtv-8, Mtv-9, and Mtv-11 in individual mice was correlated with deletion of V $\beta$ 5, 11, and 12 as summarized.

The Czech II strain was derived from feral mice and has previously been shown to lack expression of any integrated MMTV provirus when assayed by Southern analysis.<sup>27</sup> Mice of this strain were

therefore analyzed for their TCR V $\beta$  expression to determine whether they might express non-MMTV ligands capable of mediating V $\beta$ -specific deletion. V $\beta$  expression in CD4<sup>+</sup> and CD8<sup>+</sup> Czech T cells



**Figure 2.** T cell receptor V $\beta$  expression by CD4<sup>+</sup> T cells in MMTV provirus-negative Czech II mice, in their F<sub>1</sub> offspring, and in B10.A mice foster-nursed by Czech II females. Results are expressed as the percent of CD4<sup>+</sup> spleen cells that express the indicated TCR V $\beta$  product. ■, B10.A; ▨, Czech II; ▤, (B10.A x Czech) F<sub>1</sub>; ▩, B10.A-F-Czech.

was compared with expression in the strain C57BL/10 (B10), which undergoes no known  $V\beta$ -specific deletions due to its failure to express an MMTV product with a permissive class II MHC product, or with the strain B10.A which does exhibit a number of MMTV-dependent  $V\beta$ -specific deletions. Czech T cells expressed a wide profile of  $V\beta$  products, in general at levels similar to those observed in B10 T cells. However,  $V\beta 3^+$  T cells were expressed at significantly lower levels on both  $CD4^+$  (Figure 2A) and  $CD8^+$  (data not shown) Czech T cells than on the corresponding B10 or B10.A populations. Two approaches were taken to determine whether this reduced level of  $V\beta 3$  expression was the result of negative selection during T cell development. First, the expression of  $V\beta 3$  during intrathymic T cell development was compared in thymocytes from Czech, from non- $V\beta 3$ -deleting B10.A, or from known Mls<sup>c</sup> + Mtv-1 + Mtv-6 + C3H/HeJ.<sup>10</sup> As previously reported,<sup>9</sup> superantigen-mediated  $V\beta 3$  deletion in C3H/HeJ thymocytes occurred at the transition from immature cells expressing a low density of TCR to mature cells expressing a high TCR density, so that  $V\beta 3$ -bright cells were selectively deleted whereas there was a much smaller effect on  $V\beta 3$ -dull thymocytes. In contrast, Czech thymocytes exhibited similarly reduced frequencies in  $V\beta 3$ -bright and  $V\beta 3$ -dull populations, suggesting that the reduced levels of  $V\beta 3$  expression did not result from clonal deletion during intrathymic maturation (data not shown).

In order to further assess the expression of a  $V\beta 3$ -deleting ligand in the Czech genome,  $F_1$  mice were generated by breeding Czech males with females of the B10.A strain, a strain which is known to have an MHC type permissive for  $V\beta 3$  deletion, but which does not express a  $V\beta 3$ -specific MMTV ligand. These  $F_1$  mice failed to delete  $V\beta 3$ , indicating that the Czech genome does not encode a ligand capable of mediating  $V\beta 3$  deletion in these animals (Figure 2B). In contrast, when  $F_1$ s were constructed between Czech males and females from the Mls<sup>c</sup> +, Mtv-1 + Mtv-6 + strain C3H/HeJ, these (C3H/HeJ  $\times$  Czech)  $F_1$ s deleted  $V\beta 3$  at the nearly complete level of deletion characteristic of the C3H/HeJ parental strain (data not shown). To test the possibility that a milk-borne agent (such as an infectious MMTV) carried by Czech mice, and not a genomically transmitted product, could mediate  $V\beta 3$  deletion, B10.A mice were foster-nursed on Czech females and the resulting mice analyzed. No  $V\beta 3$  deletion was observed in these foster-nursed animals, thus failing

to detect a milk-borne  $V\beta 3$ -deleting agent (Figure 2B). Collectively, these data indicate that MMTV-free Czech mice fail to express a detectable  $V\beta$ -deleting ligand and are consistent with the conclusion that  $V\beta$ -specific T cell clonal deletion is a unique property of MMTV-dependent ligands.

### Effect of rat bone marrow-derived cells on the mouse T cell receptor $V\beta$ repertoire

The existence of  $V\beta$ -specific clonal deletion has to date been demonstrated only in the mouse. Specific attempts to identify a similar phenomenon in the rat have failed to detect  $V\beta$ -specific deletion.<sup>28</sup> In man, individual differences in  $V\beta$  expression have been identified, but have not been shown to result from ligand-specific events in T cell selection. To test the possibility that endogenous products of mammalian species other than mouse might mediate  $V\beta$ -specific deletion, an approach was designed using mixed radiation bone marrow chimeras. Lethally irradiated C57BL/10 (B10) mice were reconstituted with either B10 mouse bone marrow alone or with a mixture of B10 and rat bone marrow.<sup>29</sup> Mice reconstituted with B10 mouse bone marrow alone did not exhibit any  $V\beta$ -specific deletions in their reconstituted T cell repertoires (Figure 3). In contrast the B10 bone marrow-derived T cells of chimeras which had been reconstituted with a mixture of B10 and WF strain rat bone marrow demonstrated specific deletion of  $V\beta 5^+$  and  $V\beta 11^+$  T cells; chimeras which had been reconstituted with F344 strain rat marrow did not show such deletion (Figure 3). Thus, rat bone marrow derived cells were able to contribute in a strain-specific manner to the ligand for  $V\beta$ -specific deletion in mouse T cells. The identity of the rat gene product involved in this specific deletion of mouse T cells was not established in these experiments. All of the  $V\beta$ -specific deleting ligands described to date have both MHC class II components and non-MHC-linked superantigen components. Thus, rat bone marrow-derived cells may have contributed an I-E-like class II molecule which functioned to present mouse MMTV proviral products capable of mediating  $V\beta 5$  and  $V\beta 11$  deletion; alternatively, WF strain rat cells may express distinct and presumably non-MMTV ligands capable of mediating  $V\beta 5$  and  $V\beta 11$  deletion. The presence or absence of  $V\beta$ -specific deletions in the TCR repertoire expressed by rat cells in these chimeras is now under investigation.




Figure 3. Contribution of rat bone marrow-derived cells to TCR V $\beta$ -specific T cell deletion in radiation bone marrow chimeric mice. Radiation bone marrow chimeras were prepared by injecting lethally irradiated recipients with bone marrow of the indicated rat and/or mouse strain. Mouse bone marrow-derived Th1.2<sup>+</sup> T cells were analyzed for expression of the indicated TCR V $\beta$  product. Figure reproduced from the *Journal of Experimental Medicine*, 1992, 175: 147-157 by copyright permission of the Rockefeller University Press. ▨, B10; □, B10 + F344→B10; ▩, B10 + WF→B10; ■, BR→BR.

### Influence of TCR V $\alpha$ expression on Mls<sup>a</sup> reactivity

As noted above, Mls and Mls-like superantigens have been identified primarily by their ability to mediate V $\beta$ -specific clonal deletion of developing T cells or V $\beta$ -specific clonal activation of mature T cells. The analysis of natural and transfected V $\beta$  variants has further analyzed the role of V $\beta$  sequence in recognition of Mls superantigens, and has suggested that superantigen recognition is dependent upon regions of the TCR that are distant from those involved in recognition of conventional antigens.<sup>30</sup> However, two lines of evidence have suggested that TCR specificity for endogenous Mls superantigen is not determined by the expressed V $\beta$  product alone. Yui *et al* reported that some but not all T cell clones expressing a transgenic V $\beta$ 8.1  $\beta$  chain were Mls<sup>a</sup> (Mtv-7) reactive,<sup>31</sup> and Pullen *et al* found that the ability of a transfected V $\beta$ 8.2  $\beta$  chain to mediate Mls<sup>a</sup> reactivity in hybridomas was influenced by the

$\alpha$  chain co-expressed.<sup>32</sup> In order to formally assess the role of  $\alpha$  chain expression in Mls<sup>a</sup> reactivity, the pairing of specific V $\alpha$  products with V $\beta$ 6 was analyzed in populations of Mls<sup>a</sup> responsive T cells. T cells from Mls<sup>a</sup>-negative mice were stimulated *in vitro* with Mls<sup>a</sup>-positive cells, or with control allogeneic but Mls<sup>a</sup>-negative cells, or with anti-TCR antibodies. These T cell populations were then analyzed by two-color flow cytometry for their simultaneous expression of V $\beta$  and V $\alpha$  products. When T cells from Mls<sup>a</sup>-negative strains were activated *in vitro* for 6 days with Mls<sup>a</sup>-expressing allogeneic stimulator cells, a marked selective expansion in V $\beta$ 6<sup>+</sup> cells was observed, reflecting the dominance of V $\beta$ 6-expressing T cells in the response to Mls<sup>a</sup>.<sup>33</sup> When V $\alpha$  expression was analyzed on these expanded V $\beta$ 6<sup>+</sup> T cells, it was found that the distribution of specific V $\alpha$  products was substantially different from that on V $\beta$ 6<sup>+</sup> T cells which had not been selected for Mls<sup>a</sup> responsiveness.<sup>34</sup> Mls<sup>a</sup>-responsive V $\beta$ 6<sup>+</sup> T cells showed significantly decreased frequencies of




Figure 4. Influence of TCR Vα expression on reactivity of Vβ6<sup>+</sup> T cells to Mls<sup>A</sup>. T-enriched spleen cells were cultured with the indicated stimulus for 7 days prior to analysis for Vα expression in Vβ6<sup>+</sup> T cells and were compared to freshly isolated spleen cells. TCR expression in the cultured cells was expressed relative to that seen in uncultured Vβ6<sup>+</sup> spleen cells:

$$\% \text{ of expression in spleen} = \frac{\text{V}\alpha/\text{V}\beta 6 \text{ in 7-day cultures}}{\text{V}\alpha/\text{V}\beta 6 \text{ in spleen cells}} \times 100.$$

Figure reproduced from the *Journal of Experimental Medicine*, 1992, 175:1405-1408 by copyright permission of the Rockefeller University Press. ■, Vα2/Vβ6; ▨, Vα3/Vβ6; ▩, Vα8/Vβ6; □, Vα11/Vβ6.

expression of Vα2, 3, and 8 and an increase in expression of Vα11 when compared with unstimulated Vβ6<sup>+</sup> T cells (Figure 4). This pattern was consistent in the responses of different strains to Mls<sup>A</sup> stimulators and was specific for Mls<sup>A</sup> responses in that no similar skewing of Vα usage was observed after stimulation of T cells with non-Mls alloantigens or with anti-Vβ6 antibody. Thus, T cell responsiveness to Mls<sup>A</sup> is not a function of Vβ usage

alone, but is substantially influenced by expression of specific Vα products as well.

Since Vα expression influences the *in vitro* response of mature T cells to Mls<sup>A</sup>, attempts were made to determine whether Vα expression also exerts an influence on Vβ-specific negative selection by Mls superantigens. Attempts to study Vα expression on any residual Vβ6<sup>+</sup> cells which might remain undeleted in Mls<sup>A</sup>-expressing mice were unsuccessful

due to the very small number of such cells. The completeness of this deletion itself suggests that essentially all  $V\beta 6^+$  cells, regardless of  $V\alpha$  expression, are deleted in the presence of endogenous  $Mls^a$  product. Partial  $V\beta$  deletion has been described in the case of  $V\beta 11$  deletion in A/J mice, apparently mediated by the Mtv-8 proviral ligand.  $V\alpha$  expression was therefore analyzed in the  $V\beta 11^+$  T cells found in A/J and in the non- $V\beta 11$ -deleting MHC congenic strain A.BY. No difference in  $V\alpha$  expression was observed with the two  $V\alpha$ -specific antibodies reactive with this strain. There was therefore no indication that  $V\alpha$  plays a role in the incomplete negative selection of  $V\beta 11$  in A/J mice.

These findings suggest something of a paradox in that there is an apparent influence of  $V\alpha$  expression on the *in vitro* response of mature T cells to an  $Mls$  superantigen, but there may not be a similar influence on negative selection mediated by the same ligand. It should be noted, however, that the systems in which these two effects are measured are quite different. The expansion of  $V\beta 6^+$  cells during *in vitro* T cell responses to  $Mls^a$  reflects the selective survival and/or expansion of certain  $Mls^a$ -reactive T cells. These cells may, for example, represent those with an optimal TCR affinity for the  $Mls^a$  ligand. The selective  $V\alpha$  expression in these cells is therefore characteristic of the selected population of cells which are most capable of proliferation or survival in response to  $Mls^a$ . In contrast, it is possible that negative selection *in vivo* does not reflect the same requirements; a T cell may be deleted after intrathymic interaction with an  $Mls^a$  product at some threshold level of affinity, despite the fact that the TCR expressed by this cell might not mediate a competitively strong proliferative response to that ligand *in vitro*. The existence of differential requirements for negative selection versus peripheral T cell responsiveness mediated by the same TCR has in fact been reported in mice transgenic for a TCR specific for lymphocytic choriomeningitis virus plus class I D<sup>b</sup>.<sup>35</sup>

### Signal transduction events mediating T cell responses to $Mls^a$

The precise nature of T cell stimulation by endogenous  $Mls$ -like superantigens is unknown. Substantial evidence exists supporting a role for the  $\alpha\beta$  TCR and for CD4 expressed by the T cell and for class II MHC product on the stimulatory APC. However,

as discussed above, the molecular characteristics of endogenous superantigens are unknown, and it is unclear whether signals transmitted via the TCR in response to these superantigens are identical to those transmitted in response to conventional peptide antigens presented by self MHC class II molecules. O'Rourke *et al* have in fact concluded, on the basis of studies of heterogeneous T cells and T cell clones, that stimulation of T cells by  $Mls^a$  induces proliferation by a pathway that does not involve detectable phosphatidyl inositol (PI) hydrolysis, whereas stimulation of the same T cells by MHC-encoded alloantigens does induce PI hydrolysis, leading to the conclusion that  $Mls^a$  stimulation is mediated by a distinct signal transduction pathway.<sup>36</sup> Similar conclusions were reported by Gaugler *et al*<sup>37</sup> and Patarca *et al*.<sup>38</sup> To characterize the signaling events which are involved in T cell recognition of  $Mls^a$ , the response of  $Mls^a$ -specific T cell clones to splenic stimulator cells derived from  $Mls^a$ -positive or negative stimulators were first studied. Cloned T cells were loaded with the  $[Ca^{2+}]$ -sensitive dye indo-1 and stimulator cells were labeled with the dye Dil<sub>22</sub>(3). T cells and stimulator cells were then mixed, and the cell mixture was analyzed by flow cytometry. It was observed that  $Mls^a$ -specific T cells formed a greater frequency of cell-cell conjugates with  $Mls^a$ -positive than with  $Mls^a$ -negative cells, indicating that  $Mls^a$ -specific conjugates were being formed.<sup>39</sup> In addition, those cloned T cells which were conjugated with  $Mls^a$ -positive cells demonstrated a rapid increase in intracellular  $[Ca^{2+}]$  after warming to 37 °C (Table 1). The same comparison also demonstrated a specific increase in PI hydrolysis after interaction of cloned T cells with  $Mls^a$  stimulators.

Table 1. Intracellular  $[Ca^{2+}]$  response of cloned T cells to  $Mls^a$  stimulation

| Stimulator cells | H-2 | $Mls^a$ (Mtv-7) | $[Ca^{2+}]_i$ increase* |
|------------------|-----|-----------------|-------------------------|
| AKR.J            | k   | +               | +                       |
| CBA/J            | k   | +               | +                       |
| D1.LP            | b   | +               | +                       |
| DBA/2            | d   | +               | +                       |
| CBA/CaH          | k   | -               | -                       |
| C3H/HeJ          | k   | -               | -                       |
| B10.BR           | k   | -               | -                       |
| C57BL/6          | b   | -               | -                       |
| B10.Q            | q   | -               | -                       |

\*Positive responses were increases in  $[Ca^{2+}]_i$  of greater than 200 nM compared to pre-stimulation values. Negative responses were increases less than 15 nM.



These results indicated that T cell recognition of Mls<sup>a</sup> results in both increased intracellular [Ca<sup>2+</sup>] and PI hydrolysis, suggesting that the PLC-dependent pathway of activation can be induced by Mls<sup>a</sup> recognition in cloned Mls<sup>a</sup>-specific T cells. In order to determine whether similar signaling occurs in the recognition of Mls<sup>a</sup> by normal heterogeneous T cells, peripheral T cells were prepared from Mls<sup>a</sup>-positive AKR/J mice, from Mls<sup>a</sup>-negative CBA/CaH mice, and from transgenic CBA/CaH mice expressing an Mls<sup>a</sup>-specific TCR V $\beta$ 8.1 transgene product on greater than 95% of peripheral T cells. These T cells were indo-1 loaded and mixed with labeled Mls<sup>a</sup>-positive or negative spleen cells. It was observed that T cells from Mls<sup>a</sup>-negative mice formed cell-cell conjugates with Mls<sup>a</sup>-positive stimulator cells, and that a high proportion of T cells in these conjugates underwent a significant increase in intracellular [Ca<sup>2+</sup>]. T cells from normal Mls<sup>a</sup>-negative mice formed reactive conjugates in a proportion consistent with that expected for Mls<sup>a</sup>-reactive cells in heterogeneous populations; T cells from V $\beta$ 8.1 transgenic mice responded in a correspondingly high proportion.<sup>39</sup> Thus, for cloned and normal heterogeneous T cell populations, recognition of the Mls<sup>a</sup> superantigen appears capable of inducing increased intracellular [Ca<sup>2+</sup>] as well as increased PI hydrolysis, potentially reflecting TCR-mediated PLC activation.

## References

1. Festenstein H (1973) Immunogenetic and biological aspects of *in vitro* lymphocyte allotransformation (MLR) in the mouse. *Transp Rev* 15:62-88
2. Abe R, Ryan JJ, Finkelman FD, Hodes RJ (1987) T cell recognition of Mls. T cell clones demonstrate polymorphism between Mls<sup>a</sup>, Mls<sup>b</sup>, and Mls<sup>d</sup>. *J Immunol* 138:373-379
3. Abe R, Ryan JJ, Hodes RJ (1987) Clonal analysis of the Mls system. A reappraisal of polymorphism and allelism among Mls<sup>a</sup>, Mls<sup>b</sup>, and Mls<sup>d</sup>. *J Exp Med* 165:1113-1129
4. Abe R, Ryan JJ, Hodes RJ (1987) Mls is not a single gene, allelic system: Mls<sup>a</sup> and Mls<sup>b</sup> are the products of non-allelic, unlinked genes. *J Exp Med* 166:1150-1155
5. Ryan J, Mond JJ, Finkelman FD (1987) The Mls<sup>d</sup>-defined primary mixed lymphocyte reaction: a composite response to Mls<sup>a</sup> and Mls<sup>b</sup> determinants. *J Immunol* 138:4085-4092
6. Abe R, Hodes RJ (1989) Properties of the Mls system: a revised formulation of Mls genetics and an analysis of T cell recognition of Mls determinants. *Immunol Rev* 107:5-28
7. Abe R, Hodes RJ (1989) T cell recognition of minor lymphocyte stimulating (Mls) gene products. *Annu Rev Immunol* 7:683-708
8. Abe R, Foo-Phillips M, Hodes RJ (1989) Analysis of Mls<sup>c</sup> genetics: a novel instance of genetic redundancy. *J Exp Med* 170:1059-1073
9. Pullen AM, Marrack P, Kappler JW (1988) The T cell repertoire is heavily influenced by tolerance to polymorphic self-antigens. *Nature* 335:796-801
10. Frankel WN, Rudy C, Coffin JM, Huber BT (1991) Linkage of Mls genes to endogenous mammary tumor viruses of inbred mice. *Nature* 349:526-528
11. Woodland DL, Happ MP, Gollob KJ, Palmer E (1991) An endogenous retrovirus mediating deletion of  $\alpha\beta$  T cells? *Nature* 349:529-530
12. Dyson PJ, Knight AM, Fairchild S, Simpson E, Tomonari K (1991) Genes encoding ligands for deletion of V $\beta$ 11 T cells cosegregate with mammary tumor virus genomes. *Nature* 349:531-533
13. Pullen AM, Choi Y, Kushnir E, Kappler J, Marrack P (1992) The open reading frames in the 3' long terminal repeats of several mouse mammary tumor virus integrants encode V $\beta$ 3-specific superantigens. *J Exp Med* 175:41-47
14. Wilson EK, Lai E, Concannon P, Barth RK, Hood LE (1988) Structure, organization and polymorphism of murine and human T cell receptor  $\alpha$  and  $\beta$  chain gene families. *Immunol Rev* 101:149-172
15. Matis LA (1990) The molecular basis of T cell specificity. *Annu Rev Immunol* 8:63-82
16. Kappler JW, Staerz U, White J, Marrack PC (1988) Self-tolerance eliminates T cells specific for Mls-modified products of the major histocompatibility complex. *Nature* 332:35-40
17. MacDonald HR, Schneider R, Lees RK, Howe RC, Acha-Orbea H, Festenstein H, Zinkernagel RM, Hengartner H (1988) T cell receptor V $\beta$  use predicts reactivity and tolerance to Mls<sup>a</sup>-encoded antigens. *Nature* 332:40-45
18. Abe R, Vacchio MS, Fox B, Hodes RJ (1988) Preferential expression of the T cell receptor V $\beta$ 3 gene by Mls<sup>c</sup> reactive T cells. *Nature* 335:827-830
19. Vacchio MS, Ryan JJ, Hodes RJ (1990) Characterization of the ligand(s) responsible for negative selection of V $\beta$ 11- and V $\beta$ 12-expressing T cells: effects of a new Mls determinant. *J Exp Med* 172:807-813
20. Abe R, Kanagawa O, Sheard MA, Malissen B, Foo-Phillips M (1991) Characterization of a new minor lymphocyte stimulatory system. I. Cluster of self antigens recognized by 'I-E-Reactive' V $\beta$ s, V $\beta$ 5, V $\beta$ 11, and V $\beta$ 12 T cell receptors for antigen. *J Immunol* 147:739-749
21. Fry AM, Matis LA (1988) Self tolerance alters T cell receptor expression in an antigen-specific MHC restricted immune response. *Nature* 335:830-832
22. Bill J, Appel VB, Palmer E (1988) An analysis of T cell receptor variable gene expression in major histocompatibility complex disparate mice. *Proc Natl Acad Sci USA* 85:9184-9188
23. Vacchio MS, Hodes RJ (1989) Selective decreases in T cell receptor V $\beta$  expression: decreased expression of specific V $\beta$  families is associated with expression of multiple MHC and non-MHC gene products. *J Exp Med* 170:1335-1346
24. Bill J, Kanagawa O, Woodland DL, Palmer E (1989) The MHC molecule I-E is necessary but not sufficient for the clonal deletion of V $\beta$ 11-bearing T cells. *J Exp Med* 169:1405-1419
25. Acha-Orbea H, Palmer E (1991) Mls—a retrovirus exploits the immune system. *Immunol Today* 12:356-361
26. Janeway CA (1991) Mls: makes a little sense. *Nature* 349:459-460
27. Callahan R, Drohan W, Gallahan D, D'Hoostelaere L, Potter M (1982) Novel class of mouse mammary tumor virus-related DNA sequences found in all species of Mus, including mice lacking the virus proviral genome. *Proc Natl Acad Sci USA* 79:4113-4117
28. Smith LR, Kono DH, Kammuller ME, Balderas RS, Theofilopoulos AN (1992) V $\beta$  repertoire in rats and implications for endogenous superantigens. *Eur J Immunol* 22:641-645

29. Ildstad ST, Vacchio MS, Markus PM, Hronakes ML, Wren SM, Hodes RJ (1992) Cross species transplantation tolerance: rat bone marrow derived cells can contribute to the ligand for negative selection of mouse TCR-V $\beta$  in chimeras tolerant to xenogenic antigens (mouse + rat-mouse). *J Exp Med* 175:147-157
30. Pullen AM, Wade T, Marrack P, Kappler JW (1990) Identification of the region of the T cell receptor  $\beta$  chain that interacts with the self-superantigen Mls-1<sup>a</sup>. *Cell* 61:1365-1374
31. Yui K, Komori S, Katsumata M, Siegel RM, Greene MI (1990) Self-reactive T cells can escape clonal deletion in T cell receptor V $\beta$ 8.1-transgenic mice. *Proc Natl Acad Sci USA* 87:7135-7139
32. Pullen AM, Bill J, Kubo RT, Marrack P, Kappler JW (1991) Analysis of the interaction site for the self superantigen Mls-1<sup>a</sup> on T cell receptor V $\beta$ . *J Exp Med* 173:1183-1192
33. Waanders GA, MacDonald HR (1992) Hierarchy of responsiveness *in vivo* and *in vitro* among T cells expressing distinct Mls-1<sup>a</sup>-reactive V $\beta$  domains. *Eur J Immunol* 22:291-293
34. Vacchio MS, Kanagawa O, Tomonari K, Hodes RJ (1992) Influence of T cell receptor V $\alpha$  expression on Mls<sup>a</sup> superantigen-specific T cell responses. *J Exp Med* 175:1405-1408
35. Pircher H, Hoffmann Rohrer U, Moskophidis D, Zinkernagel RM, Hengartner H (1991) Lower receptor avidity required for thymic clonal deletion than for effector T cell function. *Nature* 351:482-485
36. O'Rourke AM, Mescher M, Webb SR (1990) Activation of polyphosphoinositide hydrolysis in T cells by H-2 allo-antigen but not by Mls determinants. *Science* 249:171-174
37. Gaugler B, Langlet C, Schmitt-Verhulst AM, Guimezanes A (1991) Evidence for quantitative and qualitative differences in functional activation of Mls-reactive T cell clones and hybridomas by antigen or TcR/CD3 antibodies. *Eur J Immunol* 21:2581-2589
38. Patarca R, Wei F-Y, Iregui MV, Cantor H (1991) Differential induction of interferon  $\gamma$  gene expression after activation of CD4<sup>+</sup> T cells by conventional antigen and Mls superantigen. *Proc Natl Acad Sci USA* 88:2736-2739
39. Abe R, Ishida Y, Yui K, Katsumata M, Chused TM (1992) T cell receptor-mediated recognition of self ligand induces signaling in immature thymocytes before negative selection. *J Exp Med* 176:459-468