

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

AD-A266 660



Information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering the necessary data, completing and reviewing the collection of information, Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. REPORT DATE
2. REPORT TYPE AND DATES COVERED
Final 1 Sep 89-31 Dec 92

5. FUNDING NUMBERS

DAAL03-89-G-0107

Functional Differential Equations with Piecewise Continuous Argument

6. AUTHOR(S)

Ganzaram S. Ladde

7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)

Department of Mathematics
The University of Texas at Arlington
Arlington, Texas 76019

8. PERFORMING ORGANIZATION REPORT NUMBER

9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)

U. S. Army Research Office
P. O. Box 12211
Research Triangle Park, NC 27709-2211

10. SPONSORING/MONITORING AGENCY REPORT NUMBER

ARO 26739.16-MA-SAH

11. SUPPLEMENTARY NOTES

The view, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy, or decision, unless so designated by other documentation.

12a. DISTRIBUTION/AVAILABILITY STATEMENT

Approved for public release; distribution unlimited.

12b. DISTRIBUTION CODE

13. ABSTRACT (Maximum 200 words)

By developing certain auxiliary results, a modified version of the stochastic averaging principle is formulated to investigate stochastic dynamic systems consisting of fast and slow phenomena. Using the properties of optimal feedback control laws, the robust control problem of distributed parameter systems is analyzed. In addition, the stability analysis of a large-scale distributed parameter system is also investigated. Applications are given to illustrate the results.

93 7 08 103

93-15521



35 PK

14. SUBJECT TERMS

15. NUMBER OF PAGES

16. PRICE CODE

17. SECURITY CLASSIFICATION OF REPORT

UNCLASSIFIED

18. SECURITY CLASSIFICATION OF THIS PAGE

UNCLASSIFIED

19. SECURITY CLASSIFICATION OF ABSTRACT

UNCLASSIFIED

20. LIMITATION OF ABSTRACT

UL

Functional Differential Equations with Piecewise
Continuous Argument

FINAL REPORT

Dr. Joseph Wiener

15 April 1993

U. S. Army Research Office

Contract/Grant Number
DAAL03-89-G-0107

University of Texas-Pan American
1201 W University Drive, Edinburg TX
E-mail address: jw6bdb@panam.edu

APPROVED FOR PUBLIC RELEASE;
DISTRIBUTION UNLIMITED.

FUNCTIONAL DIFFERENTIAL EQUATIONS WITH PIECEWISE CONTINUOUS ARGUMENT

JOSEPH WIENER

April 15, 1993

CONTENTS

1. Statement of the Problem Studied	2
2. Summary of the Most Important Results	6
2.1. Boundary-Value Problems	
2.2. Initial-Value Problems	
2.3. Wave Equations with Discontinuous Time Delay	
3. List of Publications and Technical Reports	32
References	33

DTIC QUALITY INSPECTED 6

Accession For	
NTIS GRA&I	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By	
Distribution/	
Availability Codes	
Dist.	Avail and/or Special
A-1	

1. STATEMENT OF THE PROBLEM STUDIED

Functional differential equations (FDE) with delay provide a mathematical model for a physical or biological system in which the rate of change of the system depends upon its past history. Although the general theory and basic results for FDE have by now been thoroughly investigated, the literature devoted to this area of research continues to grow very rapidly. The number of interesting works is very large, so that our knowledge of FDE has been substantially enlarged in recent years. Naturally, new important problems and directions arise continually in this intensively developing field.

The report summarizes the results in the study and addresses the need for further investigation of generalized solutions to broad classes of FDE. The project concentrated on differential equations with piecewise continuous arguments (EPCA), the exploration of which has been initiated in our papers a few years ago. These equations arise in an attempt to extend the theory of FDE with continuous arguments to differential equations with discontinuous arguments. This task is also of considerable applied interest since EPCA include, as particular cases, impulsive and loaded equations of control theory and are similar to those found in some biomedical models. A typical EPCA contains arguments that are constant on certain intervals. A solution is defined as a continuous, sectionally smooth function that satisfies the equation within these intervals. Continuity of a solution at a point joining any two consecutive intervals leads to recursion relations for the solution at such points. Hence, the solutions are determined by a finite set of initial data, rather than by an initial function as in the case of general FDE. Therefore, underlying each EPCA is a dynamical system governed by a difference equation of a discrete argument which describes its stability, oscillation, and periodic properties. It is not surprising then that recent work on EPCA has caused a new surge in the study of difference equations. Of significant interest is the exploration of partial differential equations (PDE) with piecewise continuous delays. Boundary and initial-value problems for some EPCA with partial derivatives were considered and the behavior of their solutions investigated. The results were also extended to equations with positive definite operators

in Hilbert spaces. This topic is of great theoretical, computational, and applied value since it opens the possibility of approximating complicated problems of mathematical physics by simpler EPCA.

It is well known that profound and close links exist between functional and functional differential equations. Thus the study of the first often enables one to predict properties of differential equations of neutral type. On the other hand, some methods for the latter in the special case when the argument deviation vanishes at individual points have been used to investigate functional equations. Functional equations are directly related to difference equations of a discrete argument, and bordering on difference equations are impulsive FDE with impacts and switching and loaded equations (that is, those including values of the unknown solution for given constant values of the argument). *The argument deviations of the EPCA considered in the project vanish at countable sets of points, and it would be interesting to investigate the relationship between EPCA and functional equations. Another deserving direction of future research is the exploration of hybrid systems consisting of EPCA and functional equations. Furthermore, EPCA are intrinsically closer to difference rather than to differential equations. Equations with piecewise constant delay can be used to approximate differential equations that contain discrete delays. It would be useful to draw a detailed comparison of the qualitative and asymptotic properties of differential equations with continuous arguments and their EPCA approximations, which has been widely used for ordinary differential equations and their difference approximations. Since the arguments of an EPCA have intervals of constancy we must relinquish smoothness of the solutions, but we still retain their continuity. This enables us to derive a homogeneous difference equation for the values of a solution at the endpoints of the intervals of constancy and to employ it in the study of the original EPCA, thus revealing remarkable asymptotic, oscillatory, and periodic properties of this type of FDE. Of course, it is possible to further generalize the definition of a solution for an EPCA, by abandoning its continuity, and to include in the framework of EPCA the impulsive functional differential equations.*

Along with mildly weakened solutions of EPCA, the project also ex-

plored generalized-function solutions of ordinary differential and functional differential equations. The unifying theme of the project is the development of theoretically meaningful and potentially applicable generalized concepts of solutions for important classes of FDE. A common feature of these equations is that their arguments have a fixed point. Thus, the argument of a typical EPCA is the greatest-integer function, and we also focus on FDE with linearly transformed arguments. Hence, it is natural to pose the initial-value problem for such equations not on an interval but at a number of individual points. Contrary to general functional differential equations, EPCA of all types (retarded, advanced, mixed, neutral) have two-sided solutions, and FDE with linearly transformed arguments possess, under certain conditions, analytic or entire solutions. Some methods in the theory of entire solutions are applied to prove stability theorems for linear EPCA with variable coefficients. Integral transformations establish close connections between entire and generalized functions (distributions). Therefore, a unified approach may be used in the study of both distributional and entire solutions to some classes of linear ordinary and functional differential equations.

Recently there has been considerable interest in problems concerning the existence of solutions to differential and functional differential equations in various spaces of generalized functions. Many important areas in mathematics and theoretical physics employ the methods of distribution theory. Generalized functions are continuous linear functionals on spaces of infinitely smooth functions with certain conditions of decay at infinity. They provide a suitable framework where major analytical operations such as differentiation can be performed. Furthermore, the importance of the class of generalized functions stems from the fact that it includes the set of regular distributions represented by locally integrable functions. There is an abundance of singular distributions, and the Dirac delta function is one of them. It is well known that normal linear homogeneous systems of ordinary differential equations (ODE) with infinitely smooth coefficients have no singular distributional solutions. However, these solutions may appear in the case of equations whose coefficients have singularities. We de-

velop the methods of study and establish some major results for linear ODE in the space of finite-order distributions (finite linear combinations of the delta function and its derivatives). An existence criterion of such solutions for any linear ODE is found. Necessary and sufficient conditions are discovered for the simultaneous existence of solutions to linear ODE in the form of rational functions and finite-order distributions. The results are also used in the study of polynomial solutions to some important classical equations. Then distributional solutions of certain classes of ODE and FDE are presented as infinite series of the delta function and its derivatives. Existence and nonexistence theorems in spaces of infinite-order distributions are obtained for linear equations with polynomial coefficients and used to explore their entire solutions. We emphasize and investigate the conditions when linear FDE with polynomial coefficients and linearly transformed arguments have entire solutions of zero order. This is a remarkable dissimilarity between the behavior of FDE and ODE since first-order algebraic ODE have no entire transcendental solutions of order less than $\frac{1}{2}$. An equally striking phenomenon is the existence of distributional solutions for linear homogeneous FDE without singularities in the coefficients. In other words, distributional solutions to linear homogeneous FDE may be originated either by singularities of their coefficients or by argument deviations. Recent studies have shown that nonexistence of infinite order distributional solutions for linear time-dependent delay equations with real analytic coefficients implies nonexistence of small solutions (approaching zero faster than any exponential as t tends to infinity), which is important in the qualitative theory of FDE. *It would be nice to extend the results on distributional and entire solutions and their interplay to partial differential equations (PDE).* As a first step in this direction, one could take a linear PDE in two independent variables with polynomial coefficients that admits separation of variables, then consider a series whose terms are products of distributional solutions of the ordinary differential equations arising after separation.

2. SUMMARY OF THE MOST IMPORTANT RESULTS

We shall describe now some of the work that has been done in the project on the differential equations that we call equations with piecewise continuous arguments, or EPCA. A brief survey of the present status of this research is given in [5]

A typical EPCA is of the form

$$x'(t) = f(t, x(t), x(h(t))),$$

where the argument $h(t)$ has intervals of constancy. For example, in [4] equations with $h(t) = [t]$, $[t - n]$, $t - n[t]$ were investigated, where n is a positive integer and $[\cdot]$ denotes the greatest-integer function. Note that $h(t)$ is discontinuous in these cases, and although the equation fits within the general paradigm of delay differential or functional differential equations, the delays are discontinuous functions. Also note that the equation is nonautonomous, since the delays vary with t . Moreover, as we have mentioned, the solutions are determined by a finite set of initial data, rather than by an initial function, as in the case of general FDE. In fact, EPCA have the structure of continuous dynamical systems within intervals of certain lengths. Continuity of a solution at a point joining any two consecutive intervals then implies recursion relations for the solution at such points. Therefore, EPCA represent a hybrid of continuous and discrete dynamical systems and combine the properties of both differential and difference equations.

An equation in which $x'(t)$ is given by a function of x evaluated at t and at arguments $[t], \dots, [t - N]$, where N is a non-negative integer, may be called of retarded or delay type. If the arguments are t and $[t + 1], \dots, [t + N]$, the equation is of advanced type. If both these types of arguments appear in the equation, it may be called of mixed type. If the derivative of highest order appears at t and at another point, the equation is generally said to be of neutral type. All types of EPCA share similar characteristics. First of all, it is natural to pose the initial-value problem for such equations not on an interval but at a number of individual points. Secondly, for ordinary differential equations with a continuous vector field the solution exists to the right and left of the initial t -value. For retarded FDE, this is not necessarily the case [6].

Furthermore, it appears that advanced equations, in general, lose their margin of smoothness, and the method of successive integration shows that after several steps to the right from the initial interval the solution may even not exist. However, two-sided solutions do exist for all types of EPCA. Finally, the problems for EPCA studied so far are closely related to ordinary difference equations and indeed have stimulated new work on these.

It is important to note that EPCA provide the simplest examples of differential equations capable of displaying chaotic behavior. For instance, following Ladas [7], one can see that the unique solution of the initial-value problem

$$x'(t) = 3x([t]) - x^2([t]), \quad x(0) = c_0$$

where $[t]$ is the greatest-integer function, has the property that

$$x(n+1) = 4x(n) - x^2(n), \quad n = 0, 1, \dots$$

If we choose $c_0 = 4 \sin^2(\pi/9)$, then the unique solution of this difference equation is

$$x(n) = 4 \sin^2 \left(2^n \frac{\pi}{9} \right),$$

which has period three. By the well-known result [8] which states that "period three implies chaos," the solution of the above differential equation exhibits chaos. Furthermore, the equation of Carvalho and Cooke

$$x'(t) = ax(t)(1 - x([t]))$$

is analogous to the famous logistic differential equation, but t in one argument has been replaced by $[t]$. As a result, the equation has solutions that display complicated dynamics [2]. It seems likely that other simple nonlinear EPCA may display other interesting behavior.

The numerical approximation of differential equations can give rise to EPCA in a natural way, although it is unusual to take this point of view. For example, the simple Euler scheme for a differential equation $x'(t) = f(x(t))$ has the form $x_{n+1} - x_n = hf(x_n)$, where $x_n = x(nh)$ and h is the step size. This is equivalent to the EPCA

$$x'(t) = f(x([t/h]h)).$$

Impulsive differential equations and loaded equations of control theory fit within the general paradigm of EPCA. Another potential application of EPCA is the stabilization of hybrid control systems with feedback delay. By a hybrid system we mean one with a continuous plant and with discrete (sampled) controller. Some of these systems may be described by EPCA [3].

EPCA have only been researched for a few years. In each of the areas — existence, asymptotic behavior, periodic and oscillating solutions, approximation, application to control theory, biomedical models, and problems of mathematical physics — there appears to be ample opportunity for extending the known results.

2.1. Boundary-Value Problems. The main emphasis in the project was the study of partial differential equations (PDE) with piecewise continuous delay. The first fundamental paper [9] in this direction appeared in 1991. It has been shown in [9] that these equations naturally arise in the process of approximating PDE by using piecewise constant arguments. Thus, if in the equation

$$u_t = a^2 u_{xx} - bu,$$

which describes heat flow in a rod with both diffusion $a^2 u_{xx}$ along the rod and heat loss (or gain) across the lateral sides of the rod, the lateral heat change is measured at discrete times, then we get an equation with piecewise constant argument (EPCA)

$$\begin{aligned} u_t(x, t) &= a^2 u_{xx}(x, t) - bu(x, nh), \\ t &\in [nh, (n+1)h], \quad n = 0, 1, \dots \end{aligned}$$

where $h > 0$ is some constant. This equation can be written in the form

$$u_t(x, t) = a^2 u_{xx}(x, t) - bu(x, [t/h]h), \quad (1)$$

where $[\cdot]$ designates the greatest-integer function.

The diffusion-convection equation

$$u_t = a^2 u_{xx} - ru_x$$

describes, for instance, the concentration $u(x, t)$ of a pollutant carried along in a stream moving with velocity r . The term $a^2 u_{xx}$ is the diffusion contribution and $-ru_x$ is the convection component. If the convection part is measured at discrete times nh , the process results in the equation

$$u_t(x, t) = a^2 u_{xx}(x, t) - ru_x(x, [t/h]h). \quad (2)$$

These examples indicate at the considerable potential of EPCA as an analytical and computational tool in solving some complicated problems of mathematical physics. Therefore, it is important to investigate boundary-value problems (BVP) and initial-value problems (IVP) for EPCA in partial derivatives, and explore the influence of certain discontinuous delays on the behavior of solutions to some typical problems of mathematical physics.

The topic of [9] is the BVP consisting of the equation

$$\frac{\partial u(x, t)}{\partial t} + P\left(\frac{\partial}{\partial x}\right) u(x, t) = Q\left(\frac{\partial}{\partial x}\right) u\left(x, \left[\frac{t}{h}\right]h\right), \quad (3)$$

where P and Q are polynomials of the highest degree m with coefficients that may depend only on x , the boundary conditions

$$L_j u = \sum_{k=1}^m (M_{jk} u^{(k-1)}(0) + N_{jk} u^{(k-1)}(1)) = 0, \quad (4)$$

(M_{jk} and N_{jk} are constants, $j = 1, \dots, m$)

and the initial condition

$$u(x, 0) = u_0(x). \quad (5)$$

Here $[\cdot]$ designates the greatest-integer function, $(x, t) \in [0, 1] \times [0, \infty)$, and $h = \text{const.} > 0$. Conditions (4) will be written briefly as

$$Lu = 0.$$

An important result has been established that BVP (3), (4), (5) has a solution in $[0, 1] \times [nh, (n+1)h]$, if the following hypotheses hold true:

(i) The boundary-value problem

$$P\left(\frac{d}{dx}\right)X - \lambda X = 0, \quad LX = 0$$

is self-adjoint, all its eigenvalues λ_j are positive.

(ii) For each λ_j , the roots of the equation $P(z) - \lambda_j = 0$ have non-positive real parts.

(iii) The initial function $u_0(x) \in C^m[0, 1]$ satisfies (4).

The solution $u_n(x, t)$ of BVP (3), (4), (5) on the interval $nh \leq t < (n+1)h$ is represented in the form of a Fourier series

$$u_n(x, t) = \sum_{j=1}^{\infty} X_j(x) T_{nj}(t), \quad (6)$$

where $X_j(x)$ are the eigenvalues of the operator P . The functions $T_{nj}(t)$ are solutions of ordinary EPCA that arise after separation of variables.

For instance, in $[0, 1] \times [nh, (n+1)h]$, the solution $u_n(x, t)$ of Eq. (1) with boundary conditions $u_n(x, nh) = u_n(x)$ is sought in form (6). Separation of variables produces

$$X_j(x) = \sqrt{2} \sin(\pi j x), \quad T'_{nj}(t) + a^2 \pi^2 j^2 T_{nj}(t) = -b T_{nj}(nh),$$

whence

$$T_{nj}(t) = C_{nj} e^{-a^2 \pi^2 j^2 (t-nh)} - \frac{b}{a^2 \pi^2 j^2} T_{nj}(nh).$$

We put $t = nh$ in this equation and get

$$C_{nj} = \left(1 + \frac{b}{a^2 \pi^2 j^2}\right) T_{nj}(nh),$$

that is,

$$T_{nj}(t) = E_j(t - nh) T_{nj}(nh),$$

where

$$E_j(t) = e^{-a^2 \pi^2 j^2 t} - \left(1 - e^{-a^2 \pi^2 j^2 t}\right) \frac{b}{a^2 \pi^2 j^2}. \quad (7)$$

At $t = (n+1)h$ we have

$$T_{nj}((n+1)h) = E_j(h) T_{nj}(nh)$$

and since

$$T_{nj}((n+1)h) = T_{n+1,j}((n+1)h),$$

then

$$T_{n+1,j}((n+1)h) = E_j(h)T_{nj}(nh)$$

and

$$T_{nj}(nh) = E_j^n(h)T_{0j}(0).$$

Therefore,

$$T_{nj}(t) = E_j(t - nh)E_j^n(h)T_{0j}(0)$$

and

$$u_n(x, t) = \sum_{j=1}^{\infty} \sqrt{2} E_j^n(h) T_{0j}(0) E_j(t - nh) \sin(\pi j x). \quad (8)$$

Putting $t = 0, n = 0$ gives

$$u_0(x) = \sum_{j=1}^{\infty} T_{0j}(0) \sqrt{2} \sin(\pi j x) dx$$

and

$$T_{0j}(0) = \sqrt{2} \int_0^1 u_0(x) \sin(\pi j x) dx.$$

If $|E_j(h)| < 1$, then solution (8) decays exponentially as $t \rightarrow \infty$, uniformly with respect to x . From (7) it follows that this is true if

$$-a^2 \pi^2 < b < a^2 \pi^2 \frac{e^{a^2 \pi^2 h} + 1}{e^{a^2 \pi^2 h} - 1}.$$

Furthermore, from the equations

$$T_{nj}(nh) = E_j^n(h)T_{0j}(0), \quad T_{nj}((n+1)h) = E_j^{n+1}(h)T_{0j}(0)$$

we see that $T_{nj}(nh)T_{nj}((n+1)h) < 0$ if $E_j(h) < 0$. The latter inequality holds true if

$$b > \frac{a^2 \pi^2}{e^{a^2 \pi^2 h} - 1}. \quad (9)$$

Hence, under condition (9) each function $T_{nj}(t)$ ($j = 1, 2, \dots$) has a zero in the interval $[nh, (n+1)h]$, in sharp contrast to the functions $T_j(t)$ in the Fourier expansion for the solution of the equation $u_t = a^2 u_{xx} - bu$ without time delay. Moreover, the inequality $E_j(h) < 0$ takes place for sufficiently large j and any $b > 0$. Therefore, for $b > 0$ and sufficiently large j , the functions $T_{nj}(t)$ are oscillatory.

Eq. (2) on $nh \leq t < (n+1)h$ becomes

$$\frac{\partial u_n(x, t)}{\partial t} = a^2 \frac{\partial^2 u_n(x, t)}{\partial x^2} - ru'_n(x),$$

and we differentiate the latter with respect to t to obtain the equation

$$\frac{\partial y_n}{\partial t} = a^2 \frac{\partial^2 y_n}{\partial x^2}, \quad y_n = \frac{\partial u_n}{\partial t},$$

whose solution is sought in form (6). Separation of variables leads to the equations

$$X''(x) + \lambda X(x) = 0, \quad T'_n(t) + a^2 \lambda T_n(t) = 0,$$

and the boundary conditions $u_n(0, t) = u_n(1, t) = 0$ give $\lambda_j = j^2 \pi^2$ and

$$y_n(x, t) = \sum_{j=1}^{\infty} \sqrt{2} T_{nj}(nh) e^{-a^2 \pi^2 j^2 (t-nh)} \sin(\pi j x).$$

Since

$$y_n(x, nh) = a^2 u''_n(x) - ru'_n(x), \quad u_n(x) = u_n(x, nh),$$

then

$$a^2 u''_n(x) - ru'_n(x) = \sum_{j=1}^{\infty} \sqrt{2} T_{nj}(nh) \sin(\pi j x)$$

and

$$\begin{aligned} T_{nj}(nh) = & -a^2 \pi^2 j^2 \sqrt{2} \int_0^1 u_n(x) \sin(\pi j x) dx \\ & + r \pi j \sqrt{2} \int_0^1 u_n(x) \cos(\pi j x) dx. \end{aligned}$$

Finally

$$u_n(x, t) = u_n(x) + \sum_{j=1}^{\infty} \frac{\sqrt{2} T_{nj}(nh) (1 - e^{-a^2 \pi^2 j^2 (t-nh)}) \sin(\pi j x)}{a^2 \pi^2 j^2}.$$

Given the initial function $u(x, 0) = u_0(x)$, we can find the coefficients $T_{0j}(0)$ and the solution $u_0(x, t)$ on $0 \leq t \leq h$. Since $u_0(x, h) = u_1(x)$, we can calculate the coefficients $T_{1j}(h)$ and the solution $u_1(x, t)$ on $h \leq t \leq 2h$. By the method of steps the solution can be extended to any interval $[nh, (n+1)h]$.

The equation

$$iq \frac{\partial u(x, t)}{\partial t} = -\frac{q^2}{2m_0} \frac{\partial^2 u(x, t)}{\partial x^2} + V(x) u\left(x, \left[\frac{t}{h}\right] h\right)$$

is a piecewise constant analogue of the one-dimensional Schrödinger equation

$$iq \psi_t(x, t) = \frac{-q^2}{2m_0} \psi_{xx}(x, t) + V(x) \psi(x, t).$$

If $u(x, t)$ satisfies conditions (4) and (5), with $m = 2$, then separation of variables produces a formal solution

$$u_n(x, t) = \sum_{j=1}^{\infty} C_{nj} e^{-i\lambda_j(t-nh)/q} X_j(x) + P^{-1} Q u_n(x),$$

for $nh \leq t \leq (n+1)h$. Here, $X_j(x)$ are the eigenfunctions of the operator $q^2(d^2/dx^2)/2m_0$, and $P^{-1} Q u_n(x)$ is the solution $v_n(x)$ of the equation

$$q^2 v_n''(x) = 2m_0 V(x) u_n(x)$$

that satisfies (4).

The Fourier method was also used to find weak solutions of the boundary-value problem (3), (4), (5) and it is easily generalized to similar problems in Hilbert space. First, we remind a few well known definitions. Let H be a Hilbert space and let P be a linear operator in H (additive and homogeneous but, possibly, unbounded) whose domain $\mathcal{D}(P)$ is dense in H , that is $\overline{\mathcal{D}(P)} = H$. The operator P is called symmetric if $(Pu, v) = (u, Pv)$, for any $u, v \in \mathcal{D}(P)$. If P is symmetric, then (Pu, v)

is a symmetric bilinear functional and (Pu, u) is a quadratic form. A symmetric operator P is called positive if $(Pu, u) \geq 0$ and $(Pu, u) = 0$ if and only if $u = 0$. A symmetric operator P is called positive definite if there exists a constant $\gamma^2 > 0$ such that $(Pu, u) \geq \gamma^2 \|u\|^2$. With every positive operator P a certain Hilbert space H_P can be associated, which is called the energy space of P . It is the completion of $\mathcal{D}(P)$, with the inner product $(u, v)_P = (Pu, v)$; $u, v \in \mathcal{D}(P)$. This product induces a new norm $\|u\|_P = (Pu, u)^{1/2}$, $u \in \mathcal{D}(P)$, and if P is positive definite, then $\|u\| \leq \gamma^{-1} \|u\|_P$. Since $\mathcal{D}(P)$ is dense in H , it follows by using the latter inequality that the energy space H_P of a positive definite operator P is dense in the original space H .

Assuming P is positive definite, we may consider the solution $u(x, t)$ of (3), (4), (5) for a fixed t as an element of H_P . If $\mathcal{D}(Q) \subset H$, then $Qu(x, [t/h]h)$ may be treated as an abstract function $Qu([t/h]h)$ with the values in H . Therefore, the given BVP is reduced to the abstract Cauchy problem

$$\frac{du}{dt} + Pu = Qu \left(\left[\frac{t}{h} \right] h \right), \quad t > 0, u|_{t=0} = u_0 \in H. \quad (10)$$

If (10) has a solution, we multiply each term by an arbitrary function $g(t) \in H_P$ in the sense of inner product in H and get on the interval $nh \leq t < (n+1)h$ the equation

$$\left(\frac{du}{dt}, g \right) + (u, g)_P = (Qu_n, g), \quad (11)$$

where $u_n = u(nh)$. Conversely, if $u \in C^1((nh, (n+1)h); \mathcal{D}(P))$ for all integers $n \geq 0$ and satisfies (11), then it also satisfies Eq. (10). Indeed, if $u \in \mathcal{D}(P)$, then $(u, g)_P = (Pu, g)$, and (11) can be written

$$\left(\frac{du}{dt} + Pu - Qu_n, g \right) = 0, \quad nh \leq t < (n+1)h.$$

Since H_P is dense in H , then $u(t)$ is a solution of Eq. (10).

Definition. An abstract function $u(t): [0, \infty) \rightarrow H$ is called a *weak solution* of problem (10) if it satisfies the conditions:

- (i) $u(t)$ is continuous for $t \geq 0$ and strongly continuously differentiable for $t > 0$, with the possible exception of the points $t = nh$ where one-sided derivatives exist.
- (ii) $u(t)$ is continuous for $t > 0$ as an abstract function with the values in H_P and satisfies Eq. (11) on each interval $nh \leq t < (n+1)h$, for any function $g(t): [0, \infty) \rightarrow H_P$.
- (iii) $u(t)$ satisfies the initial condition (10), that is,

$$\lim_{t \rightarrow 0} \|u(t) - u_0\|_H = 0.$$

A weak solution $u(t)$ is also an ordinary solution if $u(t) \in \mathcal{D}(P)$, for any $t > 0$, and $u(x, t) \rightarrow u_0(x)$ as $t \rightarrow 0$ not only in the norm of H but uniformly as well. It is said that a symmetric operator P has a discrete spectrum if it has an infinite sequence $\{\lambda_j\}$ of eigenvalues with a single limit point at infinity and a sequence $\{X_j\}$ of eigenfunctions which is complete in H . Suppose the operator P in (11) is positive definite and has a discrete spectrum and assume the existence of a solution $u(t) = u(x, t)$ to Eq. (11) with the condition $u(0) = u_0$. On the interval $nh \leq t < (n+1)h$ this solution can be expanded into series (6), where $T_j(t) = (u(t), X_j)$. To find the coefficients $T_j(t)$, we put $g(t) = X_k$ in (11) and since X_k does not depend on t , then

$$\left(\frac{du(t)}{dt}, X_k \right) = \frac{d}{dt} (u(t), X_k) = T'_k(t),$$

$$(u, X_k)_P = (Pu, X_k) = (u, PX_k) = \lambda_k(u, X_k) = \lambda_k T_k(t),$$

which leads to the equation

$$T'_{nj}(t) + \lambda_j T_{nj}(t) = (Qu_n, X_j)$$

By selecting a proper space H , a weak solution corresponding to conditions (4) can be constructed. A theorem has been stated in [10] that if P and Q are linear operators in a Hilbert space and P is positive definite with a discrete spectrum, then there exists a unique weak solution of problem (10).

2.2. Initial-Value Problems. This topic has been explored recently by Wiener and Debnath [10]. Eq. (3) with constant coefficients and initial condition (5) has been considered in the domain

$$(x, t) \in \Omega = (-\infty, \infty) \times [0, \infty).$$

Let $u_n(x, t)$ be the solution of the given problem on $nh \leq t < (n+1)h$, then

$$\frac{\partial u_n(x, t)}{\partial t} + Pu_n(x, t) = Qu_n(x), \quad (12)$$

where

$$u_n(x) = u_n(x, nh). \quad (13)$$

Write

$$u_n(x, t) = w_n(x, t) + v_n(x),$$

which gives the equation

$$\frac{\partial w_n}{\partial t} + Pw_n + Pv_n(x) = Qu_n(x),$$

and require that

$$\frac{\partial w_n}{\partial t} + Pw_n = 0, \quad (14)$$

$$Pv_n(x) = Qu_n(x). \quad (15)$$

If $v_n(x)$ is a solution of ODE (15), then at $t = nh$ we have

$$w_n(x, nh) = u_n(x) - v_n(x), \quad (16)$$

and it remains to consider Eq. (14) with initial condition (16). It is well known that the solution $E(x, t)$ of the problem

$$\frac{\partial w}{\partial t} + Pw = 0, \quad w(x, 0) = w_0(x), \quad (17)$$

with $w_0(x) = \delta(x)$, where $\delta(x)$ is the Dirac delta functional, is called its *fundamental solution*. The solution of IVP (17) is given by the convolution

$$w(x, t) = E(x, t) * w_0(x). \quad (18)$$

Hence, the solution of problem (14), (16) can be written as

$$w_n(x, t) = E(x, t - nh) * w_n(x, nh), \quad (19)$$

and the solution of (12), (13) is

$$u_n(x, t) = E(x, t - nh) * (u_n(x) - v_n(x)) + v_n(x), \quad (20)$$

$$(nh \leq t < (n+1)h).$$

Continuity of the solution at $t = (n+1)h$ implies

$$u_n(x, (n+1)h) = u_{n+1}(x, (n+1)h) = u_{n+1}(x),$$

that is,

$$u_{n+1}(x) = E(x, h) * (u_n(x) - v_n(x)) + v_n(x). \quad (21)$$

Formulas (20), (21) successively determine the solution of IVP (3), (5) on each interval $nh \leq t \leq (n+1)h$. Indeed, from $Pv_0(x) = Qu_0(x)$ we find $v_0(x)$ and substitute both $u_0(x)$ and $v_0(x)$ in (20) and (21) to obtain $u_0(x, t)$ and $u_1(x)$. Then we use $u_1(x)$ in (15) to find $v_1(x)$ and substitute $u_1(x)$ and $v_1(x)$ in (20) and (21), which yields $u_1(x, t)$ and $u_2(x)$. Continuing this procedure leads to $u_n(x, t)$, the solution of (3) on $[nh, (n+1)h]$. The solution $v_n(x)$ of (15) is defined to within an arbitrary polynomial $q(x)$ of degree $< m$. Since $q(x)$ is a solution of Eq. (17) with the initial condition $w(x, 0) = q(x)$, then $q(x) = E(x, t) * q(x)$, and $q(x)$ cancels in the formulas (20), (21). This proves that if Eq. (17) with $w(x, 0) = u_0(x)$ has a unique solution on $t \in (0, \infty)$, then there exists a unique solution of IVP (3), (5) on $(0, \infty)$ and it is given by (20), for each interval $nh \leq t < (n+1)h$. Thus, there exist unique solutions of Eqs. (1) and (2), with $u(x, 0) = u_0(x)$, in the class of functions that grow to infinity slower than $\exp(x^2)$ as $|x| \rightarrow \infty$. For Eqs. (1) and (2) we have

$$v_n(x) = a^{-2}b \int_0^x (x-s)u_n(s) ds \quad \text{and} \quad v_n(x) = a^{-2}r \int_0^x u_n(s) ds,$$

respectively, and $E(x, t) = \exp(-x^2/4a^2t)/2a\sqrt{\pi t}$.

Formula (20) for the solution of the problem

$$u_t(x, t) = a^2 u_{xx}(x, t) - bu_{xx} \left(x, \left[\frac{t}{h} \right] h \right), \quad u(x, 0) = u_0(x)$$

on $nh \leq t < (n+1)h$ becomes

$$u_n(x, t) = \left(1 - \frac{b}{a^2}\right) E(x, t - nh) * u_n(x) + \frac{b}{a^2} u_n(x),$$

where $E(x, t)$ is the same as in Eqs. (1) and (2).

The above method may also be used to solve IVP for PDE of any order in t with piecewise constant delay or systems of such equations. In the latter case, P and Q in (3) are square matrices of linear differential operators and $u(x, t)$ is a vector function. Thus, the solution $u_n(x, t)$ of the problem

$$\begin{aligned} u_{tt}(x, t) &= a^2 u_{xx}(x, t) - bu_{xx}(x, [t]), \\ u(x, 0) &= f_0(x), \quad u_t(x, 0) = g_0(x) \end{aligned}$$

on $n \leq t < n+1$ is sought in the form $u_n(x, t) = w_n(x, t) + v_n(x)$ whence $a^2 v_n''(x) - bu_n''(x, n) = 0$ and $\partial^2 w_n / \partial t^2 = a^2 \partial^2 w_n / \partial x^2$. Setting $u(x, n) = f_n(x)$, $u_t(x, n) = g_n(x)$ gives

$$v_n(x) = a^{-2} b f_n(x), \quad w(x, n) = (1 - a^{-2} b) f_n(x), \quad w_t(x, n) = g_n(x),$$

and

$$\begin{aligned} u_n(x, t) &= \frac{b}{a^2} f_n(x) + \left(1 - \frac{b}{a^2}\right) \frac{f_n(x - a(t - n)) + f_n(x + a(t - n))}{2} \\ &\quad + \frac{1}{2a} \int_{x-a(t-n)}^{x+a(t-n)} g_n(s) ds. \end{aligned}$$

Putting $t = n+1$ produces the recursion relations

$$\begin{aligned} f_{n+1}(x) &= \frac{b}{a^2} f_n(x) + \left(1 - \frac{b}{a^2}\right) \frac{f_n(x - a) + f_n(x + a)}{2} \\ &\quad + \frac{1}{2a} \int_{x-a}^{x+a} g_n(s) ds, \end{aligned}$$

$$\begin{aligned} g_{n+1}(x) &= \left(1 - \frac{b}{a^2}\right) \frac{a f_n'(x + a) - a f_n'(x - a)}{2} \\ &\quad + \frac{1}{2} (g_n(x + a) + g_n(x - a)). \end{aligned}$$

Loaded partial differential equations have properties similar to those of equations with piecewise constant delay. The IVP for the following class of loaded equations

$$\frac{\partial u(x, t)}{\partial t} = P\left(\frac{\partial}{\partial x}\right) u(x, t) + \sum_{j=1}^q Q_j\left(\frac{\partial}{\partial x}\right) u(x, t_j), \quad (22)$$

$$u(x, 0) = u_0(x)$$

was considered in [1] and [10], where $(x, t) \in \mathbb{R}^n \times [0, T]$, the $t_j \in (0, T]$ are given, $P(s)$ and $Q_j(s)$ are polynomials in $s = (s_1, \dots, s_n)$, and $\sum |Q_j(s)| \not\equiv 0$. Eq. (22) arises in solving certain inverse problems for systems with elements concentrated at specific moments of time. The Fourier transform $U(s, t)$ of $u(x, t)$ satisfies the equation

$$U_t(s, t) = P(is)U(s, t) + \sum_{j=1}^q Q_j(is)U(s, t_j),$$

whence,

$$U(s, t) = U_0(s)e^{P(is)t} + k(P(is), t) \sum_{j=1}^q Q_j(is)U(s, t_j), \quad (23)$$

where $U_0(s)$ is the Fourier transform of $u_0(x)$ and

$$k(P(is), t) = \int_0^t e^{P(is)y} dy.$$

Denote

$$A_j = U_0(s)e^{P(is)t_j}, \quad k_j = k(P(is), t_j), \quad B = \sum_{j=1}^q Q_j(is)U(s, t_j), \quad (24)$$

then multiply by $Q_j(is)$ each of the equations

$$U(s, t_j) = A_j + k_j B, \quad j = 1, \dots, q$$

and add them. Hence,

$$B = \sum_{j=1}^q A_j Q_j(is) + B \sum_{j=1}^q k_j Q_j(is)$$

or

$$\left(1 - \sum_{j=1}^q k_j Q_j(is)\right) B = \sum_{j=1}^q A_j Q_j(is). \quad (25)$$

The equation

$$\Delta(s) \equiv 1 - \sum_{j=1}^q Q_j(is) k(P(is), t_j) = 0 \quad (26)$$

is called the characteristic equation for (22) and its solution set $\mathbf{Z}$ is called the characteristic variety of (22). It is said [1] that (22) is absolutely nondegenerate if $\mathbf{Z} = \emptyset$, nondegenerate of type a if

$$a = \inf |\operatorname{Im} s| < \infty, \quad s \in \mathbf{Z} \neq \mathbf{C}^n,$$

and degenerate if $\mathbf{Z} = \mathbf{C}^n$. The case $\mathbf{Z} = \emptyset$ implies $\Delta(s) = \text{const.}$, since $\Delta(s)$ is meromorphic, and a meromorphic function that is not constant assumes every complex value with at most two exceptions. The equation $\Delta(s) = C$ can be written as

$$P(is) + \sum_{j=1}^q Q_j(is) - \sum_{j=1}^q Q_j(is) e^{P(is)t_j} = CP(is)$$

and is possible for $q > 1$ only if $P(s) = \text{const.}$, otherwise $\exp(P(is)t_j)$ would grow faster than any polynomial, which breaks the latter equality. For $q = 1$ we have

$$\Delta(s) = \frac{P(is) + Q_1(is) - Q_1(is) e^{P(is)t_1}}{P(is)},$$

and in this case $\mathbf{Z} = \emptyset$ is equivalent to $P(is) + Q_1(is) \equiv 0$. On the other hand, $\Delta(s) \equiv 0$ is equivalent to

$$P(is) + \sum_{j=1}^q Q_j(is) - \sum_{j=1}^q Q_j(is) e^{P(is)t_j} \equiv 0,$$

which implies $P(s) = \text{const.}$ This establishes the following proposition which was stated in [1] without proof, namely Eq. (22) is absolutely

nondegenerate if and only if either of the following conditions holds true:

$$(i) \quad P(s) \equiv C_1, \quad \sum_{j=1}^q Q_j(s)k(C_1, t_j) \equiv C_2 \neq 1;$$

or

$$(ii) \quad q = 1, \quad P(s) + Q_1(s) \equiv 0.$$

Eq. (22) is degenerate if and only if

$$P(s) \equiv C_1, \quad \sum_{j=1}^q Q_j(s)k(C_1, t_j) \equiv 1.$$

Substituting B from (25) in (23) leads to the proof that the uniqueness classes for the solution of the Cauchy problem for an absolutely nondegenerate equation (22) are the same as those for the equation (without "loads") $u_t(x, t) = Pu(x, t)$. The homogeneous degenerate IVP (22) ($u_0(x) = 0$) has nontrivial solutions, with compact support. Suppose that (22) is of finite type a ($0 < a < \infty$) and that $u(x, t)$ is a solution of (22) with $u_0(x) \equiv 0$. If

$$|u(x, t)| \leq Ce^{\alpha|x|}, \quad x \in \mathbb{R}^n, \quad t \in [0, T], \quad (27)$$

and $\alpha < a$, then $u(x, t) \equiv 0$. For any $\alpha > a$ there exists a solution $u(x, t) \not\equiv 0$ of (22) with $u_0(x) \equiv 0$ satisfying (27). Integral transformations have also been used in the study of EPCA.

Consider the nonlinear initial-value problem

$$\begin{aligned} \frac{\partial u}{\partial t} &= A(D)u(x, t) + f(t, u(x, [t])), \\ u(x, 0) &= u_0(x), \end{aligned} \quad (28)$$

where $u(x, t)$ and $u_0(x)$ are m -vectors, $x = (x_1, x_2, \dots, x_N) \in \mathbb{R}^N$,

$$A(D) = \sum_{|\alpha| \leq r} A_\alpha D^\alpha,$$

$$\begin{aligned} \alpha &= (\alpha_1, \alpha_2, \dots, \alpha_N), \quad |\alpha| = \alpha_1 + \alpha_2 + \dots + \alpha_N, \\ D^\alpha &= D_1^{\alpha_1} \dots D_N^{\alpha_N}, \quad D_k = i\partial/\partial x_k \quad (k = 1, 2, \dots, N), \end{aligned}$$

the coefficients A_α are given constant matrices of order $m \times m$, and the m -vector $f \in C^1([n, n+1) \times \mathfrak{L}^2(\mathbb{R}^N), \mathfrak{L}^2(\mathbb{R}^N))$, $n = 0, 1, 2, \dots$. The number r is called the order of the system. It is assumed that $u_0 \in \mathfrak{L}^2(\mathbb{R}^N)$, and the solutions sought are such that $u(x, t) \in \mathfrak{L}^2(\mathbb{R}^N)$, for every $t \geq 0$. Let $\mu_1(s), \mu_2(s), \dots, \mu_m(s)$ be the eigenvalues of the matrix $A(s)$. The system

$$\frac{\partial u}{\partial t} = A(D)u \quad (29)$$

is said to be parabolic by Shilov if

$$\operatorname{Re} \mu_j(s) \leq -c|s|^h + b, \quad j = 1, \dots, m$$

where $h > 0$, $c > 0$, and b are constants. For a fixed t we may consider the solution $u(x, t)$ as an element of $\mathfrak{L}^2(\mathbb{R}^N)$, and $f(t, u(x, [t]))$ may be treated as an abstract function $f(t, u([t]))$ with the values in $\mathfrak{L}^2$. Therefore, IVP (28) is reduced to the abstract Cauchy problem

$$\frac{du}{dt} = Au + f(t, u([t])), \quad u|_{t=0} = u_0 \in \mathfrak{L}^2. \quad (30)$$

Applying to (29), with the initial condition $u(x, 0) = u_0(x)$, the Fourier transformation $\mathcal{F}$ in x produces the system of ordinary differential equations

$$U_t(s, t) = A(s)U(s, t), \quad (31)$$

with the initial condition $U(s, 0) = U_0(s)$, where $U(s, t) = \mathcal{F}(u(x, t))$, $U_0(s) = \mathcal{F}(u_0(x))$, and $A(s)$ is a matrix with polynomial entries depending on $s = (s_1, s_2, \dots, s_N)$. The solution of (31) is given by the formula

$$U(s, t) = e^{tA(s)}U_0(s).$$

Parabolicity of (29) by Shilov implies that the semigroup $T(t)$ of operators of multiplication by $e^{tA(s)}$, for $t > 0$, is an infinitely smooth semigroup of operators bounded in $\mathfrak{L}^2(\mathbb{R}^N)$. Together with the requirement $h = r$, this ensures that the Cauchy problem for (29) is uniformly correct in $\mathfrak{L}^2(\mathbb{R}^N)$ and all its solutions are infinitely smooth functions

of t , for $t > 0$. Since f is continuously differentiable, problem (28) has on $[0, 1)$ a unique solution

$$u(t) = T(t)u_0 + \int_0^t T(t-s)f(s, u_0) ds.$$

Denoting $u_1 = u(1)$, we can find the solution

$$u(t) = T(t-1)u_1 + \int_1^t T(t-s)f(s, u_1) ds$$

of (28) on $[1, 2)$ and continue this procedure successively. If

$$f(t, u([t])) = Bu([t]),$$

where B is a constant matrix, the solution of (28) for $t \in [0, \infty)$ is given by

$$u(t) = \left(T(t-[t]) + \int_{[t]}^t T(t-s)B ds \right) \times \prod_{k=[t]}^1 \left(T(1) + \int_{k-1}^k T(k-s)B ds \right) u_0,$$

This proves that problem (28) has a unique solution on $\mathbb{R}^N \times [0, \infty)$ if system (29) is parabolic by Shilov, the index of parabolicity h coincides with its order r , and $f \in C^1([n, n+1]) \times \mathfrak{L}^2(\mathbb{R}^N), \mathfrak{L}^2(\mathbb{R}^N)$, $n = 0, 1, 2, \dots$

2.3. Wave Equations with Discontinuous Time Delay. The influence of terms with piecewise constant time on the behavior of the solutions, especially their oscillatory properties, of the wave equation was initiated in 1991 by Wiener and Debnath [11, 12].

First, we shall discuss separation of variables in systems of PDE. Consider the BVP consisting of the equation

$$U_t(x, t) = AU_{xx}(x, t) + BU_{xx}(x, [t]), \quad (32)$$

the boundary conditions

$$U(0, t) = U(1, t) = 0, \quad (33)$$

and the initial condition

$$U(x, 0) = U_0(x). \quad (34)$$

Here, $U(x, t)$ and $U_0(x)$ are real $m \times m$ matrices, A and B are real constant $m \times m$ matrices and $[\cdot]$ denotes the greatest-integer function. Looking for a solution in the form

$$U(x, t) = T(t)X(x) \quad (35)$$

gives

$$T'(t)X(x) = AT(t)X''(x) + BT([t])X''(x),$$

whence

$$(AT(t) + BT([t]))^{-1}T'(t) = X''(x)X^{-1}(x) = -P^2,$$

which generates the BVP

$$\begin{aligned} X''(x) + P^2X(x) &= 0, \\ X(0) &= X(1) = 0 \end{aligned} \quad (36)$$

and the equation with piecewise constant argument

$$T'(t) = -AT(t)P^2 - BT([t])P^2. \quad (37)$$

The general solution of Eq. (36) is

$$X(x) = \cos(xP)C_1 + \sin(xP)C_2,$$

where

$$\cos(xP) = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n} P^{2n}}{(2n)!}, \quad \sin(xP) = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+1} P^{2n+1}}{(2n+1)!}$$

and C_1, C_2 are arbitrary constant matrices. From $X(0) = 0$ we conclude that $C_1 = 0$, and the condition $X(1) = 0$ enables us to choose $\sin P = 0$ (although this is not the necessary consequence of the equation $(\sin P)C_2 = 0$). This can be written $e^{iP} - e^{-iP} = 0$, $e^{2iP} = I$. Assuming that all eigenvalues $p_1, p_2, \dots, p_m$ of P are distinct and $S^{-1}PS = \mathcal{D} = \text{diag}(p_1, p_2, \dots, p_m)$, we have $\exp(2iS\mathcal{D}S^{-1}) = I$, $Se^{2i\mathcal{D}}S^{-1} = I$, and $e^{2i\mathcal{D}} = I$. Therefore, $\mathcal{D} = \text{diag}(\pi j_1, \pi j_2, \dots, \pi j_m)$, where the j_k are integers, and $P = S\mathcal{D}S^{-1}$,

$$P^2 = S\mathcal{D}^2S^{-1} = S \text{diag}(\pi^2 j_1^2, \pi^2 j_2^2, \dots, \pi^2 j_m^2)S^{-1},$$

$\sin(xP) = S \sin(x\mathfrak{D})S^{-1} = S \operatorname{diag}(\sin \pi j_1 x, \dots, \sin \pi j_m x)S^{-1}$. Furthermore, we can put

$$P_j = \operatorname{diag}(\pi(m(j-1)+1), \dots, \pi mj), \quad (38)$$

$$(j = 1, 2, \dots)$$

in (36) and obtain the following result:

There exists an infinite sequence of matrix eigenfunctions for BVP (36),

$$X_j(x) = \sqrt{2} \operatorname{diag}(\sin \pi(m(j-1)+1)x, \dots, \sin \pi mjx), \quad (39)$$

$$(j = 1, 2, \dots)$$

which is complete and orthonormal in the space $\mathfrak{L}^2[0, 1]$ of $m \times m$ matrices, that is,

$$\int_0^1 X_j(x) X_k(x) ds = \begin{cases} 0, & j \neq k \\ I, & j = k \end{cases}$$

where I is the identity matrix.

Let $E(t)$ be the solution of the problem

$$T'(t) = -AT(t)P^2, \quad T(0) = I \quad (40)$$

and let

$$M(t) = E(t) + (E(t) - I)A^{-1}B. \quad (41)$$

If the matrix A is nonsingular, then Eq. (37) with the initial condition $T(0) = C_0$ has on $[0, \infty)$ a unique solution

$$T(t) = M(t - [t])M^{[t]}(1)C_0. \quad (42)$$

If $\|M(1)\| < 1$, then $\|T(t)\|$ exponentially tends to zero as $t \rightarrow +\infty$.

For the scalar parabolic equation

$$u_t(x, t) = a^2 u_{xx}(x, t) + bu_{xx}(x, [t])$$

we have $m = 1$ and $P_j = \pi j$, according to (38). For Eq. (40) with $A = a^2$ and $P = P_j$, we have $E_j(t) = e^{-a^2 \pi^2 j^2 t}$ and

$$M_j(t) = e^{-a^2 \pi^2 j^2 t} - \frac{b}{a^2}(1 - e^{-a^2 \pi^2 j^2 t}).$$

Hence, the inequality $|M_j(1)| < 1$ is equivalent to

$$-1 < e^{-a^2\pi^2j^2} - \frac{b}{a^2}(1 - e^{-a^2\pi^2j^2}) < 1,$$

whence

$$-a^2 < b < a^2 \frac{1 + e^{-a^2\pi^2j^2}}{1 - e^{-a^2\pi^2j^2}}.$$

Since the function $(1 + e^{-t})/(1 - e^{-t})$ is decreasing, all functions $T_j(t)$ exponentially tend to zero as $t \rightarrow \infty$ if and only if

$$-a^2 < b \leq a^2. \quad (43)$$

If $b < -a^2$, then all $T_j(t)$ monotonically tend to infinity as $t \rightarrow \infty$; and if

$$b > a^2 \frac{1 + e^{-a^2\pi^2}}{1 - e^{-a^2\pi^2}},$$

then all $T_j(t)$ are unbounded and oscillatory. For any $b > a^2$, there exists a positive integer j_0 such that the $T_j(t)$ are unbounded and oscillatory, for $j > j_0$. Indeed, letting $b = a^2 + \epsilon$ and solving the inequality

$$a^2 + \epsilon > a^2 \frac{1 + e^{-a^2\pi^2j^2}}{1 - e^{-a^2\pi^2j^2}}$$

gives

$$e^{-a^2\pi^2j^2} < \frac{\epsilon}{2a^2 + \epsilon},$$

which holds for any positive ϵ and sufficiently large j and implies $M_j(1) < -1$. If $b = -a^2$, then $M_j(t) = 1$, $T_j(t) = T_j(0)$, and $u(x, t) = u_0(x)$, for all t . Therefore, the condition $|b| \leq a^2$ is necessary and sufficient for the series

$$u(x, t) = \sum_{j=1}^{\infty} T_j(t) X_j(x) \quad (44)$$

to be a solution of the scalar BVP (32)–(34), with $A = a^2$ and $B = b$, if $u_0(x)$ is three times continuously differentiable. The coefficients $T_j(0)$

are given by

$$T_j(0) = \int_0^1 u_0(x) X_j(x) dx,$$

where $X_j(x) = \sqrt{2} \sin(\pi j x)$ and $u_0(x) \in C^3[0, 1]$ satisfies

$$u_0(0) = u_0(1) = 0.$$

The solution $T = 0$ of Eq. (37) is globally asymptotically stable as $t \rightarrow +\infty$ if and only if the eigenvalues λ_r of the matrix $M(1)$ satisfy the inequalities

$$|\lambda_r| < 1, \quad r = 1, \dots, m. \quad (45)$$

If all eigenvalues of A have positive real parts and $U_0(x) \in C^3[0, 1]$, $\|A^{-1}B\| < 1$, then BVP (32)–(34) has a solution (44). This series and all its term-by-term derivatives converge uniformly.

Separation of variables in the equation with constant coefficients

$$u_{tt}(x, t) = a^2 u_{xx}(x, t) - b u_{xx}(x, [t]) \quad (46)$$

and boundary conditions (33) yields $X_j(x) = \sqrt{2} \sin(\pi j x)$ and leads to the EPCA

$$T_j''(t) + a^2 \pi^2 j^2 T_j(t) = b \pi^2 j^2 T_j([t]). \quad (47)$$

For brevity, omit the subindex j and use the substitution $T'(t) = V(t)$, which changes (47) to a vector EPCA

$$w'(t) = Aw(t) + Bw([t]), \quad (48)$$

where $w = \text{col}(T, V)$ and

$$A = \begin{pmatrix} 0 & 1 \\ -a^2 \pi^2 j^2 & 0 \end{pmatrix}, \quad B = \begin{pmatrix} 0 & 0 \\ b \pi^2 j^2 & 0 \end{pmatrix}.$$

Eq. (48) on the interval $n \leq t < n+1$ becomes

$$w'(t) = Aw(t) + Bc_n, \quad c_n = w(n)$$

with the solution

$$w(t) = M(t - n)c_n,$$

where

$$M(t) = e^{At} + (e^{At} - I)A^{-1}B. \quad (49)$$

Therefore, Eq. (48) with the initial condition $w(0) = c_0$ has on $[0, \infty)$ a unique solution given by the right-hand side of (42), where $M(t)$ is defined in (49).

For $b < 0$, the solution $w = 0$ of Eq. (48) is unstable. Indeed, computations show that

$$e^{At} = \cos(\omega t)I + \omega^{-1} \sin(\omega t)A$$

and

$$e^{At} - I = \begin{pmatrix} \cos \omega t - 1 & \omega^{-1} \sin \omega t \\ -\omega \sin \omega t & \cos \omega t - 1 \end{pmatrix},$$

where $\omega = a\pi j$. Also

$$(e^{At} - I)A^{-1}B = \begin{pmatrix} b(1 - \cos \omega t)/a^2 & 0 \\ (b\omega \sin \omega t)/a^2 & 0 \end{pmatrix}$$

Hence,

$$M(t) = \begin{pmatrix} \cos \omega t + ba^{-2}(1 - \cos \omega t) & \omega^{-1} \sin \omega t \\ (ba^{-2} - 1)\omega \sin \omega t & \cos \omega t \end{pmatrix}$$

and

$$\det M(1) = 1 - \frac{b}{a^2} + \frac{b}{a^2} \cos \omega.$$

The condition $b < 0$ implies $\det M(1) > 1$ and shows that at least one of the eigenvalues λ of $M(1)$ satisfies $|\lambda| > 1$. Therefore, $\|w(t)\| \rightarrow \infty$ as $t \rightarrow +\infty$, for some initial vector $c_0 \neq 0$.

For $b > a^2$, the solution $w = 0$ of Eq. (48) is unstable. Calculations yield

$$\det(M(1) - \lambda I) = \lambda^2 - 2 \left(\cos \omega + \frac{b}{a^2} \sin^2 \frac{\omega}{2} \right) \lambda + 1 - \frac{b}{a^2} + \frac{b}{a^2} \cos \omega$$

and the expressions $\lambda_1 = s + d$, $\lambda_2 = s - d$ for the eigenvalues λ_1 , λ_2 of $M(1)$, where

$$s = \cos \omega + \frac{b}{a^2} \sin^2 \frac{\omega}{2}, \quad d^2 = \left(\frac{b}{a^2} - 1 \right) \sin^2 \omega + \frac{b^2}{a^4} \sin^4 \frac{\omega}{2}.$$

The condition $b > a^2$ shows that $d^2 > 0$ and $\lambda_1 > 1$. The latter inequality implies $\|w(t)\| \rightarrow \infty$ as $t \rightarrow +\infty$, for some initial vector $c_0 \neq 0$.

The solution $w = 0$ of Eq. (48) is asymptotically stable as $t \rightarrow +\infty$ if and only if

$$0 < b < a^2, \quad (50)$$

and $\omega \neq 2\pi n$, $n = 0, 1, 2, \dots$. The condition $d^2 < 0$, which means that the eigenvalues of $M(1)$ are complex, leads to

$$\cos^2 \frac{\omega}{2} > \frac{b^2}{(2a^2 - b)^2},$$

whence

$$b < a^2 \left(1 - \tan^2 \frac{\omega}{4} \right) \quad \text{or} \quad b < a^2 \left(1 - \cot^2 \frac{\omega}{4} \right).$$

Since $|\lambda_1| = |\lambda_2|$ and $\det M(1) = \lambda_1 \lambda_2$, the inequality $|\lambda_1| < 1$ is equivalent to $\det M(1) < 1$, that is, to $b > 0$. Therefore, in the case of complex eigenvalues, a criterion for asymptotic stability is

$$0 < b < \max \left(a^2 \left(1 - \tan^2 \frac{\omega}{4} \right), a^2 \left(1 - \cot^2 \frac{\omega}{4} \right) \right).$$

The inequality $d^2 > 0$ in the case of distinct real eigenvalues leads to

$$b > \max \left(a^2 \left(1 - \tan^2 \frac{\omega}{4} \right), a^2 \left(1 - \cot^2 \frac{\omega}{4} \right) \right),$$

and the inequalities $\lambda_1 < 1$, $\lambda_2 > -1$ yield $b < a^2$. Hence, in this case a criterion of asymptotic stability is

$$\max \left(a^2 \left(1 - \tan^2 \frac{\omega}{4} \right), a^2 \left(1 - \cot^2 \frac{\omega}{4} \right) \right) < b < a^2.$$

Finally, if

$$b = \max \left(a^2 \left(1 - \tan^2 \frac{\omega}{4} \right), a^2 \left(1 - \cot^2 \frac{\omega}{4} \right) \right),$$

then $d = 0$ and $\lambda_1 = \lambda_2 = \cos \omega + ba^{-2} \sin^2 \omega/2$, whence

$$\cos \omega < \lambda_1 < \cos^2 \omega/2$$

and $|\lambda_1| < 1$. According to (45), this implies asymptotic stability and completes the proof of criterion (50).

If $b = a^2$, then $\lambda_1 = 1$, $\lambda_2 = \cos \omega$, and the solutions of (48) are bounded but not asymptotically stable. If $\omega = 2\pi n$, then $\lambda_1 = \lambda_2 = 1$, which leads to the existence of unbounded solutions for (48). If the coefficient a is irrational, then (50) is a criterion of asymptotic stability of the solutions to (47) for all j , since recalling that $\omega = \omega_j = a\pi j$, we note that the equality $a\pi j = 2\pi n$ is impossible for any irrational a . For any rational a , there exist infinitely many integers j such that the corresponding solutions $w_j(t)$ of (48) are unbounded. Furthermore, each component of every solution of Eq. (48) oscillates if and only if either of the following conditions holds true:

$$(i) \quad b < \max \left(a^2 \left(1 - \tan^2 \frac{\omega}{4} \right), a^2 \left(1 - \cot^2 \frac{\omega}{4} \right) \right),$$

$$(ii) \quad \max \left(a^2 \left(1 - \tan^2 \frac{\omega}{4} \right), a^2 \left(1 - \cot^2 \frac{\omega}{4} \right) \right) < b < \frac{a^2}{2 \sin^2 \frac{\omega}{2}}$$

$$\text{and } \cos \omega < -\frac{1}{2}.$$

In conclusion, it is worth noting that the asymptotic properties of Eq. (47) depend on the algebraic nature of the coefficient a . For $b < 0$, all solutions of Eq. (47) are unstable and oscillatory; for $b > a^2$ all solutions of Eq. (47) are unstable and nonoscillatory. These two cases hold true for both rational and irrational values of a . For

$$0 < b < \max \left(a^2 \left(1 - \tan^2 \frac{\omega}{4} \right), a^2 \left(1 - \cot^2 \frac{\omega}{4} \right) \right),$$

all solutions of (47) are asymptotically stable and oscillatory, provided that $\omega \neq 2\pi n$. However, for any rational a , there exist infinitely many integers j such that $\omega_j = 2\pi n$, which leads to the existence of unbounded solutions for (47). Furthermore, since $\omega = \omega_j = a\pi j$ the inequality $\cos \omega < -1/2$ breaks down for infinitely many integers j . Therefore, under the above hypothesis (ii), there are infinitely many solutions of Eq. (47) which are asymptotically stable and oscillatory, as well as infinitely many solutions which are asymptotically stable and nonoscillatory ($\omega \neq 2\pi n$). Also, for $\omega \neq 2\pi n$ and $a^2/2 \sin^2(\omega/2) < b < a^2$, the solutions of (47) are asymptotically stable and nonoscillatory. Problems of this nature deserve further investigation.

3. LIST OF PUBLICATIONS AND TECHNICAL REPORTS

Proposal Number 26739-MA Funding Document DAAL03-89-G-0107

Boundary-Value Problems for Partial Differential Equations with Piecewise Constant Delay, by Joseph Wiener

A Survey of Differential Equations with Piecewise Continuous Arguments, by Joseph Wiener and Kenneth L. Cooke

Resonance in Linear Differential Equations and L'Hospital's Rule, by Joseph Wiener

Averaging Principle and Systems of Singularly Perturbed Stochastic Differential Equations, by J. Golec and G. Ladde

Partial Differential Equations with Piecewise Constant Delay, by Joseph Wiener and Lokenath Debnath

A Calculus Exercise for the Sums of Integer Powers, by Joseph Wiener

Modeling of Dynamic Systems by Ito-Type Systems of Stochastic Differential Equations, by G. S. Ladde

Stochastic Delay Differential Systems, by G. S. Ladde

A Parabolic Differential Equation with Unbounded Piecewise Constant Delay, by Joseph Wiener and Lokenath Debnath

Singularly Perturbed Linear Boundary-Value Problems, by M. Kathirkamanayagan and G. S. Ladde

Coexistence of Analytic and Distributional Solutions for Linear Differential Equations, by J. Wiener et al.

The Fourier Method for Partial Differential Equations with Piecewise Continuous Delay, by J. Wiener and L. Debnath

A Wave Equation with Discontinuous Time Delay, by Joseph Wiener and Lokenath Debnath

Ordinary and Delay Differential Equations, by Joseph Wiener and Jack K. Hale (editors)

Partial Differential Equations, by Joseph Wiener and Jack K. Hale (editors)

REFERENCES

1. V. M. Borok and Ya. I. Zhitomirskii, *The Cauchy problem for a certain class of loaded equations*, Uspekhi Mat. Nauk **34** (1979), 221-222.
2. L. A. V. Carvalho and K. L. Cooke, *A nonlinear equation with piecewise continuous argument*, Differential and Integral Equations **1** (1988), 359-367.
3. K. L. Cooke, J. Turi, and G. Turner, *Stabilization of hybrid systems in the presence of feedback delays*, Preprint Series 906, Inst. Math. and Its Appls., University of Minnesota, 1991.
4. K. L. Cooke and J. Wiener, *Retarded differential equations with piecewise constant delays*, J. Math. Anal. Appl. **99** (1984), 26-39.
5. ———, *A survey of differential equations with piecewise continuous arguments*, in Delay Differential Equations and Dynamical Systems, eds. S. Busenberg and M. Martelli (Lecture Notes in Math. **1475** Springer-Verlag, Berlin-Heidelberg-New York, 1991) pp. 1-15.
6. J. K. Hale, *Theory of Functional Differential Equations* (Springer-Verlag, New York-Heidelberg-Berlin, 1977).
7. G. Ladas, *Oscillations of equations with piecewise constant mixed arguments*, in Differential Equations and Applications II, ed. A. R. Aftabizadeh (Ohio University Press, Athens, 1988) pp. 64-69.
8. T.-Y. Li and J. A. Yorke, *Period three implies chaos*, Amer. Math. Monthly **82** (1975), 985-992.
9. J. Wiener, *Boundary-value problems for partial differential equations with piecewise constant delay*, Internat. J. Math. Math. Sci. **14** (1991), 301-321.
10. J. Wiener and L. Debnath, *Partial differential equations with piecewise constant delay*, Internat. J. Math. Math. Sci. **14** (1991), 485-496.
11. ———, *The Fourier method for partial differential equations with piecewise continuous delay*, in Contemporary Mathematics **129**, eds. J. R. Graef and J. K. Hale (Amer. Math. Soc., Providence, 1992) pp. 241-263.
12. ———, *A wave equation with discontinuous time delay*, Internat. J. Math. Math. Sci. **15** (1992), 781-792.