

AD-A268 178



Naval Research Laboratory

Stennis Space Center, MS 39529-5004



NRL/MR/7323--93-7023

Report Detailing Modifications to the 1/8° Global Bathymetry

WESLEY J. YOUTSEY

Sverdrup Technology, Inc.
Stennis Space Center, MS 39529

May 1993

DTIC
ELECTE
AUG 19 1993
S E D

93-8-18-025


93-19205



135AY

Approved for public release; distribution is unlimited.

DISCLAIMER NOTICE



THIS DOCUMENT IS BEST QUALITY AVAILABLE. THE COPY FURNISHED TO DTIC CONTAINED A SIGNIFICANT NUMBER OF COLOR PAGES WHICH DO NOT REPRODUCE LEGIBLY ON BLACK AND WHITE MICROFICHE.

REPORT DOCUMENTATION PAGE*Form Approved*
OBM No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. Agency Use Only (Leave blank).		2. Report Date. May 1993	3. Report Type and Dates Covered. Final
4. Title and Subtitle. Report Detailing Modifications to the 1/8° Global Bathymetry			5. Funding Numbers. <i>Program Element No.</i> 0603207N <i>Project No.</i> X0513 <i>Task No.</i> <i>Accession No.</i> DN259002 <i>Work Unit No.</i> 73509403
6. Author(s). Wesley J. Youtsey*			8. Performing Organization Report Number. NRL/MR/7323--93-7023
7. Performing Organization Name(s) and Address(es). Naval Research Laboratory Detachment Ocean Dynamics and Prediction Branch Stennis Space Center, MS 39529-5004			10. Sponsoring/Monitoring Agency Report Number. NRL/MR/7323--93-7023
9. Sponsoring/Monitoring Agency Name(s) and Address(es). Space and Naval Warfare Systems Command PMW 165 Washington, DC 20363-5200			11. Supplementary Notes. *Sverdrup Technology Stennis Space Center, MS 39529
12a. Distribution/Availability Statement. Approved for public release; distribution is unlimited.			12b. Distribution Code.
13. Abstract (Maximum 200 words). This report describes the modifications made to the 1/8° global topography used in the Navy Layered Ocean Model. The regions, which were modified, are listed and figures are provided to show the Earth Topography 5-minute (ETOPO5) global topography, the 1/8° global topography prior to modification, and both the smoothed and unsmoothed versions of the 1/8° global topography after modification.			
14. Subject Terms. Ocean Models, Military Oceanography, Data Assimilation			15. Number of Pages. 129
			16. Price Code.
17. Security Classification Unclassified	18. Security Classification of Report. Unclassified	19. Security Classification of This Page. Unclassified	20. Limitation of Abstract of Abstract. SAR

Report Detailing Modifications to the 1/8° Global Bathymetry

Introduction

This report details the changes made to the 1/8° global topography used in the Navy Layered Ocean Model using GEBCO (General Bathymetric Chart of the Oceans) as a reference. Changes were made to the following regions:

PACIFIC OCEAN

- Aleutian Islands
- Hawaiian Islands (including the shoals and islands to the west)
- Channel Islands west of Los Angeles
- Galapagos Islands
- Tahiti and the Marquesas
- Line Islands
- Samoa Islands
- New Hebride Islands
- Fiji Islands (including Northern New Caledonia)
- TOGA/COARE region (especially the Gilbert Islands)
- Indonesian throughflow region
- Philippine Islands
- South China Sea
- East China Sea
- Izu Ridge (including the Mariana Islands)
- Tsushima Strait
- Sea of Japan
- Soya Strait

Accession For	
NTIS CRA&I	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
A-1	

DTIC QUALITY INSPECTED 1

- Kuril Islands

ATLANTIC OCEAN

- GIN Sea region
- Azores
- Canary Islands
- Cape Verde Islands
- Barbados
- Lesser Antilles
- West Indies
- Bahama Islands
- Bermuda

INDIAN OCEAN

- Maldiv Islands
- Andaman Islands
- Selat Mentawai
- Indonesian throughflow region
- Chagos Laccadive Ridge and Diego Garcia Island
- Mascarene Plateau
- Shoals and Islands to the north and west of Madagascar Island
- Seychelle Islands

MEDITERRANEAN SEA

- Balearic Islands
- Sicilian Channel

The new global topography has been named top117a. This topography incorporates the 1/8° N Atlantic (na632k) and 1/8° N Pacific (p133d) topographies. The 1/8° global topography, which preceded top117a, was named top08_200g_global. The modifications made to generate the new topography include the addition of missing islands and shoals, the removal of nonexistent islands, the shifting of islands and shoals to undersea ridges, the conversion between model land/plot sea points and land points, and the modification of straits.

The Pacific Ocean (Figure 1) required the most changes. Some of the regions, such as the Aleutian Islands, required little change. Most, however, required major revision particularly the Hawaiian Islands, the TOGA/COARE region, the Kuril Islands and the Indonesian throughflow region.

Soya and Tsushima Straits were modified to provide a more realistic geometry that more closely resembled the actual straits. This required setting depths that were less than 200 m in the ETOPO5 topography to 201 m in the p133d topography. A separate file was then generated for each strait that contains the depth for that gridpoint from the ETOPO5 bathymetry. These two files can then be used as inputs to the model to generate friction values to dampen the flow through these straits.

Plots are included that show the N Pacific (Figure 2), N Atlantic (Figure 3), ETOPO5 bathymetry, the unmodified global bathymetry, the modified global bathymetry, and the twice-smoothed modified global bathymetry (Figures 1-100). Also included are the sill depths (Tables 1-4) for the ETOPO5, modified global, and twice-smoothed modified global bathymetries for selected straits of interest. Plots are also included that show the locations of the sill depths (Figures 101-115).

The changes to the Atlantic Ocean (Figure 4) were concentrated in the West Indies and the Lesser Antilles. Most of the changes in the West Indies were in the small islands and shoals to the north of Cuba and Hispanola. Extensive changes were made to the Lesser Antilles along with sill depths being modified to be in closer agreement with the ETOPO5 depths. Also, Bermuda and Barbados Islands were added to the topography.

The Azores, Canary, and Cape Verde Islands of the eastern Atlantic Ocean required the addition and shifting of many islands in the groups.

Changes in the Indian Ocean were made mostly to the Maldive Islands, the shoals to the south of the equator in the central and western Indian Ocean, and the islands and shoals in the eastern Indian Ocean along the Andaman Sea and the Selat Mentawai region.

Minor changes were made to other regions of the world. These were concentrated in the Drake Passage/Falkland Islands region, New Zealand, the Loyalty Islands and the Mediterranean Sea. Some smoothing and removal of small bays in Antarctica and peninsulas was also performed.

Acknowledgments

This work has been performed as part of the Naval Research Laboratory (NRL) 6.2/6.3 Global Ocean Prediction System projects. These are components of the Navy Ocean Modeling Program managed by Robert Peloquin under program elements 62435N and 63207N.

Table 1: Comparison of sill depths for the p133d, p133d smoothed and ETOPO5 topographies for the N Pacific Ocean.

	p133d	p133d smoothed	ETOPO5
Strait	Depth (m)	Depth (m)	Depth (m)
Aleutian Islands 1	2056	2260	2072
Aleutian Islands 2	1413	1512	NA
Aleutian Islands 3	959	996	797
Aleutian Islands 4	747	812	NA
Aleutian Islands 5	752	892	921
Aleutian Islands 6	1029	1201	1018
Aleutian Islands 7	398	461	414
Vitiaz Strait	925	710	741
SW Pacific Ocean 1	276	288	286
SW Pacific Ocean 2	254	304	272
SW Pacific Ocean 3	397	373	444
SW Pacific Ocean 4	481	500	490
SW Pacific Ocean 5	1806	1816	1768
SW Pacific Ocean 6	534	631	524
East China Sea 1	787	838	785
East China Sea 2	414	486	NA
East China Sea 3	243	277	245
East China Sea 4	201	268	242
East China Sea 5	393	390	344
East China Sea 6	1043	986	1040
East China Sea 7	366	652	806
East China Sea 8	684	649	NA
East China Sea 9	314	419	265
East China Sea 10	551	548	513
Izu Ridge 1	412	441	401
Izu Ridge 2	212	274	NA
Izu Ridge 3	253	269	NA
Izu Ridge 4	891	865	NA
Izu Ridge 5	477	502	NA
Kuril Islands 1	201	200	NA
Kuril Islands 2	469	517	478
Kuril Islands 3	2066	2359	2000
Kuril Islands 4	400	601	540
Kuril Islands 5	810	1068	NA
Kuril Islands 6	887	1195	NA
Kuril Islands 7	1035	1241	NA
Kuril Islands 8	220	253	266

Table 2: Comparison of sill depths for the na632k, na632k smoothed and ETOPO5 topographies for the N Atlantic Ocean.

	na632k	na632k smoothed	ETOPO5
Strait	Depth (m)	Depth (m)	Depth (m)
Anegada Passage	1480	1360	1923
St. Christopher Passage	390	424	409
Guadeloupe Passage	550	625	495
Dominica Channel	552	573	464
Martinique Channel	807	880	900
St. Lucia Channel	924	936	855
St. Vincent Channel	642	739	541
Grenada Passage	468	516	511
Mona Passage	507	542	545
Windward Passage	2134	2142	2134
Old Bahama Passage	273	279	293
Florida Strait	566	510	578
Northwest Providence Channel	530	488	546
Crooked Island Passage	1817	1636	1780
Mayaguana Passage	1558	1344	NA
Caicos Passage	1712	2201	2554
Turks Islands Passage	524	537	NA
Mouchoir Passage	579	770	NA
Silver Bank Passage	2719	2256	2737
GIN Sea 1	547	402	594
GIN Sea 2	690	837	936
GIN Sea 3	300	296	289

Table 3: Comparison of sill depths for the $1/8^\circ$ world ocean model topography (wom08), wom08 smoothed and ETOPO5 topographies for the Mediterranean Sea, the Falkland Islands and New Zealand.

	wom08	wom08 smoothed	ETOPO5
Strait	Depth (m)	Depth (m)	Depth (m)
Strait of Gibraltar	201	201	NA
Mediterranean Sea 1	680	627	784
Mediterranean Sea 2	322	355	388
Mediterranean Sea 3	332	349	351
Mediterranean Sea 4	615	734	992
Mediterranean Sea 5	682	640	738
Falkland Island	393	398	399
New Zealand	508	521	496

Table 4: Comparison of sill depths for the $1/8^\circ$ world ocean model topography (w117a), w117a smoothed topography, ETOPO5 topography, $1/8^\circ$ world ocean model topography (w116e), and w116e smoothed topography for the Indian Ocean.

	w117a	w117a	ETOPO5	w116e	w116e
		smoothed			smoothed
Strait	Depth (m)	Depth (m)	Depth (m)	Depth (m)	Depth (m)
Socotra Island	685	680	718	598	843
Maldiv Islands 1	600	623	1923	NA	NA
Maldiv Islands 2	510	426	409	NA	NA
Maldiv Islands 3	230	232	495	NA	NA
Maldiv Islands 4	353	555	464	225	668
Maldiv Islands 5	1414	1403	900	NA	NA
Maldiv Islands 6	1997	1857	1896	1990	1972
Andaman Sea 1	377	484	399	635	759
Andaman Sea 2	1110	1271	NA	1262	1191
Andaman Sea 3	971	970	827	943	1099
Andaman Sea 4	482	760	228	NA	NA
Andaman Sea 5	236	521	207	NA	NA
Andaman Sea 6	1737	1598	1408	1709	1610
Andaman Sea 7	327	405	512	325	730
Indo. throughflow 1	249	514	230	201	465
Indo. throughflow 2	588	801	578	767	721
Indo. throughflow 3	1063	1065	1045	1063	1236
Indo. throughflow 4	1007	949	1020	1007	1099
Indo. throughflow 5	1452	1291	1530	1458	1212
SW Indian Ocean 1	480	619	NA	NA	NA
SW Indian Ocean 2	483	846	309	576	890
SW Indian Ocean 3	904	1264	903	1227	1191
SW Indian Ocean 4	395	411	NA	450	699
SW Indian Ocean 5	624	827	785	624	848
SW Indian Ocean 6	640	807	1454	840	964

Figure Captions

Figure 1: Plot of the Pacific Ocean from the new $1/8^\circ$ global topography which has been smoothed by two passes of a 9-point smoother.

Figure 2: Plot of the North Pacific Ocean from the new $1/8^\circ$ global topography which has been smoothed by two passes of a 9-point smoother.

Figure 3: Plot of the North Atlantic Ocean from the new $1/8^\circ$ global topography which has been smoothed by two passes of a 9-point smoother.

Figure 4: Plot of the Atlantic Ocean from the new $1/8^\circ$ global topography which has been smoothed by two passes of a 9-point smoother.

Figure 5: Plot of the Aleutian Islands from the unmodified $1/8^\circ$ Pacific topography.

Figure 6: Plot of the Aleutian Islands from the $1/12^\circ$ ETOPO5 topography.

Figure 7: Plot of the Aleutian Islands from the new $1/8^\circ$ Pacific topography.

Figure 8: Plot of the Aleutian Islands from the new $1/8^\circ$ Pacific topography which has been smoothed by two passes of a 9-point smoother.

Figure 9: Plot of the Hawaiian Islands and the shoals to the northwest from the unmodified $1/8^\circ$ Pacific topography.

Figure 10: Plot of the Hawaiian Islands and the shoals to the northwest from the $1/12^\circ$ ETOPO5 topography.

Figure 11: Plot of the Hawaiian Islands and the shoals to the northwest from the new $1/8^\circ$ Pacific topography.

Figure 12: Plot of the Hawaiian Islands and the shoals to the northwest from the new $1/8^\circ$ Pacific topography which has been smoothed by two passes of a 9-point smoother.

Figure 13: Plot of the Channel Islands west of Los Angeles, California from the unmodified $1/8^\circ$ Pacific topography.

Figure 14: Plot of the Channel Islands west of Los Angeles, California from the $1/12^\circ$ ETOPO5 topography.

Figure 15: Plot of the Channel Islands west of Los Angeles, California from the new $1/8^\circ$ Pacific topography.

Figure 16: Plot of the Channel Islands west of Los Angeles, California from the new $1/8^\circ$ Pacific topography which has been smoothed by two passes of a 9-point smoother.

Figure 17: Plot of the Galapagos Islands from the unmodified $1/8^\circ$ Pacific topography.

Figure 18: Plot of the Galapagos Islands from the $1/12^\circ$ ETOPO5 topography.

Figure 19: Plot of the Galapagos Islands from the new $1/8^\circ$ Pacific topography.

Figure 20: Plot of the Galapagos Islands from the new $1/8^\circ$ Pacific topography which has been smoothed by two passes of a 9-point smoother.

Figure 21: Plot of the Line Islands from the unmodified $1/8^\circ$ Pacific topography.

Figure 22: Plot of the Line Islands from the $1/12^\circ$ ETOPO5 topography.

Figure 23: Plot of the Line Islands from the new $1/8^\circ$ Pacific topography.

Figure 24: Plot of the Line Islands from the new $1/8^\circ$ Pacific topography which has been smoothed by two passes of a 9-point smoother.

Figure 25: Plot of the Samoan Islands from the unmodified $1/8^\circ$ Pacific topography.

Figure 26: Plot of the Samoan Islands from the $1/12^\circ$ ETOPO5 topography.

Figure 27: Plot of the Samoan Islands from the new $1/8^\circ$ Pacific topography.

Figure 28: Plot of the Samoan Islands from the new $1/8^\circ$ Pacific topography which has been smoothed by two passes of a 9-point smoother.

Figure 29: Plot of the New Caledonia, New Hebrides, and the Fiji Islands from the unmodified $1/8^\circ$ Pacific topography.

Figure 30: Plot of the New Caledonia, New Hebrides, and the Fiji Islands from the $1/12^\circ$ ETOPO5 topography.

Figure 31: Plot of the New Caledonia, New Hebrides, and the Fiji Islands from the new $1/8^\circ$ Pacific topography.

Figure 32: Plot of the New Caledonia, New Hebrides, and the Fiji Islands from the new $1/8^\circ$ Pacific topography which has been smoothed by two passes of a 9-point smoother.

Figure 33: Plot of the TOGA/COARE region from the unmodified $1/8^\circ$ Pacific topography.

Figure 34: Plot of the TOGA/COARE region from the 1/12° ETOPO5 topography.

Figure 35: Plot of the TOGA/COARE region from the new 1/8° Pacific topography.

Figure 36: Plot of the TOGA/COARE region from the new 1/8° Pacific topography which has been smoothed by two passes of a 9-point smoother.

Figure 37: Plot of the Indonesian Throughflow region from the unmodified 1/8° Pacific topography.

Figure 38: Plot of the Indonesian Throughflow region from the 1/12° ETOPO5 topography.

Figure 39: Plot of the Indonesian Throughflow region from the new 1/8° Pacific topography.

Figure 40: Plot of the Indonesian Throughflow region from the new 1/8° Pacific topography which has been smoothed by two passes of a 9-point smoother.

Figure 41: Plot of the Philippine Islands and the S China Sea from the unmodified 1/8° Pacific topography.

Figure 42: Plot of the Philippine Islands and the S China Sea from the 1/12° ETOPO5 topography.

Figure 43: Plot of the Philippine Islands and the S China Sea from the new 1/8° Pacific topography.

Figure 44: Plot of the Philippine Islands and the S China Sea from the new 1/8° Pacific topography which has been smoothed by two passes of a 9-point smoother.

Figure 45: Plot of the E China Sea from the unmodified 1/8° Pacific topography.

Figure 46: Plot of the E China Sea from the 1/12° ETOPO5 topography.

Figure 47: Plot of the E China Sea from the new 1/8° Pacific topography.

Figure 48: Plot of the E China Sea from the new 1/8° Pacific topography which has been smoothed by two passes of a 9-point smoother.

Figure 49: Plot of the Mariana Islands from the unmodified 1/8° Pacific topography.

Figure 50: Plot of the Mariana Islands from the 1/12° ETOPO5 topography.

Figure 51: Plot of the Mariana Islands from the new 1/8° Pacific topography.

Figure 52: Plot of the Mariana Islands from the new 1/8° Pacific topography which has been smoothed by two passes of a 9-point smoother.

Figure 53: Plot of the Izu Ridge from the unmodified $1/8^\circ$ Pacific topography.

Figure 54: Plot of the Izu Ridge from the $1/12^\circ$ ETOPO5 topography.

Figure 55: Plot of the Izu Ridge from the new $1/8^\circ$ Pacific topography.

Figure 56: Plot of the Izu Ridge from the new $1/8^\circ$ Pacific topography which has been smoothed by two passes of a 9-point smoother.

Figure 57: Plot of the Sea of Japan from the unmodified $1/8^\circ$ Pacific topography.

Figure 58: Plot of the Sea of Japan from the $1/12^\circ$ ETOPO5 topography.

Figure 59: Plot of the Sea of Japan from the new $1/8^\circ$ Pacific topography.

Figure 60: Plot of the Sea of Japan from the new $1/8^\circ$ Pacific topography which has been smoothed by two passes of a 9-point smoother.

Figure 61: Plot of the Sea of Okhotsk from the unmodified $1/8^\circ$ Pacific topography.

Figure 62: Plot of the Sea of Okhotsk from the $1/12^\circ$ ETOPO5 topography.

Figure 63: Plot of the Sea of Okhotsk from the new $1/8^\circ$ Pacific topography.

Figure 64: Plot of the Sea of Okhotsk from the new $1/8^\circ$ Pacific topography which has been smoothed by two passes of a 9-point smoother.

Figure 65: Plot of the GIN Sea from the unmodified $1/8^\circ$ global topography.

Figure 66: Plot of the GIN Sea from the $1/12^\circ$ ETOPO5 topography.

Figure 67: Plot of the GIN Sea from the new $1/8^\circ$ Atlantic topography.

Figure 68: Plot of the GIN Sea from the new $1/8^\circ$ Atlantic topography which has been smoothed by two passes of a 9-point smoother.

Figure 69: Plot of the Canary and Azore Islands from the unmodified $1/8^\circ$ global topography.

Figure 70: Plot of the Canary and Azore Islands from the $1/12^\circ$ ETOPO5 topography.

Figure 71: Plot of the Canary and Azore Islands from the new $1/8^\circ$ Atlantic topography.

Figure 72: Plot of the Canary and Azore Islands from the new $1/8^\circ$ Atlantic topography which has been smoothed by two passes of a 9-point smoother.

Figure 73: Plot of the Caribbean Sea and Gulf of Mexico from the unmodified $1/8^\circ$ global topography.

Figure 74: Plot of the Caribbean Sea and Gulf of Mexico from the $1/12^\circ$ ETOPO5 topography.

Figure 75: Plot of the Caribbean Sea and Gulf of Mexico from the new $1/8^\circ$ Atlantic topography.

Figure 76: Plot of the Caribbean Sea and Gulf of Mexico from the new $1/8^\circ$ Atlantic topography which has been smoothed by two passes of a 9-point smoother.

Figure 77: Plot of the Mediterranean Sea from the unmodified $1/8^\circ$ global topography.

Figure 78: Plot of the Mediterranean Sea from the $1/12^\circ$ ETOPO5 topography.

Figure 79: Plot of the Mediterranean Sea from the new $1/8^\circ$ Atlantic topography.

Figure 80: Plot of the Mediterranean Sea from the new $1/8^\circ$ Atlantic topography which has been smoothed by two passes of a 9-point smoother.

Figure 81: Plot of the Drake Passage and the Falkland Islands from the unmodified $1/8^\circ$ global topography.

Figure 82: Plot of the Drake Passage and the Falkland Islands from the $1/12^\circ$ ETOPO5 topography.

Figure 83: Plot of the Drake Passage and the Falkland Islands from the new $1/8^\circ$ global topography.

Figure 84: Plot of the Drake Passage and the Falkland Islands from the new $1/8^\circ$ global topography which has been smoothed by two passes of a 9-point smoother.

Figure 85: Plot of the Arabian Sea from the unmodified $1/8^\circ$ global topography.

Figure 86: Plot of the Arabian Sea from the $1/12^\circ$ ETOPO5 topography.

Figure 87: Plot of the Arabian Sea from the new $1/8^\circ$ global topography.

Figure 88: Plot of the Arabian Sea from the new $1/8^\circ$ global topography which has been smoothed by two passes of a 9-point smoother.

Figure 89: Plot of the Southwestern Indian Ocean from the unmodified $1/8^\circ$ global topography.

Figure 90: Plot of the Southwestern Indian Ocean from the $1/12^\circ$ ETOPO5 topography.

Figure 91: Plot of the Southwestern Indian Ocean from the new $1/8^\circ$ global topography.

Figure 92: Plot of the Southwestern Indian Ocean from the new $1/8^\circ$ global topography which has been smoothed by two passes of a 9-point smoother.

Figure 93: Plot of the Andaman Sea from the unmodified $1/8^\circ$ global topography.

Figure 94: Plot of the Andaman Sea from the $1/12^\circ$ ETOPO5 topography.

Figure 95: Plot of the Andaman Sea from the new $1/8^\circ$ global topography.

Figure 96: Plot of the Andaman Sea from the new $1/8^\circ$ global topography which has been smoothed by two passes of a 9-point smoother.

Figure 97: Plot of the Indonesian Throughflow region from the unmodified $1/8^\circ$ global topography.

Figure 98: Plot of the Indonesian Throughflow region from the $1/12^\circ$ ETOPO5 topography.

Figure 99: Plot of the Indonesian Throughflow region from the new $1/8^\circ$ global topography.

Figure 100: Plot of the Indonesian Throughflow region from the new $1/8^\circ$ global topography which has been smoothed by two passes of a 9-point smoother.

Figure 101: Plot of New Zealand from the unmodified $1/8^\circ$ global topography.

Figure 102: Plot of New Zealand from the $1/12^\circ$ ETOPO5 topography.

Figure 103: Plot of New Zealand from the new $1/8^\circ$ global topography.

Figure 104: Plot of New Zealand from the new $1/8^\circ$ global topography which has been smoothed by two passes of a 9-point smoother.

Figure 105: Plot of Aleutian Island sill depth locations listed in table 1.

Figure 106: Plot of Southwest Pacific Ocean sill depth locations listed in table 1.

Figure 107: Plot of E China Sea sill depth locations listed in table 1.

Figure 108: Plot of Izu Ridge sill depth locations listed in table 1.

Figure 109: Plot of Kuril Island sill depth locations listed in table 1.

Figure 110: Plot of Caribbean Sea and Leeward Island sill depth location listed in table 2.

Figure 111: Plot of GIN Sea sill depth location listed in table 2.

Figure 112: Plot of Mediterranean Sea sill depth locations listed in table 3.

Figure 113: Plot of New Zealand sill depth location listed in table 3.

Figure 114: Plot of Falkland Island sill depth location listed in table 3.

Figure 115: Plot of Arabian Sea sill depth location listed in table 4.

Figure 116: Plot of Andaman Sea sill depth location listed in table 4.

Figure 117: Plot of Indonesian Throughflow sill depth location listed in table 4.

Figure 118: Plot of Southwest Indian Ocean sill depth location listed in table 4.

w117a2 topography 00117****.4

DX,DY = 0.176,0.125 DEG DB = 500.0 M



NRL 323 19-Oct-92

Figure 1: Plot of the Pacific Ocean from the new $1/8^\circ$ global topography which has been smoothed by two passes of a 9-point smoother.

w117a2 topography

00117:***.A

DX,DY = 0.176,0.125 DEG DB = 500.0 M





Figure 2: Plot of the North Pacific Ocean from the new $1/8^\circ$ global topography which has been smoothed by two passes of a 9-point smoother.

w117a topography

DX,DY = 0.176,0.125 DEG DB = 500.0 M

00117:777.4



NRL 323 20-Oct-92

Figure 3: Plot of the North Atlantic Ocean from the new $1/8^\circ$ global topography which has been smoothed by two passes of a 9-point smoother.

DATA SOURCE: GEOSAT
 DX, DY = 0.176, 0.125 DEG DB = 500.0 M




Figure 4: Plot of the Atlantic Ocean from the new $1/8^\circ$ global topography which has been smoothed by two passes of a 9-point smoother.





Figure 5: Plot of the Aleutian Islands from the unmodified $1/8^\circ$ Pacific topography.

ETOPO5 topography
 DX,DY = 0.083,0.083 DEG DB = 500.0 M 00999: 0.0



NRL 323 11-Jun-92

Figure 6: Plot of the Aleutian Islands from the 1/12° ETOPO5 topography.




Figure 7: Plot of the Aleutian Islands from the new $1/8^\circ$ Pacific topography.




Figure 9: Plot of the Hawaiian Islands and the shoals to the northwest from the unmodified $1/8^\circ$ Pacific topography.




Figure 10: Plot of the Hawaiian Islands and the shoals to the northwest from the 1/12° ETOPO5 topography.

p133d topography

00133: 0.0

DX,DY = 0.176,0.125 DEG DB = 500.0 M





Figure 11: Plot of the Hawaiian Islands and the shoals to the northwest from the new 1/8° Pacific topography

p133d2 topography

00133: 0.0

DX,DY = 0.176,0.125 DEG DB = 500.0 M



MAX DEPTH = 6109.6 M

NRL 323 5-Jun-92

Figure 12: Plot of the Hawaiian Islands and the shoals to the northwest from the new 1/8° Pacific topography which has been smoothed by two passes of a 9-point smoother.




Figure 13: Plot of the Channel Islands west of Los Angeles, California from the unmodified $1/8^\circ$ Pacific topography.




Figure 14: Plot of the Channel Islands west of Los Angeles, California from the 1/12° ETOPO5 topography.




Figure 15: Plot of the Channel Islands west of Los Angeles, California from the new 1/8° Pacific topography.




Figure 16: Plot of the Channel Islands west of Los Angeles, California from the new $1/8^\circ$ Pacific topography which has been smoothed by two passes of a 9-point smoother.




Figure 17: Plot of the Galapagos Islands from the unmodified 1/8° Pacific topography.




Figure 18: Plot of the Galapagos Islands from the 1/12° ETOPO5 topography.




Figure 19: Plot of the Galapagos Islands from the new 1/8° Pacific topography.




Figure 20: Plot of the Galapagos Islands from the new $1/8^\circ$ Pacific topography which has been smoothed by two passes of a 9-point smoother.





Figure 21: Plot of the Line Islands from the unmodified 1/8° Pacific topography.

ETOPO5 topography

DX,DY = 0.083,0.083 DEG DB = 500.0 M

00999: 0.0







Figure 23: Plot of the Line Islands from the new 1/8° Pacific topography.

p133d2 topography

DX,DY = 0.176,0.125 DEG

DR = 500.0 M

00133: 0.0



MAX DEPTH = 5873.0 M

NRL 323 5-Jun-92

Figure 24: Plot of the Line Islands from the new 1/8° Pacific topography which has been smoothed by two passes of a 9-point smoother.




Figure 25: Plot of the Samoan Islands from the unmodified $1/8^\circ$ Pacific topography.




Figure 26: Plot of the Samoan Islands from the 1/12° ETOPO5 topography.




Figure 27: Plot of the Samoan Islands from the new $1/8^\circ$ Pacific topography.




Figure 28: Plot of the Samoan Islands from the new $1/8^\circ$ Pacific topography which has been smoothed by two passes of a 9-point smoother.





Figure 29: Plot of the New Caledonia, New Hebrides, and the Fiji Islands from the unmodified 1/8° Pacific topography.

ETOPO5 topography

00999: 0.0

DX,DY = 0.083,0.083 DEG DB = 500.0 M



MAX DEPTH = 7889.5 M

NRL 323 11-Jun-92

Figure 30: Plot of the New Caledonia, New Hebrides, and the Fiji Islands from the 1/12° ETOPO5 topography.

p133d topography

00133: 0.0

PX, PY = 0.176, 0.125 DEG DB = 500.0 M




Figure 31: Plot of the New Caledonia, New Hebrides, and the Fiji Islands from the new 1/8° Pacific topography.





Figure 32: Plot of the New Caledonia, New Hebrides, and the Fiji Islands from the new $1/8^\circ$ Pacific topography which has been smoothed by two passes of a 9-point smoother.

p133c topography

00133: 0.0

DX,DY = 0.176,0.125 DEG DB = 500.0 M



MAX DEPTH = 6500.0 M


NRL 323 11-Jun-92

Figure 33: Plot of the TOGA/COARE region from the unmodified 1/8° Pacific topography.

ETOP05 topography

00999: 0.0

DX,DY = 0.083,0.083 DEG DB = 500.0 M



MAX DEPTH = 7918.0 M


NRL 323 3-Jun-92

Figure 34: Plot of the TOGA/COARE region from the 1/12° ETOP05 topography.

p133d topography

00133: 0.0

DX,DY = 0.176,0.125 DEG DB = 500.0 M



MAX DEPTH = 6500.0 M

NRL 323 3-Jun-92

Figure 35: Plot of the TOGA/COARE region from the new 1/8° Pacific topography.




Figure 37: Plot of the Indonesian Throughflow region from the unmodified 1/8° Pacific topography.




Figure 38: Plot of the Indonesian Throughflow region from the 1/12° ETOPO5 topography.




Figure 39: Plot of the Indonesian Throughflow region from the new $1/8^\circ$ Pacific topography.




Figure 40: Plot of the Indonesian Throughflow region from the new $1/8^\circ$ Pacific topography which has been smoothed by two passes of a 9-point smoother.




Figure 41: Plot of the Philippine Islands and the S China Sea from the unmodified 1/8° Pacific topography.




Figure 42: Plot of the Philippine Islands and the S China Sea from the 1/12° ETOP05 topography.





Figure 43: Plot of the Philippine Islands and the S China Sea from the new 1/8° Pacific topography.

p133d2 topography

DX,DY = 0.176,0.125 DEG DB = 500.0 M

00133: 0.0



NRI 323 5-Jun-92

Figure 44: Plot of the Philippine Islands and the S China Sea from the new $1/8^\circ$ Pacific topography which has been smoothed by two passes of a 9-point smoother.




Figure 45: Plot of the E China Sea from the unmodified 1/8° Pacific topography.




Figure 46: Plot of the E China Sea from the 1/12° ETOPO5 topography.




Figure 47: Plot of the E China Sea from the new 1/8° Pacific topography.




Figure 48: Plot of the E China Sea from the new $1/8^\circ$ Pacific topography which has been smoothed by two passes of a 9-point smoother.

P133c topography

DX,DY = 0.176,0.125 DEG

DB = 500.0 M

00133: 0.0



NRL 323 11 Jun-92

Figure 49: Plot of the Mariana Islands from the unmodified 1/8° Pacific topography.




Figure 50: Plot of the Mariana Islands from the 1/12° ETOPO5 topography.




Figure 51: Plot of the Mariana Islands from the new 1/8° Pacific topography.




Figure 52: Plot of the Mariana Islands from the new $1/8^\circ$ Pacific topography which has been smoothed by two passes of a 9-point smoother.




Figure 53: Plot of the Izu Ridge from the unmodified 1/8° Pacific topography.




Figure 54: Plot of the Izu Ridge from the 1/12° ETOP05 topography.




Figure 55: Plot of the Izu Ridge from the new $1/8^\circ$ Pacific topography.

p133d2 topography
 DX,DY = 0.176,0.125 DEG DB = 500.0 M 00117: 0.0




Figure 56: Plot of the Izu Ridge from the new $1/8^\circ$ Pacific topography which has been smoothed by two passes of a 9-point smoother.




Figure 57: Plot of the Sea of Japan from the unmodified $1/8^\circ$ Pacific topography.




Figure 58: Plot of the Sea of Japan from the 1/12° ETOPO5 topography.




Figure 59: Plot of the Sea of Japan from the new $1/8^\circ$ Pacific topography.




Figure 60: Plot of the Sea of Japan from the new $1/8^\circ$ Pacific topography which has been smoothed by two passes of a 9-point smoother.




Figure 61: Plot of the Sea of Okhotsk from the unmodified $1/8^\circ$ Pacific topography.




Figure 62: Plot of the Sea of Okhotsk from the 1/12° ETOPO5 topography.




Figure 63: Plot of the Sea of Okhotsk from the new 1/8° Pacific topography.





Figure 64: Plot of the Sea of Okhotsk from the new $1/8^\circ$ Pacific topography which has been smoothed by two passes of a 9-point smoother.

global 1/8 deg topography

DX,DY = 0.176,0.125 DEG DB = 500.0 M

00800: 0.0



MAX DEPTH = 4941.9 M

NRL 323 10-Aug-92

Figure 65: Plot of the GIN Sea from the unmodified 1/8° global topography.




Figure 66: Plot of the GIN Sea from the 1/12° ETOP05 topography.





Figure 67: Plot of the GIN Sea from the new 1/8° Atlantic topography.

w117a2 topography

00117: 0.0

DX,DY = 0.176,0.125 DEG DB = 500.0 M



NRL 323 7-Aug-92

Figure 68: Plot of the GIN Sea from the new $1/8^\circ$ Atlantic topography which has been smoothed by two passes of a 9-point smoother.






Figure 69: Plot of the Canary and Azore Islands from the unmodified 1/8° global topography.



NRI 323 3-Aug-92

Figure 70: Plot of the Canary and Azore Islands from the 1/12° ETOPO5 topography.



NRL 323 6-Aug-92

Figure 71: Plot of the Canary and Azore Islands from the new $1/8^\circ$ Atlantic topography.




Figure 72: Plot of the Canary and Azore Islands from the new 1/8° Atlantic topography which has been smoothed by two passes of a 9-point smoother.




Figure 73: Plot of the Carribean Sea and Gulf of Mexico from the unmodified 1/8° global topography.




Figure 74: Plot of the Caribbean Sea and Gulf of Mexico from the 1/12° ETOPO5 topography.




Figure 75: Plot of the Caribbean Sea and Gulf of Mexico from the new 1/8° Atlantic topography.




Figure 76: Plot of the Caribbean Sea and Gulf of Mexico from the new $1/8^\circ$ Atlantic topography which has been smoothed by two passes of a 9-point smoother.




Figure 77: Plot of the Mediterranean Sea from the unmodified 1/8° global topography.





Figure 78: Plot of the Mediterranean Sea from the 1/12° ETOPO5 topography.

w117a topography

00117: 0.0

DX,DY = 0.176,0.125 DEG DB = 500.0 M



MAX DEPTH = 4971.9 M

NRI 323 3-Aug-92


Figure 79: Plot of the Mediterranean Sea from the new 1/8° Atlantic topography.

w117a2 topography

DX,DY = 0.176,0.125 DEG

DB = 500.0 M

00117: 0.0



MAX DEPTH = 4919.1 M

NRL 323 3 Aug-92

Figure 80: Plot of the Mediterranean Sea from the new 1/8° Atlantic topography which has been smoothed by two passes of a 9-point smoother.




Figure 81: Plot of the Drake Passage and the Falkland Islands from the unmodified 1/8° global topography.




Figure 82: Plot of the Drake Passage and the Falkland Islands from the 1/12° ETOP05 topography.





Figure 83: Plot of the Drake Passage and the Falkland Islands from the new $1/8^\circ$ global topography.



NRL 323 3-Aug-92

Figure 84: Plot of the Drake Passage and the Falkland Islands from the new $1/8^\circ$ global topography which has been smoothed by two passes of a 9-point smoother.





Figure 85: Plot of the Arabian Sea from the unmodified 1/8° global topography.



NRL 323 3 Aug-92

Figure 86: Plot of the Arabian Sea from the 1/12° ETOP05 topography.




Figure 87: Plot of the Arabian Sea from the new 1/8° global topography.




Figure 88: Plot of the Arabian Sea from the new $1/8^\circ$ global topography which has been smoothed by two passes of a 9-point smoother.




Figure 89: Plot of the Southwestern Indian Ocean from the unmodified 1/8° global topography.




Figure 90: Plot of the Southwestern Indian Ocean from the 1/12° ETOPO5 topography.




Figure 91: Plot of the Southwestern Indian Ocean from the new $1/8^\circ$ global topography.




Figure 92: Plot of the Southwestern Indian Ocean from the new $1/8^\circ$ global topography which has been smoothed by two passes of a 9-point smoother.




Figure 93: Plot of the Andaman Sea from the unmodified 1/8° global topography.




Figure 94: Plot of the Andaman Sea from the 1/12° ETOPO5 topography.




Figure 95: Plot of the Andaman Sea from the new $1/8^\circ$ global topography.




Figure 96: Plot of the Andaman Sea from the new $1/8^\circ$ global topography which has been smoothed by two passes of a 9-point smoother.




Figure 97: Plot of the Indonesian Throughflow region from the unmodified 1/8° global topography.




Figure 98: Plot of the Indonesian Throughflow region from the 1/12° ETOPO5 topography.

w117a topography
 DX,DY = 0.176,0.125 DEG DB = 500.0 M 00117:***.4





Figure 99: Plot of the Indonesian Throughflow region from the new 1/8° global topography.

w117a2 topography

00117:***.4

DX,DY = 0.176,0.125 DEG DB = 500.0 M



MAX DEPTH = 6530.1 M

NRL 323 29-Oct-92

Figure 100: Plot of the Indonesian Throughflow region from the new 1/8° global topography which has been smoothed by two passes of a 9-point smoother.




Figure 101: Plot of New Zealand from the unmodified 1/8° global topography.





Figure 102: Plot of New Zealand from the 1/12° ETOPO5 topography.

w117a topography
 DX,DY = 0.176,0.125 DEG DB = 500.0 M 00117: 0.0



NRL 323 3 AUG 92

Figure 103: Plot of New Zealand from the new $1/8^\circ$ global topography.





Figure 104: Plot of New Zealand from the new $1/8^\circ$ global topography which has been smoothed by two passes of a 9-point smoother.

Aleutian Islands

DX,DY = 0.176,0.125 DEG CB = 0.0 M



MAX DEPTH = 6500.0 M

NRL 323 21-Sep-92

Figure 105: Plot of Aleutian Island sill depth locations listed in table 1.

Southwest Pacific Ocean

DX,DY = 0.176,0.125 DEG

DB = 0.0 M





Figure 106: Plot of Southwest Pacific Ocean sill depth locations listed in table 1.

East China Sea

DX,DY = 0.176,0.125 DEG

DB = 0.0 M



NRL 323 21-Sep-92

Figure 107: Plot of E China Sea sill depth locations listed in table 1.

Izu Ridge




Figure 108: Plot of Izu Ridge sill depth locations listed in table 1.

Kuril Islands





Figure 109: Plot of Kuril Island sill depth locations listed in table 1.

Caribbean Sea

DX,DY = 0.176,0.125 DEG

DB = 0.0 M




NRL 323 21-Sep-92

Figure 110: Plot of Caribbean Sea and Leeward Island sill depth location listed in table 2.

GIN Sea

DX,DY = 0.176,0.125 DEG DB = 0.0 M




MAX DEPTH = 5464.5 M

NRL 323 21-Sep-92

Figure 111: Plot of GIN Sea sill depth location listed in table 2.

Mediterranean Sea

DX,DY = 0.176,0.125 DEG DB = 0.0 M



NRL 323 21-Sep-92

Figure 112: Plot of Mediterranean Sea sill depth locations listed in table 3.

New Zealand





Figure 113: Plot of New Zealand sill depth location listed in table 3.

Falkland Islands

DX,DY = 0.176,0.125 DEG

DB = 0.0 M



NRL 323 21-Sep-92

Figure 114: Plot of Falkland Island sill depth location listed in table 3.

Arabian Sea





Figure 115: Plot of Arabian Sea sill depth location listed in table 4.

Andaman Sea

DX,DY = 0.176,0.125 DEG

DB = 0.0 M




NRL 323 21-Sep-92

Figure 116: Plot of Andaman Sea sill depth location listed in table 4.

Indonesian Throughflow Region

DX,DY = 0.176,0.125 DEG DB = 0.0 M



MAX DEPTH = 6837.8 M

NRL 323 13-Oct-92

Figure 117: Plot of Indonesian Throughflow sill depth location listed in table 4.

Southwest Indian Ocean




Figure 118: Plot of Southwest Indian Ocean sill depth location listed in table 4.