

AD-A268 491




TION PAGE

Form Approved
OMB No. 0704-0188


age 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Avenue, Washington, DC 20503, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE 1993		3. REPORT TYPE AND DATES COVERED Reprint	
4. TITLE AND SUBTITLE (see title on reprint)				5. FUNDING NUMBERS PE: NWED QAXM WU: 00129	
6. AUTHOR(S) Dubois et al.					
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Armed Forces Radiobiology Research Institute 8901 Wisconsin Ave. Bethesda, MD 20889-5603				8. PERFORMING ORGANIZATION REPORT NUMBER SR93-13	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Defense Nuclear Agency 6801 Telegraph Road Alexandria, VA 22310-3398				10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES					
12a. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited.				12b. DISTRIBUTION CODE	
Accession For 13. ABSTRACT (Maximum 200 words) DTIC TAB ☐ Unannounced ☐ Justification ☐ By _____ Distribution / Availability Codes Dist Avail and/or Special A-1 20 DTIC QUALITY INSPECTED 3 DTIC ELECTE AUG 24 1993 E D 93-19542 93 8 23 035					
14. SUBJECT TERMS				15. NUMBER OF PAGES 8	
				16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED		18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED		19. SECURITY CLASSIFICATION OF ABSTRACT	
				20. LIMITATION OF ABSTRACT	

DISCLAIMER NOTICE



THIS DOCUMENT IS BEST QUALITY AVAILABLE. THE COPY FURNISHED TO DTIC CONTAINED A SIGNIFICANT NUMBER OF PAGES WHICH DO NOT REPRODUCE LEGIBLY.



Hematopoietic growth factors and glucocorticoids synergize to mimic the effects of IL-1 on granulocyte differentiation and IL-1 receptor induction on bone marrow cells in vivo

Claire M. Dubois,¹ Ruth Neta,² Jonathan R. Keller,³ Sten E.W. Jacobsen,¹ Joost J. Oppenheim,¹ Francis Ruscetti¹

¹The Laboratory of Immunoregulation, NCI-Frederick Cancer Research & Development Center, Frederick, MD; ²Department of Experimental Hematology, Armed Forces Radiobiological Research Institute, Bethesda, MD; ³Biological Carcinogenesis and Development Program, Program Resources/DynCorp Inc., NCI-Frederick Cancer Research & Development Center, Frederick, MD
Offprint requests to: Dr. Francis W. Ruscetti, Laboratory of Molecular Immunoregulation, Bldg. 560, Rm. 21-89A, Frederick Cancer Research & Development Center, Frederick, MD 21702-1201

(Received 18 December 1991; in revised form 23 June 1992; accepted 31 August 1992)

Abstract. The mechanisms by which interleukin-1 (IL-1) stimulates hematopoiesis are not clear. We have previously shown that in vivo administration of IL-1 indirectly increases IL-1 receptor (IL-1R) expression on both immature and mature bone marrow (BM) cells, partly due to IL-1-induced hematopoietic growth factor (HGF) production. Because IL-1 also stimulates the hypothalamic pituitary-adrenal axis resulting in the production of glucocorticoids (GC), we assessed whether in vivo treatment with HGF and glucocorticoids upregulates IL-1R. Administration of IL-1 to adrenalectomized mice reduces by 53% IL-specific binding on light density bone marrow (LDBM) cells compared to sham-operated mice. The administration of dexamethasone (dex) alone induced only a slight increase in IL-1R expression but synergized with granulocyte colony-stimulating factor (G-CSF), granulocyte-macrophage CSF (GM-CSF), IL-3 and IL-6 to upregulate IL-1R expression. Flow cytometry analysis using the RB6-8C5 antibody, which is differentially expressed on myeloid cells, indicated that combined G-CSF and dex treatment acts to promote increased numbers of differentiated myeloid progenitors in the bone marrow. Autoradiographic analysis confirmed that while G-CSF and dex increased IL-1R expression on all myeloid cells, it was particularly pronounced for myelocytes, promyelocytes and metamyelocytes. These results suggest that the ability of IL-1 to enhance granulocyte differentiation in vivo is partly due to its ability to induce a cascade of cytokines and steroids which in turn regulate IL-1 receptor expression.

Key words: IL-1 receptor—CSFs—Glucocorticoids—Myelopoiesis

Introduction. Interleukin-1 (IL-1) is a pleiotropic cytokine affecting the immune, inflammatory, neuroendocrine and hematopoietic systems [1,2]. Much of this diversity is based on the ability of IL-1 to induce production of other biologic mediators such as corticotropin releasing factor, corticosteroids, adrenocorticotrophic hormone (ACTH), insulin and hematopoietic growth factors (G-CSF, GM-CSF, IL-6, IL-8, transforming growth factor beta [TGF- β] and tumor necrosis factor alpha [TNF- α]) from multiple cell types.

The ability of IL-1 to synergize with HGFs to promote the growth and differentiation of primitive progenitor cells in

vitro [3-5] suggests that IL-1 also plays an important role in the regulation of hematopoiesis [6]. In vivo administration of IL-1 induces an initial rapid mobilization of neutrophils from the bone marrow, followed by cycling of hematopoietic progenitor cells resulting in the expansion of the granulocytic compartment [7-11]. These effects presumably contribute to the ability of IL-1 to accelerate the recovery of hematopoietic stem cells and blood neutrophils following myelosuppression by chemotherapeutic drugs or exposure to lethal radiation [12-16].


In order to better understand the mechanism of action of IL-1 in vivo, we demonstrated that the IL-1 mediated upregulation of IL-1 receptors on bone marrow cells after in vivo administration of IL-1 occurs by an indirect mechanism [17]. In contrast, several hematopoietic growth factors, such as GM-CSF, G-CSF, IL-3 and IL-6 but not IL-1, act in vitro to upregulate IL-1R on both progenitor cells [18] and mature myeloid cells [18,19]. In vivo administration of HGFs, however, only partially mimic the ability of IL-1 to upregulate IL-1R expression [17]. It is, therefore, likely that other factors also contribute to the in vivo upregulation of IL-1 on hematopoietic progenitor cells. Since the in vivo administration of IL-1 also results in an elevation of corticosteroids in plasma [20] and glucocorticoids enhance in culture the expression of IL-1R on monocytes, B lymphocytes and fibroblasts [21,22], GCs might participate in the upregulation of IL-1R on bone marrow cells observed in response to IL-1. In this study, evidence is presented for a role for GCs as well as HGFs in IL-1R regulation.

Materials and methods

Mice. CD2F1 male mice were purchased from the Animal Genetics and Production Branch, National Cancer Institute, NIH (Frederick, MD). Animals were handled as previously described [23]. Animal care was provided in accordance with the procedures outlined in the *Guide for Care and Use of Laboratory Animals* (NIH Publication #86-23, 1985).

Reagents. Human recombinant IL-1 α was supplied by Hoffmann-La Roche (Nutley, NJ). Human recombinant G-CSF was supplied by Amgen Corp. (Thousand Oaks, CA). Human recombinant IL-6 (5×10^6 U/mL) in pyrogen-free solution was kindly provided by Dr. Menachem Rubinstein (Interpharm

Fig. 1. Comparative effect of *in vivo* IL-1 on IL-1R expression on BM cells from adrenalectomized (removal of adrenal gland) mice vs. sham-adrenalectomized (similar surgical procedure without removal of the adrenal gland) mice. Mice were adrenalectomized 14 days before injection of 100 ng IL-1. Sixteen to 18 hours after IL-1 administration, BM cells were harvested and tested for the expression of IL-1R as described in Materials and methods. Each bar represents the mean $\pm$ SE of 2 experiments from which the level of background binding (280 ± 41) was subtracted. The numbers in parentheses represent the number of mice receiving each treatment.



Laboratory, Ness-Ziona, Israel). Murine IFN- γ was kindly provided by Genentech (San Francisco, CA). Murine recombinant IL-3 and GM-CSF were generously provided by Dr. Steven Gillis (Immunex Corp., Seattle, WA). Human recombinant TGF- β was generously provided by Dr. Tony Purchio (Bristol-Myers/Squibb Pharmaceuticals, Seattle, WA). Dexamethasone sodium phosphate was purchased from LymphoMed Inc. (Rosemont, IL).

In vivo procedures. Cytokines and dexamethasone were diluted in pyrogen-free saline on day of injection. Predetermined optimal doses of cytokines [17,23,24] were given intraperitoneally (i.p.) at the same time as dex administered at 50 μ g/mouse or at the dose indicated in the text. Bone marrow cells were tested for the expression of IL-1R 16 to 18 hours after treatment. Adrenalectomy was performed under anesthesia 14 days before the experiment. Adrenalectomized mice were given 1.0% (wt/vol) NaCl in drinking water.

Measurement of CSF activity in serum. CSF titer in serum was measured as follows. Briefly, mice were bled 2 to 3 hours after IL-1 injection and serum was collected by centrifugation after clot formation. CSF activity was determined using bone marrow colony assay for CSF activity as previously described [20]. Briefly, BM cells were suspended in 1.0 mL IMDM, 10% FCS, in 0.3% Seaplaque agarose (Rockland, ME) in the presence or absence of serum (serial 2-fold dilutions). The cells were plated in 35 mm Lux petri dishes (Miles Laboratories Inc., Naperville, IL) and incubated at 37°C in 5.0% CO₂ and scored for colonies (>50 cells) growth after 7 days of incubation. CSF activity was expressed as colony-forming units per millimeter, based on the colony count at 50% of maximum response (Ed50).

Preparation of bone marrow cells. Murine bone marrow cells were aspirated from femurs and low-density mononuclear cells were isolated by separation on Lymphocyte Separation Medium (Organon Teknika Corp., Durham, NC).

Flow cytometry analysis of bone marrow cells. BM cells from saline-, dex- and/or cytokine-treated mice were labeled with either monoclonal antibody (MAB) RB6-8C5, Thy-1, L3T4, F480 or control IgG in an indirect immunofluorescence assay. Briefly, 10⁶ BM cells were resuspended in RPMI with 10% fetal calf serum and incubated for 30 minutes at 4°C with 1.0 μ g of

the appropriate MAB. The cells were washed and then incubated with fluorescein-labeled goat antirat antibody for 30 minutes at 4°C. The cells were then washed 2 times with PBS alone, fixed with 1.0% paraformaldehyde in PBS and analyzed using Coulter Profile II.

Preparation of iodinated IL-1. Human rIL-1 α was labeled with ¹²⁵I using chloramine-T reagent as described previously [25]. The radiolabeled IL-1 α had a specific activity that ranged from 1 to 3x10¹⁵ cpm/mmol. There was no significant loss of biological activity of radiolabeled IL-1 α as determined by the thymocyte comitogenic activity assay.

Receptor binding assay. Fractionated bone marrow cell suspensions were washed once with cold medium and cell pellets were treated for 1 minute on ice with 50 mM glycine-HCl (pH 3.0) to remove potentially bound cytokines. Subsequently, the cells were washed twice with binding medium (RPMI 1.0% BSA supplemented with 0.1% sodium azide and 10 mM Hepes) and incubated at 4°C with 500 pm ¹²⁵I-labeled IL-1 α in a final volume of 0.2 mL. After 1 to 2 hours of incubation, cell-bound radioactivity was separated from unbound ¹²⁵I-IL-1 α by centrifugation of the sample through a mixture of 1.5:1 (vol/vol) dibutyl phthalate/bis(2-ethylhexyl)-phthalate (Eastman Kodak Co., Rochester, NY). Nonspecific binding was determined by incubating bone marrow cells with labeled IL-1 α in the presence of 50-fold excess of unlabeled ligand.

Autoradiography. LDBM cells from mice treated 16 to 18 hours with IL-1, G-CSF and/or dex were prepared as described for receptor binding assay and incubated at 4°C with 1.0 nM ¹²⁵I-IL-1 α . After 1 hour of incubation, cell-bound radioactivity was separated from unbound IL-1 by centrifugation of the sample through a layer of cold FBS. The autoradiography was performed using a modification of a previously described technique [26]. Briefly, 2x10⁵ cells were centrifuged onto microscope slides coated with 0.5% gelatin, fixed in methanol for 10 minutes, coated with Kodak NTB2 photographic emulsion and exposed at 4°C for 4 weeks. Slides were developed with Kodak D-19 developer, fixed with Kodak fixer, stained with Jenner-Giemsa. The number of grains was determined for over 50 cells per slide for 2 slides.

Results

Endogenous corticosteroid production is involved in the upregula-

Table 1. IL-1-stimulated CSF production in adrenalectomized and sham-adrenalectomized mice

Mice ^a	IL-1 injection (ug/mice) ^b	CSF titer (U/mL) ^c
Sham-ADX	None	< 20
Sham-ADX	0.1	590
Sham-ADX	1.0	705
ADX	None	< 20
ADX	0.1	750
ADX	1.0	853

^aMice were either adrenalectomized or sham-adrenalectomized as described [23].


^bMice were bled 2 to 3 hours after IL-1 injection as described in Materials and methods.

^cSerum was assayed for CSF activity as described in Materials and methods. The data represent CSF titers of pooled serum from 3 mice.

tion of IL-1R. To examine the role of endogenous corticosteroids in the upregulation of IL-1R on bone marrow cells after IL-1 injection, mice were adrenalectomized. The highest dose of IL-1 (100 ng/mouse) that adrenalectomized mice could tolerate was given i.p. 14 days after surgery. As already demonstrated in normal animals [17], the administration of 100 ng of IL-1 to sham-adrenalectomized mice resulted in a 15-fold increase in IL-1-specific binding compared to saline-treated sham-adrenalectomized mice at 16 hours after injection of IL-1 (Fig. 1). This time was chosen based on our previous studies showing maximal IL-1R expression at 16 to 18 hours [17]. The specific binding of IL-1 on BM cells from IL-1-treated adrenalectomized mice was reduced by 53% compared with IL-1-treated sham-adrenalectomized mice. No specific IL-1 binding was detectable on bone marrow cells from adrenalectomized mice. These data suggest that endogenous corticosteroids participate in the constitutive and IL-1-induced expression of IL-1R.

Since it has been established that IL-1 induces HGF production [2,23], we studied whether adrenalectomy influences IL-stimulated HGF production. For this, adrenalectomized and control (sham-adrenalectomized) mice were bled 2 to 3 hours after IL-1 injection. IL-1 induced similar levels of CSF production in all mice tested (Table 1), indicating that adrenalectomy does not impair IL-stimulated HGF production. We next examined the ability of exogenous addition of dex to synergize with IL-1 in the regulation of IL-1R expression. The injection of optimal concentrations of dex (50 µg) [27] in the presence of optimal amounts of IL-1 (1.0 µg) [17] did not increase IL-1R expression (Fig. 2). At concentrations of 0.1 and 0.3 µg of IL-1 per mouse, however, dex increased IL-1R expression 2-fold. The magnitude of IL-1 upregulation in these cases, however, did not equal the magnitude seen with optimal IL-1 concentrations.

Dexamethasone synergizes with HGFs in the upregulation of IL-1R. Since we have recently determined that increased IL-1R expression after IL-1 administration to mice was mediated, in part, through endogenous HGF production [17], we examined the effect of dexamethasone in combination with HGFs on IL-1R expression on BM cells. Mice were injected with equivalent doses of hematopoietic growth factors such as G-CSF, GM-CSF, IL-3 and IL-6 as well as TGF-β, a negative regu-

**Fig. 2.** Dose-dependent interaction between dex and IL-1 administration for increase in IL-1R expression

Mice were injected i.p. with either saline, the indicated doses of IL-1 and maximal dose (50 µg) of dex. Sixteen to 18 hours after treatment radioreceptor assays for the expression of IL-1R on BM cells were done as described in Materials and methods. The data represent the mean ± SE of duplicate determinations using pooled cells from 3 animals. Background was 259 ± 34, which was subtracted from the data shown.

lator of hematopoiesis. Among the cytokines tested, G-CSF, GM-CSF, IL-3 or IL-6 treatment in combination with dex (50 µg) results in a synergistic effect on IL-1R expression on bone marrow cells (Fig. 3). At equivalent doses, rG-CSF was slightly more potent than GM-CSF, followed by IL-6 and IL-3 (Fig. 3). G-CSF plus dex and GM-CSF plus dex were equal to or better than IL-1. TGF-β alone did not increase IL-1R expression and did not synergize with dexamethasone.

Using G-CSF, the dose-dependence of the synergistic interaction between dex and HGFs was studied. The administration of a previously determined [17] optimal dose of G-CSF significantly increased the expression of IL-1R on BM cells (4.3-fold) (Fig. 4). The administration of dex alone induced a 2-fold increase in IL-1R expression but synergized with G-CSF to upregulate IL-1R in vivo in a dose-dependent fashion (Fig. 4). In all experiments performed (n=7), the administration of G-CSF (5.0 µg) and dex (50 µg) upregulated IL-1R expression to a greater extent than seen with an optimal dose (1.0 µg) of IL-1 alone (Fig. 4). This synergy with G-CSF in increasing IL-1R expression was dose-dependent on dex (Fig. 4).

To determine if G-CSF and dex can act on isolated hematopoietic cells in vitro, normal BM cells were incubated in vitro with G-CSF and/or different concentrations of dex. G-

Fig. 3. Hematopoietic growth factors synergize with dexamethasone to upregulate IL-1R expression

Mice were injected i.p. with either saline, IL-1 (1.0 μ g), G-CSF (5.0 μ g), GM-CSF (5.0 μ g), IL-3 (5.0 μ g), IL-6 (5.0 μ g), TGF- β (5.0 μ g) and/or dex (50 μ g). Sixteen to 18 hours after treatment, BM cells were harvested and tested for the expression of IL-1R using radioreceptor assay. The data represent the mean $\pm$ SE of determinations using 3 to 15 mice as indicated in parentheses, 2 to 4 experiments for each cytokine. The level of background binding was 315 ± 52 , which was subtracted from the data shown here.





Fig. 4. Dose-dependent synergistic interaction between dexamethasone and G-CSF administration for the increase in IL-1R expression

Mice were injected i.p. with either saline, maximal dose of IL-1 (1.0 μ g), maximal dose of G-CSF (5.0 μ g) and/or increasing doses of dex. Sixteen to 18 hours after the treatments, radioreceptor assay for the expression of IL-1R on BM cells was performed as described in Materials and methods. The data represent the mean $\pm$ SE of duplicate determinations of at least 2 experiments using pooled cells from 3 animals. The level of background binding was 259 ± 34 , which was subtracted from the data shown here.

CSF and dex also synergize in vitro to increase IL-1-specific binding (Table 2), whereas IL-1 by itself had no effect.

Distribution of IL-1R on bone marrow cells from G-CSF- and dex-treated mice. We next evaluated whether the in vivo administration of G-CSF and dex increased expression of IL-1R on a specific population or subpopulation of BM cells. Mice received a single injection of saline or a combination of G-CSF and dex. IL-1 binding to BM cells was determined by autoradiography, 16 to 18 hours after injection. Autoradiographic analysis of cells from saline-treated animals showed that most of the labeled cells belonged to the granulocytic series (Table 3, Fig. 5). Seven percent of the undifferentiated blast/early cells were labeled with 8 grains per cell. After treatment with G-CSF and dex, these cells were 19% positive with 28 grains per cell. Promyelocytes and myelocytes (43% positive with 19 grains per cell) were the most heavily labeled cells after G-CSF and dex treatment (87% positive with 57 grains per cell). Thirteen and 16% of eosinophilic and monocytic cells exhibited a similar pattern of labeling with 7 to 8

specific grain per cell. More monocytic cells were labeled after G-CSF and dex treatment (31%) with a small increase in the number of grains per cell (8 to 17). This treatment had no effect on eosinophilic cells. No specific IL-1 labeling was observed on erythroid cells. These results clearly demonstrate that G-CSF and dex treatment results in an increase in IL-1R expression along the myelocytic series and is particularly pronounced for myelocytes followed by metamyelocytes and segmented neutrophils.

G-CSF and dexamethasone promote granulopoiesis. Because the administration of IL-1 induces an initial rapid mobilization of neutrophils from the bone marrow followed by increased cycling of hematopoietic progenitor cells, resulting in the expansion of granulocytes in the marrow [7-11] we examined whether the combination of G-CSF and dex also promoted an expansion of myeloid cells. Bone marrow cells were analyzed by fluorescence-activated cell sorting according to the differential expression of RB6-8C5 antigen on myeloid cells [28].

Table 2. In vitro interaction between G-CSF and dexamethasone in the upregulation of IL-1R on bone marrow cells

Factor added ^a	IL-1-specific binding (CPM) ^b
None	798 ± 56
DEX 10 ⁻⁹ M	1341 ± 30
G-CSF	4195 ± 273
G-CSF + DEX 10 ⁻¹⁰ M	5985 ± 409
G-CSF + DEX 10 ⁻⁹ M	8685 ± 22
IL-1	747 ± 220

^aLDBM cells obtained from normal mice by Ficoll separation were incubated 24 hours in the presence or absence of the 20 ng/mL of IL-1 or G-CSF and the indicated concentrations of dex.

^bBone marrow cells were treated for IL-1 binding as described in Materials and methods. The data represent the mean ± SEM of duplicate determination of a representative experiment of 2 experiments. The level of background binding was 402 ± 168 cpm which was subtracted from the total cpm to give the data shown here.

The RB6-8C5^{hi} cells are enriched for the end stage (segmented) neutrophils (>75%) while the RB6-8C5^{lo} cells are enriched for myeloblasts, promyelocytes and myelocytes (>80%). The RB6-8C5^{lo} population represents 19 to 24% of total bone marrow and contains 50% of CFU-GM progenitors [28]. As previously demonstrated [28], the administration of IL-1 to mice results in a 206% increase in the RB6-8C5^{lo} population and a concomitant 41% loss of RB6-8C5^{hi} (Fig. 6, Table 4). Treatment of mice with G-CSF and dex induced a 406% increase in the RB6-8C5^{lo} immature myeloid population and no reduction in the RB6-8C5^{hi} population. In comparison, dex and G-CSF promoted a 206% and 218% increase in RB6-8C5^{lo} population, respectively, and a 137% increase and a 45% reduction in the RB6-8C5^{hi} population, respectively.

Discussion

We have previously demonstrated that injection of mice with IL-1 results in considerable upregulation of type II IL-1R on myeloid-enriched progenitors [17]. By the administration of antibody against type I IL-1R not present on bone marrow progenitors, we have clearly demonstrated that this upregulation occurs through an indirect mechanism. Administration of IL-1 in vivo stimulates the hypothalamic-pituitary-adrenal axis, resulting in the production of GC [20] as well as HGF production by type I IL-1R expressing accessory cells [29-32]. Both GC and HGF are elevated rapidly, with maximal levels reached within 2 hours after IL-1 administration [20,29]. GC have been previously shown to upregulate IL-1R on human monocytes and B cells in vitro [21,22].

In this report, evidence is presented that IL-1-induced endogenous GC synergize with HGF to mediate IL-1 responsiveness on bone marrow cells. The injection of IL-1 to adrenalectomized mice reduced the upregulation of IL-1R by 53% compared with sham-adrenalectomized mice. This effect, which is associated with diminished GC production, is not due to impaired IL-1-induced HGF production in adrenalectomized animals (Table 1). The concomitant administration of GC and HGF such as G-CSF, GM-CSF, IL-3 and IL-6, to normal mice synergistically increased functional IL-1R on bone marrow cells. In addition, this synergy between GC and HGF was also seen in vitro. The observations concerning glu-

Table 3. Distribution of IL-1R on BM cells from G-CSF plus dexamethasone-treated mice

Cell type	Specific IL-1 binding			
	% labeled		mean grain count	
	saline	G-CSF + Dex	saline	G-CSF + Dex
Blasts/early cells	7	19	8	28
Promyelocytes/myelocytes	43	87	19	57
Metamyelocytes	29	72	11	30
Later neutrophils	23	39	7	23
Eosinophils	13	13	7	8
Monocyte	16	31	8	17
Nucleated erythroid	0	0	0	0

Data represent background-subtracted grain count over more than 20 cells of each type. The level of background binding of >5 grains was subtracted from the total grain count to give the data shown here.

corticoid modulation of IL-1R in vivo using adrenalectomized mice and the synergy between G-CSF and dex are in agreement with a recent observation from Shieh et al. [19]. While the data indicate a role for GC in IL-1 receptor regulation, other mechanisms cannot be excluded.

In addition, we found that GM-CSF and G-CSF were equally potent in synergizing with dex while IL-6 and IL-3 were approximately 50% as potent. Unlike IL-1, HGF and GCs can stimulate IL-1R expression on hematopoietic cells in vitro [33, 34]. Whether this is a direct effect must await the results of technically difficult binding and antibody-blocking assays on single cells.

Because administration of IL-1 rapidly induces an initial mobilization of neutrophils from the bone marrow followed by increased cycling of hematopoietic progenitor cells, resulting in the expansion of granulocytic compartment in the bone marrow [20,23], we examined whether G-CSF and dex have the same effect. Bone marrow cells were analyzed by fluorescence-activated cell sorting according to the differential expression of RB6-8C5 antigen on myeloid cells [28]. We had previously shown that RB6-8C5^{hi} cells are enriched for the end stage (segmented) neutrophils (>75%) while the RB6-8C5^{lo} cells are enriched for myeloblasts, promyelocytes and myelocytes (>80%). While the administration of IL-1 to mice results in a 206% increase in the RB6-8C5^{lo} population and a concomitant 41% loss of RB6-8C5^{hi}, treatment of mice with G-CSF and dex induced a 406% increase in the RB6-8C5^{lo} immature myeloid population and no reduction in the RB6-8C5^{hi} population. In addition, autoradiographic analysis of IL-1 binding on bone marrow cells after G-CSF and dex treatment showed that while cells in all stages of granulocytic development had increased IL-1 binding, the most dramatic increase in terms of number of cells positive and grains per cell were the myelocytes, promyelocytes and metamyelocytes.




Fig. 5. Autoradiography of ^{125}I -IL-1-labeled LDBM cells from control and G-CSF plus dex-treated mice. Mice received a single i.p. injection of G-CSF (5.0 μg) and dex (50 μg). After 16 hours, bone marrow cells were harvested and radioautography of BMC labeled with ^{125}I -IL-1. Emulsion films were developed after 4 weeks. Panels **A**, **C** and **D** without cold IL-1; panel **B** with an excess of cold IL-1.

Whether the increased IL-1R binding leads to increased IL-1 responsiveness is being studied. Thus, it is shown that G-CSF and dex in combination *in vivo* mimic the effects of IL-1 in granulocyte differentiation.

In addition, we have previously shown that an antibody against type I IL-1R not expressed on neutrophils blocked most of the initial mobilization of neutrophils together with HGF production by type I-expressing stromal cells [17,23]. This confirms that chemotactic response of neutrophils to IL-1 is indirect, probably mediated through IL-1-induced potent chemotactic cytokines such as IL-8 [35,36]. In this report, we show that G-CSF alone can mimic the extent of the initial mobilization of bone marrow neutrophils due to IL-1 (Fig 6, Table 4). Unlike IL-1, G-CSF is directly chemotactic for neutrophils *in vitro* [37]. It is therefore likely that G-CSF participates with other cytokines induced by IL-1 in the mobilization of neutrophils observed after IL-1 administration.

In general, the amplitude of the response to IL-1 correlates with cell surface receptor expression. For example, positive regulators such as PDGF increased the number of IL-1R together with the capacity of the cell to respond to IL-1 [38]. In addition, treatment of hematopoietic progenitor cells with negative regulators such as TGF- β blocked the ability of IL-1 to promote high proliferative potential (HPP) colony forma-




Fig. 6. Differential expression of RB6-8C5 antigen on BM cells from mice treated with G-CSF and/or dexamethasone. Mice were injected i.p. with either saline, IL-1 (1.0 μg), G-CSF (5.0 μg) and/or dex (50 μg). Sixteen to 18 hours after treatment, BM cells were labeled in indirect immunofluorescence assay by using the MAB RB6-8C5 as outlined in Materials and methods. The cells were gated according to fluorescent intensity into 8CS^{neg} (solid region) showing fluorescence between channels 0 and 60, 8CS^{lo} (hatched region) between 60 and 175 and 8CS^H (open region) between 175 and 250. The background staining of the isotype-matched control antibody was less than 3.5% for each treatment.

tion as well as greatly reduced the expression of IL-1R expression [39]. Clinically, increased IL-1R expression has been noted in sepsis, organ failure and acute disseminated inflammation [40]. In this report, concomitant injection of HGF and GC increased IL-1R expression on myeloid cells in the bone marrow with the most increase seen on the myelocyte and promyelocytes followed by metamyelocytes and segmented neutrophils. Such an increase in IL-1R on a premitotic population would serve to promote cellular differentiation and/or cell division, resulting in an amplification of granulocyte differentiation. Therefore, the ability of IL-1 to enhance granulopoiesis in normal [7-11] as well as in myelosuppressed [12-16] mice may be partly due to the unique ability of IL-1 to induce a complex cascade of cytokines and steroids, which can then act to regulate IL-1 receptor expression.

In conclusion, these results provide new insights into the mechanism of IL-1 restorative effects in the marrow. IL-1 stimulates production of HGF and GC which in turn upregulate the expression of IL-1R and render the cells more responsive to IL-1. This accounts for the initial burst of granu-

Table 4. Flow cytometric analysis of BM cells from dex- and G-CSF-treated mice using RB6-8C5 antibody

Treatments ^a	Populations ^b	
	RB6-8C5 ^{lo}	RB6-8C5 ^{hi}
Saline	7.7 ± 2.4 (100)	15.8 ± 1.1 (100)
IL-1	16.3 ± 4.6 (210)	9.8 ± 1.3 (62)
Dex	15.4 ± 5.6 (200)	23.3 ± 0.9 (147)
G-CSF	23.6 ± 2.2 (306)	11.8 ± 1.6 (74)
G-CSF + Dex	31.3 ± 7.4 (406)	18.0 ± 6.0 (114)

^aMice were injected i.p. with either saline, maximal doses of IL-1 (1 µg), G-CSF (5 µg) and/or dex (50 µg). Sixteen to 18 hours after treatment, 10⁶ BM cells were labeled in an indirect immunofluorescence assay by using the MAB RB6-8C5 as outlined in Materials and methods. Background staining of the isotype-matched control antibody was >3.5% for each treatment. Data represent mean ± SEM for 2 experiments using pooled cells from 3 mice.

^bThe numbers in parentheses represent the percent of stimulation and inhibition of controls (saline-treated). Using an established protocol (28), 8C5^{neg} showed fluorescence between channels 0 and 60, 8C5^{lo} between 60 and 175 and 8C5^{hi} between 175 and 250.

lopoiesis seen after administration of even nanogram amounts of IL-1.

Acknowledgments

This project was funded in part by federal funds from the Department of Human Health and Services under contract #N01-CO-74102. The content of this publication does not necessarily reflect the views or policies of the Department of Health and Human Services, nor does mention of trade names, commercial products or organizations imply endorsement by the U.S. government. In addition, the work was supported by the Armed Forces Radiobiology Research Institute, Defense Nuclear Agency, under work unit 00129. No endorsement by the Defense Nuclear Agency or the Department of Defense has been given or should be inferred. Claire M. Dubois is a recipient of a research fellowship from the Fonds de la Recherche en Santé du Québec.

We thank Dr. Dan L. Longo for helpful discussions and review of this manuscript.

References

- Oppenheim JJ, Kovacs EJ, Matsushima K, Durum SK (1986) There is more than one interleukin-1. *Immunol Today* 7:45
- Dinarello CA (1991) Interleukin-1 and interleukin-1 antagonism. *Blood* 77:1627
- Stanley ER, Bartocci A, Patinkin D, Rosendall M, Bradley TR (1986) Regulation of very primitive, multipotent, hemopoietic cells by hemopoietin-1. *Cell* 4:669
- Mochizuki DY, Eisenman JR, Conlon PJ, Larsen AD, Tushinski RJ (1987) Interleukin-1 regulates hematopoietic activity, a role previously ascribed to hemopoietin-1. *Proc Natl Acad Sci USA* 84:5267
- Zsebo KM, Wypych J, Yuschenkoff VN, Lu H, Hunt P, Dukes PP, Langley KE (1988) Effects of hematopoietin-1 and interleukin-1 activities on early hematopoietic cells of the bone marrow. *Blood* 71:962
- Moore MAS (1989) Role of interleukin-1 in hematopoiesis. *Immunol Res* 8:165
- Kampschmidt RF, Upchurch HF (1977) Possible involvement of leukocytic endogenous mediator in granulopoiesis. *Proc Soc Exp Biol Med* 155:89
- Stork LC, Peterson VM, Rundus CH, Robinson WA (1988) Interleukin-1 enhances murine granulopoiesis in vivo. *Exp Hematol* 16:163
- Neta R, Szein MB, Oppenheim JJ, Gillis S, Douches SD (1987) The in vivo effect of interleukin-1. I. Bone marrow cells are induced to cycle after administration of interleukin-1. *J Immunol* 139:1861
- Johnson CJ, Douglas JK, Topper MI, Braunschweiger PG, Furmanski P (1989) In vivo hematopoietic effects of recombinant interleukin-1α in mice: Stimulation of granulocytic, monocytic, megakaryocytic, and early erythroid progenitors, suppression of late-stage erythropoiesis, and reversal of erythroid suppression with erythropoietin. *Blood* 73:678
- Newton RC, Sandlin G, Pezzella K, Huang J, McKearn J (1989) Flow cytometric analysis of the effect of interleukin-1 administration on bone marrow populations in mice. *J Biol Response Mod* 8:155
- Moore MAS, Warren DJ (1987) Synergy of interleukin-1 and granulocyte colony-stimulating factor: in vivo stimulation of stem-cell recovery and hematopoietic regeneration following 5-fluorouracil treatment of mice. *Proc Natl Acad Sci USA* 84:7134
- Benjamin WR, Tare NS, Hayes TJ, Becker JM, Anderson TD (1989) Regulation of hemopoiesis in myelosuppressed mice by human recombinant IL-1α. *J Immunol* 142:792
- Moreb J, Zucali JR, Gross MA, Weiner RS (1989) Protective effects of IL-1 on human hematopoietic progenitor cells treated in vitro with 4-hydroperoxycyclophosphamide. *J Immunol* 142:1937
- Futami H, Jansen R, MacPhee MJ, Keller J, McCormick K, Longo DL, Oppenheim JJ, Ruscetti FW, Wiltrott RH (1990) Chemoprotective effects of rHL-1α in cyclophosphamide-treated normal and tumor-bearing mice: protection from acute toxicity, hematological effects, development of late mortality and enhanced therapeutic efficacy. *J Immunol* 145:4121
- Neta R, Douches S, Oppenheim JJ (1986) Interleukin-1 is a radioprotector. *J Immunol* 136:2483
- Dubois CM, Ruscetti FW, Keller JR, Oppenheim JJ, Hestdal K, Chizzonite R, Neta R (1991) In vivo IL-1 administration indirectly promotes type II IL-1R expression on hematopoietic bone marrow cells: novel mechanism for the hematopoietic effects of IL-1. *Blood* 78:2841
- Dubois CM, Ruscetti FW, Oppenheim JJ, Keller JR (1990) Induction of interleukin-1 receptor (IL-1R) on normal hematopoietic progenitor cells by colony-stimulating factor. *Exp Hematol* 18:615 (abstr)
- Shieh J-H, Peterson RHF, Moore MAS (1991) IL-1 modulation of cytokine receptors on bone marrow cells. *J Immunol* 147:1273
- Besedovski H, Adriana DR, Sorkin E, Dinarello C (1986) Immunoregulatory feedback between interleukin-1 and glucocorticoid hormones. *Science* 233:652
- Akahoshi T, Oppenheim JJ, Matsushima K (1988) Interleukin-1 stimulates its own receptor on human fibroblasts through the endogenous production of prostaglandins. *J Clin Invest* 82:1219
- Akahoshi T, Oppenheim JJ, Matsushima K (1988)

- Induction of high affinity interleukin-1 receptor on human peripheral blood lymphocytes by glucocorticoid hormones. *J Exp Med* 167:924
23. Neta R, Vogel SN, Plocinski JM, Tare NS, Benjamin W, Chizzonite R, Pilcher M (1990) In vivo modulation with anti-interleukin-1 (IL-1) receptor (p80) antibody 35F5 of the response to IL-1: the relationship of radioprotection, colony-stimulating factor, and IL-6. *Blood* 76:57
24. Neta R, Vogel SN, Sipe JD, Wong GG, Nordan RP (1987) Comparison of in vivo effects of human recombinant IL-1 and human recombinant IL-6 in mice. *Lymphokine Res* 7:403
25. Palaszynski EW, Ihle JN (1984) Evidence for specific receptors for interleukin-3 on lymphokine-dependent cell lines established from long-term bone marrow cultures. *J Immunol* 132:1872
26. Nicola NA, Metcalf D (1985) Binding of 125 I-labeled granulocyte colony-stimulating factor to normal murine hematopoietic cells. *J Cell Physiol* 124:313
27. Gustavson LE, Benet LZ (1985) Pharmacokinetics of natural and synthetic glucocorticoids. In: Anderson DC, Winder JSD (eds) *The adrenal cortex*. Butterworth, NY: Wiley-Liss, 321
28. Hestdal K, Ruscetti FW, Ihle N, Jacobsen SEW, Dubois CM, Kopp WC, Longo DL, Keller JR (1991) Characterization and regulation of RB6-8C5 antigen expression on murine bone marrow cells. *J Immunol* 147:22
29. Vogel SN, Douches SD, Kaufman EN, Neta R (1987) Induction of colony-stimulating factor in vivo by recombinant interleukin-1 α and recombinant tumor necrosis factor. *J Immunol* 138:2143
30. Zucali JR, Dinarello CA, Oblon DJ, Gross MA, Anderson L, Weiner RS (1986) Interleukin-1 stimulates fibroblasts to produce granulocyte-macrophage colony-stimulating activity and prostaglandin E₂. *J Clin Invest* 77:1857
31. Bagby GC, Dinarello CA, Wallace P, Wagner C, Hefeneider S, McCall E (1986) Interleukin-1 stimulates granulocyte-macrophage colony-stimulating activity release by vascular endothelial cells. *J Clin Invest* 78:1316
32. Fibbe WE, Van Damme J, Billiau A, Voogt PJ, Duinkerken N, Kluck PMC, Falkenburg JHF (1986) Interleukin-1 (22 K factor) induces release of granulocyte-macrophage colony-stimulating activity from human mononuclear phagocytes. *Blood* 68:1316
33. Shieh J-H, Peterson RHF, Moore MAS (1991) Granulocyte colony-stimulating factor modulation of cytokine receptors on bone marrow cells. *J Immunol* 147:2984
34. Dubois CM, Neta R, Ruscetti FW, Jacobsen SEW, Oppenheim JJ, Keller JK (1990) Upregulation of p65 IL-1 receptor (IL-1R) on bone marrow cells by G-CSF and glucocorticoids. *J Cell Biochem suppl* 15F:109 (abstr)
35. Yoshimura T, Matsushima K, Oppenheim JJ, Leonard EJ (1987) Neutrophil chemotactic factor produced by lipopolysaccharide (LPS)-stimulated human blood mononuclear leukocytes: partial characterization and separation from interleukin-1 (IL-1). *J Immunol* 139:788
36. Matsushima K, Oppenheim JJ (1989) Interleukin-8 and MCAF: novel inflammatory cytokines inducible by IL-1 and TNF. *Cytokine* 1:2
37. Wang JM, Chen ZG, Colella S, Bonilla MA, Welte K, Bordignon C, Mantovani A (1988) Chemotactic activity of recombinant human granulocyte colony-stimulating factor. *Blood* 72:1456
38. Bonin PD, Sings JP (1988) Modulation of interleukin-1 receptor expression and interleukin-1 response in fibroblasts by platelet-derived growth factor. *J Biol Chem* 263:11052
39. Dubois CM, Ruscetti FR, Palaszynski EW, Falk LA, Oppenheim JJ, Keller JR (1990) Transforming growth factor β is a potent inhibitor of interleukin-1 (IL-1) receptor expression: proposed mechanism of inhibition of IL-1 action. *J Exp Med* 172:737
40. Fasano MB, Cousart S, Neal S, McCall C (1991) Increased Expression of the interleukin-1 receptor on blood neutrophils with the sepsis syndrome. *J Clin Invest* 88:1452