

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor Battelle Memorial Institute, nor any of their employees, makes any warranty, expressed or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or Battelle Memorial Institute. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

PACIFIC NORTHWEST LABORATORY
operated by
BATTELLE MEMORIAL INSTITUTE
for the
UNITED STATES DEPARTMENT OF ENERGY
under Contract DE-AC06-76RLO 1830

Printed in the United States of America

*Available to DOE and DOE contractors from the
Office of Scientific and Technical Information, P.O. Box 62, Oak Ridge, TN 37831;
prices available from (615) 576-8401. FTS 626-8401.*

*Available to the public from the National Technical Information Service,
U.S. Department of Commerce, 5285 Port Royal Rd., Springfield, VA 22161.*

ADA 269657

**AN EVALUATION OF THE ENVIRONMENTAL
FATE AND BEHAVIOR OF MUNITIONS MATERIEL
(TETRYL AND POLAR METABOLITES OF TNT) IN
SOIL AND PLANT SYSTEMS**

Preliminary Evaluation of TNT-Polar Metabolites in Plants

Project Order No. 88PP8853

**R. J. Fellows
S. D. Harvey
D. A. Cataldo**

November 1992

**Prepared for
U.S. Army Medical Research and Development Command
under a Related Services Agreement
with the U.S. Department of Energy
Contract DE-AC06-76RLO 1830**

Final Report

**Pacific Northwest Laboratory
Richland, Washington 99352**

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)

2. REPORT DATE

November 1992

3. REPORT TYPE AND DATES COVERED

Final Report

4. TITLE AND SUBTITLE An Evaluation of the Environmental Fate and Behavior of Munitions Materiel (Tetryl and Polar Metabolites of TNT) in Soil and Plant Systems -- Preliminary Evaluation of TNT-Polar Metabolites in Plants

5. FUNDING NUMBERS

88PP8853

62720A

3E162720A835 00

DA314939

6. AUTHOR(S)

R. J. Fellows, S. D. Harvey, D. A. Cataldo

7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)

Pacific Northwest Laboratory
P. O. Box 999
Richland, WA 99352

8. PERFORMING ORGANIZATION
REPORT NUMBER

9. SPONSORING, MONITORING AGENCY NAME(S) AND ADDRESS(ES)

U.S. Army Biomedical Research and Development Laboratory
Fort Detrick
Frederick, MD 21702-5010

10. SPONSORING, MONITORING
AGENCY REPORT NUMBER

11. SUPPLEMENTARY NOTES

Subtitle: Preliminary Evaluation of TNT-Polar Metabolites in Plants

12a. DISTRIBUTION, AVAILABILITY STATEMENT

Approved for publication; distribution unlimited

12b. DISTRIBUTION CODE

13. ABSTRACT (Maximum 200 words)

A preliminary study was conducted to elucidate the nature of the polar metabolites reported for plant tissues and xylem exudates following root accumulation of trinitrotoluene. Studies focused on the nature of the proposed conjugates of TNT-residues in xylem exudates of bush bean plants. Use of enzymatic methods failed to demonstrate that the conjugates were carbohydrate based, but acid hydrolysis indicated that the conjugates may be protein based. Of the five polar conjugates isolated from exudates, the presence of aminodinitrotoluene isomers and one unknown TNT residue was demonstrated.

14. SUBJECT TERMS

munitions, bioavailability, plant uptake and metabolism,
environmental fate, RA3

15. NUMBER OF PAGES

16. PRICE CODE

17. SECURITY CLASSIFICATION
OF REPORT

unclassified

18. SECURITY CLASSIFICATION
OF THIS PAGE

unclassified

19. SECURITY CLASSIFICATION
OF ABSTRACT

unclassified

20. LIMITATION OF ABSTRACT

unlimited

FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the U.S. Army.

X Where copyrighted material is quoted, permission has been obtained to use such material.

X Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

X Citations of commercial organizations and trade names in this report do not constitute an official Department of the Army endorsement or approval of the products or services of these organizations.

X In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and Use of Laboratory Animals of the Institute of Laboratory Animal Resources, National Research Council (NIH Publication No. 86-23, Revised 1985.)

X For the protection of human subjects, the investigator(s) have adhered to policies of applicable Federal Law 45-CFR-46.

DA Bello 9/10/83
PI Signature Date

Accession For	
NTIS	CRA&I ☒
DTIC	LAB ☐
Unannounced	☐
Justification	
By	
Distribution /	
Availability Codes	
Dist	Availability / or Special
A-1	

EXECUTIVE SUMMARY

Trinitrotoluene (TNT), unlike hexahydro-1,3,5-trinitro-1,3,5-triazine (RDX), undergoes rapid conversion or metabolism to polar forms following accumulation by terrestrial plants. This results in some uncertainty as to the potential food chain risk that may be imposed by TNT when it is released to the environment. A prior study at the same laboratory suggested the presence of potential environmentally important polar TNT and TNT-metabolite conjugates following root accumulation of trinitrotoluene by plants. To begin to address this question, a preliminary study was conducted to clarify and elucidate the nature of these polar metabolites present in plant tissues and xylem exudates.

The study focused on the nature of the proposed conjugates of TNT-residues contained in xylem exudates of bush bean plants, which represent the transport form(s) for TNT residues between roots and shoot tissues. Use of enzymatic methods failed to demonstrate that the conjugates were carbohydrate based. However, acid hydrolysis of TNT-residue-containing components isolated from xylem fluids indicated that the conjugates may be protein based. Of the five polar conjugates isolated from exudates, the presence of aminodinitrotoluene isomers and one unknown TNT residue (more polar than the aminodinitrotoluene isomers) was demonstrated. Determining the importance of these findings awaits further elucidation of the role of protein and possibly DNA adduction.

CONTENTS

REPORT DOCUMENT PAGE	iii
FOREWORD	v
EXECUTIVE SUMMARY	vii
1.0 INTRODUCTION	1
1.1 REVIEW OF RELATED LITERATURE	2
1.1.1 Toxicity of TNT	2
1.1.2 Chemistry and Analytic Methods	2
1.1.3 Plant Uptake and Metabolism	2
1.2 STUDY OBJECTIVES	6
1.3 TECHNICAL APPROACH	6
2.0 MATERIALS AND METHODS	7
2.1 PURITY AND ANALYSIS OF TNT SOURCES	7
2.1.1 Chemicals and Standards	7
2.1.2 Chemical Analysis of ¹⁴ C-TNT by HPLC	7
2.2 PLANT CULTIVATION AND SAMPLING	8
2.2.1 Plant Selection and Cultivation	8
2.2.2 Xylem Exudate Collection	9
2.3 CHEMICAL FRACTIONATION AND ANALYSES	9
2.3.1 Enzymatic Digestion for Conjugates	9
2.3.2 Residue Analysis	10
2.3.3 Radioanalyses	11
3.0 RESULTS AND DISCUSSION	12
3.1 CHARACTERIZATION OF PLANT METABOLITES OF TNT	12
3.1.1 Enzymatic Hydrolysis of Xylem Exudate	12
3.1.2 Radiochromatographic Studies of Bush Bean Exudate	13
4.0 SUMMARY AND CONCLUSIONS	19
5.0 LITERATURE CITED	20
DISTRIBUTION	22

FIGURES

1.1.	Radiochromatogram of Bush Bean Exudate	11
1.2.	Radiochromatogram of Acid-Hydrolyzed Bush Bean Exudate	12
2.1.	Radiochromatogram of ¹⁴ C-TNT Illustrating the High Degree of Purity	13
2.2.	Representative Standard Curve for TNT	14
3.1.	Representative HPLC Chromatogram with UV detection of Stem Exudate from a Bush Bean Plant Maintained in TNT-Amended Hydroponic Culture	20
3.2.	Radiochromatogram of Bush Bean Stem Exudate from a Plant Maintained in a TNT-Amended Hydroponic Culture	21
3.3.	Radiochromatogram of Acid-Hydrolyzed Peak 2.....	22
3.4.	Radiochromatogram of Acid-Hydrolyzed Peak 3.....	23
3.5.	Radiochromatogram of Acid-Hydrolyzed Peak 4.....	23
3.6.	Radiochromatogram of Acid-Hydrolyzed Peak 5.....	24

TABLES

2.1.	Buffer Compositions Utilized for the Enzymatic Hydrolyzes.....	16
3.1.	Percentage of Diethyl Ether-Extractable Radiolabel After Enzymatic Hydrolysis of Bush Bean Exudate Collected from a Plant Exposed Hydroponically to TNT-Amended Hydroponic Solution	19

1.0 INTRODUCTION

Munitions materials currently used as propellants or as explosive charges include trinitrotoluene (TNT), hexahydro-1,3,5-trinitro-1,3,5-triazine (RDX), octahydro-1,3,5,7-tetranitro-1,3,5,7-tetrazocine (HMX), and 2,4,6-trinitrophenylmethylnitramine (tetryl). These materials and their decomposition or combustion products may enter the environment as a result of both production and manufacturing activities, and field usage and disposal (Small and Rosenblatt 1974; Kitchens et al. 1978; Spanggord et al. 1983; Ryon et al. 1984).

The presence of these specific munitions-related components in the environment is not, however, indicative of the existence or the severity of an environmental impact. Persistence, and therefore accumulation of the contaminant, can be further influenced by physical processes such as its relative chemical stability and/or the presence of biologically mediated degradative processes.

These physical processes may be dependent on the sorptive processes of the soil and the contaminant's relative water solubility. Organic contaminants, such as the munitions material, may undergo chemical partitioning as they enter terrestrial and aquatic environments, thus affecting their short-term accumulation and mobility.

Biotic processes have been clearly indicated as being effective in the degradation and detoxification of a variety of organic xenobiotic compound classes, and they may operate in a similar fashion with munitions. With this in mind, biotic processes important for assessing the relative long-term behavior of munitions-related material released to the environment have been the subject of prior reports. Important factors include 1) the extent to which soil microbes can degrade and/or modify the contaminant; 2) the extent to which the parent compounds and their major decomposition products are accumulated by food-chain plants; and 3) the extent to which plant-accumulated contaminants are metabolized and degraded. The third of these is the focus of this preliminary study to elucidate the chemical fate of TNT residues once they are accumulated by terrestrial plants.

1.1 REVIEW OF RELATED LITERATURE

1.1.1 Toxicity of TNT

There are several recorded instances of human toxicity associated with TNT exposure. Documented cases over the last 60 years report instances of liver damage and anemia among chronically exposed munitions workers (Sax and Lewis 1989). In the environment, TNT and its derivatives have been reported to inhibit growth of freshwater algae at 2 to 15 $\mu\text{g/g}$ (Smock et al. 1976) and to negatively affect other green algae and oyster larvae; to inhibit the growth and metabolism of microorganisms generally (Klausmeier et al. 1973; Nay et al. 1974) as well as, among the vertebrates, fish (Nay et al. 1974), rats and mice (Lee et al. 1975). Furthermore, TNT was found to be toxic to lower plants (duckweed) at levels in excess of 1 $\mu\text{g/g}$ (Schott and Worthley 1974). With the exception of studies by Smock et al. (1976) and Schott and Worthley (1974), no chemical analyses were performed on the materials accumulated by the organisms. However, little if any information is available as to the the chemical fate of metabolized TNT residues in higher plants, let alone the subsequent toxicity of these metabolites to other trophic levels along the food chain.

1.1.2 Chemistry and Analytic Methods

Palazzo and Leggett (1986) were the first to address the analysis of explosive residues in plant tissues. Since their initial report, methods described by Cataldo and co-workers (Cataldo et al. 1989, 1990; Harvey et al. 1990, 1991) represent significant advances in the analytical methodology for determining explosives residues in plant tissues. Of particular interest is the use of methodologies that indicate a rapid conversion of TNT and related residues to more polar, although unidentified, chemical forms.

1.1.3 Plant Uptake and Metabolism

The literature contains few studies related to higher plant soil/plant fate and bioavailability of TNT. One such study was performed using hydroponically grown yellow nutsedge to assess the uptake, toxicity, and metabolic transformations of TNT (Palazzo and Leggett 1986). This study showed shoot and particularly root growth to be inhibited at 5 $\mu\text{g/g}$.

Although this point was not noted by Palazzo and Leggett, these toxicity symptoms are characteristic of dinitroaniline herbicide damage, which might be expected based on the chemical structures of TNT and the aminodinitrotoluenes. Chemical analysis showed that >90% of all tissue-extractable material was present as 2- and 4-ADNT, with only a small amount of TNT being recovered. Because these species were not observed in the nutrient solutions, it is assumed that they were metabolic detoxification products.

Detailed plant fate studies related to TNT have been conducted recently (Cataldo et al. 1989; Harvey et al. 1990). In bush beans grown in hydroponic solutions and exposed to ^{14}C -TNT, TNT-derived ^{14}C was found to be localized primarily in the roots, with only a small amount (<9% of the total radiolabel in the plant) having been transported to the shoot and leaves even after 7 days of exposure. When the leaf tissue from these plants, which contained approximately 8.6% of the total label in the plant, was homogenized with water and subsequently extracted with diethyl ether, only 4% of the label present in this tissue (less than 0.3% of total plant label) was found to partition into the ether layer; 57% of the leaf's label (less than 4.9% of total plant label) remained in the water layer, and the remaining 40% (3.4% of total plant label) was retained in the pellet. This indicated that the TNT that had been transported to the leaf from the root was metabolized to a very polar, and perhaps conjugated, form in this tissue.

In an attempt to free possible conjugates from the tissue and identify the water-extractable components, Cataldo et al. (1989) implemented acid hydrolysis immediately after homogenization of the tissue. When a subsample of the leaf tissue was hydrolyzed with 1 N HCl for 1 h prior to extraction with diethyl ether, 23% of the leaf tissue radiolabel (1.9% of total plant label) was ether-extractable, with 41% (3.5% of total plant label) remaining in the HCl layer. This experiment illustrated that approximately 19% of the TNT or TNT-derived compounds (~1.6% of total plant label) in the leaves of these plants had been metabolized to HCl hydrolyzable conjugates after only 7 days. Interestingly, acid hydrolysis did not increase the amount of radiolabel that was solubilized from the pellet. In both the acid hydrolysis and the water experimental conditions, slightly over 60% of the total radiolabel in the leaf tissue (>5% of total plant label) was consistently found to be solubilized in the aqueous or ether layers after 7 days of exposure.

When the stem and roots of the bush bean were analyzed in a similar manner, it was observed that almost twice as much of the total label in the roots was contained in the ether-extractable fractions (TNT and aminodinitrotoluene isomers) as in the HCl and water fractions (Cataldo et al. 1989). Stem tissue contained concentrations of TNT and the aminodinitrotoluene isomers that were intermediate between the large concentrations in the roots and the scant concentrations in the leaves. Furthermore, the proportion of the total label remaining in the pellet following extraction for these two tissues was consistently lower, at $28 \pm 10\%$. This distribution indicates progressive formation of polar compounds as TNT metabolites were transported from the root to the leaves.

In an attempt to identify the primary transport forms of TNT, additional experiments were conducted to examine the xylem exudate of bush beans grown in hydroponic solutions containing ^{14}C -TNT (Cataldo et al. 1989). Analysis of stem exudate has several inherent advantages over the analysis of stem tissue. Paramount among these is the fact that stem exudate contains only xylem transport products, or those metabolites being transported from root to shoot tissues, whereas stem tissue contains both transport products and storage forms. Additional advantages include the non-viscosity of the exudates, the absence of pigments, and the relatively simple chemical composition of exudates.

In that study, the roots of an intact 27-day-old bush bean were submerged in a 25-ppm TNT solution containing $10\ \mu\text{Ci}$ of radiolabeled TNT. The stem was severed and 1.3 mL of exudate collected over a period of 5 h. This sample was divided into two aliquots. The first of these were lyophilized and then reconstituted with $100\ \mu\text{l}$ of 50:50 MeOH:H₂O. A radiochromatogram of this sample is presented in Figure 1.1. The overwhelming majority of radiolabel eluted in the dead volume of the reverse-phase column, indicating the highly polar nature of the transport products. A very small amount of radiolabeled compound did not elute until after a retention time of 13 min. This corresponds to the retention time of the aminodinitrotoluene isomers.

The second 0.65-mL aliquot of exudate was acid-hydrolyzed with HCl at $100\ ^\circ\text{C}$ for 1 h prior to extraction with diethyl ether. The aqueous layer was made basic by the addition of




Figure 1.1. Radiochromatogram of Bush Bean Exudate (Cataldo et al. 1989)

NH₄OH and again extracted with diethyl ether. The radiochromatogram obtained on the acid-hydrolyzed material (Figure 1.2) represented a surprisingly simple chromatographic profile that contained only two major constituents. The highest peak (retention time of 13 min) was due to the aminodinitrotoluene isomers; the second-highest peak (retention time of 22 min) was due to a compound of unknown composition. This experiment clearly demonstrated that TNT is transported from the root to leaf tissue almost exclusively as conjugates of the aminodinitrotoluene isomers. Unfortunately, this information was not pursued during the duration of this contract and the characterization of the conjugates could not be completed.




Figure 1.2. Radiochromatogram of Acid-Hydrolyzed Bush Bean Exudate (Cataldo et al. 1989)

1.2 STUDY OBJECTIVES

The objectives of the study described in this report were to assess the chemical composition of the TNT-conjugated polar transport moiety (or moieties) in bush beans. Particular emphasis was placed on elucidating 1) the composition of the transport conjugate (carbohydrate or protein) and 2) the enumeration of the number of possible TNT-derived transport forms within the xylem of the plant.

1.3 TECHNICAL APPROACH

Ongoing research with TNT (Cataldo et al. 1989), RDX (Cataldo et al. 1990), and other organic xenobiotics for U.S. Army Medical Research and Development Command has clearly shown the need to understand the critical aspects controlling the behavior and chemical fate of environmental contaminants. This study continues to address the environmental fate and behavior of TNT in plants. The methods and approaches previously developed for TNT are being applied to understanding the composition of the principal TNT-metabolites that may have significant environmental effects.

2.0 MATERIALS AND METHODS

2.1 PURITY AND ANALYSIS OF TNT SOURCES

2.1.1 Chemicals and Standards

A series of standards, including 2,4,6-trinitrotoluene, 2-amino-4,6-dinitrotoluene, and 4-hydroxylamino-2,6-dinitrotoluene, were obtained from Chem Service (West Chester, Pennsylvania), the U.S. Army THAMA SARM Repository, and Oak Ridge National Laboratory, respectively. Purity was >98% for each compound.

2.1.2 Chemical Analysis of ^{14}C -TNT by HPLC

Uniformly ring-labeled ^{14}C -TNT was obtained from New England Nuclear (E.I. du Pont de Nemours & Co., Boston, Massachusetts). The purity of the TNT radiolabel to be used for these experiments was determined by radiochromatography. The radiochromatogram, presented in Figure 2.1, indicates a purity of 99.26 %. This purity was sufficient for our metabolic studies, as shown in the example of a standard concentration curve (Figure 2.2), and the radiolabel was used without further purification.




Figure 2.1. Radiochromatogram of ^{14}C -TNT Illustrating the High Degree of Purity




Figure 2.2. Representative Standard Curve for TNT

2.2 PLANT CULTIVATION AND SAMPLING

2.2.1 Plant Selection and Cultivation

The chemical fate of TNT-polar metabolites in plants was evaluated using bush beans (*Phaseolus vulgaris*, cv. Blue Lake Bush). All plants used were grown from seed. Plants were grown for 18 to 26 days in 600-mL beakers containing 500 mL of hydroponic nutrient solutions described previously (Garland et al. 1981). These contained 150 mg KCl, 120 mg MgSO_4 , 946 mg $\text{Ca}(\text{NO}_3)_2 \cdot 4 \text{H}_2\text{O}$, 68 mg KH_2PO_4 , 0.06 mg $\text{ZnSO}_4 \cdot 7 \text{H}_2\text{O}$, 0.69 mg H_3BO_3 , 0.017 mg $\text{CuCl}_2 \cdot 2 \text{H}_2\text{O}$, 0.024 mg $\text{Na}_2 \text{MoO}_4 \cdot 2 \text{H}_2\text{O}$, 0.022 mg $\text{MnCl}_2 \cdot 4 \text{H}_2\text{O}$, and 0.60 mg Fe^{+3} (as ethylenediaminetetraacetic acid ferric sodium salt) per liter. The solution pH was adjusted to 5.8 and brought to volume daily with fresh solution. Plants were maintained in a controlled-environment chamber with a 16/8-h light/dark cycle ($\sim 450 \mu\text{E}/\text{m}^2$ per sec, photosynthetically active radiation, at leaf surface or $850 \text{ W}/\text{m}^2$), a day/night temperature cycle of $28/22^\circ\text{C}$, and 50% RH. All solutions were bubbled with filtered air to provide aeration and mixing as described by Cataldo et al. (1978). The beakers were jacketed in opaque sheaths to protect the roots from light, and to minimize the photolysis of TNT.

2.2.2 Xylem Exudate Collection

In experiments aimed at elucidating the transportation forms of TNT, bush bean exudate was collected from plants grown in hydroponic solutions containing ^{14}C -TNT, using the method described by Fellows et al. (1989). For each collection, ten 29-day-old bush bean plants were placed overnight (14 h) in 500-mL beakers containing 400 mL of a TNT-amended nutrient solution. Each 400 mL of nutrient solution contained 20 μCi of radiolabeled TNT at a final concentration of 25 $\mu\text{g/g}$. The following morning, the plants were taken from the growth chambers within the first 30 min of the start of the light cycle (the time of highest exudation rate) and the stem was severed with a razor blade approximately 6 to 8 cm above the solution level. The first 100 μL of stem exudate was blotted away and the severed stem fitted with a latex tube connected to Teflon tubing that led to a scintillation vial. The vial was maintained at 4 °C during exudate collection (2 to 4 h). The exudates from all of the plants were pooled, yielding 4 to 5 mL of exudate having an activity of ~ 10 dpm/ μL , and analyzed as described below.

2.3 CHEMICAL FRACTIONATION AND ANALYSES

2.3.1 Enzymatic Digestion for Conjugates

The enzymes β -glucuronidase, β -glucosidase, protease, papain, and ficin were obtained from Sigma Chemical (St. Louis, Missouri). The buffers used for the incubations are listed in Table 2.1. Enzymes were incubated with bush bean exudates at 37 °C for 24 h; the mixtures were subsequently extracted with diethyl ether and the organic and aqueous phases separated. The diethyl ether layer was evaporated to dryness and the residue reconstituted with methanol. Volumes of the methanol solution and the aqueous layer were determined gravimetrically prior to removal of 100 μL of each solution for liquid scintillation counting.

Table 2.1. Buffer Compositions Used for the Enzymatic Hydrolysis

Enzyme Treatment	Buffer Used for Hydrolysis	Units Added (as defined by Sigma ^(a))
β -glucuronidase	0.008 M phosphate buffer at pH = 6.8	200
β -glucosidase	0.5 M acetate buffer at pH = 5.0	110
Protease	50 mM phosphate buffer at pH = 7.5	35
Papain	0.05 M sodium acetate pH 4.5	554
Ficin	2.0 M NaCl and 0.03 M cysteine pH = 5.0	41

(a) Sigma Chemical, St. Louis, Missouri.

2.3.2 Residue Analysis

The high-performance liquid chromatography (HPLC) system consisted of a Waters Model 600E pump and system controller. The methanol extract (20 μ L) of soil or plant tissue extracts was injected by a Waters WISP 710 automatic injector onto a Beckman Ultrasphere 5- μ m octadecyl silica column and the components were separated by a linear solvent program at a flow rate of 1.0 mL/min. The solvent system was water/acetonitrile, with a 20-min gradient from 40% to 100% acetonitrile. Components were detected by ultraviolet (UV) absorption at 254 nm (Waters Model 490E detector), with a detector sensitivity of 0.008 AUFS. Peak areas obtained from a Hewlett-Packard 3390 integrator were used for quantitative measurements.

Radiochromatographic detection was extensively used for unambiguous identification of transformation products derived from TNT. During selected chromatographic runs, the column eluate was collected in 0.5-mL increments for a total of 30 min. Each fraction was assayed for radioactivity by liquid scintillation spectrometry. Radiochromatograms were generated by plotting the disintegrations per min (dpm) in each successive aliquot as a function of retention time. Transformation products and/or metabolites identified in this manner were collected by repetitive HPLC runs to accumulate enough material for subsequent mass-spectral studies.

2.3.4 Radioanalyses

Samples for radiocarbon analysis were oxidized by total combustion in a Packard Model 306 oxidizer (Packard Instrument Co., Downers Grove, Illinois) to determine the amount of radiocarbon associated with each sample. Combusted samples and liquid samples were counted using a Beckman 9800 Liquid Scintillation Spectrometer (Beckman Instruments, Palo Alto, California) with appropriate quench correction.

3.0 RESULTS AND DISCUSSION

Results of xylem exudate studies with bush bean (Cataldo et al. 1989; Harvey et al. 1990) indicated that TNT is transported as polar conjugates to aerial portions of the plant, where it undergoes further metabolic alteration. Subsequent polar TNT metabolites are sequestered within the plant and are not appreciably transpired as volatile organics or $^{14}\text{CO}_2$. Chemical fractionation of plant tissues grown in TNT-containing hydroponic solution indicated that less than 12% of the incorporated radiolabel was due to TNT, 2-amino-4,6-dinitrotoluene, or 4-amino-2,6-dinitrotoluene. Most of the radiolabel was found in either a polar ether-extractable fraction or a more polar non-ether-extractable fraction. The preponderance of these previously unknown, polar ether-extractable TNT transformation products was a major finding of the previous studies. The objective of the present effort was to further elucidate the chemical nature of the polar residues.

3.1 CHARACTERIZATION OF PLANT METABOLITES OF TNT

In view of the chemical complexity of TNT polar metabolites in plants, exploratory experiments were initiated to optimize the chances of identifying potential classes of conjugates. These experiments involved analysis and characterization of polar xylem transport forms.

3.1.1 Enzymatic Hydrolysis of Xylem Exudate

The xylem exudate of hydroponically grown bush beans exposed to ^{14}C -TNT was subjected to enzymatic hydrolysis followed by solvent extraction and HPLC analysis of the organic phase for the aminodinitrotoluene isomers. Several enzymes were used to clarify the nature of the conjugates; these included β -glucuronidase, β -glucosidase, protease, papain, and ficin.

If a particular enzyme is active in cleaving polar conjugates from the TNT metabolites, the resulting non-polar radiolabel would be expected to partition into the diethyl ether layer on extraction. Results of the enzymatic experiments are provided in Table 3.1. In general, the enzymes failed to effectively cleave the conjugates recovered from xylem exudates. In the

Table 3.1. Percentage of Diethyl Ether-Extractable Radiolabel After Enzymatic Hydrolysis of Bush Bean Exudate Collected from a Plant Exposed Hydroponically to TNT-Amended Hydroponic Solution

Enzyme Treatment	Aqueous Layer	Diethyl Ether Layer	Mass Balance
β -glucuronidase	72.9	12.9	85.8
β -glucosidase #1	47.2	31.8	79.0
β -glucosidase #2	55.9	27.4	83.3
Protease	76.6	8.6	85.2
Papain	65.2	22.9	88.1
Ficin	66.1	17.8	83.9
0.5 M Acetate Buffer (pH = 5.0)	60.3	30.4	90.7

duplicate analyses performed with the β -glucosidase, 31.8 and 27.4% of the radiolabel was ether-extractable after hydrolysis. Incubation of bush bean exudate in acetate buffer that did not contain enzyme gave 30.4% ether-extractable radiolabel. It is likely that this is due to the formation of extractable ion pairs formed with the acetate buffer. As can be seen from Table 3.1, enzyme treatments resulted in very little partitioning of the radiolabel into the diethyl ether layer. These results indicate that cleavage of the polar TNT conjugates did not occur to an appreciable degree with the enzymes investigated. Thus the conjugates are likely neither carbohydrate nor protein based.

3.1.2 Radiochromatographic Studies of Bush Bean Exudate

Further investigation focused on radiochromatographic examination of the bush bean stem exudate. For this experiment, the exudate used came from bush bean plants exposed to high-specific-activity hydroponic solutions. Hydroponic solutions contained 25 μ g/g TNT and a total of 100 μ Ci 14 C-TNT per 500-ml beaker. Plants were placed in these solutions overnight

and exudate collections made 12 h later. Exudate used for these experiments was collected during the first 2 h following decapitation.

To characterize the polar conjugates contained within the bush bean stem exudate, gradient reversed-phase chromatography was employed with a weaker solvent system than was previously used (Cataldo et al. 1989). Conditions commonly employed by our laboratory for profiling TNT and the aminodinitrotoluene isomers start with a mobile phase composition of 40% acetonitrile. To profile the polar TNT-conjugates, the initial mobile phase composition was changed to 100% water. The gradient ramp was from pure water to 50% acetonitrile over 25 min. At this point, the gradient was continued at a faster rate to 100% acetonitrile in an additional 15 min. Direct injection of the bush bean exudate under these conditions resulted in the chromatogram shown in Figure 3.1. A radiochromatogram corresponding to the UV trace is shown in Figure 3.2. The principal peaks that contain radioactivity are numbered 1 through 5 in both these figures. This experiment clearly demonstrated that TNT is transported from the roots to shoots as a number of discrete polar metabolites.




Figure 3.1. Representative HPLC Chromatogram with UV detection (254 nm) of Stem Exudate from a Bush Bean Plant Maintained in TNT-Amended Hydroponic Culture




Figure 3.2. Radiochromatogram of Bush Bean Stem Exudate from a Plant Maintained in a TNT-Amended Hydroponic Culture

As previous experiments demonstrated (Cataldo et al. 1989), acid hydrolysis of the polar conjugates results in the formation of aminodinitrotoluene isomers and an unknown TNT metabolite that elutes with a retention time of 22 min. Next, the column eluate was pooled for each individual peak shown in Figures 3.1 and 3.2. The acetonitrile solvent was then removed by a gentle stream of nitrogen. Acid hydrolysis was accomplished by addition of 5.0 mL of 1 M HCl followed by heating at 100 °C for 1 h. After cooling, the hydrolysis mixture was extracted with 5.0 ml of diethyl ether. The aqueous layer was made basic and again extracted with diethyl ether. The ether layers were pooled and evaporated to dryness, and the residue was dissolved in methanol. A 20- μ L aliquot of this solution was counted by liquid scintillation spectrometry; the remainder of the solution was analyzed by radiochromatography.

Radiochromatograms of the acid-hydrolyzed peaks are shown in Figures 3.3 through 3.6 for Peaks 2 through 5, respectively. The quantity of radiolabel collected for peak 1 was not sufficient for radiochromatographic characterization.

The radiochromatograms presented in Figures 3.3 and 3.4 show hydrolysis products from peaks 2 and 3 (Figure 3.2) and indicate the presence of aminodinitrotoluene isomers. The amount of radiolabel collected for peak 4 was only sufficient to give an indication of the components liberated by acid hydrolysis. The radiochromatogram of acid-hydrolyzed peak 4 is shown in Figure 3.5. This radiochromatogram suggests that the acid-hydrolyzed peak 4 contains both the aminodinitrotoluene isomers and an unknown TNT metabolite that is more polar than the aminodinitrotoluene isomers (retention time of 8.0 min). Finally, acid hydrolysis of peak 5 yielded the same unknown metabolite, as well as lesser amounts of the aminodinitrotoluene isomers (Figure 3.6). This unknown metabolite is not a diaminonitrotoluene isomer as indicated by the fact that injection of a 2,6-diamino-4-nitrotoluene standard had a retention time of 4.36 min when analyzed under the same chromatographic conditions.




Figure 3.3. Radiochromatogram of Acid-Hydrolyzed Peak 2




Figure 3.4. Radiochromatogram of Acid-Hydrolyzed Peak 3




Figure 3.5. Radiochromatogram of Acid-Hydrolyzed Peak 4




Figure 3.6. Radiochromatogram of Acid-Hydrolyzed Peak 5

A quantity of the unknown metabolite that elutes at 22 min was collected from acid-hydrolyzed bush bean exudate and has been archived for further chemical characterization.

4.0 SUMMARY AND CONCLUSIONS

The results of this preliminary study failed to elucidate the nature of the conjugates associated with the TNT-residue transport forms in xylem exudates. However, results clearly show that

- 1) the majority of the TNT and its residues that are accumulated by plants are polar, and not simple transformation products,
- 2) the transport and likely tissue storage forms are conjugated in some consistent fashion,
- 3) the TNT residues can be released from the supposed conjugates by acid hydrolysis to yield one known metabolite (aminodinitrotoluene isomers) and one unknown TNT metabolite (retention time of 8.0 min) that is more polar than the aminodinitrotoluene isomers, and
- 4) the TNT-residue conjugates are likely not simple carbohydrate or related conjugates.

The results, while not definitive, do suggest that the polar conjugates may be small peptide-based molecules, since they are readily released by acid hydrolysis. This suggests that their ultimate chemical fate and potential impacts in plants and other trophic levels of the food chain may reside in their adduction with proteins and possibly DNA. If so, their potential to adversely impact the environment and humans requires further study.

5.0 LITERATURE CITED

- Cataldo, D. A., T. R. Garland, and R. E. Wildung. 1978. "Nickel in Plants 1. Uptake Kinetics Using Intact Soybean Seedlings." Plant Physiol. 62:563-565.
- Cataldo, D. A., S. D. Harvey, R. J. Fellows, R. M. Bean, and B. D. McVeety. 1989. An Evaluation of the Environmental Fate and Behavior of TNT in Soil and Plant Systems. AD-A223546, PNL-7370, Pacific Northwest Laboratory, Richland, Washington.
- Cataldo, D. A., S. D., Harvey, and R. J. Fellows. 1990. An Evaluation of the Environmental Fate and Behavior of RDX. PNL-7529, Pacific Northwest Laboratory, Richland, Washington.
- Fellows, R. J., D. A. Cataldo, and R. M. Bean. 1989. "Uptake and Metabolism of Exogenously Applied Indole by Soybeans Grown in Solution Culture and Soils." J. Agric. and Food Chem. 37:1444-1454.
- Garland, T. R., D. A. Cataldo, and R. E. Wildung. 1981. "Absorption, Transport, and Chemical Fate of Plutonium in Soybean Plants." J. Agric. Food Chem. 29:915-920.
- Harvey, S. D., R. J. Fellows, D. A. Cataldo, and R. M. Bean. 1990. "Analysis of 2,4,6-Trinitrotoluene (TNT) and Its Transformation Product in Soils and Plant Tissues by High-Performance Liquid Chromatography." J. Chromatog. 518:361-374.
- Harvey, S. D., R. J. Fellows, D. A. Cataldo, and R. M. Bean. 1991. "Fate of the Explosive Hexahydro-1,3,5-Trinitro-1,3,5 triazine (RDX) in Soil and Bioaccumulation in Bush Bean Hydroponic Plants." Environ. Toxicol. and Chem. 10:845-855.
- Kitchens, J. F., S. G. Brownlee, W. E. Harward, R. G. Hyde, W. E. Jones, D. A. Price, R. S. Wentzel, and R. S. Valentine. 1978. Preliminary Problem Definition Study on Munitions-Related Chemicals. Final Report. DAMD17-77-C-7057, Atlantic Research Corp.
- Klausmeier, R. E., J. L. Osmon, and D. R. Wells. 1973. "The Effect of Trinitrotoluene on Microorganisms." Dev. Ind. Microbiol. 15:309-317.
- Lee, C. C., J. V. Dilley, J. R. Hodgson, D. O. Helton, W. J. Weigand, D. N. Roberts, B. S. Anderson, L. M. Halfpapp, and L. D. Kurtz. 1975. Mammalian Toxicity of Munition Compounds. Phase I. Acute Oral Toxicity. Primary Skin and Eye Irritation. Dermal Sensitization, and Disposition and Metabolism. No. 3900-B. Contract No. DAMD17-74-C-4073, Midwest Research Institute, Kansas City, Missouri.
- Nay, M. W., C. W. Randall, and P. H. King. 1974. "Biological Treatability of Trinitrotoluene Manufacturing Wastewater." J. Water Pollut. Control Fed. 46:485-497.

Palazzo, A. J. and D. C. Leggett. 1986. "Effect and Disposition of TNT in a Terrestrial Plant." J. Environ. Qual. 15:49-52.

Ryon, M. G., B. C. Pal, S. S. Talamage, and R. H. Ross. 1984. Database Assessment of the Health and Environmental Effects of Munition Production Waste Products. Final Report ORNL-6018, Oak Ridge National Laboratory, Oak Ridge, Tennessee.

Sax, N. I., and R. J. Lewis. 1989. Dangerous Properties of Industrial Chemicals. 7th ed. Reinhold Publishing, New York.

Schott, C. D., and E. G. Worthley. 1974. The Toxicity of TNT and Related Wastes to an Aquatic Flowering Plant: *Lemna perpusilla* Torr. Edgewood Arsenal Technical Rep. EB-TR-74016, Edgewood Arsenal, Aberdeen Proving Ground, Maryland.

Small, M. J., and D. H. Rosenblatt. 1974. Munitions Production Products of Potential Concern - Phase II. Technical Report 7404, AD919031, U.S. Army Medical Bioengineering Laboratory. Fort Detrick, Frederick, Maryland.

Smock, L. A., D. L. Stoneburner, and J. R. Clark. 1976. "The Toxic Effects of TNT and its Primary Degradation Products on Two Species of Algae and Fathead Minnow." Water Res. 10:534-543.

Spanggord, R. J., W. R. Mabey, T. Mill, T.-W. Chou, J. H. Smith, S. Lee, and D. Roberts. 1983. Environmental Fate Studies on Certain Munitions Wastewater Constituents. Phase IV-Lagoon Model Studies. Final Report Phase IV. DAMD17-78-C-8081, SRI International, Menlo Park, California.

DISTRIBUTION

OFFSITE

- 2 DOE - Office of Scientific and Technical Information
- 30 Dr. Wayne Mitchell
U.S. Army Biomedical Research and Development Laboratory
ATTN: SGRD-UBG-E
Building 568
Fort Detrick
Frederick, Maryland 21701-5010
- 3 Commander
U. S. Army Medical Research and Development Command
ATTN: SGRD-RMI-S
Fort Detrick
Frederick, Maryland 21702-5012

ONSITE

- 2 DOE Richland Operations Office

R. B. Goranson

Pacific Northwest Laboratory
- 36 D. A. Cataldo (25)
R. J. Fellows
L. K. Grove
S. D. Harvey
L. E. Rogers
Publishing Coordination
Technical Report Files (5)

Routing
- 1 R. M. Ecker
M. J. Graham
P. M. Irving
R. L. Skaggs
C.S. Sloan
P. C. Hays (last)

Distr.1