

AD-A274 064

STATION PAGE

Form Approved
OMB No. 0704-0188



ted to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE 1 February, 1992		3. REPORT TYPE AND DATES COVERED Final Report 2/1/88-4/1/91	
4. TITLE AND SUBTITLE Molecular Regulation of Light Adaptation and Nitrogen Assimilation in Marine Diatoms				5. FUNDING NUMBERS N00014-88-K-0258	
6. AUTHOR(S) Dr. G. Jason Smith / Dr. Hewson Swift					
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Department of Molecular Genetics and Cell Biology EBC 101 The University of Chicago 1102 E. 57 th Street Chicago, IL 60637				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Office of Naval Research 800 N. Quincy St. Arlington, VA 22217-5000				10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES					
12a. DISTRIBUTION/AVAILABILITY STATEMENT Distribution Unlimited				12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) The objective of this project was to investigate the molecular and cellular processes that regulate the rate of adaptation of photosynthetic light harvesting and nitrogen assimilation capacity of marine diatoms to fluctuating light and nutrient environments. The initial tasks were to develop and verify nucleic acid and antibody probes for assessing gene presence, transcriptional activity and posttranslational modulation of the expression of nitrate reductase (NR), glutamine synthetase (GS) and the fucoxanthin-chlorophyll a-c pigment-protein (FCP) in diatoms. Given the distinct evolutionary history of diatoms we independently cloned segments of these genes using the polymerase chain reaction and developed antisera specific to the diatom forms of these enzymes. Sequence analysis indicates that NR and GS in diatoms exhibit limited homology with the forms occurring in land plants. Concurrent application of the NR nucleic acid probe and antisera indicated that the expression of NR is under strong transcriptional and post translational control when cells are exposed to realistic environmental shifts. NR and FCP were observed to be inversely regulated by shifts in light availability.					
14. SUBJECT TERMS gene expression, nitrate reductase, glutamine synthetase, fucoxanthin-chlorophyll a-c pigment-protein, diatoms				15. NUMBER OF PAGES 3	
				16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT UL		

DTIC
ELECTE
DEC 23 1993

93 12 21 1 32

93-30759

SPJ



Accession For	
NTIS	CRA&I
DTIC	TAB
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
A-1	

FINAL TECHNICAL REPORT

GRANT # N00014-88-K-0258

R&T CODE: 4412040

PRINCIPAL INVESTIGATOR: Dr. Hewson Swift/ Dr. G. Jason Smith

INSTITUTION: The University of Chicago

GRANT TITLE: Molecular Regulation of Light Adaptation and Nitrogen Assimilation in Marine Diatoms

AWARD PERIOD: 1 February 1988 - 1 April 1991

OBJECTIVE: To investigate the molecular and cellular processes that regulate the rate of adaptation of photosynthetic light harvesting and nitrogen assimilation capacity of marine diatoms to fluctuating light and nutrient environments.

ACCOMPLISHMENTS: A major goal of the research was to develop cDNA probes for the genes encoding the rate-limiting enzymes for nitrate and ammonium assimilation, nitrate reductase (NR) and glutamine synthetase (GS), respectively as well as a probe for the diatom light harvesting complex a, fucoxanthin chlorophyll a-c pigment-protein (FCP). Prior attempts to isolate genes encoding NR & GS from diatoms by screening diatom Southern or northern blots with heterologous (ie. higher plant) probes proved unsuccessful; western analysis indicated that heterologous antisera would also not be suitable for screening diatom expression cDNA libraries or for the study of the regulation of these genes by environmental factors. Therefore, we initiated a cloning strategy based on polymerase chain reaction (PCR) amplification of conserved domains of nucleic acid sequences encoding these proteins. Amino acid sequence comparisons of higher plant, fungal and bacteria forms of NR and GS were performed and revealed sets of 6 - 10 conserved amino acid residues which served as suitable primer sites for PCR. Degenerate oligonucleotides corresponding to all possible codon usages for these domains were synthesized and used in PCR amplification of genomic DNA and cDNA from higher plant, green algal, diatom and other chromophytic macroalgal species. Products of the size expected for the regions bounded by the primers were obtained and hybridized strongly to diatom Southern and northern blots. Sequence analysis of these products indicate that although the green algal "control" PCR products generally exhibit greater than 50% identity at the amino acid level to published plant sequences, the diatom and brown algal sequences exhibit less than 40% identity, suggesting that in chromophytes these proteins are significantly divergent from green algae and higher plants. Native gel electrophoresis was used to isolate pure FCP from diatoms and kelp. V8 protease treatment of these proteins followed by amino acid microsequencing

was used to determine internal amino acid domains for designing PCR and hybridization oligonucleotides for FCP. The NR, GS, and FCP clones are currently being used as hybridization probes to examine the dynamics of the transcripts in response to physiologically realistic transients in light and nitrogen availability. The probes are also being used to isolate full length coding sequences for these genes. The NR probe has been used successfully to demonstrate that the accumulation of NR transcripts is light and nitrate dependent (Smith *et al.* 1992).

We are also interested in the post-transcriptional processes that may regulate the expression of NR, GS and FCP and towards this end have been developing antibody probes against these proteins. An anti-FCP serum, already available, has been used to verify that FCP accumulates under low light conditions, but that this accumulation is not dependent upon the nitrogen source used for growth. A heterologous antisera raised against squash NR was used to demonstrate that in diatoms, enhancement of NR activity in response to light and nitrate enrichment occurs independently of any net synthesis of NR protein, suggesting that light can directly activate existing enzyme (Smith *et al.* 1992). Due to the low cross-reactivity of this antisera with diatom NR we undertook the purification, characterization and antibody production of diatom NR. Antisera for Skeletonema costatum NR has been obtained and affinity purified; western blot analysis indicates that this antisera is specific to the diatom form of NR. The purified NR will also be used to further examine the nature of the light activation response. A protocol for purification of GS from chromophytic algae based on ion exchange chromatography has been developed and purification for antibody production is underway. In addition, one of our graduate students has developed a protocol for the isolation of high molecular weight and restrictable DNA from Prochloron, the prochlorophyte symbiont of didemnid ascidians (Urbach *et al.* 1992). This development will enable us to examine the genomic organization of GS and NR in this evolutionary divergent algal group.

SIGNIFICANCE: The development of these molecular and immunological probes specific for marine diatoms will enable us to examine the influence of light and nitrogen availability in modulating gene expression. These studies will determine the relationships between molecular level changes and the physiological expression of these changes as they relate to adaptation to variable light and nitrogen regimes. Comparisons of light and nitrogen dependent gene expression between diatoms and higher plants may also provide us with insights as to the evolution of gene expression control circuits in photosynthetic organisms. Additionally, our development of "universal" PCR primers for NR, GS, and FCP will enable rapid characterization of these genes in a variety of marine algae.

FUTURE WORK: We plan to complete characterizing the NR, GS, and FCP genes in diatoms with the immediate goal of isolating full length genomic sequences. These clones will be used to map and sequence the putative

promoter elements associated with the 5'- regions of these genes and sequence searches will be conducted to determine if these regions are homologous to light responsive promoter elements characterized in higher plants. Application of nucleic acid probes and antibodies in conjunction with measurement of the associated physiological processes in diatom cells subjected to environmentally realistic transitions in their light and nitrogen environment will enable us to define the cellular levels of control on the expression of that physiological process. Measurement of transients in other cellular metabolite pools (amino acids, adenine nucleotides, photosynthetic pigments) will be conducted to determine whether these components play a role in the transduction of environmental signals to gene expression events.

PUBLICATONS:

- Smith, G.J. and Alberte, R.S. 1991. Characterization of photosystem I-associated polypeptides from the chlorophyll *b*-rich alga Tetraselmis spp. (Pleurostrophyceae) and other chlorophyte algae. J. Phycol. 27:92-101.
- Swift, H. and Robertson, D.L. 1991. Structural aspects of the prochloron-tunicate symbiosis. Symbiosis 10: 95-113.
- Gao, Y., Smith, G.J. and Alberte, R.S. 1992. Light regulation of nitrate reductase in Ulva fenestrata (Chlorophyceae): I. The influence of light regimes on nitrate reductase activity. Mar. Biol. in press.
- Smith, G.J., Zimmerman, R.C. and Alberte R.S. 1992. Molecular and physiological responses of diatoms to variable levels of irradiance and nitrogen availability. I. Growth of Skeletonema costatum in simulated upwelling conditions. Limnol. Oceanogr. in press.
- Urbach, E., Robertson, D.L. and Chisholm, S.W. 1992. Multiple evolutionary origins of prochlorophytes within the cyanobacterial radiation. Nature 355:267-270.

Distribution List for Final Reports

Attach a copy of the REPORT DOCUMENTATION PAGE (DO FORM 1473) to your final report as the first page and mail two copies (including the postcard labelled DTIC FORM 50) to:

Defense Technical Information Center
Building 5, Cameron Station
Alexandria, VA 22314

This allows other investigators to obtain copies of your report directly from DTIC. DTIC will fill out the postcard DTIC ACCESSION NOTICE (DTIC FORM 50) and return it to you with their number for your report. When you refer people to DTIC to get a copy of your report, give this number to expedite the request.

Mail one copy to each of the following and attach this very page to the back of your report - otherwise the folks below will think they have mistakenly received a copy meant for the Molecular Biology Program):

- | | |
|---|--|
| (a) Dr. Michael Marron
ONR Code 1141
Molecular Biology Program
800 N. Quincy Street
Arlington, VA 22217-5000 | (e) Director
Chemical and Biological Sci Div
Army Research Office
P. O. Box 12211
Research Triangle Park, NC 27709 |
| (b) Administrative Grants Officer
ONR Resident Representative
(address varies - see copy of your
grant/contract) | (f) Life Sciences Directorate
Air Force Office of Scientific Res
Bolling Air Force Base
Washington, DC 20332 |
| (c) Director,
Applied Research Directorate
ONR Code 12
800 N. Quincy Street
Arlington, VA 22217-5000 | (g) Director
Naval Research Laboratory
Technical Information Div
Code 2627
Washington, DC 20375 |
| (d) Director
Office of Naval Technology
Code 22
800 N. Quincy Street
Arlington, VA 22217-5000 | |