


NASA Contractor Report 191560

AD-A274 823

ICASE Report No. 93-82



ICASE



ON THE GIBBS PHENOMENON III: RECOVERING EXPONENTIAL ACCURACY IN A SUB-INTERVAL FROM A SPECTRAL PARTIAL SUM OF A PIECEWISE ANALYTIC FUNCTION

David Gottlieb

Chi-Wang Shu

BEST AVAILABLE COPY

DTIC
ELECTE
JAN 19 1994
S E D

NASA Contract No. NAS1-19480
November 1993

Institute for Computer Applications in Science and Engineering
NASA Langley Research Center
Hampton, Virginia 23681-0001

Operated by the Universities Space Research Association



Approved for public release
Distribution unlimited

94-01651



National Aeronautics and
Space Administration
Langley Research Center
Hampton, Virginia 23681-0001

94 1 14 04

**ON THE GIBBS PHENOMENON III:
RECOVERING EXPONENTIAL ACCURACY IN A SUB-INTERVAL
FROM A SPECTRAL PARTIAL SUM OF A PIECEWISE ANALYTIC FUNCTION**

David Gottlieb and Chi-Wang Shu¹
Division of Applied Mathematics
Brown University
Providence, RI 02912

ABSTRACT

We continue the investigation of overcoming Gibbs phenomenon, i.e., obtaining exponential accuracy at all points including at the discontinuities themselves, from the knowledge of a spectral partial sum of a discontinuous but piecewise analytic function.

We show that if we are given the first N expansion coefficients of an L_2 function $f(x)$ in terms of either the trigonometrical polynomials or the Chebyshev or Legendre polynomials, we can construct an exponentially convergent approximation to the point values of $f(x)$ in any sub-interval in which it is analytic.

DTIC QUALITY INSPECTED B

Accession For	
NTIS CRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification _____	
By _____	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
A-1	

¹This research was supported in part by the National Aeronautics and Space Administration under NASA Contract No. NAS1-19480 while the authors were in residence at the Institute for Computer Applications in Science and Engineering (ICASE), NASA Langley Research Center, Hampton, VA 23681. Research also supported by AFOSR grant 93-0090, ARO grant DAAL03-91-G-0123, DARPA grant N00014-91-J-4016, NSF grant DMS-9211820, and NASA grant NAG1-1145.

1 Introduction

In this paper we continue our investigation of overcoming Gibbs phenomenon, i.e., recovering pointwise exponential accuracy at all points including at the discontinuities themselves, from the knowledge of a spectral partial sum of a discontinuous but piecewise analytic function, which we started in [5] and [6].

Spectral approximations based upon trigonometric polynomials (Fourier, for periodic problems) or polynomials (Chebyshev or Legendre, for non-periodic problems) are exponentially accurate for analytic functions [4], [3]. However, for discontinuous but piecewise analytic functions, the spectral partial sum approximates the function poorly throughout the domain. Away from the discontinuity only first order accuracy is achieved. Near the discontinuity there are $O(1)$ oscillations which do not decrease with N , the number of terms retained in the spectral sum. This is known as Gibbs phenomenon.

In [5] and [6] we treated a representative problem :

- Let $f(x)$ be an *analytic but nonperiodic function*. Suppose that we are given its first $-N \leq k \leq N$ Fourier coefficients. Can one obtain exponentially convergent pointwise approximations in the maximum norm?

To solve this problem we used the Gegenbauer polynomials $C_n^\lambda(x)$, which are orthogonal in $[-1, 1]$ with the weight function $(1 - x^2)^{\lambda - \frac{1}{2}}$.

The procedure of overcoming the Gibbs phenomenon consists of two steps:

1. Given the Fourier partial sum of the first N terms, we first recover the first $m \sim N$ Gegenbauer expansion coefficients with exponential accuracy. This can be achieved for any L_2 function, as long as we choose λ in the weight function of Gegenbauer polynomials to be proportional to N . The error incurred at this stage is called the truncation error.
2. The next step is to prove, *for an analytic function*, the exponential convergence of its Gegenbauer expansion, when the parameter λ in the weight function is

proportional to the number of terms retained in the expansion. The error at this stage is labelled the regularization error.

Thus we have shown how to overcome the Gibbs phenomenon for an analytic but non-periodic function. By a simple shift this procedure covers the case of any analytic function with one discontinuity.

In this paper we treat a broader class of problems. Here we assume that $f(x)$ is an L_2 function on $[-1, 1]$ and analytic in a subinterval $[a, b] \subset [-1, 1]$. We assume that the spectral partial sum (based on either the Fourier or Chebyshev or Legendre expansion) of a function over the full interval $[-1, 1]$ is known, and try to recover exponentially accurate point values over a subinterval $[a, b]$.

We will follow the same path as in [5]. Basically we will show that the first N Fourier (or Chebyshev or Legendre) expansion coefficients contain enough information such that a rapidly converging Gegenbauer expansion in the subinterval $[a, b]$ can be constructed. As before, we will separate the analysis of the error into two parts: truncation error and regularization error. Truncation error measures the difference between the exact Gegenbauer coefficients and those obtained by using the spectral partial sum. These will be investigated in Section 3 for the Fourier case and in Section 4 for the Chebyshev and Legendre cases. Regularization error then measures the difference between the Gegenbauer expansion using the first few Gegenbauer coefficients and the function itself in a sub-interval $[a, b]$, in which the function is assumed analytic. The estimation of the regularization error is in Section 5. In Section 2 we will provide some useful properties of Gegenbauer polynomials to be used later. Section 6 contains a summary theorem and some remarks.

Throughout this paper, we will use A to denote a generic constant or at most a polynomial in the growing parameters. It may not be the same at different locations.

The message of this paper is that the knowledge of the partial spectral sum of an L_2 function in $[-1, 1]$ furnishes enough information so that an *exponentially convergent approximation* can be constructed in any subinterval in which $f(x)$ is analytic.

In particular, if $f(x)$ is *piecewise analytic*, then an exponentially convergent approximation in the maximum norm can be recovered from its partial spectral expansion.

2 Preliminaries

This section is devoted to a collection of results about the Gegenbauer polynomials. Though the results are classical, they are not widely known and certainly not in our context. We rely heavily on the standardization in Bateman [2].

We start by defining the Gegenbauer polynomials $C_n^\lambda(x)$ in the following

Definition 2.1

The Gegenbauer polynomial $C_n^\lambda(x)$ is defined by

$$(1-x^2)^{\lambda-\frac{1}{2}} C_n^\lambda(x) = \frac{(-1)^n}{2^n n!} G(\lambda, n) \frac{d^n}{dx^n} [(1-x^2)^{n+\lambda-\frac{1}{2}}] \quad (2.1)$$

where $G(\lambda, n)$ is given by

$$G(\lambda, n) = \frac{\Gamma(\lambda + \frac{1}{2}) \Gamma(n + 2\lambda)}{\Gamma(2\lambda) \Gamma(n + \lambda + \frac{1}{2})} \quad (2.2)$$

□

Under this definition we have

$$C_n^\lambda(1) = \frac{\Gamma(n + 2\lambda)}{n! \Gamma(2\lambda)} \quad (2.3)$$

and

$$|C_n^\lambda(x)| \leq C_n^\lambda(1), \quad -1 \leq x \leq 1 \quad (2.4)$$

The Gegenbauer polynomials are orthogonal under their weight function, in fact we have

Lemma 2.2

The Gegenbauer polynomials satisfy the following orthogonality condition:

$$\int_{-1}^1 (1-x^2)^{\lambda-\frac{1}{2}} C_k^\lambda(x) C_n^\lambda(x) dx = \delta_{k,n} h_n^\lambda \quad (2.5)$$

where

$$h_n^\lambda = \pi^{\frac{1}{2}} C_n^\lambda(1) \frac{\Gamma(\lambda + \frac{1}{2})}{\Gamma(\lambda)(n + \lambda)} \quad (2.6)$$

□

We will need to use heavily the asymptotics of the Gegenbauer polynomials for large n and λ . For this we need

Lemma 2.3 (Stirling)

For any number x such that $x \geq 1$ we have

$$(2\pi)^{\frac{1}{2}} x^{x+\frac{1}{2}} e^{-x} \leq \Gamma(x+1) \leq (2\pi)^{\frac{1}{2}} x^{x+\frac{1}{2}} e^{-x} e^{\frac{1}{12x}} \quad (2.7)$$

□

Lemma 2.4

There exists a constant A independent of λ and n such that

$$A^{-1} \frac{\lambda^{\frac{1}{2}}}{(n+\lambda)} C_n^\lambda(1) \leq h_n^\lambda \leq A \frac{\lambda^{\frac{1}{2}}}{(n+\lambda)} C_n^\lambda(1) \quad (2.8)$$

The proof follows from (2.6) and Stirling's formula (2.7).

□

In the analysis we will extensively use the relationship between the Fourier functions $e^{in\pi x}$ and the Gegenbauer polynomials $C_n^\lambda(x)$. In particular, we will focus our attention on the *Legendre polynomials*

$$P_n(x) = C_n^{\frac{1}{2}}(x) \quad (2.9)$$

and the *Chebyshev polynomials*

$$T_n(x) = \frac{n}{2} C_n^0(x) \quad (2.10)$$

Our basic formula is taken from [2], page 213:

Lemma 2.5

$$\frac{1}{h_l^\lambda} \int_{-1}^1 (1-x^2)^{\lambda-\frac{1}{2}} e^{in\pi x} C_l^\lambda(x) dx = \Gamma(\lambda) \left(\frac{2}{\pi n}\right)^\lambda i^l (l+\lambda) J_{l+\lambda}(\pi n) \quad (2.11)$$

where $J_\nu(x)$ is the Bessel function.

□

Lemma 2.5 gives us the Gegenbauer expansion coefficients of the *Fourier functions*. However, what we really want are the *Fourier coefficients* of the Gegenbauer polynomials. We have those coefficients, luckily, for the Legendre and Chebyshev polynomials.

Lemma 2.6

Let a_k^N be the Fourier coefficients of the Legendre polynomial $P_N(x)$ defined in (2.9) and b_k^N be the Fourier coefficients of the Chebyshev polynomial $T_N(x)$ defined in (2.10), i.e.

$$P_N(x) = \sum_{k=-\infty}^{\infty} a_k^N e^{ik\pi x} \quad (2.12)$$

and

$$T_N(x) = \sum_{k=-\infty}^{\infty} b_k^N e^{ik\pi x} \quad (2.13)$$

Then

$$a_k^N = \frac{i^N}{\sqrt{2k}} J_{N+\frac{1}{2}}(\pi k) \quad (2.14)$$

and

$$b_k^N = \frac{(-i)^N \pi}{2} J_N(\pi k) \quad (2.15)$$

□

Equation (2.14) follows from (2.11) upon substituting $\lambda = \frac{1}{2}$ and using (2.6) and (2.3). Equation (2.15) can be found in [7], page 836.

□

Lemma 2.6 provides us with a better understanding of the terms “small scales” and “large scales”. The question is the following: it is clear that $e^{iN\pi x}$ is small scale for large N , intuitively $T_N(x)$ and $P_N(x)$ are also small scale functions, however the expansions (2.12) and (2.13) contain *all* the scales. So, in what sense are $T_N(x)$ and $P_N(x)$ small scales?

The answer is given in the following Lemma:

Lemma 2.7

Let a_k^N and b_k^N be the Fourier coefficients of the Legendre polynomial $P_N(x)$ and the Chebyshev polynomial $T_N(x)$ respectively, given in (2.12)-(2.15). Then

$$|a_k^N| \leq A \min \left(1, \left(\frac{c\pi k}{2N} \right)^N \right) \quad (2.16)$$

$$|b_k^N| \leq A \min \left(1, \left(\frac{c\pi k}{2N} \right)^N \right) \quad (2.17)$$

where A is a constant independent of k or N .

Proof:

We start with the following estimate for the Bessel function $J_n(nz)$ (see [1], page 362):

$$|J_n(nz)| \leq \left(\frac{ze^{\sqrt{1-z^2}}}{1 + \sqrt{1-z^2}} \right)^n, \quad 0 \leq z \leq 1 \quad (2.18)$$

From (2.18) we can deduce trivially that

$$|J_n(nz)| \leq \left(\frac{ez}{2} \right)^n \quad 0 \leq z \leq 1 \quad (2.19)$$

We also have the uniform bound on Bessel functions

$$|J_n(z)| \leq 1 \quad (2.20)$$

Using (2.19) and (2.20) in (2.14) and (2.15), with $z = \frac{\pi k}{N}$ and $n = N$, we obtain (2.16) and (2.17).

□

Lemma 2.7 provides the answer why $P_N(x)$ and $T_N(x)$ (for large N) can be considered as small scale functions. Basically only the terms that are greater than N appear in their Fourier expansions (2.12) and (2.13). The lower terms decay *exponentially* with N .

3 Truncation Error in a Sub-interval—Fourier Method

Consider an arbitrary L_2 function $f(x)$ defined in $[-1, 1]$. Suppose that the first $2N + 1$ Fourier coefficients of this function are given

$$\hat{f}(k) = \frac{1}{2} \int_{-1}^1 f(x) e^{-ik\pi x} dx \quad |k| \leq N \quad (3.1)$$

In [5] we showed that we can recover (with exponential accuracy) the first $m \sim N$ terms in the Gegenbauer expansion of $f(x)$ based on the interval $[-1, 1]$.

Here we are interested to find the Gegenbauer expansion of $f(x)$ based on a sub-interval $[a, b] \subset [-1, 1]$. This will allow us to handle multiple discontinuities and discontinuities of unknown locations. We will show that in this case too, we can get exponential accuracy. We start by introducing the local variable ξ :

Definition 3.1

The local variable ξ is defined by

$$x = x(\xi) = \epsilon\xi + \delta \quad (3.2)$$

where

$$\epsilon = \frac{b-a}{2}, \quad \delta = \frac{b+a}{2} \quad (3.3)$$

Thus when $a \leq x \leq b$, $-1 \leq \xi \leq 1$.

□

Denote now the Fourier partial sum by $f_N(x)$, namely

$$f_N(x) = \sum_{k=-N}^N \hat{f}(k) e^{ik\pi x} \quad (3.4)$$

As mentioned before, we are trying to recover the Gegenbauer expansion coefficients based on the sub-interval $[a, b]$, i.e. we would like to find the first $m \sim N$ coefficients $\hat{f}_\epsilon^\lambda(l)$ in the expansion

$$f(x(\xi)) = \sum_{l=0}^{\infty} \hat{f}_\epsilon^\lambda(l) C_l^\lambda(\xi) \quad (3.5)$$

where

$$\hat{f}_\epsilon^\lambda(l) = \frac{1}{h_l^\lambda} \int_{-1}^1 (1 - \xi^2)^{\lambda - \frac{1}{2}} f(x(\xi)) C_l^\lambda(\xi) d\xi \quad (3.6)$$

Of course we do not have $\hat{f}_\epsilon^\lambda(l)$ at our disposal but only an approximation based on the Fourier partial sum $f_N(x)$, thus we have

$$\hat{g}_\epsilon^\lambda(l) = \frac{1}{h_l^\lambda} \int_{-1}^1 (1 - \xi^2)^{\lambda - \frac{1}{2}} f_N(x(\xi)) C_l^\lambda(\xi) d\xi \quad (3.7)$$

The *Truncation Error* is the measure of deviation of the approximate Gegenbauer expansion from the true expansion:

$$TE(\lambda, m, N, \epsilon) = \max_{-1 \leq \xi \leq 1} \left| \sum_{l=0}^m (\hat{f}_\epsilon^\lambda(l) - \hat{g}_\epsilon^\lambda(l)) C_l^\lambda(\xi) \right| \quad (3.8)$$

We can show that the truncation error is exponentially small:

Theorem 3.2:

Let $f(x)$ be an L_2 function whose first $2N + 1$ Fourier coefficients $\hat{f}(k)$ defined in (3.1) are known. Let $\hat{g}_\epsilon^\lambda(l)$ be the Gegenbauer coefficients, based on the sub-interval $[a, b]$ expansion of the Fourier partial sum $f_N(x)$ defined in (3.7), and $\hat{f}_\epsilon^\lambda(l)$ be those of $f(x)$ defined in (3.6). Assume that

$$\lambda = \alpha \epsilon N \quad m = \beta \epsilon N \quad (3.9)$$

Where ϵ is the half length of the sub-interval $[a, b]$ defined in (3.3). Then the truncation error is bounded by

$$TE(\alpha \epsilon N, \beta \epsilon N, N, \epsilon) \leq A q^{\epsilon N}, \quad (3.10)$$

where

$$q = \frac{(\beta + 2\alpha)^{\beta + 2\alpha}}{(2\pi\epsilon)^\alpha \alpha^\alpha \beta^\beta} \quad (3.11)$$

and A grows at most quadratically with N . In particular, if $\alpha = \beta \approx \frac{1}{4}$, then

$$q \approx 0.8$$

Proof:

The proof follows exactly the one in [5], Theorem 3.1 and 3.3, with proper accounting for the interval length ϵ .

□

4 Truncation Error in a Sub-interval—Chebyshev and Legendre Methods

In this section we assume that we have the first $N + 1$ Chebyshev or Legendre coefficients in the expansion of an L_2 function $f(x)$ based on the interval $-1 \leq x \leq 1$. We show how to get its Gegenbauer expansion based on a sub-interval $[a, b] \subset [-1, 1]$ with exponential accuracy in N . The proof makes use of Lemma 2.7, which states that the *low modes* of the Fourier expansion of the Chebyshev or Legendre polynomial of degree N decay exponentially with N . Thus only the high modes count, and those are shown in Section 2 to have small Gegenbauer coefficients.

For simplicity of notation we will deal with $C_k^{\frac{1}{2}}(x)$, which is the Legendre polynomial $P_k(x)$ (see (2.9)), and $C_k^0(x)$, which is $\frac{2}{k}$ times the Chebyshev polynomial $T_k(x)$ (see (2.10)). The Chebyshev or Legendre expansion coefficients of $f(x)$ are given by

$$\hat{f}^\mu(k) = \frac{1}{h_k^\mu} \int_{-1}^1 (1-x^2)^{\mu-\frac{1}{2}} f(x) C_k^\mu(x) dx \quad (4.1)$$

for $\mu = 0$ or $\mu = \frac{1}{2}$.

In this section we consider functions $f(x)$ satisfying

Assumption 4.1:

$$|\hat{f}^\mu(k)| \leq A \text{ for all } k.$$

□

We remark that if $f(x)$ is an L_2 function the assumption is fulfilled.

We assume that we know the first $N + 1$ Chebyshev or Legendre coefficients, $\hat{f}^\mu(k)$ for $0 \leq k \leq N$, and define

$$f_N^\mu(x) = \sum_{k=0}^N \hat{f}^\mu(k) C_k^\mu(x) \quad (4.2)$$

Note that $f_N^\mu(x)$ does not converge fast to $f(x)$ if there exist discontinuities inside the domain.

The function $f(x)$ has also a Gegenbauer expansion in a sub-interval $[a, b]$. With ξ , ϵ and δ defined in (3.2)-(3.3), we have

$$f(\epsilon\xi + \delta) = \sum_{l=0}^{\infty} \hat{f}_l^\lambda C_l^\lambda(\xi) \quad (4.3)$$

where the Gegenbauer coefficients \hat{f}_l^λ are defined by

$$\hat{f}_l^\lambda = \frac{1}{h_l^\lambda} \int_{-1}^1 (1 - \xi^2)^{\lambda - \frac{1}{2}} f(\epsilon\xi + \delta) C_l^\lambda(\xi) d\xi \quad (4.4)$$

As before, we do not have \hat{f}_l^λ at our disposal, but only an approximation based on the partial Chebyshev or Legendre sum $f_N^\mu(x)$, thus we have

$$\hat{g}_l^\lambda = \frac{1}{h_l^\lambda} \int_{-1}^1 (1 - \xi^2)^{\lambda - \frac{1}{2}} f_N^\mu(\epsilon\xi + \delta) C_l^\lambda(\xi) d\xi \quad (4.5)$$

How well do \hat{g}_l^λ approximate \hat{f}_l^λ ? To answer this question we define

Definition 4.2:

The truncation error is defined by

$$TE(\lambda, m, N, \epsilon) = \max_{-1 \leq \xi \leq 1} \left| \sum_{l=0}^m (\hat{f}_l^\lambda - \hat{g}_l^\lambda) C_l^\lambda(\xi) \right| \quad (4.6)$$

where \hat{f}_l^λ are defined by (4.4) and \hat{g}_l^λ are defined by (4.5).

□

The truncation error is the measure of the distance between the true Gegenbauer expansion in the interval $[a, b]$ and its approximation based on the Chebyshev or Legendre partial sum in $[-1, 1]$.

We first have the following lemma:

Lemma 4.3:

$$TE(\lambda, m, N, \epsilon) \leq \sum_{q=N+1}^{\infty} |\hat{f}^\mu(q)| \sum_{l=0}^m \left| \frac{C_l^\lambda(1)}{h_l^\lambda} \int_{-1}^1 (1 - \xi^2)^{\lambda - \frac{1}{2}} C_q^\mu(\epsilon\xi + \delta) C_l^\lambda(\xi) d\xi \right| \quad (4.7)$$

Proof:

From (4.4) and (4.5) we have

$$\left| \hat{f}_\epsilon^\lambda(l) - \hat{g}_\epsilon^\lambda(l) \right| = \frac{1}{h_l^\lambda} \left| \int_{-1}^1 (1 - \xi^2)^{\lambda - \frac{1}{2}} (f(\epsilon\xi + \delta) - f_N^\mu(\epsilon\xi + \delta)) C_l^\lambda(\xi) d\xi \right| \quad (4.8)$$

Substituting (4.8) into (4.6), recalling (2.4) and

$$f(\epsilon\xi + \delta) - f_N^\mu(\epsilon\xi + \delta) = \sum_{q=N+1}^{\infty} \hat{f}^\mu(q) C_q^\mu(\epsilon\xi + \delta) \quad (4.9)$$

we obtain (4.7). □

For simplicity of notations we denote:

$$F_{q,l,\lambda} = \left| \frac{C_l^\lambda(1)}{h_l^\lambda} \int_{-1}^1 (1 - \xi^2)^{\lambda - \frac{1}{2}} C_q^\mu(\epsilon\xi + \delta) C_l^\lambda(\xi) d\xi \right|, \quad q > N \quad (4.10)$$

In order to estimate this term, we consider the Fourier series approximation of $C_q^\mu(\epsilon\xi + \delta)$ from (2.12) and (2.13) to get

Lemma 4.4:

$$F_{q,l,\lambda} \leq A C_l^\lambda(1) \Gamma(\lambda) \sum_{k=1}^{\infty} \left(\frac{2}{k\pi\epsilon} \right)^\lambda |J_{l+\lambda}(k\pi\epsilon)| |a_k^q| \quad (4.11)$$

where a_k^q is given either by (2.14) or by (2.15), and A grows at most as fast as a linear polynomial in λ and l .

Proof:

We expand $C_q^\mu(x)$ using (2.12) or (2.13):

$$C_q^\mu(x) = \sum_{k=-\infty}^{\infty} a_k^{q,\mu} e^{ik\pi x} = \sum_{k=-\infty}^{\infty} a_k^{q,\mu} e^{ik\pi\epsilon\xi} e^{ik\pi\delta} \quad (4.12)$$

where $a_k^{q,\frac{1}{2}} = a_k^q$ is given by (2.14) and $a_k^{q,0} = \frac{2}{q} b_k^q$ is given by (2.15).

We now use (2.11) to get

$$\left| \frac{1}{h_l^\lambda} \int_{-1}^1 (1 - \xi^2)^{\lambda - \frac{1}{2}} e^{ik\pi\epsilon\xi} e^{ik\pi\delta} C_l^\lambda(\xi) d\xi \right| = \Gamma(\lambda) \left(\frac{2}{|k|\pi\epsilon} \right)^\lambda (l + \lambda) |J_{l+\lambda}(k\pi\epsilon)| \quad (4.13)$$

Substituting (4.12) and (4.13) into (4.10) gives us (4.11).

□

We are ready now to estimate directly $F_{q,l,\lambda}$ defined in (4.10).

Lemma 4.5:

Consider $F_{q,l,\lambda}$ defined in (4.10). Let $\lambda = \frac{2}{27\epsilon}\epsilon N$ and $l = \gamma\lambda$ where $\gamma < 1$. Then for every $q > N$ we have

$$F_{q,l,\lambda} \leq A e^{-\lambda} \left(\frac{N}{q}\right)^\lambda. \quad (4.14)$$

where A grows at most quadratically with N .

Proof:

Let $k_q = \left[\frac{q}{\pi\epsilon}\right]$ be the integer part of $\frac{q}{\pi\epsilon}$. We split the sum on the right hand side of (4.11) into

$$F_{q,l,\lambda} \leq A \left(\sum_{k=1}^{k_q} + \sum_{k=k_q+1}^{\infty} \right) \left[C_l^\lambda(1) \Gamma(\lambda) \left(\frac{2}{k\pi\epsilon}\right)^\lambda |J_{l+\lambda}(k\pi\epsilon)| |a_k^q| \right] = A (F_{q,l,\lambda}^{(1)} + F_{q,l,\lambda}^{(2)}) \quad (4.15)$$

We first use (2.3) and (2.7) to obtain:

$$C_l^\lambda(1) \leq A \left(\frac{(\gamma+2)^{\gamma+2}}{4\gamma^\gamma}\right)^\lambda \leq A \left(\frac{27}{4}\right)^\lambda \quad (4.16)$$

and

$$\Gamma(\lambda) \leq A \lambda^\lambda e^{-\lambda} \quad (4.17)$$

We start with an estimation of the second sum

$$F_{q,l,\lambda}^{(2)} = \sum_{k=k_q+1}^{\infty} C_l^\lambda(1) \Gamma(\lambda) \left(\frac{2}{k\pi\epsilon}\right)^\lambda |J_{l+\lambda}(k\pi\epsilon)| |a_k^q|$$

We use (4.16), (4.17), the fact that $|J_{l+\lambda}(x)| \leq 1$ and $|a_k^q| \leq 1$, and $k > k_q = \left[\frac{q}{\pi\epsilon}\right]$ and $\lambda = \frac{2}{27\epsilon}\epsilon N$ to get

$$F_{q,l,\lambda}^{(2)} \leq A \left(\frac{27\lambda}{2\epsilon q}\right)^\lambda \leq A e^{-\lambda} \left(\frac{N}{q}\right)^\lambda. \quad (4.18)$$

where A grows at most linearly with N , the growth coming from the summation. For the first sum

$$F_{q,l,\lambda}^{(1)} = \sum_{k=1}^{k_q} C_l^\lambda(1) \Gamma(\lambda) \left(\frac{2}{k\pi\epsilon}\right)^\lambda |J_{l+\lambda}(k\pi\epsilon)| |a_k^q|,$$

we still use (4.16), (4.17) and the bound $|J_{l+\lambda}(x)| \leq 1$. However, for the estimate on $|a_k^q|$, we use (2.16), (2.17):

$$|a_k^q| \leq A \left(\frac{e\pi k}{2q} \right)^q \quad (4.19)$$

and the fact $q > N > \lambda$ to get

$$F_{q,l,\lambda}^{(1)} \leq A \sum_{k=1}^{k_q} \left(\frac{e\pi k}{2q} \right)^q \left(\frac{27\lambda}{2e\pi k\epsilon} \right)^\lambda \leq Ak_q \left(\frac{e\pi k_q}{2q} \right)^q \left(\frac{27\lambda}{2e\pi k_q\epsilon} \right)^\lambda \quad (4.20)$$

and, upon substituting for $k_q = \left\lfloor \frac{q}{\pi\epsilon} \right\rfloor$ and $\lambda = \frac{2}{27e}\epsilon N$:

$$F_{q,l,\lambda}^{(1)} \leq Ak_q \left(\frac{1}{2} \right)^q e^{-\lambda} \left(\frac{N}{q} \right)^\lambda \leq Ae^{-\lambda} \left(\frac{N}{q} \right)^\lambda \quad (4.21)$$

Combining (4.18) and (4.21) we obtain (4.14). □

We are now ready for the main theorem of this section:

Theorem 4.6:

Let the truncation error be defined in (4.6). Let $\lambda = \frac{2}{27e}N$ and $m \leq \lambda$, then

$$TE(\lambda, m, N, \epsilon) \leq Ae^{-\lambda} \quad (4.22)$$

where A grows at most as a fourth order polynomial in N .

Proof:

The theorem follows from (4.7), the Assumption 4.1, and (4.14). □

5 Regularization Error for a Sub-interval

In this section we study the second part of the error, which is caused by using a finite Gegenbauer expansion based on a sub-interval $[a, b] \subset [-1, 1]$, to approximate a function $f(x)$ which is assumed analytic in this sub-interval $[a, b]$. Since in previous sections we have

established exponentially small truncation errors if λ and m are both growing linearly with N , in this section we will consider the case of $\lambda \sim m$.

We will assume that $f(x)$ is an analytic function on $[a, b]$ satisfying

Assumption 5.1

There exists constants $\rho \geq 1$ and $C(\rho)$ such that, for every $k \geq 0$,

$$\max_{a \leq x \leq b} \left| \frac{d^k f}{dx^k}(x) \right| \leq C(\rho) \frac{k!}{\rho^k} \quad (5.1)$$

□

This is a standard assumption for analytic functions. ρ is the distance from $[a, b]$ to the nearest singularity of $f(x)$ in the complex plane (see, for example, [8]). This assumption and the following proof can be modified to allow for an arbitrary ellipse region around $[a, b]$ for the analyticity of $f(x)$ (H. Vandeven, private communication).

Let us consider the Gegenbauer partial sum of the first m terms for the function $f(\epsilon\xi + \delta)$:

$$f_m^{\lambda, \epsilon}(\xi) = \sum_{l=0}^m \hat{f}_l^\lambda(l) C_l^\lambda(\xi) \quad (5.2)$$

with ξ , ϵ and δ defined by (3.2) and (3.3), and the Gegenbauer coefficients based on $[a, b]$ defined by

$$\hat{f}_l^\lambda(l) = \frac{1}{h_l^\lambda} \int_{-1}^1 (1 - \xi^2)^{\lambda - \frac{1}{2}} f(\epsilon\xi + \delta) C_l^\lambda(\xi) d\xi \quad (5.3)$$

We want to estimate the regularization error in the maximum norm:

$$RE(\lambda, m, \epsilon) = \max_{-1 \leq \xi \leq 1} \left| f(\epsilon\xi + \delta) - \sum_{l=0}^m \hat{f}_l^\lambda(l) C_l^\lambda(\xi) \right| \quad (5.4)$$

We have the following result for the estimation of the regularization error, when $\lambda \sim m$:

Theorem 5.2:

Assume $\lambda = \gamma m$ where γ is a positive constant. If $f(x)$ is analytic in $[a, b] \subset [-1, 1]$ satisfying the Assumption 5.1, then the regularization error defined in (5.4) can be bounded by

$$RE(\gamma m, m, \epsilon) \leq Aq^m \quad (5.5)$$

where q is given by

$$q = \frac{\epsilon(1+2\gamma)^{1+2\gamma}}{\rho^{2^{1+2\gamma}}\gamma(1+\gamma)^{1+\gamma}} \quad (5.6)$$

which is always less than 1. In particular, if $\gamma = 1$ and $m = \beta N$ where β is a positive constant, then

$$RE(\beta N, \beta N, \epsilon) \leq Aq^N \quad (5.7)$$

with

$$q = \left(\frac{27\epsilon}{32\rho}\right)^\beta \quad (5.8)$$

Proof:

The proof follows in exactly in the same fashion as in [5], Lemma 4.2, Theorems 4.3 and 4.4, with proper accounting for the transformation $x(\xi) = \epsilon\xi + \delta$.

□

6 Concluding Remarks

In this Section, we combine the estimates for truncation errors and regularization errors in previous sections, to obtain the main theorem of this paper:

Theorem 6.1: Removal of the Gibbs Phenomenon for the sub-interval case

Consider an L_2 function $f(x)$ on $[-1, 1]$, which is analytic in a sub-interval $[a, b] \subset [-1, 1]$ and satisfies

$$\max_{a \leq x \leq b} \left| \frac{d^k f}{dx^k}(x) \right| \leq C(\rho) \frac{k!}{\rho^k}, \quad \rho \geq 1 \quad (6.1)$$

Assume that either the first $2N + 1$ Fourier coefficients

$$\hat{f}(k) = \frac{1}{2} \int_{-1}^1 f(x) e^{-ik\pi x} dx, \quad -N \leq k \leq N$$

or the first $N + 1$ Chebyshev or Legendre coefficients

$$\hat{f}^\mu(k) = \frac{1}{h_k^\mu} \int_{-1}^1 (1-x^2)^{\mu-\frac{1}{2}} f(x) C_k^\mu(x) dx, \quad \mu = 0 \text{ or } \mu = \frac{1}{2}$$

are known. Let $\hat{g}_i^\lambda(l)$, $0 \leq l \leq m$ be the Gegenbauer expansion coefficients, based on the sub-interval $[a, b]$, of the spectral partial sum $f_N(x)$ in (3.4) or $f_N^\mu(x)$ in (4.2). These Gegenbauer coefficients are defined in (3.7) for the Fourier case and in (4.5) for the Chebyshev or Legendre case. Then for $\lambda = m = \beta \epsilon N$ where $\beta < \frac{2\pi\epsilon}{27}$ (Fourier case) or $\beta \leq \frac{2}{27\epsilon}$ (Chebyshev or Legendre case), we have

$$\max_{-1 \leq \xi \leq 1} \left| f(\epsilon\xi + \delta) - \sum_{l=0}^m \hat{g}_i^\lambda(l) C_l^\lambda(\xi) \right| \leq A (q_T^{\epsilon N} + q_R^{\epsilon N}) \quad (6.2)$$

where

$$q_T = \left(\frac{27\beta}{2\pi\epsilon} \right)^\beta < 1, \quad q_R = \left(\frac{27\epsilon}{32\rho} \right)^\beta < 1$$

for the Fourier case and

$$q_T = e^{-\beta} < 1, \quad q_R = \left(\frac{27\epsilon}{32\rho} \right)^\beta < 1$$

for the Chebyshev or Legendre case, and A is at most a fourth order polynomial in N .

Proof:

Just combine the results of Theorems 3.2, 4.6 and 5.2.

□

We finally give two remarks:

Remark 6.2

It can be seen from (6.2), that the truncation error gets bigger when the interval half-length ϵ decreases, because the parameters m and λ must decrease with ϵ . This in turn also affects the regularization error, although there is a factor ϵ in q_R which offsets this effect.

□

Remark 6.3

No attempt has been made in this paper to optimize the parameters.

□

References

- [1] M. Abramowitz and I. A. Stegun, *Handbook of Mathematical Functions*, Dover, 1970.
- [2] H. Bateman, *Higher Transcendental Functions*, vol. 2, McGraw-Hill, 1953.
- [3] C. Canuto, M.Y. Hussaini, A. Quarteroni and T. A. Zang, *Spectral Methods in Fluid Dynamics*, Springer-Verlag, 1988.
- [4] D. Gottlieb and S. Orszag, *Numerical Analysis of Spectral Methods: Theory and Applications*, SIAM-CBMS, Philadelphia, 1977.
- [5] D. Gottlieb, C.-W. Shu, A. Solomonoff and H. Vandeven, *On The Gibbs Phenomenon I: recovering exponential accuracy from the Fourier partial sum of a nonperiodic analytic function*, J. Comput. Appl. Math., v43, 1992, pp.81-92.
- [6] D. Gottlieb and C.-W. Shu, *Resolution properties of the Fourier method for discontinuous waves*, ICASE Report No. 92-27. June 1992.
- [7] I. Gradshteyn and I. Ryzhik, *Tables of Integrals, Series, and Products*, Academic Press, 1980.
- [8] F. John, *Partial Differential Equations*, Springer-Verlag, 1982.

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE November 1993	3. REPORT TYPE AND DATES COVERED Contractor Report	
4. TITLE AND SUBTITLE ON THE GIBBS PHENOMENON III: RECOVERING EXPONENTIAL ACCURACY IN A SUB-INTERVAL FROM A SPECTRAL PARTIAL SUM OF A PIECEWISE ANALYTIC FUNCTION			5. FUNDING NUMBERS C NAS1-19480 WU 505-90-52-01	
6. AUTHOR(S) David Gottlieb Chi-Wang Shu				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Institute for Computer Applications in Science and Engineering Mail Stop 132C, NASA Langley Research Center Hampton, VA 23681-0001			8. PERFORMING ORGANIZATION REPORT NUMBER ICASE Report No. 93-82	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) National Aeronautics and Space Administration Langley Research Center Hampton, VA 23681-0001			10. SPONSORING/MONITORING AGENCY REPORT NUMBER NASA CR-191560 ICASE Report No. 93-82	
11. SUPPLEMENTARY NOTES Langley Technical Monitor: Michael F. Card Final Report Submitted to SIAM Journal on Numerical Analysis				
12a. DISTRIBUTION/AVAILABILITY STATEMENT Unclassified-Unlimited Subject Category 64			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) We continue the investigation of overcoming Gibbs phenomenon, i.e., obtaining exponential accuracy at all points including at the discontinuities themselves, from the knowledge of a spectral partial sum of a discontinuous but piecewise analytic function. We show that if we are given the first N expansion coefficients of an L_2 function $f(x)$ in terms of either the trigonometrical polynomials or the Chebyshev or Legendre polynomials, we can construct an exponentially convergent approximation to the point values of $f(x)$ in any sub-interval in which it is analytic.				
14. SUBJECT TERMS Gibbs phenomenon; Fourier series; Chebyshev series; Legendre series; Gegenbauer polynomials; exponential accuracy			15. NUMBER OF PAGES 19	
			16. PRICE CODE A03	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT	20. LIMITATION OF ABSTRACT	

NSN 7540-01-280-5500

Standard Form 298 (Rev. 2-89)
Prescribed by ANSI Std. Z39-18
298-102