

AD-A276 584



DOCUMENTATION PAGE

Form Approved
OMB No 0704-0188

2

on is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and reviewing the collection of information, sending comments regarding this burden estimate or any other aspect of this collection of information, including this burden estimate, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188) Washington, DC 20503

2. REPORT DATE 25-2-94		3. REPORT TYPE AND DATES COVERED Technical Report	
4. TITLE AND SUBTITLE Synthesis, Characterization and Molecular Structure of a Compound with a Ring of Group 13, 15 and 16 Atoms, (Me ₃ CCH ₂) ₂ InP(t-Bu) ₂ In(CH ₂ CMe ₃) ₂ SePh		5. FUNDING NUMBERS Grant: N00014-90-J-1530 R&T Code: 4135002	
6. AUTHOR(S) O. T. Beachley, Jr., Sun-Hua L. Chao, Melvyn Rowen Churchill and Charles H. Lake		8. PERFORMING ORGANIZATION REPORT NUMBER Technical Report No. 39	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Department of Chemistry State University of New York at Buffalo Buffalo, NY 14214-3094		10. SPONSORING/MONITORING AGENCY REPORT NUMBER n/a	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Office of Naval Research 800 N. Quincy Street Arlington, VA 22217-5000		11. SUPPLEMENTARY NOTES Accepted for publication - Organometallics	
12a. DISTRIBUTION/AVAILABILITY STATEMENT This document has been approved for public release and sale; its distribution is unlimited. Reproduction in whole or in part is permitted for any purpose of the United States government.		12b. DISTRIBUTION CODE n/a	
13. ABSTRACT (Maximum 200 words) The compound (Me ₃ CCH ₂) ₂ InP(t-Bu) ₂ In(CH ₂ CMe ₃) ₂ SePh has been prepared from equimolar quantities of [(Me ₃ CCH ₂) ₂ InP(t-Bu) ₂] ₂ and [(Me ₃ CCH ₂) ₂ InSePh] ₂ in pentane solution, recrystallized and fully characterized. This mixed-bridge organoindium species crystallized in the centrosymmetric space group P $\bar{1}$ (No. 2) with $a = 10.069(2)\text{\AA}$, $b = 11.627(4)\text{\AA}$, $c = 18.452(4)\text{\AA}$, $\alpha = 100.74(2)^\circ$, $\beta = 99.33(2)^\circ$, $\gamma = 103.38(2)^\circ$, $V = 2016.9(8)\text{\AA}^3$, and $Z = 2$. The structure was solved and refined to $R = 2.51\%$ and $wR = 2.52\%$ for those 3747 unique data with $ F_o < 6\sigma(F_o)$ ($R = 4.51\%$ for all 5304 data). The two indium atoms are bridged by both a bis(tert-butyl)-phosphido ligand and a phenylselenido ligand. The four-membered In ₂ SeP core has a puckered arrangement such that the fold angle about the In...In axis is 20.9° .			
14. SUBJECT TERMS Indium, phosphorus, selenium compound, X-ray structural study		15. NUMBER OF PAGES 21	
17. SECURITY CLASSIFICATION OF REPORT Unclassified		16. PRICE CODE n/a	
18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT UL	

SDTC
ELECTE
MAR 03 1994
S E D

466
352

94-07071

NSN 7540-01-280-5500

Standard Form 298 (Rev 2-89)
Prescribed by ANSI Std Z39-18
298-102

94 3 03 010

OFFICE OF NAVAL RESEARCH

Contract N-00014-90-J-1530

R&T Code 4135002

TECHNICAL REPORT NO. 39

Synthesis, Characterization and Molecular Structure of a
Compound with a Ring of Group 13, 15 and 16 Atoms.

$(\text{Me}_3\text{CCH}_2)_2\text{InP}(\text{t-Bu})_2\text{In}(\text{CH}_2\text{CMe}_3)_2\text{SePh}$

by

O. T. Beachley, Jr., Sun-Hua L. Chao, Melvyn Rowen Churchill
and Charles H. Lake

Prepared for Publication

in

Organometallics

State University of New York at Buffalo
Department of Chemistry
Buffalo, New York 14214

25 February 1994

Reproduction in whole or in part is permitted for
any purpose of the United States Government


*This document has been approved for public release
and sale; its distribution is unlimited

DTIC QUALITY INSPECTION

Accession For	
NTIS CRA&I	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
A-1	

Contribution from the Department of Chemistry,
State University of New York at Buffalo, Buffalo, NY 14214

Synthesis, Characterization and Molecular Structure
of a Compound with a Ring of
Group 13, 15 and 16 Atoms,




by

O. T. Beachley, Jr.*, Sun-Hua L. Chao, Melvyn Rowen Churchill*
and Charles H. Lake

Summary: The compound $\overline{(\text{Me}_3\text{CCH}_2)_2\text{InP}(\text{t-Bu})_2\text{In}(\text{CH}_2\text{CMe}_3)_2\text{SePh}}$ has been prepared from equimolar quantities of $[(\text{Me}_3\text{CCH}_2)_2\text{InP}(\text{t-Bu})_2]_2$ and $[(\text{Me}_3\text{CCH}_2)_2\text{InSePh}]_2$ in pentane solution, recrystallized and fully characterized. This mixed-bridge organoindium species crystallized in the centrosymmetric space group $P\bar{1}$ (No. 2) with $a = 10.069(2)\text{\AA}$, $b = 11.627(4)\text{\AA}$, $c = 18.452(4)\text{\AA}$, $\alpha = 100.74(2)^\circ$, $\beta = 99.33(2)^\circ$, $\gamma = 103.38(2)^\circ$, $V = 2016.9(8)\text{\AA}^3$, and $Z = 2$. The structure was solved and refined to $R = 2.51\%$ and $wR = 2.52\%$ for those 3747 unique data with $|F_o| > 6\sigma(|F_o|)$ ($R = 4.51\%$ for all 5304 data). The two indium atoms are bridged by both a bis(tert-butyl)phosphido ligand and a phenylselenido ligand. The four-membered In_2SeP core has a puckered arrangement such that the fold angle about the $\text{In}\cdots\text{In}$ axis is 20.9° .

Compounds of the type $[R_2MER'_2]_n$ ($M = Al, Ga, In$; $E = N, P, As$) typically have four- and six-membered cyclic structures.¹ If these rings are envisioned to involve a series of donor-acceptor bonds, it should be possible to replace a group 15 moiety (ER'_2) with a Lewis base moiety from another or replace a group 13 moiety (MR_2) with a different Lewis acid. Thus, a series of gallium and indium compounds which have two different bridging atoms, a phosphorus or arsenic and a chlorine or bromine, have been prepared and fully characterized. These compounds have the general formula $R_2ME(SiMe_3)_2MR_2X$ ($M = Ga, E = As, R = Ph, X = Cl,^2 Br;^2 M = Ga, E = As, R = CH_2CMe_3,^3 CH_2SiMe_3,^4 CMe_3,^5 X = Cl; M = Ga, E = P, R = Ph, X = Cl;^6 M = In, E = P, R = CH_2CMe_3,^7 CH_2SiMe_3,^8 X = Cl$ and $M = In, E = As, R = CH_2SiMe_3, X = Cl^9$). This interesting class of mixed bridge compounds has now been extended to include a ring with group 13, 15 and 16 atoms, $(Me_3CCH_2)_2\overline{InP(t-Bu)_2In(CH_2CMe_3)_2SePh}$.

The compound $(Me_3CCH_2)_2\overline{InP(t-Bu)_2In(CH_2CMe_3)_2SePh}$ was prepared from equimolar quantities of the two dimeric reactants, $[(Me_3CCH_2)_2InP(t-Bu)_2]_2$ ¹⁰ and $[(Me_3CCH_2)_2InSePh]_2$ ¹¹ in pentane solution at room temperature. The ease of formation



of the new ring structure demonstrates the lability of the original four-membered indium-phosphorus and/or indium-selenium rings. The mixed bridge product was recrystallized from pentane solution in approximately 75% yield and fully characterized in solution and in the solid state including a single crystal X-ray structural study.

The unit cell consists of two discrete molecules of $(\text{Me}_3\text{CCH}_2)_2\text{InP}(\text{t-Bu})_2\text{In}(\text{CH}_2\text{CMe}_3)_2\text{SePh}$ which are separated by normal van der Waals' distances. There are no anomalously short intermolecular contacts. A list of interatomic distances and angles is included in Table I. A diagram of the molecule, showing the atomic labeling scheme, is shown in Figure 1.

The four-membered In_2SeP core (Figure 2) has a puckered arrangement with the following In-Se and In-P distances: $\text{In}(1) - \text{Se}(1) = 2.769(1)\text{\AA}$ and $\text{In}(2) - \text{Se}(1) = 2.748(1)\text{\AA}$ (average = $2.759\text{\AA}$) and $\text{In}(1) - \text{P}(1) = 2.653(2)\text{\AA}$ and $\text{In}(2) - \text{P}(1) = 2.651(2)\text{\AA}$ (average = $2.652\text{\AA}$). These In-P distances are shorter than those associated with the symmetrically bridged compound $[(\text{Me}_3\text{CCH}_2)_2\text{InP}(\text{t-Bu})_2]_2$, in which the average In-P distance is $2.701\text{\AA}$.¹⁰ The In-Se distances are slightly elongated relative to those in the symmetrically bridged compound $[(\text{Me}_3\text{CCH}_2)_2\text{InSePh}]_2$, which has an average In-Se distance of $2.744\text{\AA}$.¹¹ The fold angle about the $\text{In}(1)\cdots\text{In}(2)$ axis, as defined by the angle of intersection of the $\text{In}(1) - \text{Se}(1) - \text{In}(2)$ and $\text{In}(1) - \text{P}(1) - \text{In}(2)$ planes, is 20.9° . The fold angle about the $\text{Se}(1)\cdots\text{P}(1)$ axis, which is defined by the angle of intersection of the planes $\text{Se}(1) - \text{In}(1) - \text{P}(1)$ and $\text{Se}(1) - \text{In}(2) - \text{P}(1)$, is 19.2° . These fold angles are much smaller than the corresponding fold angles in $[(\text{Me}_3\text{CCH}_2)_2\text{InSePh}]_2$, in which that about the $\text{In}\cdots\text{In}$ axis is 30.8° and that about the $\text{Se}\cdots\text{Se}$ axis is 31.7° .¹¹ In comparison, $(\text{Me}_3\text{CCH}_2)_2\text{InP}(\text{SiMe}_3)_2\text{In}(\text{CH}_2\text{CMe}_3)_2\text{Cl}$ is planar,⁷ whereas $(\text{Me}_3\text{SiCH}_2)_2\text{InP}(\text{SiMe}_3)_2\text{In}(\text{CH}_2\text{SiMe}_3)_2\text{Cl}$ has a puckered ring with a fold angle about the $\text{In}\cdots\text{In}$ axis of 33.3° . The nonbonding distances across the core in the present molecule are $\text{In}(1)\cdots\text{In}(2) = 3.954\text{\AA}$ and $\text{Se}(1)\cdots\text{P}(1) = 3.631\text{\AA}$. These distances are intermediate between those for symmetrically bridged bis(*t*-butyl)phosphido and phenylselenide

compounds. Thus, both In...In distances and (bridging ligand)···(bridging ligand) distance vary in the order $[(\text{Me}_3\text{CCH}_2)_2\text{InP}(\text{t-Bu})_2]_2 > (\text{Me}_3\text{CCH}_2)_2\text{InP}(\text{t-Bu})_2\text{In}(\text{CH}_2\text{CMe}_3)_2\text{SePh} > [(\text{Me}_3\text{CCH}_2)_2\text{InSePh}]_2$. Both the present mixed-bridge compound and the symmetrically bridged phenylselenide compound crystallize with a puckered core. This conformation is adopted because of the low symmetry of the bridging ligands which allows the ring to fold and reduce the total amount of steric interference between groups attached to the central ring.¹¹ The symmetrically bridged bis(t-butyl)phosphido derivative has symmetrical ligands bridging the two indium atoms and has a planar In_2P_2 core.

The bridging phenylselenide ligand is associated with a Se(1) - C(1) bond length of 1.922(5) Å and bond angles of In(1) - Se(1) - C(1) = 112.6(1)° and In(2) - Se(1) - C(1) = 101.3(2) Å. The Se-C distance is identical to the Se-C distances in $[(\text{Me}_3\text{CCH}_2)_2\text{InSePh}]_2$. The phenyl group is bent out of the In(1) - Se(1) - In(2) plane by 25.9°. The orientation of this ligand is determined by intermolecular packing rather than intramolecular forces.

The bridging bis(tert-butyl)phosphido ligand is associated with the following phosphorus-carbon distances: P(1) - C(311) = 1.895(5) Å, P(1) - C(321) = 1.891(5) Å (average = 1.893 Å); the interligand angle C(311) - P(1) - C(321) is 112.2(2)°. These P-C distances are very similar to those associated with $[(\text{Me}_3\text{CCH}_2)_2\text{InP}(\text{t-Bu})_2]_2$. The plane formed by C(311) - P(1) - C(321) intersects the In(1) - P(1) - In(2) plane at an angle of 89.2°. The equivalent angle in $[(\text{Me}_3\text{CCH}_2)_2\text{InP}(\text{t-Bu})_2]_2$ is 88.6°.

There are two neopentyl ligands associated with each indium atom. In-C distances are In(1) - C(111) = 2.194(4) Å, In(1) - C(121) = 2.194(5) Å, In(2) - C(211) = 2.181(6) Å and In(2) - C(221) = 2.193(4) Å (average = 2.191(5) Å). Interligand angles are C(111) - In(1)

- C(121) = 119.5(2)° and C(211) - In(2) - C(221) = 122.3(2)° (average = 120.9°). These angles are intermediate between those in [(Me₃CCH₂)₂InP(t-Bu)₂]₂ (average interligand angle = 115.2°) and [(Me₃CCH₂)₂InSePh]₂ (average interligand angle = 133.6°).

The NMR spectroscopic data confirm the existence of a single, unique compound in benzene and toluene solutions rather than an equilibrium mixture of reactants and products. A comparison of the ¹H and ³¹P NMR chemical shift data for (Me₃CCH₂)₂ $\overline{\text{InP(t-Bu)}_2\text{In(CH}_2\text{CMe}_3)_2\text{SePh}}$, [(Me₃CCH₂)₂InP(t-Bu)₂]₂¹⁰ and [(Me₃CCH₂)₂InSePh]₂¹¹ is provided in Table II. It is significant that the resonance for the t-butyl group protons bound to phosphorus in (Me₃CCH₂)₂ $\overline{\text{InP(t-Bu)}_2\text{In(CH}_2\text{CMe}_3)_2\text{SePh}}$ is a doublet at 1.36 ppm, whereas in [(Me₃CCH₂)₂InP(t-Bu)₂]₂ the resonance for the t-butyl group protons is a triplet at 1.46 ppm.¹⁰ The trans coupling of the protons of the t-butyl group to phosphorus in [(Me₃CCH₂)₂InP(t-Bu)₂]₂ is not possible for the mixed-bridge compound with one phosphorus atom. The chemical shifts of the phosphorus atoms in the ³¹P NMR spectra of the reactant and product differ by 11.0 ppm. The X-ray structural study identified the presence of unique neopentyl and t-butyl groups due to the relative position of the phenyl group on selenium but these were not observed in solution by NMR spectroscopy. The two neopentyl groups bound to indium are equivalent as are the two t-butyl groups on phosphorus at room temperature and at -80°C in toluene. Either the substituents on indium and phosphorus are not differentiated by the phenyl group or, more likely, an exchange process makes the protons of each group equivalent. The most likely process would appear to be rapid inversion of the puckered four-membered ring, even at -80°C. Alternatively, ring cleavage and rotation would be required.

The melting point and solubility properties of the bulk solid are consistent with the molecule identified by the X-ray structural study but not consistent with a mixture of the two starting compounds, $[(\text{Me}_3\text{CCH}_2)_2\text{InP}(\text{t-Bu})_2]_2$ and $[(\text{Me}_3\text{CCH}_2)_2\text{InSePh}]_2$. The melting point of the product (144 - 145°C) is sharp and higher than the melting point of $[(\text{Me}_3\text{CCH}_2)_2\text{InSePh}]_2$ (103.8 - 105.5°C)¹¹ but lower than the decomposition temperature of $[(\text{Me}_3\text{CCH}_2)_2\text{InP}(\text{t-Bu})_2]_2$ (200 - 202°C).¹⁰ The mixed bridge compound is also significantly more soluble in pentane and benzene than either of the two compounds used to make it. The phosphide $[(\text{Me}_3\text{CCH}_2)_2\text{InP}(\text{t-Bu})_2]_2$ is almost insoluble in pentane and has insufficient solubility in benzene for a cryoscopic molecular weight study (0.01 ~ 0.10 m).¹¹

EXPERIMENTAL

Compounds described in this investigation were extremely sensitive to oxygen and moisture and were handled in a standard vacuum line or under a purified argon atmosphere. The starting compounds $[(\text{Me}_3\text{CCH}_2)_2\text{InP}(\text{t-Bu})_2]_2$ ¹⁰ and $[\text{In}(\text{CH}_2\text{CMe}_3)_2(\text{SePh})]_2$ ¹¹ were prepared and purified by literature methods. Solvents were dried by conventional procedures. Elemental analyses were performed by E + R Microanalytical Laboratory, Inc., Corona, NY. Infrared spectra of Nujol mulls between CsI plates were recorded by means of a Perkin-Elmer 683 spectrometer. The ^1H NMR spectra were recorded at 400 MHz by means of a Varian VXR-400 S spectrometer. Proton chemical shifts were reported in δ units (ppm) and were referenced to C_6H_6 at δ 7.15 ppm. The $^{13}\text{C}\{^1\text{H}\}$ NMR spectra were recorded by means of a Varian Gemini 300 at 75 MHz. The ^{13}C spectra were reported relative to C_6D_6 at δ 128.0 ppm. The $^{31}\text{P}\{^1\text{H}\}$ and ^{31}P NMR spectra were recorded at 161.9 MHz by means of a Varian VXR-400 S spectrometer. The ^{31}P spectra were reported relative to H_3PO_4 at δ 0.0 ppm via an external standard. All samples for NMR spectra were contained in sealed NMR tubes. Melting points were observed in sealed capillaries under purified argon. Molecular weights were measured cryoscopically in benzene solution by using an instrument similar to that described by Shriver and Drezdson.¹²

Synthesis of $(\text{Me}_3\text{CCH}_2)_2\text{InP}(\text{t-Bu})_2\text{In}(\text{CH}_2\text{CMe}_3)_3\text{SePh}$. Stoichiometric amounts of $[(\text{Me}_3\text{CCH}_2)_2\text{InP}(\text{t-Bu})_2]_2$ ¹⁰ (0.251 g, 0.311 mmol) and $[\text{In}(\text{CH}_2\text{CMe}_3)_2\text{SePh}]_2$ ¹¹ (0.258 g, 0.312 mmol) were combined in pentane (25 mL) at room temperature and stirred for 2 h. The product was filtered and then recrystallized from pentane to yield X-ray quality crystals

of $(\text{Me}_3\text{CCH}_2)_2\text{InP}(\text{t-Bu})_2\text{In}(\text{CH}_2\text{CMe}_3)_3\text{SePh}$ (0.392 g, 0.481 mmol, 77% based on $[(\text{Me}_3\text{CCH}_2)_2\text{InP}(\text{t-Bu})_2]_2$). Mp: 144 - 145 °C. Anal. Calcd for $\text{C}_{29}\text{H}_{56}\text{In}_2\text{PSe}$: C, 50.08; H, 8.28. Found: C, 50.08; H, 8.38. ^1H NMR (C_6D_6) δ 1.27 (s, 36 H, InCCCH_3), 1.35 (d, 18 H, PCCH_3 , $J = 14$ Hz), 1.45 (s, 8H, InCH_2 , $w_{1/2} = 6$ Hz), 6.91 (t, 1 H, p -H of Ph, $^3J = 7$ Hz, $^4J = 2$ Hz), 6.98 (tm, 2 H, m -H of Ph, $^3J = 7$ Hz, $^4J = 2$ Hz), 7.67 (dm, 2 H, o -H of Ph, $^3J = 7$ Hz, $^4J = 1$ Hz). ^1H NMR ($\text{C}_6\text{D}_5\text{CD}_3$) δ 1.26 (s, InCCCH_3), 1.36 (d, $J = 13$ Hz, PCCH_3), 1.45 (s, InCH_2), 6.90 (t, p -H of Ph, $J = 8$ Hz), 6.97 (t, m -H of Ph, $J = 7$ Hz), 7.62 (d, o -H of Ph, $J = 8$ Hz). ^1H NMR (-80 °C, $\text{C}_6\text{D}_5\text{CD}_3$) δ 1.25 (d, $J = 13$ Hz, PCCH_3), 1.29 (s, InCCCH_3), 1.46 (s, InCH_2), 6.75 (t, p -H of Ph), 6.85 (m, m -H of Ph), 7.64 (d, o -H of Ph, $J = 7$ Hz). $^{31}\text{P}\{^1\text{H}\}$ NMR (C_6D_6) δ 44.2 (s, PCCH_3). ^{31}P NMR (C_6D_6) δ 44.2 (m, PCCH_3 , $J = 13$ Hz). $^{31}\text{P}\{^1\text{H}\}$ NMR ($\text{C}_6\text{D}_5\text{CD}_3$) δ 46.3 (s, PCCH_3). $^{31}\text{P}\{^1\text{H}\}$ NMR (-80 °C, $\text{C}_6\text{D}_5\text{CD}_3$) δ 45.4 (s, PCCH_3). $^{13}\text{C}\{^1\text{H}\}$ NMR (C_6D_6) δ 33.1 (s, $\text{InCH}_2\text{CMe}_3$), 33.4 (d, PCMe_3 , $J_{\text{CP}} = 5$ Hz), 34.8 (s, $\text{InCH}_2\text{CMe}_3$), 35.2 (d, PCMe_3 , $J_{\text{CP}} = 7$ Hz), 40.0 (d, $\text{InCH}_2\text{CMe}_3$, $J_{\text{CP}} = 11$ Hz), 126.2 (s, p -C of Ph), 128.9 (s, m -C of Ph), 135.5 (s, o -C of Ph). IR (Nujol mull) cm^{-1} 1576 (w), 1358 (vs, sh), 1297 (vw), 1231 (m), 1212 (vw), 1170 (w), 1112 (m), 1093 (w), 1066(w), 1017 (m), 1010 (m), 997 (m), 960 (vw), 926 (vw), 831 (vw), 804 (vw), 731 (vs), 688 (s), 661 (m), 569 (w), 462 (w), 446 (vw), 376 (vw). Cryoscopic molecular weight, formula weight 815.48 (obsd molality, obsd mol wt, association): 0.0512, 814.6, 1.00; 0.0379, 811.1, 0.99; 0.0239, 852.7, 1.05.

Collection of X-ray Diffraction Data for $(\text{Me}_3\text{CCH}_2)_2\text{InP}(\text{t-Bu})_2\text{In}(\text{CH}_2\text{CMe}_3)_3\text{SePh}$.

A crystal of the mixed bridge indium species was selected and sealed into a 0.3 mm thin walled capillary under strictly anaerobic conditions. The capillary was then mounted and

aligned on a Siemens R3m/V single crystal diffractometer. Data collection was performed as described previously.¹³ Details of the data collection appear in Table III.

The crystal belongs to the triclinic crystal system, the space group being $P\bar{1}$ (No. 2). This choice of space group was made on the basis of intensity statistics and was later confirmed by the successful solution of the crystal structure in the centrosymmetric space group $P\bar{1}$. All data were corrected for Lorentz and polarization effects. An empirical absorption correction was applied. A total of 10608 reflections (a complete shell with $2\theta = 5^\circ$ to 45° for Mo $K\alpha$ radiation) were collected and merged ($R_{\text{int}} = 1.69\%$) into a unique set of 5304 reflections.

Solution and Refinement of the Structure. All crystallographic calculations were carried out with the aid of the Siemens SHELXTL PLUS program set.¹⁴ The analytical scattering factors for neutral atoms were corrected for both components ($\Delta f'$ and $i\Delta f''$) of anomalous dispersion.¹⁵

The structure was solved by the use of direct methods. Positional and anisotropic thermal parameters for all non-hydrogen atoms were refined. Hydrogen atoms were placed at calculated positions with $d(\text{C-H}) = 0.96\text{\AA}$ ¹⁶ and the appropriate staggered tetrahedral geometry. The isotropic thermal parameter of each hydrogen atom was fixed equal to the U_{eq} value for that carbon atom to which it was bonded.

Refinement converged with $R = 2.51\%$ and $wR = 2.52\%$ for 344 parameters refined against those 3747 reflections with $|F_o| > 6\sigma(|F_o|)$, and $R = 4.51\%$ for all data. During refinement the structure was corrected for the effects of secondary extinction. This was done by refining the parameter χ in the expression $F^* = F [1 + 0.002\chi F^2/\sin(2\theta)]^{-1/4}$; the final value for χ was $1.4(4) \times 10^{-4}$. Following refinement, the extreme features left on the

difference-Fourier map were a peak of $0.94\text{e}\text{\AA}^{-3}$ and a negative feature of $-0.68\text{e}\text{\AA}^{-3}$. Final atomic coordinates appear in Table IV.

Acknowledgements. This work was supported in part by the Office of Naval Research and by a grant from Eastman Kodak Co. Purchase of the Siemens R3m/V diffractometer was made possible by Grant No. 89-13733 from the Chemical Instrumentation Program of the National Science Foundation.

Supplementary Material Available. Complete listings of interatomic distances and angles, anisotropic thermal parameters and calculated positions for all hydrogen atoms (4 pages). Ordering information is given on any current masthead page.


References

1. Cowley, A. H.; Jones, R. A. *Angew. Chem. Int. Ed. Engl.* **1989**, 28, 1208 and references therein.
2. Holley, W. K.; Wells, R. L.; Shafieezad, S.; McPhail, A. T.; Pitt, C. G. *J. Organomet. Chem.* **1990**, 381, 15.
3. Wells, R. L.; McPhail, A. T.; Pasterczyk, J. W.; Alvanipour, A. *Organometallics* **1992**, 11, 226.
4. Wells, R. L.; Pasterczyk, J. W.; McPhail, A. T.; Johansen, J. D.; Alvanipour, A. *J. Organomet. Chem.* **1991**, 407, 17.
5. Wells, R. L.; Pasterczyk, J. W.; McPhail, A. T.; Alvanipour, A. *Polyhedron* **1992**, 11, 839.
6. Wells, R. L.; Aubuchon, S. R.; Self, M. F.; Jasinski, J. P.; Woudenberg, R. C.; Butcher, R. J. *Organometallics* **1992**, 11, 3370.
7. Wells, R. L.; McPhail, A. T.; Self, M. F. *Organometallics* **1993**, 12, 3363.
8. Wells, R. L.; McPhail, A. T.; Self, M. F. *Organometallics* **1992**, 11, 221.
9. Wells, R. L.; Jones, L. J.; McPhail, A. T.; Alvanipour, A. *Organometallics* **1991**, 10, 2345.
10. Beachley, O. T., Jr.; Chao, S-H. L.; Churchill, M. R.; Lake, C. H. *Organometallics* **1993**, 12,
11. Beachley, O. T., Jr.; Lee, J. C., Jr.; Gysling, H. J.; Chao, S-H. L.; Churchill, M. R.; Lake, C. H. *Organometallics* **1992**, 11, 3144.

12. Shriver, D. F.; Drezdson, M. A. "The Manipulation of Air Sensitive Compounds," Wiley: New York, 1986, p. 38.
13. Churchill, M. R.; Lashewycz, R. A.; Rotella, F. J. Inorg. Chem. 1977, 16, 265.
14. Siemens SHELXTL PLUS Manual, 2nd ed. Siemens Analytical Instruments, Madison, WI, 1990.
15. International Tables for X-Ray Crystallography, Kynoch Press: Birmingham, England, 1974, Vol. 4, pp. 99-101 and 149-150.
16. Churchill, M. R. Inorg. Chem. 1973, 12, 1213.

Table I

Selected Bond Lengths and Bond Angles for



Bond lengths (Å)

In(1)-Se(1)	2.769 (1)	In(1)-P(1)	2.653 (2)
In(1)-C(111)	2.194 (4)	In(1)-C(121)	2.194 (5)
In(2)-Se(1)	2.748 (1)	In(2)-P(1)	2.651 (2)
In(2)-C(211)	2.181 (6)	In(2)-C(221)	2.193 (4)
Se(1)-C(1)	1.922 (5)	P(1)-C(311)	1.895 (5)
P(1)-C(321)	1.891 (5)		

Bond angles (°)


Se(1)-In(1)-P(1)	84.1(1)	Se(1)-In(1)-C(111)	105.9(1)
P(1)-In(1)-C(111)	117.1(1)	Se(1)-In(1)-C(121)	109.4(1)
P(1)-In(1)-C(121)	114.0(1)	C(111)-In(1)-C(121)	119.5(2)
Se(1)-In(2)-P(1)	84.5(1)	Se(1)-In(2)-C(211)	106.7(2)
P(1)-In(2)-C(211)	108.9(2)	Se(1)-In(2)-C(221)	111.0(1)
P(1)-In(2)-C(221)	116.6(1)	C(211)-In(2)-C(221)	122.3(2)
In(1)-Se(1)-In(2)	91.6(1)	In(1)-Se(1)-C(1)	112.6(1)
In(2)-Se(1)-C(1)	101.3(2)	In(1)-P(1)-In(2)	96.4(1)
In(1)-P(1)-C(311)	114.8(1)	In(2)-P(1)-C(311)	112.4(2)
In(1)-P(1)-C(321)	110.1(2)	In(2)-P(1)-C(321)	110.0(2)
C(311)-P(1)-C(321)	112.2(2)	Se(1)-C(1)-C(2)	116.5(4)
Se(1)-C(1)-C(6)	123.7(4)	C(2)-C(1)-C(6)	119.8(5)
C(1)-C(2)-C(3)	119.9(5)	C(2)-C(3)-C(4)	120.5(7)
C(3)-C(4)-C(5)	121.1(8)	C(4)-C(5)-C(6)	119.1(6)
C(1)-C(6)-C(5)	119.7(5)	In(1)-C(111)-C(112)	120.0(3)
In(2)-C(211)-C(212)	124.9(3)	In(1)-C(121)-C(122)	120.7(4)
In(2)-C(221)-C(222)	123.5(3)		

Table II
Comparison of ^1H and ^{31}P NMR Data

$(\text{Me}_3\text{CCH}_2)_2\text{InP}(\text{t-Bu})_2\text{In}(\text{CH}_2\text{CMe}_3)_2\text{SePh}$		$[(\text{Me}_3\text{CCH}_2)_2\text{InSePh}]_2$	$[(\text{Me}_3\text{CCH}_2)_2\text{InP}(\text{t-Bu})_2]_2$
InCCCH ₃	1.26 (s)	1.12 (s)	1.39 (s)
PCCH ₃	1.36 (d)	—	1.46 (t)
InCH ₂	1.45 (s)	1.47 (s)	1.54 (s)
SeC ₆ H ₅	6.91 (tt)	6.94 (m)	—
	6.98 (tm)	—	—
	7.67 (dm)	7.44 (m)	—
^{31}P	44.2	—	55.0

Table III

Experimental Data for the X-Ray Diffraction Study of

Crystal Data

Empirical Formula	$\text{C}_{34} \text{H}_{67} \text{In}_2 \text{P Se}$
Crystal System	Triclinic
Space Group	$\text{P}\bar{1}$
Unit Cell Dimensions	$\underline{a} = 10.069(2) \text{ \AA}$ $\underline{b} = 11.627(4) \text{ \AA}$ $\underline{c} = 18.452(4) \text{ \AA}$ $\alpha = 100.74(2)^\circ$ $\beta = 99.33(2)^\circ$ $\gamma = 103.38(2)^\circ$
Volume	$2016.9(8) \text{ \AA}^3$
Z	2
Formula weight	815.4
Density(calc.)	1.343 Mg/m^3
Absorption Coefficient	1.996 mm^{-1}
F(000)	836

Table III (Cont.)

Data Collection

Diffractionmeter Used	Siemens R3m/V
Radiation	MoK α ($\lambda = 0.71073 \text{ \AA}$)
Temperature (K)	297
Monochromator	Highly oriented graphite crystal
2 θ Range	5.0 to 45.0°
Scan Type	2 θ - θ
Scan Speed	Constant; 2.00°/min. in ω
Scan Range (ω)	0.50° plus K α -separation
Background Measurement	Stationary crystal and stationary counter at beginning and end of scan, each for 25.0% of total scan time
Standard Reflections	3 measured every 97 reflections
Index Ranges	$-10 \leq h \leq 10$, $-12 \leq k \leq 12$ $-19 \leq l \leq 19$
Reflections Collected	10608
Independent Reflections	5304 ($R_{\text{int}} = 1.69\%$)
Observed Reflections	3747 ($F > 6.0(F)$)
Absorption Correction	Semi-empirical
Min./Max. Transmission	0.5447 / 0.7183

Table III (Cont.)

Solution and Refinement

System Used	Siemens SHELXTL PLUS (VMS)
Solution	Direct Methods
Refinement Method	Full-Matrix Least-Squares
Quantity Minimized	$\sum w(F_o - F_c)^2$
Extinction Correction	$\chi = 0.00014(4)$, where $F^* = F [1 + 0.002\chi F^2 / \sin(2\theta)]^{-1/4}$
Hydrogen Atoms	Riding model, fixed isotropic U
Weighting Scheme	$w^{-1} = \sigma^2(F) + 0.0002F^2$
Number of Parameters refined	344
Final R indices (all data)	R = 4.51 %, wR = 3.39 %
R Indices (6 σ data)	R = 2.51 %, wR = 2.52 %
Goodness-of-Fit	1.07
Largest and Mean Δ/σ	0.002, 0.000
Data-to-Parameter Ratio	15.4:1
Largest Difference Peak	0.94 eÅ ⁻³
Largest Difference Hole	-0.68 eÅ ⁻³

Table IV

Final Atomic Parameters for
 $(\text{Me}_3\text{CCH}_2)_2\text{InP}(\text{t-Bu})_2\text{In}(\text{CH}_2\text{CMe}_3)_2\text{SePh}$

	x	y	z	U(eq)
In(1)	610(1)	2779(1)	2719(1)	41(1)
In(2)	3010(1)	5779(1)	2407(1)	38(1)
Se(1)	878(1)	5248(1)	3154(1)	45(1)
P(1)	3058(1)	3494(1)	2348(1)	39(1)
C(1)	-561(5)	5742(5)	2572(3)	50(2)
C(2)	-868(6)	6786(5)	2913(3)	70(3)
C(3)	-1835(8)	7235(7)	2515(5)	94(4)
C(4)	-2519(7)	6644(7)	1819(5)	92(4)
C(5)	-2235(6)	5596(6)	1460(4)	81(3)
C(6)	-1245(5)	5145(5)	1849(3)	58(2)
C(111)	-1239(5)	2048(4)	1791(3)	47(2)
C(112)	-1960(5)	676(4)	1567(3)	54(2)
C(113)	-928(6)	-61(5)	1431(4)	89(3)
C(114)	-3082(6)	397(5)	839(3)	86(3)
C(115)	-2680(6)	297(5)	2186(3)	67(3)
C(121)	719(5)	2011(5)	3719(3)	55(2)
C(122)	121(5)	2511(5)	4390(3)	55(2)
C(123)	-111(7)	1551(6)	4858(3)	90(3)
C(124)	-1283(5)	2762(6)	4119(3)	74(3)
C(125)	1133(6)	3687(5)	4887(3)	74(3)
C(211)	2057(6)	5922(5)	1289(3)	75(3)
C(212)	2292(7)	7032(6)	1015(3)	74(3)
C(213)	3653(8)	7774(9)	1129(5)	186(6)
C(214)	1591(12)	7851(8)	1538(5)	182(7)
C(215)	1430(8)	6863(7)	232(3)	123(4)
C(221)	4809(5)	7192(4)	3140(3)	51(2)
C(222)	4695(6)	8129(5)	3816(3)	59(2)
C(223)	6081(7)	9118(6)	4080(3)	105(3)
C(224)	3567(7)	8723(6)	3600(3)	93(3)
C(225)	4377(7)	7535(6)	4453(3)	89(3)
C(311)	3167(5)	2676(4)	1379(3)	54(2)
C(312)	3227(7)	1368(5)	1352(3)	89(3)
C(313)	4403(6)	3360(6)	1116(3)	91(3)
C(314)	1824(6)	2615(5)	837(3)	69(3)
C(321)	4517(5)	3437(5)	3116(3)	51(2)
C(322)	5959(5)	3966(6)	2976(3)	81(3)
C(323)	4410(6)	2143(5)	3232(3)	74(3)
C(324)	4346(5)	4203(5)	3832(3)	62(2)

* Equivalent isotropic U defined as one third of the trace of the orthogonalized U_{ij} tensor

Captions to Figures

Figure 1. Labelling of atoms in the $(\text{Me}_3\text{CCH}_2)_2\text{InP}(\text{t-Bu})_2\text{In}(\text{CH}_2\text{CMe}_3)_2\text{SePh}$ molecule. [ORTEP II diagram, with all hydrogen atoms omitted.]

Figure 2. The $(\text{C})_2\text{InP}(\text{C})_2\text{In}(\text{C})_2\text{Se}(\text{C})_6$ fragment of the molecule showing the puckered core.

FIG 1





FIG 2



TECHNICAL REPORT DISTRIBUTION LIST - GENERAL

Office of Naval Research (2)*
Chemistry Division, Code 1113
800 North Quincy Street
Arlington, Virginia 22217-5000

Dr. James S. Murday (1)
Chemistry Division, Code 6100
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. Robert Green, Director (1)
Chemistry Division, Code 385
Naval Air Weapons Center
Weapons Division
China Lake, CA 93555-6001

Dr. Elek Lindner (1)
Naval Command, Control and Ocean
Surveillance Center
RDT&E Division
San Diego, CA 92152-5000

Dr. Bernard E. Douda (1)
Crane Division
Naval Surface Warfare Center
Crane, Indiana 47522-5000

Dr. Richard W. Drisko (1)
Naval Civil Engineering
Laboratory
Code L52
Port Hueneme, CA 93043

Dr. Harold H. Singerman (1)
Naval Surface Warfare Center
Carderock Division Detachment
Annapolis, MD 21402-1198

Dr. Eugene C. Fischer (1)
Code 2840
Naval Surface Warfare Center
Carderock Division Detachment
Annapolis, MD 21402-1198

Defense Technical Information
Center (2)
Building 5, Cameron Station
Alexandria, VA 22314

* Number of copies to forward