

REPORT DOCUMENT

AD-A277 486

Form Approved
OMB No. 0704-0188

2

Public reporting burden for this report is estimated to be 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this information collection. Send comments regarding this burden estimate or any other aspect of this information collection, including suggestions for reducing the burden, to Washington Headquarters Service, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302.



Prescribing instructions, searching existing data sources, gathering this current estimate or any other aspect of this information collection, including suggestions for reducing the burden, to Washington Headquarters Service, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302.

1. AGENCY USE ONLY (Leave blank)

2. REPORT DATE

March 23, 1994

3. REPORT TYPE AND DATES COVERED

Reprint

4. TITLE AND SUBTITLE

High Energy Cosmic Ray Modulation in March-June 1991

5. FUNDING NUMBERS

PE 61102F

PR 2311

TA G4

WU 01

6. AUTHOR(S)

M.A. Shea, D.F. Smart, D.B. Swinson*, M.D. Wilson**

7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)

Phillips Lab/GPSG
29 Randolph Road
Hanscom AFB, MA 01731-30108. PERFORMING ORGANIZATION
REPORT NUMBER

PL-TR-94-2065

9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)

10. SPONSORING / MONITORING
AGENCY REPORT NUMBER

*Department of Physics and Astronomy, University of New Mexico,
800 Yale Blvd NE, Albuquerque, NM 87131 **Herzberg Institute of Astrophysics,
National Research Council, Ottawa, Canada K1A 0R6
Reprint from 23rd International Cosmic Ray Conference, Conference Papers, 3,
735-738, 1993

Approved for public release; Distribution unlimited

13. ABSTRACT (Maximum 200 words)

The lowest galactic cosmic ray intensity since the initiation of continuous neutron monitor measurements was recorded in June 1991. Using neutron monitors and both surface and underground muon detectors we show that the solar-terrestrial phenomena between March and June 1991 modulated the galactic cosmic ray intensity above 45 GeV.

DTIC
ELECTE
MAR 29 1994

S E D

DTIC QUALITY ASSURANCE

14. SUBJECT TERMS

Cosmic rays, Neutron monitors, Muon detectors, Solar activity, Cosmic ray modulation

15. NUMBER OF PAGES

4

16. PRICE CODE

17. SECURITY CLASSIFICATION
OF REPORT
UNCLASSIFIED18. SECURITY CLASSIFICATION
OF THIS PAGE
UNCLASSIFIED19. SECURITY CLASSIFICATION
OF ABSTRACT
UNCLASSIFIED20. LIMITATION OF ABSTRACT
SAR

Reprint from 23rd International Cosmic Ray Conference, Conference Papers, 3, 735-738, 1993.

High Energy Cosmic Ray Modulation in March-June 1991

M.A. Shea¹, D.F. Smart¹, D.B. Swinson², and M.D. Wilson³

¹Space Physics Division (GPSG), Geophysics Directorate/PL
Hanscom AFB, Bedford, MA 01731-3010, U.S.A.

²Department of Physics and Astronomy, University of New Mexico,
800 Yale Blvd., NE, Albuquerque, NM 87131, U.S.A.

³Herzberg Institute of Astrophysics, National Research Council,
Ottawa, Canada K1A 0R6

ABSTRACT

The lowest galactic cosmic ray intensity since the initiation of continuous neutron monitor measurements was recorded in June 1991. Using neutron monitors and both surface and underground muon detectors we show that the solar-terrestrial phenomena between March and June 1991 modulated the galactic cosmic ray intensity above 45 GeV.

1. INTRODUCTION

A series of Forbush decreases were recorded at the earth in March through mid-June 1991. These large decreases were sufficiently frequent that the galactic cosmic ray intensity at the earth reached the lowest value recorded since routine neutron monitor measurements commenced in 1952. Figure 1 displays the daily average cosmic ray intensity from March through June 1991 as recorded by three cosmic ray detectors. Table 1 lists these detectors and their threshold and median rigidity responses.

TABLE 1. Cosmic Ray Detectors

Name	Detector	Threshold Rigidity	Median Rigidity
Inuvik	NM	1 GV	~ 9 GV
Inuvik	Muon	5 GV	~ 50 GV
Embudo	UG Muon	19 GV	~ 145 GV
Socorro	UG Muon	45 GV	~ 305 GV

2. SOLAR AND GEOMAGNETIC ACTIVITY

The sun was extremely active from March through June 1991 with 38 solar flares of optical importance $\geq 2B$ and 29 solar flares with a soft X-ray flux magnitude $\geq 1X$. Most of the major events occurred in episodes of activity in March and June.

Major solar flares that occur near the central meridian of the sun (as viewed from the earth) often generate an interplanetary magnetic shock which, if the interplanetary conditions are favorable, interacts with the earth's magnetic field resulting in a geomagnetic storm. The arrival of an interplanetary shock is registered as a "Sudden Commencement" geomagnetic disturbance. There were 18 sudden commencement geomagnetic disturbances recorded at the earth during March-June 1991. This is another indicator of an extremely perturbed heliosphere, particularly in the immediate vicinity of the earth.

During major geomagnetic storms, the cosmic ray measurements often record a Forbush decrease - a rapid attenuation of the intensity as the interplanetary disturbance envelops and passes beyond the earth. The effect of these Forbush decreases may last for several days. Figure 1 reveals three major periods of cosmic ray attenuation: March, April and the end of May through the first part of June.

Accession For	
NTIS CRA&I	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By _____	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
A-1	20

SPG

94-09569

94 3 28 117




Figure 1. Daily averages of the cosmic ray counting rates during March through June 1991. The scale factor for the Embudo telescope is 128; the scale factor for each of the Inuvik detectors is 100. The threshold rigidity for each detector is identified.

3. MARCH 1991

The March 1991 solar activity was associated primarily with three active regions passing across the solar disk. The largest cosmic ray decrease of this period was on 24 March with the Embudo underground telescope recording a 4% decrease in the hourly values as illustrated in Figure 2. This figure also shows that the unusual spatial modulation present at that time (Shea and Smart, 1993) affected particles in excess of 19 GV. Unfortunately there was a malfunction with the camera at the Socorro underground detector at this critical time (Murphy, 1949).

4. APRIL-MAY 1991

A smaller disturbance, recorded on April 26, is evident in both Inuvik detectors, but is barely discernible in the Embudo record. All three detectors record a small decrease on 17 May although another decrease a few days later is recorded primarily by the neutron monitor.

5. MAY-JUNE 1991

The extended cosmic ray decrease during this period commenced around 28 May - well before solar region 6659 rotated over the eastern limb of the sun. Six sudden commencement geomagnetic disturbances were recorded at the earth from 4-17 June. Figure 3 illustrates the cosmic ray intensity recorded by four detectors from 11-14 June. Taking the cosmic ray intensity values at 19-21 UT on 12 June as a baseline, the cosmic ray intensity decreased by 3, 5, 8, and 17 percent at the Socorro, Embudo, Inuvik muon and Inuvik neutron monitors respectively. The lowest (hourly) intensity recorded by the Inuvik monitor was ~40% below the monthly average of May 1965 which was the highest monthly value between cycles 19 and 20.




Figure 2. Relative changes in the cosmic ray intensity on 24-25 March 1991 as recorded by three detectors. The intensity between 02-03 UT on 24 March was taken as the 100% level.

5. SUMMARY

We have presented data showing that the solar activity and interplanetary disturbances in March-June 1991 combined to establish an all-time cosmic ray low as recorded by neutron monitors (since the start of routine monitoring in 1952). The interplanetary disturbances in June were sufficient turbulent to appreciably attenuate the cosmic ray intensity above 45 GeV.




Figure 3. Cosmic ray counting rates on 11-14 June 1991. The scale factor for the Socorro telescope is 32, for the Embudo telescope is 128; the scale factor for each of the Inuvik detectors is 100. The threshold rigidity for each detector is identified.

REFERENCES

- Murphy, E.: 1949, As quoted in the book, Murphy's Law, by A. Bloch, Price/Stern/Sloan Publishers, Inc., Los Angeles.
- Shea, M.A. and Smart, D.F.: 1993, 23rd ICRC, these proceedings.