

AD-A280 333

ENTATION PAGE

Form Approved
OMB No. 0704-0188

①

estimated average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering the collection of information, Send comments regarding this burden estimate or any other aspect of this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Avenue, Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE 1994		3. REPORT TYPE AND DATES COVERED Journal article	
4. TITLE AND SUBTITLE Low risk of sexual transmission of hepatitis C virus in Somalia				5. FUNDING NUMBERS PE - 62787A PR - 3M162787A870 TA - AR WC - 1288	
6. AUTHOR(S) Watts DM, Corwin AL, Omar MA, Hyams KC				7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Medical Research Institute Commanding Officer 8901 Wisconsin Avenue Bethesda, Maryland 20889-5607	
8. PERFORMING ORGANIZATION REPORT NUMBER NMRI 94-18				9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) Naval Medical Research and Development Command National Naval Medical Center Building 1, Tower 12 8901 Wisconsin Avenue Bethesda, Maryland 20889-5606	
10. SPONSORING / MONITORING AGENCY REPORT NUMBER DN243536				11. SUPPLEMENTARY NOTES Reprinted from: Transactions of the Royal Society of Tropical Medicine and Hygiene 1994 Vol.88 pp.55-56	
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution is unlimited.				12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words)				Accession For	
				NTIS CRA&I ☒	
				DTIC TAB ☐	
				Unannounced ☐	
				Justification	
				By Distribution /	
				Availability Codes	
				Dist Avail and / or Special	
				A-1 20	
14. SUBJECT TERMS sexually transmitted diseases, viral hepatitis, hepatitis C, epidemiology, serology				15. NUMBER OF PAGES 2	
				16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified		18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified		19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	
				20. LIMITATION OF ABSTRACT Unlimited	

94-18470



380

94 6 14 145

Low risk of sexual transmission of hepatitis C virus in Somalia

Douglas M. Watts¹, Andrew L. Corwin², Mahmoud A. Omar³ and Kenneth C. Hyams¹ ¹Naval Medical Research Institute, Bethesda, Maryland, USA; ²US Naval Medical Research Unit No. 3, Cairo, Egypt; ³Ministry of Health, Mogadishu, Somalia

Abstract

The prevalence in Somalia of antibody to hepatitis C virus (anti-HCV) was determined in a survey of 236 female prostitutes, 80 sexually transmitted disease (STD) clinic patients, 79 male soldiers, and 43 tuberculosis patients. Of 98 (22%) serum samples repeatedly anti-HCV reactive by first and second generation enzyme-linked immunosorbent assay kits, only 8 (1.8%) were anti-HCV positive by immunoblot assay (RIBA-2). Anti-HCV seropositivity by immunoblot assay was not associated with any risk group or with positive syphilis serology (found in 18% of subjects) or antibody to human immunodeficiency virus 1 (in 1.4% of subjects). These data indicate that sexual transmission of hepatitis C virus is not common in Somalia among sexually active populations, including female prostitutes and other groups at high risk of STDs and the acquired immune deficiency syndrome.

Introduction

Hepatitis C virus (HCV) has been found in developed countries to be a common cause of post-transfusion hepatitis and a common infection among groups with frequent parenteral exposure, such as illicit drug abusers (ALTER *et al.*, 1989a; ESTEBAN *et al.*, 1990; VAN DEN HOEK *et al.*, 1990). The role of sexual transmission in the spread of HCV is not as well understood, particularly in developing countries, although HCV does not seem to be transmitted by sexual contact as easily as hepatitis B virus (ALTER *et al.*, 1989b; EVERHART *et al.*, 1990).

In developing countries, HCV infection has not been found as frequently as hepatitis B infection; however, high-risk groups in these countries have not been comprehensively evaluated (AL-FALEH *et al.*, 1991; JACKSON *et al.*, 1991; SAEED *et al.*, 1991). In this investigation, populations living in Somalia and at high risk of sexually transmitted diseases (STDs) and human immunodeficiency virus (HIV) infection were investigated for HCV infection.

Patients and Methods

During 1990, the Somali Ministry of Health surveyed 438 subjects living in 3 major urban centres in Somalia: Mogadishu, Merca, and Chismayu. Subjects included 236 female prostitutes, 80 STD patients (69% male), 79 male military personnel, and 43 patients (58% male) with *Mycobacterium tuberculosis* infection. The mean age of the entire population was 28 years (range 13-79 years). All subjects were chosen sequentially during clinic or hospital visits on the days that the study was conducted; no selection criterion was used.

A serum sample was obtained from all study subjects and tested for total anti-HCV antibody using both first and second generation commercial enzyme-linked immunosorbent assay (ELISA) kits (Abbott Laboratories, Abbott Park, Illinois, USA). Repeatedly reactive sera were further verified with a second-generation immunoblot assay (RIBA-2[®]; Chiron Corporation, Emeryville, California, USA). Sera reactive by both ELISA and RIBA-2 were considered to be positive for anti-HCV and to represent active infection (FOLLETT *et al.*, 1991). Sera were also screened for HIV-1 antibody by ELISA, and repeatedly reactive samples were confirmed by Western blotting. Lastly, sera were tested for syphilis infection by the rapid plasma reagin card test, with confirmation by the fluorescent treponemal antibody-absorption test (FTA-ABS).

Results

Among 438 study subjects, 83 (18.9%) had serum samples repeatedly reactive for anti-HCV by first-generation ELISA. By second generation ELISA, 74 (16.9%) sera were repeatedly reactive for anti-HCV: 59 serum samples

which were also reactive by first-generation assay and 15 additional sera which were negative by first-generation ELISA. Among all 98 repeatedly ELISA-reactive sera, only 8 (1.8%) were anti-HCV positive and 6 (1.4%) were indeterminate by RIBA-2. All 8 RIBA-positive sera were identified by the second generation ELISA; however, 4 of the 8 were missed by the first-generation ELISA. Sera from 79 subjects (18.0%) were positive by FTA-ABS and 6 (1.4%) were positive for HIV-1 antibody by Western blotting.

There was no significant association between HCV infection, determined by RIBA-2, and the age, sex, or risk group of study subjects (Table). The mean age of the 8

Table. Prevalence of antibodies to hepatitis C virus and human immunodeficiency virus 1, and positive syphilis serology, among 438 subjects living in Somalia and at high risk of sexually transmitted diseases and AIDS

Risk group	No. examined	Number positive (percentages in parentheses)		
		Anti-HCV ^a	Anti-HIV-1 ^b	FTA-ABS ^c
Female prostitutes	236	4 (1.7)	5 (2.1)	73 (30.9)
STD clinic patients	80	2 (2.5)	0 (-)	3 (3.8)
Military personnel	79	1 (1.3)	1 (1.3)	3 (3.8)
Tuberculosis patients	43	1 (2.3)	0 (-)	0 (-)

^aImmunoblot for anti-hepatitis C antibody.

^bWestern blot for human immunodeficiency virus 1.

^cFluorescent treponemal antibody absorption test.

subjects positive for anti-HCV was 32 years, compared to a mean age of 28 years for the other study subjects. None of the anti-HCV positive subjects had HIV-1 antibody, and only 3 of 8 subjects with anti-HCV were FTA-ABS positive.

Discussion

Based on currently available tests for anti-HCV, these data indicate that sexual transmission of HCV is not common in Somalia among sexually active populations, including female prostitutes and individuals with anti-HIV-1 antibody and positive syphilis serology. A low risk of HCV infection has been found in other prostitute populations in developing countries, suggesting that HCV is not readily transmitted by sexual contact even among highly sexually active groups (HYAMS *et al.*, 1992).

These data also indicate that false-positive anti-HCV ELISA serological results are common in the Somali population and that the second generation ELISA assay is much better than the first generation assay for identifying immunoblot positive sera. Other studies of African populations have found a high prevalence of false-positive anti-HCV ELISA results, which has been attributed to malaria or flavivirus infection, cross-reactive antibody to unknown antigens, and infection by HCV variants (ACETI

Address for correspondence: Dr K. C. Hyams, Naval Medical Research Institute, 12300 Washington Avenue, Rockville, MD 20852, USA.

et al., 1990; WONG et al., 1990; CHAN et al., 1991; ELLIS et al., 1991; JACKSON et al., 1991; TIBBS et al., 1991; HYAMS et al., 1993).

Our results suggest that HIV-1 infection was not common in Somalia at the time this study was conducted. A prior study in Somalia indicated that HIV-1 infection was not then widespread in that region, possibly because of minimal commerce between Somalia and the rest of Africa (SCOTT et al., 1991).

Acknowledgements

This research was supported by the US Naval Medical Research and Development Command, Bethesda, Maryland, USA, Work Units no. 3M463105H29AA335 and 3M162787A870AR1288. The opinions and assertions herein are the private ones of the authors and are not to be construed as official or as reflecting the views of the US Navy Department, the Department of Defense, or the Somali Government.

References

- Aceti, A., Taliani, G., Bac, C. & Sebastiani, A. (1990). Anti-HCV false positivity in malaria. *Lancet*, 336, 1442-1443.
- Al-Faleh, F. Z., Ayoola, E. A., Al-Jeffry, M., Al-Rashed, R., Al-Mofarreh, M., Arif, M., Ramia, S., Al-Karawi, M. & Al-Shabrawy, M. (1991). Prevalence of antibody to hepatitis C virus among Saudi Arabian children: a community-based study. *Hepatology*, 14, 215-218.
- Alter, H. J., Purcell, R. H., Shih, J. W., Melpolder, J. C., Houghton, M., Choo, Q.-L. & Kuo, G. (1989a). Detection of antibody to hepatitis C virus in prospectively followed transfusion recipients with acute and chronic non-A, non-B hepatitis. *New England Journal of Medicine*, 321, 1494-1500.
- Alter, H. J., Coleman, P. J., Alexander, W. J., Kramer, E., Miller, J. K., Mandel, E., Hadler, S. C. & Margolis, H. S. (1989b). Importance of heterosexual activity in the transmission of hepatitis B and non-A, non-B hepatitis. *Journal of the American Medical Association*, 262, 1201-1295.
- Chan, S.-W., Simmonds, P., McOmish, F., Yap, P.-L., Mitchell, R., Dow, B. & Follett, E. (1991). Serological responses to infection with three different types of hepatitis C virus. *Lancet*, 338, 1391.
- Ellis, L. A., Brown, D., Conradie, J. D., Paterson, A., Sher, R., Mollo, J. & Dusheiko, G. M. (1991). Regional prevalence of hepatitis C antibodies in South Africa: an analysis of fresh and stored serum. In: *Viral Hepatitis and Liver Disease*, Hollinger, F. B., Lemon, S. M. & Margolis, H. S. (editors). Baltimore: Williams and Wilkins, pp. 445-447.
- Esteban, J. I., Gonzalez, A., Hernandez, J. M., Viladomiu, L., Sanchez, C., Lopez-Talavera, J. C., Lucea, D., Martin-Vega, C., Vidal, W., Esteban, R. & Guardia, J. (1990). Evaluation of antibodies to hepatitis C virus in a study of transfusion-associated hepatitis. *New England Journal of Medicine*, 323, 1107-1112.
- Everhart, J. E., Di Bisceglie, A. M., Murray, L. M., Alter, H. J., Melpolder, J. J., Kuo, G. & Hoffnagle, J. H. (1990). Risk for non-A, non-B (type C) hepatitis through sexual or household contact with chronic carriers. *Annals of Internal Medicine*, 112, 544-545.
- Follett, E. A. C., Dow, B. C., McOmish, F., Lee Yap, P., Hughes, W., Mitchell, R. & Simmonds, P. (1991). HCV confirmatory testing of blood donors. *Lancet*, 338, 1024.
- Hyams, K. C., Phillips, I. A., Moran, A. Y., Tejada, A. & Wignall, F. S. (1992). Seroprevalence of hepatitis C virus antibody in Peru. *Journal of Medical Virology*, 37, 127-131.
- Hyams, K. C., Okoth, F. A., Tukei, P. M., Vallari, D. S., Morrill, J. C., Long, G., Bansal, J. & Constantine, N. (1993). Inconclusive hepatitis C virus antibody results in African sera. *Journal of Infectious Diseases*, 167, 254-255.
- Jackson, J. B., Guay, L., Goldfarb, J., Olness, K., Ndugwa, C., Mmiro, F., Kataaha, P. & Allain, J.-P. (1991). Hepatitis C virus antibody in HIV-1 infected Ugandan mothers. *Lancet*, 337, 551.
- Saeed, A. A., Alk-Admawi, A. M., Al-Rasheed, A., Fairclough, D., Bacchus, R., Ring, C. & Garson, J. (1991). Hepatitis C virus infection in Egyptian volunteer blood donors in Riyadh. *Lancet*, 338, 459-460.
- Scott, D. A., Corwin, A. L., Constantine, N. T., Omar, M. A., Guled, A., Yusef, M., Roberts, C. R. & Watts, D. M. (1991). Low prevalence of human immunodeficiency virus-1 (HIV-1), HIV-2, and human T cell lymphotropic virus-1 infection in Somalia. *American Journal of Tropical Medicine and Hygiene*, 45, 653-659.
- Tibbs, C. J., Palmer, S. J., Coker, R., Clark, S. K., Parsons, G. M., Hojvat, S., Peterson, D. & Banatvala, J. E. (1991). Prevalence of hepatitis C in tropical communities: the importance of confirmatory assays. *Journal of Medical Virology*, 34, 143-147.
- Van den Hoek, J. A. R., van Haastrecht, H. J. A., Goudsmit, J., de Wolf, F. & Coutinho, R. A. (1990). Prevalence, incidence, and risk factors of hepatitis C virus infection among drug users in Amsterdam. *Journal of Infectious Diseases*, 162, 823-826.
- Wong, D. C., Diwan, A. R., Rosen, L., Gerin, J. L., Johnson, R. G., Polito, A. & Purcell, H. (1990). Non-specificity of anti-HCV test for seroepidemiological analysis. *Lancet*, 336, 750-751.

Received 16 March 1993; revised 13 May 1993; accepted for publication 13 May 1993

Announcement

Second Tropical Neurology Congress

Limoges, France 21-23 September 1994

The official languages of this congress will be French and English. Registration fee: FF 1500 (US \$ 250) before 1 March 1994; FF 2000 (US \$ 325) after that date.

Further information from: Prof. M. Dumas, Institut d'Epidémiologie et de Neurologie Tropicale, Faculté de Médecine, 2 Rue du Docteur Marcland, 87025 Limoges Cedex, France. Telephone: (33) 55 43 58 20; fax: (33) 55 43 58 01.