

Massachusetts Institute of Technology

AD-A281 459



FINAL REPORT

DTIC
ELECTE
JUL 12 1994
S F D

Grant No.: N00014-93-1-0754

Title: "Sensitivity Study of the SPEM Model to the Assimilation of Synop Datasets
Using Two Assimilation Schemes: (1) Nudging (2) The Kalman
Filter/Smoother"

Period of Performance: May 1, 1993 through December 31, 1993

Total Amount: \$23,433

Prof. Paola Malanotte-Rizzoli
TEL: 617 253 2451 FAX: 617 253 4464
MIT, 54-1416, Cambridge, MA. 02139

This document has been approved
for public release and sale; its
distribution is unlimited

DTIC QUALITY INSPECTED 8

From two CTD surveys taken two months apart during the 1981 Ocean Tomography experiment southwest of Bermuda, a time-dependent ocean state is estimated that evolves from the first survey to the second survey.

Mesoscale motions in this region are modelled by quasi-geostrophic dynamics. The model domain is a doubly periodic 600km x 600km square centred at 26°N, 70°W. The oceanic density field is expanded in terms of a horizontal Fourier series with vertical modes. Freely propagating Rossby waves in the first and second baroclinic modes are fit to the data in a least-squares sense at the two survey times.

Linear dynamics are first used to evolve the waves in time, and are found to be too slow to evolve the ocean state to fit both surveys. Weakly nonlinear interactions between the waves are then allowed and are found to do a better job. The wave-wave interactions enable Rossby waves in the barotropic mode to enter into the baroclinic Rossby wave evolution as unknown control variables which can then be estimated.

Three techniques from optimal control theory are developed and applied to this estimation problem: the Kalman filter/smoothen, the adjoint method and dynamic programming. These methods produce estimates of the baroclinic Rossby wave amplitudes and the barotropic Rossby wave amplitudes that are consistent with the data and the weakly nonlinear dynamics. The three methods all produce the same results. The adjoint method is fastest, the Kalman filter/smoothen is slowest but is the only method that produces error estimates.

Assimilating a baroclinic wave model with CTD data, it is possible to estimate a barotropic flow that is consistent with the model and the data. The determination of the barotropic flow from observations has long been a frustrating problem for oceanographers. Here is presented a new approach to solving this problem.

Paola Malanotte-Rizzoli

Paola Malanotte-Rizzoli

Accession For	
NTIS CRA&I	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By <i>per ltr</i>	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
A-1	

94-21048



286

~~94 5 24 043~~