




REPORT DOCUMENTATION PAGE

Form Approved
 OMB No. 0704-0188

1a. REPORT SECURITY CLASSIFICATION Unclassified		1b. RESTRICTIVE MARKINGS	
2a. SECURITY CLASSIFICATION AUTHORITY		3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution is unlimited	
2b. DECL AD-A285 753		5. MONITORING ORGANIZATION REPORT NUMBER(S)	
4. PERFC 			

6a. NAME OF PERFORMING ORGANIZATION Rensselaer Polytechnic Institute	6b. OFFICE SYMBOL (if applicable) Chemical Eng.	7a. NAME OF MONITORING ORGANIZATION ONR, Code 333
6c. ADDRESS (City, State, and ZIP Code) Troy, NY 12180-3590		7b. ADDRESS (City, State, and ZIP Code) 800 North Quincy St., Room 504 Arlington, VA 22217

8a. NAME OF FUNDING/SPONSORING ORGANIZATION ONR, Code 333	8b. OFFICE SYMBOL (if applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER N-00014-94-1-0097	
8c. ADDRESS (City, State, and ZIP Code) 800 North Quincy St., Room 504 Arlington, VA 22217		10. SOURCE OF FUNDING NUMBERS	
PROGRAM ELEMENT NO. 601153N	PROJECT NO. 02402	TASK NO. 010	WORK UNIT ACCESSION NO.

11. TITLE (Include Security Classification)
 Gas-Phase Kinetics Measurements for Underwater Explosives

12. PERSONAL AUTHOR(S)
 Arthur Fontijn

13a. TYPE OF REPORT Annual	13b. TIME COVERED FROM 10/15/93 TO 10/14/94	14. DATE OF REPORT (Year, Month, Day) 10/11/94	15. PAGE COUNT 4
-------------------------------	--	---	---------------------

16. SUPPLEMENTARY NOTATION

17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB-GROUP	Explosives	Reaction Kinetics
			Fluoro-Nitramino	BO
			Boron	

19. ABSTRACT (Continue on reverse if necessary and identify by block number)

To aid in the development of boron-enhanced fluoro-nitramino explosives, the kinetics of individual reactions from those systems are investigated over wide temperature ranges. For $BO + O_2 \rightarrow BO_2 + O$ the measurements agree with earlier observation in the 400 - 1000 K range. No reaction could be observed between BO and CO_2 from 780 to 1200 K, indicating k less than $1 \times 10^{-15} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$. For $BO + N_2O \rightarrow BO_2 + N_2$ $k(305-920 \text{ K}) = 3 \times 10^{-13} \exp(-1310 \text{ K/T}) \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ was obtained.

94-32548

20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS		21. ABSTRACT SECURITY CLASSIFICATION Unclassified	
22a. NAME OF RESPONSIBLE INDIVIDUAL Dr. Richard S. Miller		22b. TELEPHONE (Include Area Code) (703) 696-4404	22c. OFFICE SYMBOL 333

PROGRESS

Our goals are:

- (i) to make accurate measurements, over wide temperature ranges, of rate coefficients for boron combustion in C/H/N/O/F environments, of importance to the development of new underwater explosives, and
- (ii) to use the measurements to obtain a further understanding, to allow predictions for additional reactions occurring with such explosives.

The measurements are made by laser-induced fluorescence, LIF, in high temperature reactors. Our efforts in this first year have concentrated on the development of methods for introducing BO in HTFFRs (high-temperature fast-flow reactors). A large number of methods were investigated. The best are: i) those by which B atoms are produced in a microwave discharge, followed by reaction with O atoms or SO₂, and ii) those where LiBO₂, HBO₂, or B₂O₃ are vaporized and passed through a microwave discharge. However, all led to rate coefficients somewhat decreasing with residence time. This appears to be due to rapid wall dimerization of BO. Nonetheless, we obtained measurements on $\text{BO} + \text{O}_2 \rightarrow \text{BO}_2 + \text{O}$ between 400 and 1000 K. These are in essential agreement with earlier measurements¹ on this reaction of $k(300\text{-}1000 \text{ K}) = 7.0 \times 10^{-12} \exp(255 \text{ K}/T) \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$. There a reactor fashioned after our real-time MHTP (metals high-temperature photochemistry) apparatus was used. Since further BO reactions show the same problem, we will use the HTFFR to determine approximate rate coefficients and make accurate measurements in the MHTP apparatus, which provides a wall-less reaction environment. The HTFFR equipment is still required for measurements from about 1000 to 1800 K, where the BO production method developed¹ for the photo-initiated reaction can not be used.

Two sets of such approximate measurements have already been made. The first of these is for the $\text{BO} + \text{CO}_2 \rightarrow \text{BO}_2 + \text{CO}$ reaction. While at room temperature a slow pressure-dependent reaction was observed, from 780 to 1200 K (the highest temperature explored thus far) no reaction was found. This indicates rate coefficient values less than about $1 \times 10^{-15} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$. This is four orders of magnitude

lower than obtained for the $\text{BO} + \text{O}_2$ reaction. Even more importantly, it is two to four orders of magnitude lower than the gestimated values in the current models.²⁻⁴ While the $\text{BO} + \text{CO}_2$ reaction and its reverse are considered to be highly sensitive for boron combustion in hydrocarbon systems,^{2,3} they are not regarded so for the envisioned fluoro-nitramino based explosives.⁴ However, such a large difference is important for those as well, once again illustrating the importance of measurements for the development of realistic models.

The measurements on the $\text{BO} + \text{N}_2\text{O} \rightarrow \text{BO}_2 + \text{N}_2$ reaction have yielded $k(305-920 \text{ K}) = 3 \times 10^{-13} \exp(-1310 \text{ K/T}) \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$.

PLANS

Preparations for work on the $\text{BO} + \text{HCl}$ and BF_3 reactions have started. Once those measurements have been made below 1200 K, work on this reaction and the O_2 and CO_2 reaction above 1200 K in the HTFFR will start. (The N_2O study is restricted to the 920 K limit because of thermal instability.) For such higher temperature work BO LIF measurements will be made at shorter wavelengths, to reduce potential wall radiation interference.

We also plan to test the compatibility of our reactor materials with HF at high temperatures for measurements with that crucial reactant. For the study of the other boron species of ONR interest, BO_2 and BF , no similar problems as encountered with BO are expected and the HTFFR technique will, at least initially, be used for their study. Additionally to the above two high-temperature techniques, where laser-induced fluorescence is used for monitoring of the boron species, our HTFFR-mass spectrometer will be used for product identification. Especially product OBF detection would be important.

PARTICIPANTS AND CONTACTS

The experimental work has been performed by G.T. Dalakos with help from D.P. Belyung. In addition to these two students, W.F. Flaherty, our group technician, has participated.

We thank R.A. Yetter of Princeton University and M. Soto and L. Pasternack of the Naval Research Laboratory for helpful discussions on the kinetics aspects of the Navy program in underwater explosives.

PUBLICATIONS AND AWARDS

None this year.

REFERENCES

1. C.T. Stanton, N.L. Garland, and H.H. Nelson, "Temperature Dependence of the Kinetics of the Reaction $BO + O_2$," *J. Phys. Chem.* **95**, 8741 (1991).
2. R.C. Brown, C.E. Kolb, S.Y. Cho, R.A. Yetter, H. Rabitz, and F.L. Dryer, "Kinetics of High Temperature, Hydrocarbon Assisted Boron Combustion," *Gas-Phase Metal Reactions*, A. Fontijn, Ed., Elsevier, Amsterdam, 1992, Chap. 27
3. L. Pasternack, "Gas-Phase Modeling of Homogeneous Boron/Oxygen/Hydrogen/Carbon Combustion," *Combust. Flame* **90**, 259 (1992).
4. R.C. Brown, C.E. Kolb, R.A. Yetter, F.L. Dryer, and H. Rabitz, "Kinetic Modeling and Sensitivity Analysis for B/H/O/C/F Combustion Systems," *Combust. Flame*, in press.

Available For	
DTIC	<input checked="" type="checkbox"/>
DAAG	<input type="checkbox"/>
DTIC	<input type="checkbox"/>
By	
Date	
Availability Codes	
Dist	Avail and/or Special
A-1	