



**EDGEWOOD**

RESEARCH, DEVELOPMENT & ENGINEERING CENTER

U.S. ARMY CHEMICAL AND BIOLOGICAL DEFENSE COMMAND

ERDEC-TR-226

**INHIBITION OF RABBIT SPERM CELL HYPERACTIVATED MOTILITY  
BY METALLIC IONS IN TOXICOLOGICAL TESTING**



Ronald J. Young  
Dale H. Heitkamp

RESEARCH AND TECHNOLOGY DIRECTORATE

March 1995

Approved for public release; distribution is unlimited.

19950531 065



Aberdeen Proving Ground, MD 21010-5423

DTIC QUALITY INSPECTED 1

#### Disclaimer

The findings in this report are not to be construed as an official Department of the Army position unless so designated by other authorizing documents.

# REPORT DOCUMENTATION PAGE

Form Approved  
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE <b>1995 March</b>	3. REPORT TYPE AND DATES COVERED <b>Final, 90 Nov - 92 Aug</b>	
4. TITLE AND SUBTITLE <b>Inhibition of Rabbit Sperm Cell Hyperactivated Motility by Metallic Ions in Toxicological Testing</b>			5. FUNDING NUMBERS <b>PR-F8J2-10-005</b>
6. AUTHOR(S) <b>Young, Ronald J.; and Heitkamp, Dale H.</b>			
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>DIR, ERDEC,* ATTN: SCBRD-RTL, APG, MD 21010-5423</b>			8. PERFORMING ORGANIZATION REPORT NUMBER <b>ERDEC-TR-226</b>
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)			10. SPONSORING / MONITORING AGENCY REPORT NUMBER
11. SUPPLEMENTARY NOTES <b>*When this study was conducted, ERDEC was known as the U.S. Army Chemical Research, Development and Engineering Center, and the authors were assigned to the Research Directorate.</b>			
12a. DISTRIBUTION / AVAILABILITY STATEMENT  <b>Approved for public release; distribution is unlimited.</b>			12b. DISTRIBUTION CODE
13. ABSTRACT (Maximum 200 words)  <b>Rabbit sperm cells developed hyperactivated motility when incubated in medium M. Hyperactivated but not normal progressive motility was inhibited by lead and cadmium, two metals implicated in fertility dysfunction. A metal of military interest, Cr<sup>+6</sup>, had a similar outcome on rabbit sperm, suggesting that it may also have the same property as cadmium and lead. Two metals, zinc and mercury, without fertility effects did not suppress hyperactivated motility. The rabbit sperm-medium M-hyperactivated motility system may have a role in the <u>in vitro</u> assessment of the antifertility effects of chemicals.</b>			
14. SUBJECT TERMS <b>Medium                      Hyperactivated                      Lead                      Zinc Rabbit                        Motility                                  Cadmium                    Mercury Sperm cells                Fertility                                Chromium</b>			15. NUMBER OF PAGES <b>18</b>
17. SECURITY CLASSIFICATION OF REPORT <b>UNCLASSIFIED</b>			16. PRICE CODE
18. SECURITY CLASSIFICATION OF THIS PAGE <b>UNCLASSIFIED</b>	19. SECURITY CLASSIFICATION OF ABSTRACT <b>UNCLASSIFIED</b>	20. LIMITATION OF ABSTRACT <b>UL</b>	

Blank

PREFACE

The work described in this report was authorized under Project No. F8J2-10-005. This work was started in November 1990 and completed in August 1992.

In conducting the research described in this report, the investigators adhered to the "Guide for the Care and Use of Laboratory Animals," National Institute of Health Publication No. 86-23, 1985, as promulgated by the committee on Revision of the Guide for Laboratory Animal Facilities and Care of the Institute of Laboratory Animal Resources, Commission of Life Sciences, National Research Council (Washington, DC). These investigations were also performed in accordance with the requirements of AR 70-18, Laboratory Animals, Procurement, Transportation, Use, Care, and Public Affairs.

The use of trade names or manufacturers' names in this report does not constitute an official endorsement of any commercial products. This report may not be cited for purposes of advertisement.

This report has been approved for release to the public. Registered users should request additional copies from the Defense Technical Information Center; unregistered users should direct such requests to the National Technical Information Service.


Accession For	
DTIC GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

## QUALITY ASSURANCE

This study, governed by Protocol Number 210910430000, was examined for compliance with Good Laboratory Practices as published by the U.S. Environmental Protection Agency in 40 CFR Part 792 (effective 18 September 1989). The dates of all inspections and the dates the results of those inspections were reported to the Study Director and management were as follows:

<u>Phase Inspected</u>	<u>Date Inspected</u>	<u>Date Reported to Study Director/Management</u>
Videotaping	05 Nov 1991	05 Nov 1991
Final Report	18 July 1994	15 Sept 1994

To the best of my knowledge, the methods described in this report were the methods followed during the study as indicated by the raw data found in the laboratory notebook. The report was determined to be an accurate reflection of the raw data recorded.

  
Kenneth P. Cameron  
Quality Specialist  
Life Sciences Department

15 Sept 1994  
Date

## CONTENTS

1.	INTRODUCTION . . . . .	7
2.	MATERIALS AND METHODS . . . . .	7
2.1	Animals . . . . .	7
2.2	Collection and Purification of Sperm Cells . . . . .	7
2.3	Incubation of Sperm Cells . . . . .	8
3.	RESULTS . . . . .	8
4.	DISCUSSION AND CONCLUSION . . . . .	10
	LITERATURE CITED . . . . .	17

## TABLES

1	Effect of $\text{CdCl}_2$ on Rabbit Sperm Cell Motility . . . . .	9
2	Effect of $\text{PbCl}_2$ on Hyperactivated Motility . . . . .	11
3	Effect of $\text{K}_2\text{Cr}_2\text{O}_7$ on Hyperactivated Motility . . . . .	12
4	Effect of $\text{HgCl}_2$ on Hyperactivated Motility . . . . .	13
5	Effect of $\text{ZnCl}_2$ on Hyperactivated Motility . . . . .	14

# INHIBITION OF RABBIT SPERM CELL HYPERACTIVATED MOTILITY BY METALLIC IONS IN TOXICOLOGICAL TESTING

## 1. INTRODUCTION

The rabbit sperm cell is a suitable system for the in vitro assay of chemical cytotoxicity.<sup>1</sup> Work in progress points to this sperm's potential use for in vitro testing of the fertility effects of chemicals. A recent study shows that T6 is the medium of choice<sup>2</sup> for rabbit sperm cell culture; but, this medium, in common with DM and other media frequently used to culture sperm cells, contains carbonate, phosphate, sulfate, and organic acids, and requires the use of an atmosphere containing 5% CO<sub>2</sub>. These conditions are not suitable for testing volatile compounds and compounds that form insoluble carbonates, phosphates, sulfates, or salts of low solubility with organic acids. In addition, rabbit sperm develop hyperactivated motility only after lengthy incubation in these media, an impediment to the application of the system for an in vitro fertility assay based on the inhibition of hyperactivated motion. A new simple salts medium (medium M) in which rabbit sperm cells develop hyperactivated motility in 0.5-1 hr and which supports cellular motion for 20 hr has been developed.<sup>3</sup> This medium contains only chlorides, glucose, and bovine serum albumin (BSA) and is buffered by TRIS-HCl. To determine the suitability of the medium for in vitro testing, the effects of PbCl<sub>2</sub>, CdCl<sub>2</sub>, HgCl<sub>2</sub>, ZnCl<sub>2</sub>, and K<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub> on the motion of rabbit sperm cells were studied. The metals Pb, Cd, Zn, Hg, and Cr were chosen for study because their effects on male reproduction are either known or are under study, and because their cations form insoluble carbonates, phosphates, or sulfates. The experiments conducted are described herein.

## 2. MATERIALS AND METHODS

### 2.1 Animals.

New Zealand white rabbits were individually housed in standard rabbit cages in a room maintained at 75 °F and 50 ± 10% relative humidity (RH) with a 12-hr light/dark cycle. Standard certified laboratory rabbit chow and water were available ad libitum.

### 2.2 Collection and Purification of Sperm Cells.

Semen was collected, and sperm cells were purified by centrifugation through a discontinuous Percoll gradient previously described.<sup>4</sup> Seminal plasma was also removed by

diluting semen with 2 mL of warm medium and centrifugating it at room temperature for 5 min at 350 g. The pellet was washed once and resuspended in warm medium.

### 2.3 Incubation of Sperm Cells.

Purified cells were resuspended in medium at a concentration of  $5-10 \times 10^6/\text{mL}$  and incubated at  $37^\circ\text{C}$  in air when medium M was used or 5%  $\text{CO}_2$  in air when medium T6 was used. Cells were exposed to three concentrations of the test compounds together with a control without a test compound. The compounds and concentrations were:  $\text{ZnCl}_2$ , 0.0002M, 0.0001M, 0.00005M;  $\text{K}_2\text{Cr}_2\text{O}_7$ , 0.000005M, 0.0000025M, 0.000001M;  $\text{CdCl}_2$ , 0.0001M, 0.00005M, 0.00002M;  $\text{HgCl}_2$ , 0.000001M, 0.0000005M, 0.0000002M, and  $\text{PbCl}_2$ , 0.000025M, 0.00001M, 0.000005M. Each experiment was carried out several times with cells from different rabbits to assess variation in response among rabbits. Bovine serum albumin formed a flocculent precipitate with  $\text{ZnCl}_2$  when used at 3 mg/mL but not at 2 mg/mL. Experiments with this compound were carried out at the lower BSA concentration. At 0, 0.5, 1, 2, and 4 hr after adding the test compounds, a 5-7- $\mu\text{L}$  drop of the cell suspension was placed either into a warm 20- $\mu\text{m}$  deep chamber or a 10- $\mu\text{L}$  drop was put into a warm 20- $\mu\text{L}$  MicroCell slide (Fertility Technologies, Natick, MA). Observation of cellular motility was carried out at 100x under negative phase contrast with an Olympus microscope equipped with a warming stage. Some experiments were videotaped<sup>1</sup> for later analysis.

## 3. RESULTS

### Changes in Motility Characteristics.

Rabbit sperm cells suspended in T6 medium moved either progressively or in circles over a 6-hr period; but, few cells developed hyperactivated motility. Adding  $\text{CdCl}_2$  at 0.0001M did not alter the motility characteristics. Motility was inhibited in 2 hr when the salt was added at a concentration of 0.001M and in 4 hr at a concentration of 0.0005M; but, cells remained motile over 6 hr at a concentration of 0.0002M. A completely different result was obtained when incubation was carried out in medium M. Motility was inhibited in 1 hr by  $\text{CdCl}_2$  at a concentration of 0.0002M, demonstrating that in T6 medium,  $\text{Cd}^{2+}$  is sequestered from the solution even though a precipitate was not readily visible. Hyperactivated motility also developed in medium M, and the extent of hyperactivated motility decreased as a function of time of exposure and concentration of  $\text{CdCl}_2$  (Table 1). The response of cells to  $\text{CdCl}_2$  varied according to the cell source. The low concentrations of  $\text{CdCl}_2$  that inhibited hyperactivated motility had no apparent effect on the motility of non-hyperactivated cells as these cells remained motile after hyperactivated motion ceased.

Table 1. Effect of CdCl<sub>2</sub> on Rabbit Sperm Cell Motility.

Rabbit	Conc.	Incubation Time (Hour)				
		0	0.5	1	2	4
038	0	≥50*,4#	≥60,5	≤45,5	≤35,5	≤20,4
	H		≤50,4	≤20,3	0	0
	M		≥60,4	≤30,4	≤25,3	0
	L		≥60,4	≤40,3	≤25,3	≤20,3
213	0	0	≥15,3	≥30,3	≤5,3	≤2,3
	H		≤10,3	≤1,3	0	0
	M		≥10,3	≤5,3	0	0
	L		≥30,3	≥5,3	0	0
967	0	≥30,3	≥50,3	≤35,4	≥30,4	≤5,3
	H		≤30,3	≤30,3	0	0
	M		≥50,3	≤35,4	≤5,3	0
	L		≥50,4	≥35,4	≤30,3	≤5,3
793	0	0	≤5,3	≥40,4	≤25,4	≤5,3
	H		≤3,3	≤15,3	≤3,3	≤2,3
	M		≤5,3	≤30,3	≥5,3	≤2,3
	L		≥3,3	≤40,3	≤20,3	≥2,3
772	0	0	≤2,3	≥10,4	≥10,3	≤3,3
	H		"	≤5,3	0	0
	M		"	≤10,3	≤3,3	0
	L		"	≤6,3	≥5,3	≤3,3
967	0	≥50,3	≥50,3	≤50,3	≤2,3	≤2,3
	H1		N.M.			

A suspension of sperm cells in medium M was incubated at 37 °C with CdCl<sub>2</sub>. At the indicated time intervals, a drop of the cell suspension was placed into a warm 20 μm deep chamber, and cell motion was observed under negative phase contrast with an Olympus microscope equipped with a heated stage.

\* Percent of motile sperm cells that are hyperactivated based on microscopic observation at 100x under negative phase contrast.

# Quality of motion of total sperm cell population ranked from 1 (low) to 5 (high).

N.M. Non motile

H = 0.1mM, M = 0.05mM, L = 0.002mM; H1 = 0.2mM

Lead chloride did not form a visible precipitate with T6 medium when the chloride was added at a concentration  $<0.0005M$ , and cell motility was not affected in this medium. However, hyperactivated motility was inhibited when sperm cells were incubated in medium M with  $PbCl_2$  (Table 2). Here also, motility of non-hyperactivated sperm cells did not appear to be changed by concentrations of  $PbCl_2$  that inhibited hyperactivated motion. Similar results were obtained when sperm cells in medium M were exposed to  $K_2Cr_2O_7$  (Table 3). As shown in Tables 2 and 3, response to the two compounds differed among rabbits.

The response of sperm cells to  $HgCl_2$  was quite varied. Hyperactivated motility of cells from some rabbits was not inhibited by this compound at a concentration of  $0.000001M$  (Table 4, rabbit 967); whereas, the same concentration completely inhibited motility in others (Table 4, rabbit 213). At a concentration in which cells remained motile,  $HgCl_2$  did not seem to inhibit hyperactivated motility (Table 4, rabbits 213 and 712).

The action of  $ZnCl_2$  was also variable. Motility, hyperactivated or non-hyperactivated, of cells from some rabbits was unaffected. Hyperactivated motility of cells from other rabbits was not reduced after short incubation; but, longer incubation inhibited both types of motion (Table 5). This result may be due, in part, either to the lower concentration of BSA used in experiments with  $ZnCl_2$ , or to BSA denaturation by  $ZnCl_2$ . A precipitate was not visible in the sperm cell suspension; but, denatured BSA, although not visible, may have affected sperm viability.

#### 4. DISCUSSION AND CONCLUSION

Results show that medium M is suitable for use in assessing the effects of metal cations on sperm cell motion. The concentrations of all the metal cations used in the study encompass the ranges found in the body fluids of human males exposed to the metals.<sup>5-8</sup> Lead is suspect as the cause of fertility problems in exposed workers.<sup>5,9,10</sup> At high exposure levels, spermatogenesis is disturbed. Cadmium is a potent testicular poison, and exposure to this metal impairs spermatogenesis. However, this does not adequately explain fertility consequences following exposure to the metals.<sup>9,10</sup> Inhibition of hyperactivated motility demonstrated with the two metals in the present study provides one explanation for fertility effects associated with the metals when motility of seminal sperm cells falls within the normal range. The ability of  $Cr^{+6}$  to inhibit hyperactivated motility suggests that exposure to this metal could result in disturbance of the reproductive function. Epidemiological studies suggest there is cause for

Table 2. Effect of  $PbCl_2$  on Hyperactivated Motility

Rabbit	Conc.	Incubation Time (Hour)				
		0	0.5	1	2	4
772	0	$\leq 5^*, 3^\#$	$\leq 10, 3$	$\leq 25, 3$	$\leq 1, 3$	0
	H		$\leq 1, 3$	$\leq 1, 3$	$\leq 1, 3$	0
	M		$\leq 1, 3$	$\leq 1, 3$	$\leq 1, 3$	0
	L		$\leq 5, 3$	$\leq 3, 3$	$\leq 1, 3$	0
712	0	$\leq 2, 3$	$\leq 15, 3$	$\geq 40, 3$	$\leq 40, 3$	$\leq 5, 3$
	H		$\leq 5, 3$	$\leq 40, 3$	$\leq 5, 3$	$\leq 5, 3$
	M		$\leq 15, 3$	$\leq 40, 3$	$\leq 30, 3$	$\leq 5, 3$
	L		$\leq 15, 3$	$\leq 40, 3$	$\leq 40, 3$	$\leq 5, 3$
864	0	$\leq 2, 2$	$\leq 15, 4$	$\leq 10, 4$	$\leq 2, 3$	0
	H		$\leq 1, 3$	0	0	0
	M		$\leq 5, 3$	$\leq 5, 3$	0	0
	L		$\leq 5, 4$	$\geq 5, 4$	0	0
779	0	$\leq 2, 2$	$\leq 2, 3$	$\leq 10, 3$	$\leq 5, 3$	0
	H		0	0	0	0
	M		-	$\leq 1, 3$	0	0
	L		-	$\leq 3, 3$	0	0
038	0		$\leq 5, 3$	$\geq 50, 5$	$\geq 35, 4$	$\leq 30, 4$
	H		$\leq 5, 2$	$\leq 50, 4$	$\leq 30, 3$	$\leq 20, 3$
	M		$\leq 5, 3$	$\leq 50, 3$	$\leq 30, 3$	$\geq 20, 3$
	L		$\leq 5, 3$	$\leq 50, 4$	$\geq 30, 3$	$\geq 20, 3$

A suspension of sperm cells in medium M was incubated at 37 °C with  $PbCl_2$ . At the indicated time intervals, a drop of the cell suspension was placed into a warm 20  $\mu$ m deep chamber, and cell motion was observed under negative phase contrast with an Olympus microscope equipped with a heated stage.

\* Percent of motile sperm cells that are hyperactivated based on microscopic observation at 100x under negative phase contrast.

# Quality of motion of total sperm cell population ranked from 1 (low) to 5 (high).

H = 0.025mM, M = 0.01mM, L = 0.005mM

Table 3. Effect of  $K_2Cr_2O_7$  on Hyperactivated Motility.

Rabbit	Conc.	Incubation Time (Hour)				
		0	0.5	1	2	4
038	O	$\leq 3^*, 2^\#$	$\geq 50, 3$	$\geq 40, 3$	$\geq 30, 4$	$\geq 30, 4$
	H	-	$\leq 10, 3$	$\leq 5, 3$	$\leq 5, 3$	$\leq 3, 3$
	M	-	$\leq 30, 3$	$\leq 10, 3$	$\leq 25, 3$	$\leq 20, 3$
	L	-	$\leq 40, 3$	$\leq 40, 3$	$\leq 30, 3$	$\leq 30, 4$
793	O	0	$\leq 5, 3$	$\geq 15, 3$	$\leq 15, 3$	0
	H	-	0	$\leq 10, 3$	$\leq 1, 3$	0
	M	-	$\leq 5, 4$	$\leq 10, 3$	$\leq 2, 3$	0
	L	-	$\leq 5, 3$	$\geq 20, 3$	$\leq 5, 3$	0
712	O	$\leq 3, 2$	$\geq 20, 4$	$\geq 20, 4$	$\geq 20, 4$	$\geq 20, 4$
	H	-	0	$\leq 5, 3$	$\leq 5, 3$	0
	M	-	-	$\leq 12, 3$	$\leq 10, 3$	0
	L	-	-	$\leq 15, 3$	$\leq 15, 4$	$\leq 10, 3$
967	O	$\leq 40, 3$	$\geq 50, 4$	$\geq 20, 3$	$\geq 20, 3$	$\geq 10, 3$
	H	-	$\leq 50, 3$	$\leq 30, 3$	$\leq 10, 3$	$\leq 2, 3$
	M	-	$\leq 50, 3$	$\leq 30, 3$	$\geq 10, 3$	$\leq 2, 3$
	L	-	$\leq 50, 3$	$\geq 35, 3$	$\geq 10, 3$	$\leq 5, 3$
772	O	$\geq 20, 4$	$\geq 25, 4$	$\geq 25, 4$	$\geq 15, 4$	$\geq 5, 4$
	H	-	$\leq 5, 4$	$\leq 5, 3$	$\leq 5, 3$	0
	M	-	$\leq 20, 3$	$\leq 20, 3$	$\geq 10, 3$	0
	L	-	$\leq 20, 3$	$\leq 20, 3$	$\geq 10, 3$	$\leq 2, 3$

A suspension of sperm cells in medium M was incubated at 37 °C with  $K_2Cr_2O_7$ . At the indicated time intervals, a drop of the cell suspension was placed into a warm 20  $\mu$ m deep chamber, and cell motion was observed under negative phase contrast with an Olympus microscope equipped with a heated stage.

\* Percent of motile sperm cells that are hyperactivated based on microscopic observation at 100x under negative phase contrast.

# Quality of motion of total sperm cell population ranked from 1 (low) to 5 (high).

H = 0.005mM, M = 0.0025mM, L = 0.001mM

Table 4. Effect of HgCl<sub>2</sub> on Hyperactivated Motility

Rabbit	Conc.	Incubation Time (Hour)				
		0	0.5	1	2	4
864	O	0	≤ 5,3	≥30,3	≤ 5,3	0
	H	-	≤ 1,1	0	0	0
	M	-	≤ 1,3	≤10,3	≤ 3,3	0
	L	-	≥ 5,3	≤ 5,3	≤ 3,3	0
038	O	0	≥60,5	≥60,5	≥35,4	≥30,4
	H	-	≤ 3,2	0	0	0
	M	-	≥30,3	≥10,3	≥10,3	≥10,3
	L	-	≥50,4	≤60,4	≤30,3	≥25,3
213	O	≤ 3*,3#	≥20,4	≥25,3	≥15,3	≥10,3
	H	-	0	0	0	0
	M	-	≤15,4	≤15,3	≥15,3	≤ 5,3
	L	-	≥20,4	≤25,3	≥15,3	≤10,3
712	O	-	≥15,2	≥25,3	≤10,3	0
	H	-	≤10,2	≥ 5,3	≤ 5,3	0
	M	-	≥20,2	≥20,3	≥ 5,3	0
	L	-	≥25,2	≤20,3	≤ 5,3	0
772	O	-	≥25,4	≥10,3	≤10,3	0
	H	-	≥15,3	≤ 3,3	≤ 2,3	0
	M	-	≥20,4	≤ 3,3	≤ 3,3	0
	L	-	≥25,4	≥ 3,3	≥ 3,3	0
967	O	≤ 2,3	≥60,4	≥50,3	≤30,3	0
	H	-	≥60,4	≤50,3	≤20,3	0
	M	-	≥60,4	≤50,3	≤20,3	0
	L	-	≥60,4	≤50,3	≤20,3	0
793	O	-	-	≥30,4	≥20,3	≤ 5,3
	H	-	-	≥30,4	≥20,3	≤ 5,3
	M	-	-	≥30,4	≥20,3	≤ 5,3
	L	-	-	≥30,4	≥20,3	≤ 5,3

A suspension of sperm cells in medium M was incubated at 37 °C with HgCl<sub>2</sub>. At the indicated time intervals, a drop of the cell suspension was placed into a warm 20 μm deep chamber, and cell motion was observed under negative phase contrast with an Olympus microscope equipped with a heated stage.

\* Percent of motile sperm cells that are hyperactivated based on microscopic observation at 100x under negative phase contrast.

# Quality of motion of total sperm cell population ranked from 1 (low) to 5 (high).

H = 0.001mM, M = 0.0005mM, L = 0.0002mM

Table 5. Effect of ZnCl<sub>2</sub> on Hyperactivated Motility

Rabbit	Conc.	Incubation Time (Hour)				
		0	0.5	1	2	4
038	0	≤ 2 <sup>*</sup> , 2 <sup>#</sup>	≥50, 3	≥50, 3	≤20, 3	≤20, 3
	H	-	≥50, 3	≥50, 3	≤20, 3	≤20, 3
	M	-	≥50, 3	≥50, 3	≤20, 3	≤20, 3
	L	-	≥50, 3	≥50, 3	≤20, 3	≤20, 3
972	0	≤ 5, 3	-	≥15, 4	≤ 5, 3	≤ 5, 3
	H	-	-	≤10, 4	≤ 5, 3	≤ 5, 3
	M	-	-	≤10, 4	≤ 5, 3	≤ 5, 3
	L	-	-	≤10, 4	≤ 5, 3	≤ 5, 3
712	0	0	≤20, 3	≥10, 3	≤ 5, 3	0
	H	-	-	≤ 5, 3	≤2, 3	0
	M	-	≤20, 3	≥10, 3	≤ 2, 3	0
	L	-	≤20, 3	≥10, 3	≤ 2, 3	0
793	0	0	≤ 5, 3	≤25, 3	≤20, 3	0
	H	-	≤ 5, 3	≤10, 3	≤10, 3	0
	M	-	≤10, 3	≤10, 3	≤10, 3	0
	L	-	≤20, 3	≤15, 3	≤10, 3	0
772	0	0	≥20, 2	≥20, 3	≥10, 3	0
	H	-	≥20, 3	≥20, 3	≤ 2, 1	0
	M	-	≤20, 3	≥20, 3	≤ 3, 3	0
	L	-	≤20, 3	≥20, 3	≤ 5, 3	0

A suspension of sperm cells in medium M was incubated at 37 °C with ZnCl<sub>2</sub>. At the indicated time intervals, a drop of the cell suspension was placed into a warm 20 μm deep chamber, and cell motion was observed under negative phase contrast with an Olympus microscope equipped with a heated stage.

\* Percent of motile sperm cells that are hyperactivated based on microscopic observation at 100x under negative phase contrast.

# Quality of motion of total sperm cell population ranked from 1 (low) to 5 (high).

H = 0.2mM, M = 0.1mM, L = 0.05mM

concern.<sup>11</sup> Zinc is present in high concentration in seminal plasma of mammals and has little effect on fertility.<sup>12</sup> It is significant that exposure to the metal generally had little consequence for hyperactivated motility of rabbit sperm cells. Hyperactivated motility was not suppressed by  $Hg^{2+}$  either in nonsensitive rabbit sperm cells or at concentrations that did not inhibit motility in sensitive cells, suggesting that at either low levels or with nonsensitive cells, inorganic mercury is relatively benign with respect to fertility impairment. The estimation of the decrease in hyperactivated motility was based on subjective visual observation by the same worker in all experiments, and the trend in hyperactivation suppression was found in the 6-7 replicates carried out with each compound studied. An objective measure of hyperactivated motility, under development, will provide a firmer basis for the above conclusions.

The inhibition of hyperactivated motility by the two metals implicated in the impairment of fertility, the overall lack of effect of another regarded as without impact on fertilization, together with the prediction of the consequence for fertilization by exposure to a fifth, demonstrate the utility of the rabbit sperm cell-medium M-hyperactivated motility system for the in vitro assessment of the fertility effects of chemicals. By adjusting the concentration range of the test compound, an assessment of the cytotoxicity of the chemical may also be obtained in the same experiment using methods previously described.<sup>1</sup>

**Blank**

## LITERATURE CITED

1. Young, R.J., Bodt, B.A., Iturralde, T.G., and Starke, W.C., "Automated Analysis of Rabbit Sperm Motility and the Effect of Chemicals on Sperm Motion Parameters," Mol. Reprod. Dev. Vol. 33, pp 347-356 (1992).
2. Young, R.J., Iturralde, T., and Starke, W.C., In vitro System for the Assessment of the Potential Antifertility Affect of Chemicals, CRDEC-TR-389, U.S. Army Chemical Research, Development and Engineering Center, Aberdeen Proving Ground, MD, July 1992, UNCLASSIFIED Report (AD A259 835).
3. Young, R.J., A New Medium for the Culture of Rabbit Sperm For Toxicology Testing, ERDEC-TR-139, U.S. Army Edgewood Research, Development and Engineering Center, Aberdeen Proving Ground, MD, December 1993, UNCLASSIFIED Report (AD A277 504).
4. Young, R.J., and Starke, W.C., A Procedure for the Removal of Vesicles and Prostate Secretions from Motile Rabbit Sperm Cells, CRDEC-TR-88129, U.S. Army Chemical Research, Development and Engineering Center, Aberdeen Proving Ground, MD, August 1988, UNCLASSIFIED Report (AD A200 288).
5. Stachel, B., Dougherty, U., Lahl, U., Schlösser, M., and Zeschmar, B., "Toxic Environmental Chemicals in Human Semen: Analytical Method and Case Studies," Andrologia Vol. 21, pp 282-291 (1989).
6. Apgar, J., "Zinc and Reproduction," Ann. Rev. Nutr. Vol. 5, pp 43-67 (1985).
7. Bonde, J.P., and Christenen, J.M., "Chromium in Biological Samples from Low-Level Exposed Stainless Steel and Mild Steel Welders," Arch. Environ. Health Vol. 46, pp 225-229 (1991).
8. Chia, S.-E., Ong, C.N., Lee, S.T., and Tsakok, F.H.M., "Blood Concentrations of Lead, Cadmium, Mercury, Zinc, and Copper and Human Semen Parameters," Arch. Androl. Vol. 29, pp 177-183 (1992).
9. Gagnon, C., "The Role of Environmental Toxins in Unexplained Male Infertility," Seminars Reprod. Endocrin. Vol. 6, pp 369-376 (1988).
10. Jockenhövel, F., Bals-Pratsch, M., Bertram, H.P., and Nieschlag, E., "Seminal Lead and Copper in Fertile and Infertile Men," Andrologia Vol. 22, pp 503-511 (1990).

11. Mortensen, J.T., "Risk for Reduced Sperm Quality Among Metal Workers with Special Reference to Welders," Scand. J. Environ. Health Vol. 14, pp 27-30 (1988).

12. Hidioglou, M., and Knipfel, J.E., "Zinc in Mammalian Sperm: A Review," J. Dairy Sci. Vol. 67, pp 1147-1156 (1984).