

90317R01
2ND COPY

LABORATORY SCREENING STUDIES ON
THE BIODEGRADATION OF ORGANICS
IN RMA GROUND WATER

Prepared by
Shell Developmnt Company
A Division of Shell Oil Company
Westhollow Research Center
Houston, Texas

Prepared for
Shell Oil Company/Holme Roberts & Owen
Denver, Colorado

August 1990

**Rocky Mountain Arsenal
Information Center
Commerce City, Colorado**

19950623 070

DTIC QUALITY INSPECTED 5

CONTENTS

	<u>Page</u>
TABLES AND FIGURES.....	ii
ABSTRACT.....	iii
INTRODUCTION.....	1
MATERIALS & METHODS.....	2
AQUIFER SEDIMENT SAMPLES.....	2
MICROCOSM TEST SYSTEM.....	2
MICROBIAL & CHEMICAL ANALYSES.....	3
RESULTS & SUMMARY.....	4
SUBSOIL ANALYSES - PHYSICAL-CHEMICAL PROPERTIES &.....	4
MICROBIAL POPULATIONS	
BIODEGRADATION OF BENZENE (B) & CHLOROFORM (CF).....	4
IN AQUIFER CORES	
FUTURE STUDIES.....	5
REFERENCES.....	5-7

TABLES

	<u>Page</u>
Table 1. Summary of Literature Data on Aerobic..... Biodegradation in Soil	8
Table 2. Subsoil Characteristics.....	9
Table 3. Bacterial Population Distribution in Sediments.....	10
Table 4. Summary of Ground Water Microcosm Results.....	11

FIGURES

	<u>Page</u>
Figure 1. Biodegradation of Benzene at 5.5-7.5 ppm (a)..... and 55-65 ppm (b) in RMA Aquifer Soil and Ground Water	12
Figure 2. Biodegradation of Chloroform at 50-80 ppb (a)..... and 450-650 ppb (b & c) in RMA Aquifer Soil and Ground Water	13

ABSTRACT

Data is presented on the potential for *in situ* bioremediation of organic compounds in ground water at RMA sites. The laboratory results demonstrate that 1) ground water microcosms containing aquifer cores from Basin A Neck can be used to assess biodegradation, and 2) benzene and chloroform (50 ppb - 50 ppm) can degrade at significant rates (2 - 20%/day) in the presence of inorganic nutrients (NH_4^+ and PO_4^{3-}) and dissolved oxygen (≥ 2 ppm). Similar laboratory experiments in soil microcosms are planned to investigate the bioremediation potential for these and other compounds (chlorobenzene, dicyclopentadiene, and bicycloheptadiene) present in the South Tank Farm plume using actual aquifer cores from that plume.

WRC TECHNICAL PROGRESS REPORT RMA 1-90

LABORATORY SCREENING STUDIES ON THE BIODEGRADATION OF
ORGANICS IN RMA GROUND WATER

by

J. P. Salanitro and H. L. Wisniewski

INTRODUCTION

An examination of the literature on the biodegradation of aromatic hydrocarbons such as benzene (B), toluene (T), xylene (X), and chlorobenzene (CB) indicates that these are readily metabolized by a variety of naturally-occurring bacteria and fungi in soil¹. Initial enzymatic attack on the ring is through cis-diol formation (oxygenases), followed by ring opening and oxidation to a dicarboxylic acid derivative. This acid is further metabolized to CO₂ through several pathways (e.g., citric acid cycle) in the cell. Table 1 gives a summary of literature on soil biodegradation of some compounds which have been observed in RMA ground water samples.

BTX concentrations of 50 - 5000 ppb are rapidly depleted at rates of 2 - 70%/day in laboratory soil/ground water microcosm test systems or bacterial cultures isolated from soil. These rates are, however, significantly higher than those obtained (ca. 1%/day) from field estimations using plume modeling techniques.

Recent evidence by Baker et al.² and Chiang et al.³ suggests that the field estimated decays for aromatic hydrocarbons is highly dependent upon the dissolved oxygen (DO) concentration in the ground water and its reaeration rate (via upstream aquifer flow and rainfall recharge events). Nevertheless, these data indicate that subsoils have the capacity to biodegrade high levels of organic compounds and factors such as DO levels, nutrient (NH₄⁺, PO₄³⁻) availability and diversity of subsoil microbial populations are probably important.

Chlorobenzene is most likely degraded similarly to benzene¹ and biotransformation rates of 0.4 - 14%/day have been observed with soil and

sewage isolates.^{6,7,8} There are reports on the biodegradability of chloroform in soil in which 0.3 and 500 ppm was metabolized at rates of 2.5 - 14%/day.^{9,10} No literature data exists on the degradation (soil or microbial cultures) of the cycloalkanes, dicyclopentadiene and bicycloheptadiene, however, they may be metabolized similarly to the cyclodiene insecticides, dieldrin, aldrin, and endrin in which an epoxide, alcohol, ketone or carboxylic acid derivative could be formed.¹¹

The following report describes some preliminary experiments on the biodegradation of benzene and chloroform in RMA ground water. Test tube microcosms were used to evaluate the degradation potential of these compounds in aquifer sediments isolated from the Basin A Neck area at the RMA site. These aquifer sediments are considered representative for measuring biodegradation potential of subsurface sediments and groundwater at the RMA site.

MATERIALS & METHODS

Aquifer Sediment Samples. Borings in the saturated zone around Basin A Neck at RMA were obtained by Morrison Knudsen Environmental Services (MKE) staff using a Waterloo saturated sand sampler described previously by Zapico et al.¹² Polybutyrate Shelby tubes (sterilized with bleach solution), measuring 3 inches by 5 feet were inserted into the stainless steel wireline piston core barrel ("Waterloo sampler"). A hollow stem auger was used to make initial boreholes just below the saturated zone of Basin A neck (Northeast Section 35). At these depths, the auger was replaced by the sampler, forced into the saturated zone and 2 foot aquifer cores retrieved. The samples were designated BANIBR-B (14.2 - 31.5 ft) placed on ice and shipped to Westhollow Research Center (WRC) for use in microcosms. The sample was from an uncontaminated area outside the Basin A plume.

Microcosm Test System. Ground water/subsoil test tube microcosms were used to simulate subsurface conditions in the degradation of benzene and chloroform. Microcosms were prepared by aseptically transferring 10g of inner core (14.2 - 16.2 ft sample) of soil from samples in polybutyrate tubes to sterile serum-stoppered test tubes fitted with Teflon®-lined rubber septa and aluminum crimp seals. Uncontaminated ground water from developed

wells was added (25 - 27ml) to microcosms and the DO's adjusted to 2 or 8 ppm by flushing with N₂ or air for varying periods. DO measurements were made with an Orbisphere Model 2111 BOD probe and Model 2607 meter. Sterile sediment slurries were used as controls and contained gamma-ray irradiated (7.5 Mrad, Neutron Products, Inc., Dickerson, Maryland) soils and filter-sterilized (0.22 micron Millipore) ground water. Duplicate sterile and non-sterile microcosms were prepared for benzene (5, 50 ppm) and chloroform (50, 500 ppb) concentrations. Slurries were placed in an anaerobic glove box (Vacuum Atmospheres Company, Hawthorne, California) containing an inert N₂ (≤ 1 ppm O₂) atmosphere. Inorganic nutrients (N & P) and trace minerals were added to microcosms at final concentrations of 5 - 10 ppm NH₄⁺ and NO₃-N and PO₄-P and 0.3-1 ppm Ca⁺⁺ and Mg⁺⁺. All test slurries were incubated at 12 - 15°C in the anaerobic chamber.

Microbial & Chemical Analyses. Microbial populations in soil and ground water were enumerated by tube extinction dilution methods. Endpoints of serial 1:10 dilutions of soil made into culture media were the highest dilution showing turbidity or microscopically discernible cell growth (aerobes, facultative anaerobes or anaerobes), FeS precipitation from H₂S production (sulfate-reducers), methane formation (methanogens) or no residual NO₃⁻ (denitrifiers). Aerobes were cultured in Trypticase Soy Broth (BBL, Becton, Dickinson & Company) for tube dilution or on R₂A medium¹³ for colony counts. Denitrifiers, sulfate-reducers and methanogens were estimated by culturing in denitrifying medium (0.8% Difco Nutrient Broth, 0.1% NaNO₃ and 0.5% succinate), modified Postgate medium E¹⁴ with 0.1% sodium acetate and Balch medium I¹⁵, respectively. Cultures were incubated at 30°C for 7 - 14 days.

At each sample point, the aqueous portion of microcosms was removed with a needle and syringe inside the anaerobic chamber and one ml sample placed into vials with no headspace. Samples were analyzed for benzene and chloroform by purge and trap gas chromatography¹⁶ methods using a Tekmar purge and trap apparatus and Varian GC (1 and 5 ppb minimum detection level for benzene and chloroform, respectively). This is essentially Method 602 proposed by the U.S.E.P.A. for volatiles in water samples.

RESULTS & SUMMARY

Subsoil Analyses - Physical-Chemical Properties & Microbial Populations. Analyses of four saturated soil cores taken from depths of 14 - 31 ft are given in Table 2 and indicate some heterogeneity in the amount of sand, silt, and clay, with clay comprising 20 - 35%. The amount of available N (NH_4^+ and NO_3^-) and P (PO_4^{3-}) also varied slightly from 3 - 7 ppm N and 15 - 25 ppm P among the cores at different depths. The samples contained low levels of soluble Fe (1 - 8 ppm) but high concentrations of total Fe (1.4 - 1.6%). These levels of available N and P would probably represent sufficient requirements for microbes to degrade 0.005 - 50 ppm benzene and chloroform in ground water (based on a balanced C/N/P cell ratio 25/5/1).

Total microbial populations of aerobes and facultative anaerobes is given in Table 3. The number of aerobic bacteria varied from 10^3 - 10^8 /g wet wt soil and appeared to increase with depth. Facultatively anaerobic bacteria were fewer in numbers (10^3 - 10^4 /g) while more strict anaerobic species (denitrifiers, sulfate-reducers, and methanogens) were essentially non-detectable. The data show that these RMA aquifer soil cores contain predominantly aerobic organisms and that any degradation potential would probably be due to aerobic metabolism.

Biodegradation of Benzene (B) & Chloroform (CF) in Aquifer Cores. Data on the potential of laboratory microcosm systems for estimating the biodegradation of B and CF in ground water are given in Figures 1 & 2 and summarized in Table 4. These represent the degradation profiles in the subsoil incubated at 12 - 15°C in the presence of 5 and 50 ppm of B, and 50 and 500 ppb CF at DO levels of 2 or 8 ppm. The biodegradation of benzene was more rapid at 50 ppm (20%/day) than at 5 ppm (2%/day) in the presence of added nutrients (NH_4^+ -N and PO_4^{3-} -P) and oxygen (DO, 2 or 8 ppm) and were substantially higher than the sterile control depletions.

It is not clear why the B depletion rate at 5 ppm is less than that at 50 ppm, but it may indicate 1) an unusually high nutrient requirement for low concentrations (not present in the experiment) or 2) that there is significant heterogeneity in microbial metabolism between soil core samples or 3) that there is significant soil sorption effects of B at

low concentrations. This effect needs to be confirmed with additional microcosm and nutrient experiments.

The biodegradation of CF at 50 and 500 ppb in ground water indicates that the rates were similar (5 - 10%/day). Although some variability was observed between duplicate microcosms (A & B) these decreases in CF were observed in the presence or absence of added nutrients and at an initial DO of 2 or 8 ppm. The potential for CF to degrade in aerobic ground water, however, may be a new finding since it has been shown that many chlorinated alkanes are primarily metabolized under strict anoxic conditions (absence of detectable O_2) by anaerobic bacteria via a reductive dehalogenation¹⁷ or aerobically in the presence of methane (as alternate co-substrate) to enhance the growth of methanotrophic bacteria in soil.¹⁸

FUTURE STUDIES

Additional experiments are planned with South Tank Farm aquifer cores to further investigate the limits and potential for the aerobic biodegradation of benzene and chloroform plus other organics, such as, chlorobenzene, dicyclopentadiene, and bicycloheptadiene. We expect the biodegradation of benzene to be similar to that observed with Basin A Neck aquifer sediments and groundwater, as well as, at other sites around the country as mentioned in the Introduction section.

REFERENCES

- ¹ Gibson, D. T. (Ed.), Microbial Degradation of Organic Compounds. Marcel Dekker, New York, 1984.
- ² Barker, J. F., G. C. Patrick, and D. Major. (1987). Natural Attenuation of Aromatic Hydrocarbons in a Shallow Sand Aquifer. Ground Water Monitoring Rev. 7:64-71.
- ³ Chiang, C. Y., J. P. Salanitro, E. Y. Chai, J. D. Colthart, and C. L. Klein. (1989). Aerobic Biodegradation of Benzene, Toluene, and Xylene in a Sandy Aquifer - Data Analysis and Computer Modeling. Ground Water. 27:823-830.

- ⁴ Mahadevaiah, B. and G. D. Miller, "Application of Microcosm Technology to Study the Biodegradation Potential of a Subsurface Alluvial Material Exposed to Selected Petroleum Hydrocarbons," Sixth Natl. Symp. & Expos. Aquifer Restoration & Ground Water Monitoring, May, 1986.
- ⁵ Jamison, V. W., R. L. Raymond, and J. D. Hudson, "Biodegradation of High-Octane Gasoline," Proc. 3rd Intl. Biodegradation Symp., Appl. Sci. Publ., 1976, pp. 187-196.
- ⁶ Wilson, J. T., C. G. Enfield, W. J. Dunlap, R. L. Crosby, D. A. Foster, and L. B. Baskin. 1981. Transport and Fate of Selected Organic Pollutants in a Sandy Soil, J. Environ. Qual. 10:501-506.
- ⁷ Haider, K., G. Jagnow, R. Kohnen, and S. U. Lim, "Degradation of Chlorinated Benzenes, Phenols, and Cyclohexane Derivatives by Benzene and Phenol-Utilizing Soil Bacteria Under Aerobic Conditions," In Decomposition of Toxic and Non Toxic Organic Compounds in Soils, R. M. Overcash (Ed.), Ann Arbor Science, pp. 207-223.
- ⁸ Reineke, K. and H-J Knackmuss. (1984). Microbial Metabolism of Haloaromatics: Isolation and Properties of a Chlorobenzene-Degrading Bacterium. Appl. Environ. Microbiol. 47:395-402.
- ⁹ Flatham, P. E. and J. R. Dahlgran. (1982). Environ. Sci. Technol. 16:130.
- ¹⁰ Henson, J. M., M. V. Yates, and J. W. Cochran. (1989). Metabolism of Chlorinated Methanes, Ethanes, and Ethylenes by a Mixed Bacterial Culture Growing on Methane. J. Indust. Microbiol. 4:29-35.
- ¹¹ Matsumura, F. and H. J. Benezet, "Microbial Degradation of Insecticides," In Pesticide Microbiology, Ed. by I. R. Hill and S. J. L. Wright, Academic Press, New York, pp. 623-667.
- ¹² Zapico, M. M., S. Vales, and J. A. Cherry. A Wireline Piston Core Barrel for Sampling Cohesionless Sand and Gravel Below the Water Table. Ground Water Monitor. Rev., Summer, 1987, pp. 74-82.

- ¹³ Reasoner, D. J. and E. E. Geldreich. (1985). A New Medium for the Enumeration and Subculture of Bacteria from Potable Water. *Appl. Environ. Microbiol.* 49:1-7.
- ¹⁴ Postgate, J. R. (1979). *The Sulfate-Reducing Bacteria*. Cambridge University Press, London.
- ¹⁵ Balch, W. E., G. E. Fox, L. J. Magrum, C. R. Woese, and R. S. Wolfe. (1979). Methanogens: Re-Evaluation of a Unique Biological Group. *Microbiol. Rev.* 43:260-296.
- ¹⁶ U. S. Environmental Protection Agency, "Guidelines Establishing Test Procedures for the Analysis of Pollutants; Proposed Regulations," Federal Register. 44:NO. 233, December 3, 1979, pp. 69474-69476.
- ¹⁷ Vogel, T. M., C. S. Criddle and P. L. McCarty. (1987). Transformations of Halogenated Aliphatic Compounds. *Environ. Sci. Technol.* 21:722-736.
- ¹⁸ Henson, J. M., M. V. Yates, J. W. Cochran, and D. L. Shockleford. (1988). Microbial Removal of Halogenated Methanes, Ethanes & Ethylenes in an Aerobic Soil Exposed to Methane. *FEMS Microbiol. Ecol.* 53:193-201.

TABLE 1: SUMMARY OF LITERATURE DATA ON AEROBIC BIODEGRADATION IN SOIL

COMPOUND	CONCENTRATIONS TESTED, ppm	TEST SYSTEM ^{a)}	DEGRADATION %/DAY	REFERENCE
Benzene	1.8 - 2.4	Soil/GW (field)	0.7 - 1.1	2,3
	0.05 - 5	Soil/GW	1.8 - 20	2,3
Toluene	1.8 - 2.6	Soil/GW (field)	1.4	2
	0.05 - 5	Soil/GW	2.5 - 20	3
Xylenes	0.05 - 5	Soil/GW	2.5 - 70	2,3,4
BTX	1 - 56	Various cultures	121.5	5
Chlorobenzene (CB)	1	Soil	1.6	6
	20	Soil	0.4	7
	660	Soil/sewage isolate	14	8
	20	<u>Nocardia/Pseudomonas</u>	6 - 10	7
Chloroform	500	Soil	2.5	9
	0.30	Soil (Methane-stimulated)	14	10
Dicyclopentadiene (DCPD)	___ ^{b)}	___	___	___
Bicycloheptadiene (BCH)	___	___	___	___

a) Most were laboratory test systems unless otherwise indicated; GW = groundwater.

b) ___, no data available.

=====

TABLE 2: SUBSOIL CHARACTERISTICS

=====

PARAMETER	<u>DEPTH (FT)</u>			
	14.2 - 16.2	19.5 - 21.5	24.5 - 26.5	29.5 - 31.5
pH	8.1	7.5	7.8	8.3
SAND/SILT/ CLAY (%)	73/8/19	54/11/35	70/5/25	64/9/27
CLASS	SL	SC	SCL	SCL
% FINES <0.1mm	59	43	42	46
AVAILABLE NUTRIENTS (ppm)				
NH ₄ -N	4.8	5.7	4.1	2.7
NO ₃ -N	2.8	.4	.7	.7
PO ₄ -P	25	16	20	21
SOLUBLE Fe (ppm)	8.1	1.1	1.6	2.5
TOTAL Fe (ppm)	14700	13860	15020	16410

OTHER PROPERTIES - range for all depths

SOLUBLE ANIONS AND CATIONS (ppm)

CL 3.8 - 5.1
 SO₄ 8.1 - 23.2
 Na-K 229 - 327
 Ca-Mg 4.3 - 13.2

BULK DENSITY (g/cc) 1.26 - 1.36

PARTICLE DENSITY (g/cc) 2.52 - 2.75

=====

* Analyzed by Soil Analytical Services, Inc., College Station, Texas, for BANIBR cores.

=====

TABLE 3: BACTERIAL POPULATION DISTRIBUTION IN SEDIMENTS^{a)}

=====

NUMBER OF BACTERIA/GRAM WET. WT.^{b)}

<u>SAMPLE DEPTH (FT)</u>	<u>AEROBES</u>		<u>FACULTATIVE ANAEROBES</u>
	<u>TUBE DILUTION</u>	<u>PLATE COUNT</u>	
	$\times 10^5$		
14.2 - 16.2	0.1 - 1	0.2 - 0.3	0.1
19.5 - 21.5	.01 - .1	.08 - 1	.01 - 1
24.5 - 26.5	100	50 - 80	.01 - 1
29.5 - 31.5	100 - 1000	30 - 100	.1

=====

a) BANBR sample (NE Section 35, Basin A Neck).

b) Number of denitrifiers, sulfate reducers and metanogens were < 100/g
(min. detection level).


 TABLE 4: SUMMARY OF GROUND WATER MICROCOSM RESULTS

<u>COMPOUND</u> (concn)	<u>% DEGRADATION</u> <u>DEGN/DAY^{a)}</u>	<u>COMMENTS</u>
Benzene (50 ppm)	20	DO, 8 ppm, + nutrients ^{b)}
(5 ppm)	2	DO, 2 or 8 ppm, ± nutrients
Chloroform (500 ppb)	5-8	DO, 2 or 8 ppm, ± nutrients
(50 ppb)	11	DO, 2 ppm, ± nutrients

a) Assumes a first-order decay, rate = $\ln(C_o/C_i)/t$

b) Effects observed at dissolved oxygen (DO) levels given and ± nutrients
 (N, NH_4^+ / NO_3^- ; P, PO_4^{3-})

FIGURE 1: BIODEGRADATION OF BENZENE AT 5.5 - 7.5 PPM (A) AND 55 - 65 PPM (B) IN RMA AQUIFER SOIL AND GROUND WATER



+ DO 2 ppm ◇ DO 2 + NUTRIENTS △ DO 8 × DO 8 + NUTRIENTS —








FIGURE 2: BIODEGRADATION OF CHLOROFORM AT 50 - 80 PPB (A) AND 450 - 650 PPB (B & C) IN RMA AQUIFER SOIL AND GROUND WATER



REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE 08/00/90	3. REPORT TYPE AND DATES COVERED	
4. TITLE AND SUBTITLE LABORATORY SCREENING STUDIES ON THE BIODEGRADATION OF ORGANICS IN RMA GROUNDWATER			5. FUNDING NUMBERS	
6. AUTHOR(S) SALANITRO, J.; WISNIEWSKI, H.				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) SHELL DEVELOPMENT COMPANY HOUSTON, TX			8. PERFORMING ORGANIZATION REPORT NUMBER 90317R01	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) SHELL OIL COMPANY DENVER, CO			10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION/AVAILABILITY STATEMENT APPROVED FOR PUBLIC RELEASE; DISTRIBUTION IS UNLIMITED				
12b. DISTRIBUTION CODE				
13. ABSTRACT (Maximum 200 words) THIS REPORT DESCRIBES PRELIMINARY EXPERIMENTS ON THE BIODEGRADATION OF C6H6 AND CHCL3 IN RMA GROUND WATER. TEST TUBE MICROCOSMS WERE USED TO EVALUATE THE DEGRADATION POTENTIAL OF THESE COMPOUNDS IN AQUIFER SEDIMENTS ISOLATED FROM THE BASIN A NECK AREA. THE LABORATORY RESULTS DEMONSTRATE THAT: 1. GROUND WATER MICROCOSMS CONTAINING AQUIFER CORES FROM BASIN A NECK CAN BE USED TO ASSESS BIODEGRADATION 2. C6H6 AND CHCL3 CAN DEGRADE AT SIGNIFICANT RATES IN THE PRESENCE OF INORGANIC NUTRIENTS (NH4 AND P04).				
14. SUBJECT TERMS CHEMICAL ANALYSIS, MICROCOSM TEST SYSTEM				15. NUMBER OF PAGES
				16. PRICE CODE
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED	18. SECURITY CLASSIFICATION OF THIS PAGE	19. SECURITY CLASSIFICATION OF ABSTRACT	20. LIMITATION OF ABSTRACT	

DTIC QUALITY INSPECTED 5