

NASA Contractor Report 198184


ICASE Report No. 95-54



ICASE

MATHEMATICAL ANALYSIS OF THE NAVIER-STOKES EQUATIONS WITH NON STANDARD BOUNDARY CONDITIONS

M. D. Tidriri



Contract No. NAS1-19480
July 1995

19950925 168

Institute for Computer Applications in Science and Engineering
NASA Langley Research Center
Hampton, VA 23681-0001

DTIC QUALITY INSPECTED 8



Operated by Universities Space Research Association

DISTRIBUTION STATEMENT A

Approved for public release;
Distribution Unlimited

Mathematical Analysis of the Navier-Stokes Equations with non standard boundary conditions

M. D. Tidriri *

Institute for Computer Applications in Science and Engineering
NASA Langley Research Center, Hampton, VA 23681-0001

Accession For	
NTIS	CRA&I <input checked="" type="checkbox"/>
DTIC	TAB <input type="checkbox"/>
Unannounced <input type="checkbox"/>	
Justification	
By	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
A-1	

Abstract

One of the major applications of the Domain Decomposition Time Marching Algorithm is the coupling of the Navier-Stokes systems with Boltzmann equations in order to compute transitional flows. Another important application, is the coupling of a global Navier-Stokes problem with a local one in order to use different modelizations and/or discretizations. Both of these applications involve a global Navier-Stokes systems with non standard boundary conditions. The purpose of this work is to prove, using the classical Leray-Schauder theory, that these boundary conditions are admissible and lead to a well posed problem.

*This work was supported by the National Aeronautics and Space Administration under NASA contract NAS1-19480 while the author was in residence at the Institute for Computer Applications in Science and Engineering, NASA Langley Research Center, Hampton, VA 23681-0001.

1 Introduction

In this paper we study the Navier-Stokes equations with the new boundary conditions introduced by the application of the Domain Decomposition Time Marching Algorithm ([6], [9], and [10]). These boundary conditions are of slip type, they could appear either by the application of the Domain Decomposition Time Marching Algorithm to a Navier-Stokes problem ([9] and [10]), or to a Navier-Stokes/Boltzmann coupling ([7] and [9]). In the latter case, they are similar to the analytical slip boundary conditions introduced in [5], which are derived from kinetic theory in order to replace solving Boltzmann equations by solving Navier-Stokes equations in the transitional regime.

We study here only the stationary problem. The treatment of the time dependent problem will follow using the same ideas developed here (see [9]), and the classical proofs for the standard boundary conditions (see [1]-[3], and [4]). We begin by describing the strong formulation of the problem in the first paragraph, then we set the preliminary results necessary for our study. In the third paragraph we show the equivalence of the strong and the weak formulations under some regularity hypothesis, from which we deduce the admissible boundary conditions. We present, then, the study of the stationary problem. Paragraph 6, deals with the uniqueness results when the data are sufficiently small. Finally, we present, in the last paragraph, some conclusions.

2 Motivations

2.1 The general coupling strategy

For coupling external Navier-Stokes equations, with local Navier-Stokes equations (dense regimes) or local Boltzmann equations (transitional regimes), we introduce two domains, a global one Ω , a local one Ω_V included in Ω , and an interface Γ_i (Fig. 1 in which Γ_e denotes Γ_∞). The global solution W on Ω and the local solution U_{loc} on Ω_V , are matched by the following boundary




Figure 1: The global geometry

conditions, inspired of Schwarz overlapping techniques :

$$\left\{ \begin{array}{l} W = \text{given imposed value on } \Gamma_\infty, \\ n \cdot \sigma(W) \cdot \tau = n \cdot \sigma(U_{loc}) \cdot \tau \text{ on the body } \Gamma_o, \text{ (equality of friction forces)} \\ q(W) \cdot n + n \cdot \sigma(W) \cdot v = q(U_{loc}) \cdot n \text{ on } \Gamma_o, \text{ (equality of total heat fluxes)} \\ v \cdot n = 0 \text{ on } \Gamma_o, \\ U_{loc} = 0 \text{ on } \Gamma_o, U_{loc} = W \text{ on the interface } \Gamma_i. \end{array} \right.$$

Above, $n \cdot \sigma \cdot n$ and $n \cdot \sigma \cdot \tau$ respectively denote the normal and the tangential force exerted by the body on the flow, with n the unit normal vector to the body oriented towards its interior.

The calculation of U_{loc} and W satisfying the above boundary conditions is then obtained by the Domain Decomposition Time Marching Algorithm, which was introduced by Le Tallec and Tidiriri ([6], [9] and [10]) and which leads to the following algorithm :

Initialization

1. Guess an initial distribution of the conservative variable W in the global domain Ω ;
2. Advance in time this distribution by using the global Navier-Stokes solver on N_1 time steps, with *Dirichlet* type boundary conditions on the body Γ_o ;

3. Deduce from this result an initial distribution of the local variable U_{loc} on the interface Γ_i and in the local domain Ω_V ;

4. Advance in time this distribution by using the local solver on N_2 time steps with Dirichlet boundary conditions on Γ_i and Γ_o .

Iterations

5. From U_{loc} , compute the friction forces $n \cdot \sigma(U_{loc}) \cdot \tau$ and heat flux $q(U_{loc}) \cdot n$ on the body Γ_o ;

6. Advance the global solution in time (N_1 steps) by using the global Navier-Stokes solver with the above viscous forces as boundary conditions on Γ_o ;

7. From W , compute the value of U_{loc} on the interface Γ_i ;

8. Using this new value as Dirichlet boundary conditions on Γ_i , advance the local solution in time (N_2 steps) and go back to step 5 until convergence is reached.

This algorithm completely uncouples the local and the global problems which can therefore be solved by independent solvers.

A parallel version is also quite possible although it is generally wiser to use parallel solvers within steps 6 and 8.

Remark 2.1 *The local problem can be either Navier-Stokes or Boltzmann equations (see [6]-[10]).*

2.2 The global Navier-Stokes problem

The global domain Ω is discretized using node centered cells defined on an unstructured grid. Then, at each time step n and for each cell i , we solve

$$\int_{C_i} \frac{W^{n+1} - W^n}{\Delta t} + \sum_{j \in V(i)} \int_{\partial C_i \cap \partial C_j} F_C(W^{n+1}) \cdot n_i + \int_{\partial C_i - \Gamma} F_D(W^{n+1}) \cdot n_i + \int_{\partial C_i \cap \Gamma_\infty} F(W^{n+1}) \cdot n_i = - \int_{\partial C_i \cap \Gamma_o} F_o \cdot n_i.$$

The fluxes F_C and F_D are computed at time step $n + 1$ and linearized, with for example F_C computed by an Osher approximate Riemann solver [9], and [10].




Figure 2: A boundary cell

On the body Γ_o , because of our special choice of boundary conditions, the flux is given by

$$\int_{\partial C_i \cap \Gamma_o} F_o \cdot n_i = \int_{\partial C_i \cap \Gamma_o} \begin{pmatrix} 0 \\ n_i \cdot \sigma(W^{n+1}) \cdot n_i \\ n_i \cdot \sigma(U_{loc}) \cdot \tau_i \\ -q(U_{loc}) \cdot n_i \end{pmatrix},$$

where the aspect of a boundary cell C_i is described in Fig. 2.

In other words, friction forces and heat flux are given explicitly as predicted by the local solver and the mass flux is imposed to zero. Then, in order to have a well-posed problem, at least in the incompressible case (see next paragraphs), the normal stress (the multiplier of the zero mass flux constraint) cannot be imposed and must be obtained from the solution W^{n+1} .

Remark 2.2 *Imposing friction forces to the global solution instead of no slip boundary conditions allows to have an accurate solution away from the boundary layer even with a coarse mesh (see [9]).*

3 Strong formulation

Let Ω be the domain occupied by the fluid, Γ its boundary as described in figure (1); we assume that Ω satisfy :

$$\left\{ \begin{array}{l} \Omega \text{ is a simply connected bounded domain in } \mathbb{R}^N, \\ \Gamma = \partial\Omega \text{ is of class } C^1, \\ \Gamma_b \cup \Gamma_\infty = \Gamma \text{ and } \Gamma_b \cap \Gamma_\infty = \emptyset, \\ \Gamma_b \text{ and } \Gamma_\infty \text{ are compacts of nonzero measure.} \end{array} \right. \quad (1)$$

In the steady incompressible case, the global problem in the coupling strategy described above consists in finding u and p satisfying the following equations :

$$-\nu\Delta u + u\nabla u + \nabla p = f \quad \text{in } \Omega, \quad (2)$$

$$\operatorname{div} u = 0 \quad \text{in } \Omega, \quad (3)$$

$$u \cdot n = 0 \quad \text{on } \Gamma_b, \quad (4)$$

$$u = h \quad \text{on } \Gamma_\infty, \quad (5)$$

$$\nu(\nabla u \cdot n)\tau = g_\tau \quad \text{on } \Gamma_b, \quad (6)$$

where n is the unit normal to Γ , τ is any unit normal to n .

We look for weak solutions of problem (2)-(6), i.e. (2), (3) are satisfied in a distribution sense and equations (4)-(6) are satisfied in some Sobolev spaces.

Remark 3.1 *From the remark (2.1) we observe that g_τ can be issued from a local kinetic model (Boltzmann equations) or a local continuous model (Navier-Stokes system).*

4 Preliminary results

Let $H^{m-\frac{1}{2}}(\Gamma)$ be a space of trace functions in $H^m(\Omega)$ with the following norm :

$$\|\psi\|_{m-\frac{1}{2},\Gamma} = \inf\{\|u\|_{m,\Omega} / u \in H^m(\Omega), u|_\Gamma = \psi\}, \quad (7)$$

and V the closed subspace of $(H^1(\Omega))^N$ defined by

$$V = \{v \in (H^1(\Omega))^N \mid \operatorname{div} v = 0, v \cdot n = 0 \quad \text{on } \Gamma_b, v = 0 \quad \text{on } \Gamma_\infty\}, \quad (8)$$

and define

$$\|v\| = \|\nabla v\|_{0,2,\Omega}. \quad (9)$$

We have the following classical lemma.

Lemma 4.1 $\|\cdot\|$ is a norm of V equivalent to the norm $\|\cdot\|_{1,2,\Omega}$.

Lemma 4.2 ([11]) Under hypothesis (1), we have

$$n : \Gamma \rightarrow \mathbb{R}^N \quad (10)$$

$$x \rightarrow n(x)$$

is C^1 and hence, the trace operator

$$(H^1(\Omega))^N \rightarrow H^{\frac{1}{2}}(\Gamma) \quad (11)$$

$$u \rightarrow u \cdot n$$

is continuous.

Lemma 4.3 ([14]). Let Ω satisfy (1). We have :

The injection

$$W^{1,p}(\Omega) \subset L^{q_1}(\Omega)$$

is compact for q_1 satisfying :

$$1 \leq q_1 < \infty \quad \text{if } p \geq N \quad (12)$$

$$1 \leq q_1 < q \quad \text{with } q \text{ given by } \frac{1}{q} = \frac{1}{p} - \frac{1}{N} \quad \text{if } 1 \leq p < N. \quad (13)$$

In particular we have for $p = 2$:

$$H^1(\Omega) \subset L^q(\Omega) \quad \text{for } 2 \leq q \leq \frac{2N}{N-2},$$

the injection being continuous. For $N = 2$, this injection takes place for any finite q such that $q \geq 2$.

Lemma 4.4 ([12]) Let Ω be a bounded connected domain. Then

- (i) The operator grad is an isomorphism from $L_0^2(\Omega)$ into V^0 ,
 - (ii) the operator div is an isomorphism from $(V^0)^\perp$ into $L_0^2(\Omega)$
- where $L_0^2(\Omega)$ and V^0 are defined by

$$V^0 = \{h \in H^{-1}(\Omega)^N, \langle h, v \rangle = 0, \forall v \in V\}.$$

$$L_0^2(\Omega) = \{h \in L^2(\Omega), \int_{\Omega} h(x) dx = 0\}.$$

We now set

$$b(u, v, w) = \sum_{i,j=1}^N \int_{\Omega} u_j \frac{\partial v_i}{\partial x_j} w_i dx, u, v, w \in V \quad (14)$$

$$\langle f, v \rangle = \int_{\Omega} f v dx, v \in V \quad (15)$$

$$\langle g_{\tau}, v \rangle_{\Gamma_b} = \int_{\Gamma_b} g_{\tau} \tau \cdot v d\gamma, v \in V \quad (16)$$

$$((u, v)) = \int_{\Omega} \nabla u \nabla v dx, u, v \in V \quad (17)$$

$$(u, v) = \int_{\Omega} u v dx, u, v \in V. \quad (18)$$

If f is in $(H^{-1}(\Omega))^N$ and g_{τ} in $H^{-\frac{1}{2}}(\Gamma_b)$ all the above operators (14)-(18) are well posed. Moreover, the operator defined by (16) and the trilinear form defined by (14) satisfy the following properties :

Lemma 4.5 *The mapping from V into \mathbb{R} defined by (16) is continuous.*

Proof: The proof follows directly from lemma 4.2.

Lemma 4.6 (i) *The trilinear form b is continuous on $V^2 \times (V \cap (L^N(\Omega))^N$*

(ii) *Let $u, v, w \in V$. Then we have the following relations :*

$$b(u, v, w) = -b(u, w, v), \quad (19)$$

$$b(u, v, v) = 0. \quad (20)$$

(iii) *Let $\{u_n\}_{n \geq 0}$ be a sequence in V such that $u_n \rightharpoonup u$ weakly in V . Then :*

$$\lim_{n \rightarrow \infty} b(u_n, u_n, v) = b(u, u, v), \forall v \in V. \quad (21)$$

Proof

We present only the proof of (i) and (ii). For the proof of (iii), see [14] or [13].

(i) Let $u \in V$ then $u_i \in H^1(\Omega)$; from lemma 4.3, we deduce :

$$u_i \in L^q(\Omega), \frac{1}{q} = \frac{1}{2} - \frac{1}{N},$$

which implies

$$\begin{aligned} \left| \int_{\Omega} u_j \frac{\partial v_i}{\partial x_j} w_i dx \right| &\leq \|u_j\|_{0,q,\Omega} \left\| \frac{\partial v_i}{\partial x_k} \right\|_{0,q,\Omega} \|w_i\|_{0,N,\Omega} \\ &\leq C \|u_j\|_V \|v_i\|_V \|w_i\|_V. \end{aligned} \quad (22)$$

Therefore we conclude to our result.

(ii) By definition

$$b(u, v, w) = \int_{\Omega} u_i v_{j,i} w_j.$$

After integration by parts, we obtain

$$\begin{aligned} b(u, v, w) &= - \int_{\Omega} v_j (u_i w_j)_{,i} + \int_{\partial\Omega} u \cdot n v \cdot w \\ &= - \int_{\Omega} v_j u_i w_{j,i} - \int_{\Omega} v_j w_j u_{i,i} + \int_{\partial\Omega} u \cdot n v \cdot w. \end{aligned}$$

Since $u \in V$, we have $u \cdot n = 0$ on $\partial\Omega$ and $u_{i,i} = \operatorname{div} u = 0$ on Ω . Therefore, we get

$$b(u, v, w) = - \int_{\Omega} v_j u_i w_{j,i} = -b(u, w, v).$$

Remark 4.1 For $N \leq 4$, it follows from lemma 4.3 that $V \cap (L^N(\Omega))^N = V$

5 Existence Result

We now go back to our initial formulation:

Find u and p such that :

$$-\nu \Delta u + u \cdot \nabla u + \nabla p = f \quad \text{in } \Omega, \quad (23)$$

$$\operatorname{div} u = 0 \quad \text{in } \Omega, \quad (24)$$

$$u \cdot n = \phi \quad \text{on } \Gamma_b, \quad (25)$$

$$u = h \quad \text{on } \Gamma_{\infty}, \quad (26)$$

$$\nu(\nabla u \cdot n)\tau = g_{\tau} \quad \text{on } \Gamma_b. \quad (27)$$

5.1 Hypothesis on ϕ and h

The data ϕ on Γ_b and h on Γ_∞ must verify some regularity and compatibility properties. More precisely we assume that :

$$h \in (H^{\frac{1}{2}}(\Gamma_\infty))^n, \quad (28)$$

$$\phi \in H^{\frac{1}{2}}(\Gamma_b), \quad (29)$$

$$\int_{\Gamma_\infty} h \cdot n = 0, \quad (30)$$

$$\int_{\Gamma_b} \phi = 0. \quad (31)$$

Lemma 5.1 *The relations (28)-(31) imply the existence of $\tilde{u} \in (H^1(\Omega))^N$ such that :*

$$\nabla \cdot \tilde{u} = 0 \quad \text{in } \Omega, \quad (32)$$

$$\tilde{u} \cdot n = \phi \quad \text{on } \Gamma_b, \quad (33)$$

$$\tilde{u} = h \quad \text{on } \Gamma_\infty. \quad (34)$$

Proof

Let ψ be a function defined on Γ such that :

$$\psi = \begin{cases} \phi \cdot n & \text{on } \Gamma_b, \\ h & \text{on } \Gamma_\infty. \end{cases} \quad (35)$$

From the assumed hypothesis, on Γ , ϕ and h we get :

$$\psi \in (H^{\frac{1}{2}}(\Gamma))^N.$$

Therefore, from the trace theorem, we can take $u \in H^1(\Omega)^N$ such that : $u|_\Gamma = \psi$. By using the relations (30) and (31) and the Green Formula, we obtain

$$\int_{\Omega} \nabla \cdot u = 0.$$

Using the regularity of u , we have $\nabla \cdot u \in L^2_0(\Omega)$.

Lemma 4.4 shows that there exists $v \in H^1_0(\Omega)^N$ such that $\nabla \cdot v = \nabla \cdot u$. Finally, by taking now $\tilde{u} = u - v$, we can see easily that \tilde{u} verify the relations (32)-(34).

Now, we are going to pick out some function ξ in $H^1(\Omega)$ such that :

$$\tilde{u} = \text{rot}\xi. \quad (36)$$

To achieve this, we will use the following proposition :

Proposition 5.1 ([13]) *Under the regularity hypothesis (1) and for $N = 3$ we have*

$$\text{rot}H^1(\Omega) = \{u \in L^2(\Omega)^N, \text{div}u = 0, \int_{\Gamma_b} u \cdot n = \int_{\Gamma_\infty} u \cdot n = 0\} \quad (37)$$

Remark 5.1 *This result is also valid for $N = 2$ ([13]).*

Under the hypothesis (30)-(31) and the relations (32)-(34) we have :

$$\int_{\Gamma_b} \tilde{u} \cdot n = \int_{\Gamma_b} \phi = 0, \quad (38)$$

$$\int_{\Gamma_\infty} \tilde{u} \cdot n = \int_{\Gamma_b} h \cdot n = 0. \quad (39)$$

Consequently, from Proposition 5.1, there exists $\xi \in H^1(\Omega)$, such that we have (36). From now on we assume that $N \leq 3$.

5.2 Weak formulation

Let $u \in V$. After multiplication of (23) by v , integration by parts and by taking into account the boundary conditions (25)-(27) and the relation (24) we obtain the following variational formulation :

$$\left\{ \begin{array}{l} \text{Find } u_o \in V \text{ such that} \\ \nu((u_o, v)) + b(\tilde{u}, u_o, v) + b(u_o, \tilde{u}, v) + b(u_o, u_o, v) \\ = -\nu((\tilde{u}, v)) - b(\tilde{u}, \tilde{u}, v) + \int_{\Gamma_b} g_\tau \tau \cdot v + \int_{\Omega} f v, \\ \forall v \in V \cap L^N(\Omega)^N \end{array} \right. \quad (40)$$

5.3 Equivalence between the two formulations

We will show here that the variational formulation (40) is equivalent to the strong formulation. This is the goal of the following proposition :

Proposition 5.2 *If (u, p) is a smooth solution of (2)-(6), then it satisfies (40). Conversely, if u_o is solution of (40), then there exists a unique function (up to a constant) $p \in L^2(\Omega)$ such that $u_o + \tilde{u}$ and p satisfy :*

- (i) (2) and (3) in a distribution sense,
- (ii) (4) in $H^{\frac{1}{2}}(\Gamma_b)$,
- (iii) (5) in $H^{\frac{1}{2}}(\Gamma_\infty)$,
- (iv) (6) in $H^{-\frac{1}{2}}(\Gamma_b)$.

Proof

The direct theorem was showed in our introduction of the weak formulation. We then show only the converse.

Let

$$\mathcal{V} = \{\psi \in (H_0^1(\Omega))^N / \text{div}\psi = 0\}.$$

Let u_o be a solution of problem (40) then after multiplication by a test function in $\mathcal{D}(\Omega)$, and integration by parts and density, we get

$$\langle -\nu\Delta u + (u \cdot \nabla)u - f, \psi \rangle = 0, \forall \psi \in \mathcal{V} \quad (41)$$

where \langle, \rangle denotes now the duality between the spaces $\mathcal{D}'(\Omega)^N$ and $\mathcal{D}(\Omega)^N$. We consider the operator $-\text{grad}(= -\nabla) \in \mathcal{L}(L_0^2(\Omega), H^{-1}(\Omega)^N)$, and $\mathcal{R}(-\text{grad})$ its image in $H^{-1}(\Omega)^N$. We have the following lemma (see [12])

Lemma 5.2 $\mathcal{R}(-\text{grad})$ is identical to \mathcal{V}° with \mathcal{V}° defined by :

$$\mathcal{V}^\circ = \{h \in H^{-1}(\Omega)^N / \langle h, \psi \rangle = 0, \forall \psi \in \mathcal{V}\}.$$

The application of this lemma implies the existence of $p \in L^2_0(\Omega)$ such that

$$-\nu\Delta u + u \cdot \nabla u - f + \nabla p = 0 \quad \text{in } (H^{-1}(\Omega))^N.$$

As $u \in V$, $u \cdot \nabla u$ and $f \in L^2$, then

$$\begin{aligned} -\nu\Delta u - \nabla p &= \operatorname{div}(-\nu\nabla u + pId) \\ &= (f - u \cdot \nabla u) \in (L^2(\Omega))^N. \end{aligned}$$

Applying the Green formula implies that there exists

$$q = (-\nu\nabla u + pId) \cdot n \in H^{-\frac{1}{2}}(\partial\Omega)$$

such that

$$\begin{aligned} &\int_{\Omega} -\operatorname{div}(-\nu\nabla u + pId) \cdot v \\ &+ \int_{\Omega} \nu\nabla u \nabla v - \int_{\Omega} p \operatorname{div} v = \int_{\partial\Omega} q \cdot v, \quad \forall v \in (H^1(\Omega))^N. \end{aligned}$$

By replacing $\operatorname{div}(-\nu\nabla u + pId)$ by $(f - u \cdot \nabla u)$ we get

$$\begin{aligned} \nu((u, v)) + b(u, u, v) - \int_{\Omega} p \operatorname{div} v - \int_{\partial\Omega} q \cdot v - \int_{\Omega} f v &= 0 \quad (42) \\ \forall v \in (H^1(\Omega))^N. \end{aligned}$$

By subtracting (43) from (40), it remains

$$\int_{\partial\Omega} q \cdot v - \int_{\Gamma_b} g_{\tau}(v \cdot \tau) = 0, \quad \forall v \in V.$$

Now let w arbitrary element of $H^{\frac{1}{2}}_{00}(\Gamma_b)$, where $H^{\frac{1}{2}}_{00}(\Gamma_b)$ is defined by :

$$H^{\frac{1}{2}}_{00}(\Gamma_b) = \{v \in L^2(\Gamma_b); \exists w \in H^1(\Omega), w = 0 \text{ on } \Gamma_{\infty} \text{ and } w = v \text{ on } \Gamma_b\}$$

(see [11] for more details about this space).

Let $v \in V$ such that

$$v \cdot n = 0 \quad \text{on } \Gamma_b,$$

$$v \cdot \tau = w \quad \text{on } \Gamma_b,$$

$$v = 0 \quad \text{on } \Gamma_\infty.$$

Applying the precedent equality, it remains

$$\int_{\Gamma_b} q \cdot \tau w = \int_{\Gamma_b} g_\tau w, \forall w \in H_{00}^{\frac{1}{2}}(\Gamma_b),$$

therefore

$$\begin{aligned} q \cdot \tau &= \tau \cdot (-\nu \nabla u + p Id) \cdot n \\ &= g_\tau \quad \text{in } H^{-\frac{1}{2}}(\Gamma_b). \end{aligned}$$

Remark 5.2 *We cannot impose the normal stress in (3)-(7) because of the presence of a pressure term in q .*

5.4 Existence result

Now, we are able to set out the main result of this section.

Theorem 5.1 *Let $f \in H^{-1}(\Omega)$; then there exists $u_o \in V$ solution of (40).*

Proof

We proceed as in the usual proof of existence for the incompressible Navier-Stokes equations. Let us define

$$\begin{aligned} [\psi_m(u), u] &= \nu \|u\|^2 + \nu((\tilde{u}, u)) + b(\tilde{u}, \tilde{u}, u) - \int_{\Gamma_b} g_\tau \tau u \\ &\quad - \int_{\Omega} f u + b(u, \tilde{u}, u). \end{aligned}$$

By using the continuity of b we obtain :

$$b(\tilde{u}, \tilde{u}, u) \leq c_1 \|\tilde{u}\|_{0,q,\Omega} \|\tilde{u}\|_{1,2,\Omega} \|u\|_{0,N} \quad (43)$$

with q satisfying (12)-(13). In addition we have :

$$b(\tilde{u}, \tilde{u}, u) \leq c_2 \|\tilde{u}\|_{1,2,\Omega}^2 \|u\|. \quad (44)$$

Using lemma 4.1 and lemma 4.2 we arrive to :

$$\begin{aligned} [\psi_m(u), u] &\geq \nu \|u\|^2 + b(u, \tilde{u}, u) \\ &\quad - \|u\| (\nu \|\tilde{u}\| + c_2 \|\tilde{u}\|_{1,2,\Omega}^2 + c_3 \|g_\tau\|_{-\frac{1}{2},\Gamma_b} + \|f\|_{-1,2,\Omega}). \end{aligned}$$

So, to be able to use the Brouwer fixed point theorem, it is sufficient to show that we have for some $\beta > 0$, the following relation :

$$\nu \|u\|^2 + b(u, \tilde{u}, u) \geq \beta \|u\|^2, \forall u \in V. \quad (45)$$

This is the goal of the following lemma :

Lemma 5.3 $\forall \gamma > 0$ we can choose \tilde{u} verifying (32)-(33) such that

$$|b(u, \tilde{u}, u)| \leq \gamma \|u\|^2. \quad (46)$$

From the above lemma, we conclude to the existence of a given k such that

$$[\psi_m(u), (u)] > 0,$$

for any $u \in V$ with $\|u\| = k$. Then the Brouwer fixed point theorem can be applied to the function $\psi_m(u)$ inside any ball of radius k belonging to any given finite dimensional approximation V_m of V . From this, we deduce the existence of a Galerkin approximation $u_m \in V_m$ of the solution u of (40). By standard compactness arguments, it follows that u_m weakly converges to a solution u_o of (40) in V . (see [5] for more details).

6 Uniqueness results

Under the assumption that $N \leq 3$, we have the following result :

Theorem 6.1 *Assume that \tilde{u} in $(L^N(\Omega))^N$ is sufficiently small so that*

$$|b(v, \tilde{u}, v)| \leq \frac{\nu}{2} \|v\|^2, \forall v \in V, \quad (47)$$

and ν is sufficiently large so that

$$\nu^2 > 4C(\|\tilde{f}\|_{V'} + \|g_\tau\|_{-\frac{1}{2}, \Gamma_b}) \quad (48)$$

where C is the constant in (22) and

$$\tilde{f} = \nu \Delta \tilde{u} - \Sigma_j \tilde{u}_j \frac{\partial}{\partial x_j} \tilde{u} + f$$

and

$$\tilde{g}_\tau = \nu \frac{\partial}{\partial n} \tilde{u} + g_\tau$$

Proof

We proceed as in the proof for the standard homogeneous boundary conditions (see for instance [13]).

Suppose that u_o is solution of the problem (40). Taking $v = u_o$ in (40) and using (20), we obtain :

$$\nu \|u_o\|_V^2 = -b(u_o, \tilde{u}, u_o) - \nu((\tilde{u}, u_o)) - b(\tilde{u}, \tilde{u}, u_o) + \int_{\Gamma_b} g_\tau \tau \cdot u_o + \int_\Omega f u_o, \quad (49)$$

Using the Green formula, we obtain

$$\nu \|u_o\|_V^2 = -b(u_o, \tilde{u}, u_o) + \langle \nu \Delta \tilde{u} - \Sigma_j \tilde{u}_j \frac{\partial}{\partial x_j} \tilde{u} + f, u_o \rangle \quad (50)$$

$$+ \langle \nu \frac{\partial}{\partial n} \tilde{u} + g_\tau, u_o \rangle_{\Gamma_b} \quad (51)$$

$$= -b(u_o, \tilde{u}, u_o) + \langle \tilde{f}, u_o \rangle + \langle \tilde{g}_\tau, u_o \rangle_{\Gamma_b} \quad (52)$$

Since $f \in (H^{-1}(\Omega))^N$ and $\tilde{u} \in (H^1(\Omega))^N$, $\Delta\tilde{u} \in (H^{-1}(\Omega))^N$ and then $\tilde{f} \in (H^{-1}(\Omega))^N$. Similarly, since $g_\tau \in (H^{-\frac{1}{2}}(\Gamma_b))^N$ and $\tilde{u} \in (H^1(\Omega))^N$ we have $\frac{\partial\tilde{u}}{\partial n} \in (H^{-\frac{1}{2}}(\Gamma_b))^N$ and then $\tilde{g}_\tau \in (H^{-\frac{1}{2}}(\Gamma_b))^N$, and the equality above is well defined. The rest of the proof is an adaptation of the proof for the standard homogeneous boundary conditions (see for instance [13]).

7 Conclusion

In this paper we have shown that the slip boundary conditions resulting from the application of the Domain Decomposition Time Marching Algorithm to either Navier-Stokes/Navier-Stokes coupling ([9] and [10]) or Navier-Stokes/Boltzmann coupling ([9], [7], and [8]) are admissible and lead to a well posed global problem at least in the incompressible case.

References

- [1] J. Leray, “*Etude de diverses équations intégrales nonlinéaires et de quelques problèmes que pose l’hydrodynamique*”, J. Math. Pures Appl., 12, 1933, p. 1-82.
- [2] J. Leray, “*Essai sur les mouvements plans d’un liquide visqueux que limitent des parois*”, J. Math. Pures Appl., 13, 1934, p. 331-418.
- [3] J. Leray, “*Essai sur le mouvement d’un liquide visqueux emplissant l’espace*”, Acta Math., 63, 1934, p. 193-248.
- [4] E. Hopf, “*On nonlinear partial differential equations*”, Lecture Series of the Symposium on partial Differential Equations, Berkeley, 1955, Ed. The Univ. of Kansas (1957), p. 1-29.
- [5] R. Gupta, C. Scott and J. Moss, “*Slip-boundary equations for multi-component nonequilibrium airflow*”. NASA Technical Paper 2452, November 1985.

- [6] P. Le Tallec and M. D. Tidriri, "*Convergence of domain decomposition algorithms with full overlapping for the advection-diffusion problems and numerical coupling of Navier-Stokes equations with Boltzmann equations*". Submitted Math. Comp.
- [7] P. Le Tallec and M. D. Tidriri, "*Coupling Navier-Stokes and Boltzmann*." Submitted to J. Comp. Phy.
- [8] P. Le Tallec, M. D. Tidriri, "*Kinetic upgrade of Navier-Stokes equations for transitional flows*." To appear as INRIA Research report.
- [9] M. D. Tidriri, "*Couplage d'approximations et de modèles de types différents dans le calcul d'écoulements externes*", PhD thesis, University of Paris IX, 1992.
- [10] M. D. Tidriri, "*Domain decomposition for compressible Navier-Stokes equations with different discretizations and formulations*." To appear in *J. Comp. Phy.*
- [11] J. Nečas, "*les méthodes directes en théorie des équations elliptiques*", Masson (1967).
- [12] V. Girault, P. A. Raviart, "*Finite element methods for Navier-stokes equations*". Springer-Verlag, Berlin Heidelberg New York Tokyo 1986.
- [13] R. Temam, "*Navier-Stokes equations*", North-Holland, (1977).
- [14] J. L. Lions, "*Quelques méthodes de résolution des problèmes aux limites non linéaires*". Dunod Gauthier-Villars, Paris 1969.
- [15] J. L. Lions, E. Magenes, "*Problèmes aux limites non homogènes et applications*". volume 1. Dunod, Paris 1968.

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE July 1995	3. REPORT TYPE AND DATES COVERED Contractor Report	
4. TITLE AND SUBTITLE MATHEMATICAL ANALYSIS OF THE NAVIER-STOKES EQUATIONS WITH NON STANDARD BOUNDARY CONDITIONS		5. FUNDING NUMBERS C NAS1-19480 WU 505-90-52-01	
6. AUTHOR(S) M.D. Tidriri			
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Institute for Computer Applications in Science and Engineering Mail Stop 132C, NASA Langley Research Center Hampton, VA 23681-0001		8. PERFORMING ORGANIZATION REPORT NUMBER ICASE Report No. 95-54	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) National Aeronautics and Space Administration Langley Research Center Hampton, VA 23681-0001		10. SPONSORING/MONITORING AGENCY REPORT NUMBER NASA CR-198184 ICASE Report No. 95-54	
11. SUPPLEMENTARY NOTES Langley Technical Monitor: Dennis M. Bushnell Final Report To be submitted to the Journal of Mathematical Analysis and its Applications			
12a. DISTRIBUTION/AVAILABILITY STATEMENT Unclassified-Unlimited Subject Category 64		12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) One of the major applications of the Domain Decomposition Time Marching Algorithm is the coupling of the Navier-Stokes systems with Boltzmann equations in order to compute transitional flows. Another important application, is the coupling of a global Navier-Stokes problem with a local one in order to use different modelizations and/or discretizations. Both of these applications involve a global Navier-Stokes system with non standard boundary conditions. The purpose of this work is to prove, using the classical Leray-Schauder theory, that these boundary conditions are admissible and lead to a well posed problem.			
14. SUBJECT TERMS Navier-Stokes Equations; Domain Decomposition Time Marching Algorithm; Slip Boundary Conditions; Friction Forces			15. NUMBER OF PAGES 19
			16. PRICE CODE A03
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT	20. LIMITATION OF ABSTRACT