

# REPORT DOCUMENTATION PAGE

Form Approved  
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)

2. REPORT DATE

9/15/94

3. REPORT TYPE AND DATES COVERED

Final

4. TITLE AND SUBTITLE

Experimental Studies of Fundamental One- and Two-Photon Lasing Processes


5. FUNDING NUMBERS

6. AUTHOR(S)

Thomas W. Mossberg

7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)

University of Oregon  
Eugene, OR 97403  
% Research and Sponsored Programs



9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)

U.S. Army Research Office  
P.O. Box 12211  
Research Triangle Park, NC 27709-2211

10. SPONSORING/MONITORING AGENCY REPORT NUMBER

11. SUPPLEMENTARY NOTES

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy, or decision, unless so designated by other documentation.

12a. DISTRIBUTION/AVAILABILITY STATEMENT

Approved for public release; distribution unlimited.

12b. DISTRIBUTION CODE

13. ABSTRACT (Maximum 200 words)

Theoretical and experimental investigations of two-photon lasing and inversionless lasing have been completed. Continuous-wave two-photon lasing has been demonstrated for the first time and its predicted bistable character verified. The role of pump bandwidth in the production of inversionless gain has been studied and the maximal efficacy of narrow bandwidth of pumps identified. Continuous-wave lasing in a Cascade-type three-level atomic system has been observed.

19951129 009

DTIC QUALITY INSPECTED 5

14. SUBJECT TERMS

two-photon laser  
inversionless lasing

15. NUMBER OF PAGES

5

16. PRICE CODE

17. SECURITY CLASSIFICATION OF REPORT

UNCLASSIFIED

18. SECURITY CLASSIFICATION OF THIS PAGE

UNCLASSIFIED

19. SECURITY CLASSIFICATION OF ABSTRACT

UNCLASSIFIED

20. LIMITATION OF ABSTRACT

UL

Experimental Studies of Fundamental One- and Two-Photon Lasing Processes

Final Report

Thomas W. Mossberg  
Professor of Physics  
Department of Physics, University of Oregon  
Eugene, Oregon 97403

September 14, 1994

U. S. Army Research Office

Accession For	
NTIS CRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification .....	
By .....	
Distribution /	
Availability Codes	
Dist	Avail and / or Special
A-1	

DAAL03-91-G-0313

Research Conducted at  
The University of Oregon  
Eugene, Oregon 97403

APPROVED FOR PUBLIC RELEASE;

DISTRIBUTION UNLIMITED

THE VIEWS, OPINIONS, AND/OR FINDINGS CONTAINED IN THIS REPORT ARE THOSE OF THE AUTHOR AND SHOULD NOT BE CONSTRUED AS AN OFFICIAL DEPARTMENT OF THE ARMY POSITION, POLICY, OR DECISION, UNLESS SO DESIGNATED BY OTHER DOCUMENTATION.

## STATEMENT OF THE PROBLEM STUDIED

Laser action is of interest as a fundamental physical process and because of its widespread importance in technology. The present project has been directed toward enhancing basic understanding of several newly discovered forms of laser action. One of these is laser action based on two-photon stimulated emission. In two-photon laser gain, the gain medium transfers energy to the light field via the simultaneous emission of photon pairs. The individual photons within a pair are highly correlated in time. Two-photon gain is non-linear and leads to qualitatively different laser performance particularly in the case of threshold. The concept of inversionless laser gain has recently become popular. In this case, laser gain occurs in the absence of a population inversion on the lasing transition of the gain medium. Naively, this situation should not occur. Inversionless gain has been suggested as a means of producing laser action in spectral regions where attainment of gain-media population inversion is technologically difficult. We have attempted to elucidate a variety of aspects of two-photon and inversionless gain and lasing using both experimental and theoretical approaches.

## SUMMARY OF THE MOST IMPORTANT RESULTS

**Two-Photon Lasing.** We have provided the first experimental demonstration of continuous-wave two-photon laser action. A novel gain medium consisting of strongly driven two-level atoms was proposed and utilized. For the first time, our experiments demonstrated the bistable operating characteristics of a two-photon laser. In stark contrast to a normal (one-photon) laser, the two-photon laser possesses two stable operating points. One of these is the "off" state. As a result of the stability of the off state, the two-photon laser has been proposed as a high energy device.

What is lasing? Investigation of the various new types of lasing process has lead us to clarify the definition of lasing. We have proposed that laser gain is defined as phase insensitive optical amplification. This definition makes it possible to distinguish lasing processes from the diverse classes of non-linear gain mechanisms.

Pump Bandwidth Effects in Inversionless gain. We have determined that narrowband pump fields lead to substantially larger inversionless gain than can be created with broad bandwidth pump fields. Previous theoretical work had emphasized broad bandwidth pump mechanisms and therefore did not predict maximal gains.

Unexpected Inversions in Three-Level Systems. We have analyzed the dynamics of doubly driven V-type three-level atomic systems and found that certain pairs of atomic states can become inverted even though there is no normal irreversible pathway available for population to enter the upper state of the transition involved. This result is of fundamental and practical interest.

Inversionless Lasing in a Cascade-Type Three-Level Atom. We have performed an experiment in an atomic beam of barium atoms that demonstrates continuous wave lasing in a driven three-level atomic system. Preliminary analysis of the experimental data indicates that the laser emission, observed on the upper transition, was produced in the absence of a population inversion on that transition. Analysis of the experimental results verifies predictions regarding inversionless gain and clarifies its physical origin.

#### LIST OF ALL PUBLICATIONS AND TECHNICAL REPORTS

Realization of a Continuous-Wave, Two-Photon Optical Laser, D. Gauthier, Q. Wu, S. Morin, and T. W. Mossberg, Phys. Rev. Lett. **68**, 464 (1992).

Dynamically Induced Irreversibility in Coherently Driven Systems, K. K. Meduri, G. A. Wilson, P. B. Sellin, and T. W. Mossberg, Phys. Rev. Lett. **71**, 4311 (1993).

Lasing without inversion - Gain Enhancement through Spectrally Colored Population Pumping, G. S. Agarwal, G. Vemuri, and T. W. Mossberg, Phys. Rev. A **48**, R4055 (1993).

Coherently driven three-level atoms: unexpected inversions, pump bandwidth effects and observation of continuous lasing, K. K. Meduri, P. B. Sellin, G. A. Wilson, and T. W. Mossberg, J. European Opt. Soc. Part B (Quantum Optics) 6, 287 (1994).

Inversionless gain in driven three-level atoms: A comparison of broadband and monochromatic excitation, G. A. Wilson, K. K. Meduri, P. B. Sellin, and T. W. Mossberg, (accepted for publication in Phys. Rev. A).

Observation of inversionless gain in Cascade-type three-level atom, P. B. Sellin, G. Wilson, K. Meduri, and T. W. Mossberg, (in preparation).

#### LIST OF PARTICIPATING SCIENTIFIC PERSONNEL

Mr. Peter B. Sellin, PhD expected 94-95 academic year

Dr. Geoffry Wilson

Dr. Krishna Meduri

Prof. Thomas W. Mossberg

#### REPORT OF INVENTIONS

None