

AD _____

GRANT NUMBER: DAMD17-94-J-4331

TITLE: Line-1 Retrotransposons as Mutagens in Human Breast
Cancer

PRINCIPAL INVESTIGATOR: Dr. Thomas G. Fanning

CONTRACTING ORGANIZATION: American Registry of Pathology
Armed Forces Institute of Pathology
Washington, DC 20306-6000

REPORT DATE: October 1995

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for public release;
distribution unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

19960129 012

DTIC QUALITY INSPECTED 1

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE October 1995	3. REPORT TYPE AND DATES COVERED Annual 30 Sep 94 - 29 Sep 95		
4. TITLE AND SUBTITLE Line-1 Retrotransposons as Mutagens In Human Breast Cancer			5. FUNDING NUMBERS DAMD17-94-J-4331	
6. AUTHOR(S) Dr. Thomas G. Fanning				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) American Registry of Pathology Armed Forces Institute of Pathology Washington, DC 20306-6000			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012			10. SPONSORING / MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) The human <i>LINE-1</i> retrotransposon (<i>LIHs</i>) is known to cause mutations by inserting into genes and inactivating them. The element is expressed in many breast tumors and breast tumor cell lines, suggesting that <i>LIHs</i> -induced mutations may play some role in this malignancy. To test this hypothesis we plan to place a "tagged" <i>LIHs</i> element in non-malignant breast epithelial cells and let it transpose. If the cells become malignant, we can isolate the gene(s) into which the element has transposed by using the unique tag. To date we have identified an appropriate cell line, MCF10A, but attempts to obtain stably transfected cells expressing <i>LIHs</i> -encoded proteins have been a failure. We are currently exploring the use of other vectors that may increase our chance of success.				
14. SUBJECT TERMS LINE-1 retrotransposon; Breast cancer			15. NUMBER OF PAGES 7	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited	

GENERAL INSTRUCTIONS FOR COMPLETING SF 298

The Report Documentation Page (RDP) is used in announcing and cataloging reports. It is important that this information be consistent with the rest of the report, particularly the cover and title page. Instructions for filling in each block of the form follow. It is important to *stay within the lines* to meet optical scanning requirements.

Block 1. Agency Use Only (Leave blank).

Block 2. Report Date. Full publication date including day, month, and year, if available (e.g. 1 Jan 88). Must cite at least the year.

Block 3. Type of Report and Dates Covered. State whether report is interim, final, etc. If applicable, enter inclusive report dates (e.g. 10 Jun 87 - 30 Jun 88).

Block 4. Title and Subtitle. A title is taken from the part of the report that provides the most meaningful and complete information. When a report is prepared in more than one volume, repeat the primary title, add volume number, and include subtitle for the specific volume. On classified documents enter the title classification in parentheses.

Block 5. Funding Numbers. To include contract and grant numbers; may include program element number(s), project number(s), task number(s), and work unit number(s). Use the following labels:

C - Contract	PR - Project
G - Grant	TA - Task
PE - Program Element	WU - Work Unit Accession No.

Block 6. Author(s). Name(s) of person(s) responsible for writing the report, performing the research, or credited with the content of the report. If editor or compiler, this should follow the name(s).

Block 7. Performing Organization Name(s) and Address(es). Self-explanatory.

Block 8. Performing Organization Report Number. Enter the unique alphanumeric report number(s) assigned by the organization performing the report.

Block 9. Sponsoring/Monitoring Agency Name(s) and Address(es). Self-explanatory.

Block 10. Sponsoring/Monitoring Agency Report Number. (If known)

Block 11. Supplementary Notes. Enter information not included elsewhere such as: Prepared in cooperation with...; Trans. of...; To be published in... When a report is revised, include a statement whether the new report supersedes or supplements the older report.

Block 12a. Distribution/Availability Statement. Denotes public availability or limitations. Cite any availability to the public. Enter additional limitations or special markings in all capitals (e.g. NOFORN, REL, ITAR).

DOD - See DoDD 5230.24, "Distribution Statements on Technical Documents."

DOE - See authorities.

NASA - See Handbook NHB 2200.2.

NTIS - Leave blank.

Block 12b. Distribution Code.

DOD - Leave blank.

DOE - Enter DOE distribution categories from the Standard Distribution for Unclassified Scientific and Technical Reports.

NASA - Leave blank.

NTIS - Leave blank.

Block 13. Abstract. Include a brief (*Maximum 200 words*) factual summary of the most significant information contained in the report.

Block 14. Subject Terms. Keywords or phrases identifying major subjects in the report.

Block 15. Number of Pages. Enter the total number of pages.

Block 16. Price Code. Enter appropriate price code (*NTIS only*).

Blocks 17 - 19. Security Classifications. Self-explanatory. Enter U.S. Security Classification in accordance with U.S. Security Regulations (i.e., UNCLASSIFIED). If form contains classified information, stamp classification on the top and bottom of the page.

Block 20. Limitation of Abstract. This block must be completed to assign a limitation to the abstract. Enter either UL (unlimited) or SAR (same as report). An entry in this block is necessary if the abstract is to be limited. If blank, the abstract is assumed to be unlimited.

Table of Contents

1. Front Cover.....	1
2. SF 298.....	2
3. Foreword.....	3
4. Table of Contents.....	4
5. Introduction.....	5
6. Experimental.....	5
7. Conclusions.....	6
8. References.....	6

Introduction

Human *LINE-1* (*LIHs*) is a transposable element that encodes a reverse transcriptase and moves *via* an RNA intermediate [1]. It therefore seems possible that cells in which *LIHs* is active may be subject to insertional mutagenesis. We have recently found that the element is expressed in a significant number of germ cell cancers [2,3] as well as in many breast tumors and breast tumor cell lines [4]. This last finding raises the possibility that the initiation or progression of some breast cancers is facilitated by *LIHs*-induced insertional mutations.

In addition to insertional mutagenesis, there are several other characteristics of the *LIHs* element that suggest its potential as an oncogenic agent. For example, *LIHs* has an internal promoter which could potentially lead to readthrough transcription and activation of downstream genes. In addition, the *LIHs*-encoded p40 protein has a leucine zipper motif, suggesting a possible interaction with other cellular proteins. Such interactions at inappropriate times might lead to the disruption of important cellular functions. Thus, *LIHs* involvement in cancer could occur by several mechanisms, either singly or in combination.

Our long range goals focus on the isolation of cellular genes that are affected by *LIHs* transposition. These genes are presumably the ones whose inactivation (by insertional mutagenesis), or activation (by readthrough transcription), is one of the steps in the pathway leading to malignancy. Specifically we have proposed to:

- (1) Place a "tagged", transpositionally competent *LIHs* element into non-malignant human breast epithelial cells.
- (2) Identify malignant cells arising from the non-malignant cell population and isolate and characterize sequences into which the tagged *LIHs* element has transposed.

Experimental

In our initial experiments we have used vectors containing only the first *LIHs* open reading frame (p40 gene). We have done this for several reasons; (1) we have an anti-p40 antibody that is capable of detecting small quantities of the protein. Thus, we can easily assay cells for *LIHs* expression. (2) Experience gained in this work will be very useful in placing the full-length *LIHs* element into the same cells.

Transfection of breast epithelial cells.

We have used three human, non-malignant, breast epithelial lines in our work: HBL100, Hs587Bst, and MCF10A. Transfection protocols have utilized (a) calcium phosphate-DNA coprecipitation, (b) a DEAE-dextran-DNA mix, and (c) several liposome-DNA complexes (LipofectAMINE, LipofectACE, and Lipofectin, all sold by Gibco-BRL). Our initial experiments utilized a β -galactosidase-containing vector, since measurement of the enzyme activity in cell extracts is very sensitive. Using this vector, the HBL100 cells were completely refractory to transfection. In addition, colleagues have informed us that this cell line is refractory to transfection by electroporation. Similarly, we were unable to obtain transfected Hs587Bst cells. Thus, we have concentrated our efforts on the MCF10A cell line.

Transfection of MCF10A cells with the β -galactosidase-containing vector suggested that transfection was possible, but the efficiency was low. Indeed, when we stained a population of transfected cells, it was apparent that less than 1% were actually taking-up DNA under our most optimal conditions. Nevertheless, this is certainly sufficient to give us stably

transfected cells that express *LIHs*, which is our goal. To obtain stably transfected cells we treat with a LipofectACE-DNA complex, then subject the cells, after an appropriate time interval, to a neomycin analog (G418) since the vector contains both the *LIHs* p40 gene and a neomycin (neo) resistance gene. Neo resistant cells are cloned, grown in large numbers, and assayed for p40 expression.

Expression of the *LIHs* p40 gene in transfected cells.

We have placed the following plasmid-based constructs into the MCF10A cell line: RSV-p40, CMV-p40 and MMTV-p40. These vectors place the *LIHs* ORF1 gene (encoding the p40 protein) behind the Rous sarcoma virus promoter/enhancer, the cytomegalovirus promoter/enhancer and the mouse mammary tumor virus promoter/enhancer, respectively. All constructs were initially tested for transient expression in COS cells, an SV40-transformed monkey kidney cell line that is easily and efficiently transfected. All constructs proved to be capable of expressing the p40 protein as shown by Western blotting extracts from the transfected COS cells.

All of the constructs produced stable, drug resistant colonies after transfection into MCF10A cells. However, in no case was p40 detected. We assume, but have not yet shown, that the promoter and/or enhancer driving the expression of the p40 gene is inactivated, probably by methylation. The promoter/enhancer for the neo gene, which is under strong selection, is obviously still functional.

Currently, we are repeating the transfection experiments with the p40 gene in several retroviral vectors: LXSX, LNSX and Elneo [5,6]. The LXSX and LNSX vectors have two promoters, one for the neo gene and one for the p40 gene. Thus, there is the possibility that they too could exhibit drug resistance, but have the p40 promoter/enhancer inactivated. The Elneo vector, however, contains only a single promoter/enhancer. With this vector the neo and p40 genes can both be expressed due to mRNA splicing. Thus, complete transcripts express the p40 protein which is located at the 5' end of the mRNA, while spliced mRNAs express the neo gene product.

Conclusions

A successful execution of the goals of the grant require that a full-length, transpositionally competent *LIHs* element be placed into a non-malignant breast epithelial cell. To date we have been unable to achieve this goal. Either the cell lines we wish to use are refractory to transfection, or the constructs we have used are rendered nonfunctional following transfection. We are currently pursuing other vectors that may offer a higher possibility of success. In addition, we are searching for other cell lines that may be more amenable than those we are currently using.

References

- [1] Fanning, T.G. and M.F. Singer (1987) *LINE-1*: a mammalian transposable element. *Biochim. Biophys. Acta* 910:203-212.
- [2] Bratthauer, G.L. and T.G. Fanning (1992) Active *LINE-1* retrotransposons in human testicular cancer. *Oncogene* 7:507-510.

[3] Bratthauer, G.L. and T.G. Fanning (1993) *LINE-1* retrotransposon expression in pediatric germ cell tumors. *Cancer* 71:2383-2386.

[4] Bratthauer, G.L., R.D. Cardiff and T.G. Fanning (1994) Expression of *LINE-1* retrotransposons in human breast cancer. *Cancer* 73:2333-2336.

[5] Miller, A.D. and G.J. Rosman (1989) Improved retroviral vectors for gene transfer and expression. *Biotechniques* 7:980-990.

[6] Heidecker, G., M. Huleihel, J.L. Cleveland, W. Kolch, T.W. Beck, P. Lloyd, T. Pawson and U.R. Rapp. Mutational activation of *c-raf-1* and definition of the minimal transforming sequence. *Mol. Cell. Biol.* 10:2503-2512.