


REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE June '94	3. REPORT TYPE AND DATES COVERED FINAL -		
4. TITLE AND SUBTITLE The Molecular and Cellular Mechanisms of Quinone Tanning of proteins			5. FUNDING NUMBERS N00014-90-J-1997	
6. AUTHOR(S) John S. Cordingley				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Univ. of Wyoming, Dept. of Molecular Biology			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) ONR			10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
 JAN 23 1995				
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION / AVAILABILITY STATEMENT unrestricted			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) The tyrosine rich eggshell protein referred to as F4 has been modelled exhaustively both by computer simulation and by physical studies of synthetic peptides. Our conclusion is that the tyrosine rich repetitive region of this protein probably forms a left handed alpha helix. I have published the suggestion, based on the established protein sequences, that the absence of phenylalanine residues may suggest a role in electron transport.				
14. SUBJECT TERMS Schistosome / Eggshell / tyrosine rich / cross-linking / tanning protein			15. NUMBER OF PAGES 4	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT undclassified	18. SECURITY CLASSIFICATION OF THIS PAGE undclassified	19. SECURITY CLASSIFICATION OF ABSTRACT undclassified	20. LIMITATION OF ABSTRACT unlimited	

NSN 7540-01-280-5500

Standard Form 298 (Rev. 2-89)
Prescribed by ANSI Std. Z39-18
298-102

DTIC QUALITY INSPECTED 1

53-85

19960117 067

FINAL REPORT

copy

Grant#: N00014-90-J-1997

R&T=441p026

PRINCIPAL INVESTIGATOR: John S. Cordingley.

INSTITUTION: University of Wyoming (Dept. of Molecular Biology).

GRANT TITLE: The Molecular and Cellular Mechanisms of
quinone tanning of proteins

AWARD PERIOD: 6/1/90 - 12/31/93

Objectives. The objective of this project was to further elucidate the mechanisms by which the schistosome eggshell becomes cross-linked. During the course of this project our aims were modified considerably in the light of our data. The most important change in my thinking was to accept the possibility, I might even say probability today, that there is no conventional phenol oxidase enzyme responsible for the final steps in the cross-linking process. This point of view has been argued in some detail in the publications listed and attached.

Accomplishments: The repetitive tyrosine rich protein which we refer to as F4, has been the major focus of our work. We have continued to refine our analysis of this unusual proteins possible secondary structure and these results have been published in detail (Middaugh et al., 1993). Our data continues to support the unusual conclusion that this protein, comprising only L-amino acids adopts a left-handed alpha helical conformation.

I have further suggested (Cordingley et al., 1993) based upon sequence data from two laboratories that the F4 protein may act as an "electron transport chain" during eggshell cross-linking. Our attempts to design experiments to test this hypothesis have been unsuccessful. The readiness with which the cross-linking reactions occur is in itself a major stumbling block to characterizing the system and we have been unsuccessful in

isolating stable intermediates of the cross-linking process.

We have begun making peptides with specific substitutions to try to create peptides in which we can measure electron transfer (or perhaps more simply "energy transfer") along the peptide. It has proved possible in other proteins to use tryptophan residues to introduce "energy" into proteins by irradiating with UV light of specific wavelengths and to detect energy transfer to other distant groups within the protein. To try to exploit this we have replaced one tyrosine residue with a tryptophan and another tyrosine residue, more C-terminal in the peptide with a cysteine residue. The cysteine residue allows us to couple reporter groups or we may be able to detect direct reduction of a disulfide bond at this position. Preliminary observations show that the peptides with the tryptophan and cysteine substitutions retain the left handed structure found in the native peptide.

The publication of another complete sequence of an F4 homologue reinforces the points made previously regarding the absence of Phe and Trp substitutions for the tyrosines in F4. There is one Phenylalanine, but it is in the signal peptide sequence. There are 120 tyrosine residues in the remainder of F4 and not a single Phenylalanine or tryptophan. The sheer number of tyrosine residues renders the lack of Trp or Phe substitutions much more significant than in a protein with only a few tyrosine residues. Clearly there must be very strong selective pressure preventing acceptance of these substitutions.

The attached reprints spell out our thinking on eggshell cross-linking in schistosomes and on the points outlined here and I see no particular advantage to repeating these ideas here. The interested reader is referred to the attached reprint collection.

Accession For	
NTIS CS&I	<input type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

Publications.

- Cordingley, J.S., Thomson, J.A., and Middaugh, C.R. (1993). Is the Tyrosine Rich Eggshell Protein of *Schistosoma mansoni* and electron transport chain? *Mat. Res. Soc. Symp. Proc.* 292, 69-76.
- Middaugh, C.R., Thomson, J.A., Burke, C.J., Mach, H., Naylor, A.M., Bogusky, M.J., Ryan, J.A., Pitzenberger, S.M., Ji, H., and Cordingley, J.S. (1993). Structure of synthetic peptides derived from an eggshell protein of *Schistosoma mansoni*. *Protein Sci.* 2, 900-914.
- Wells, K. and Cordingley, J.S. (1991). Detecting proteins containing 3,4-dihydroxyphenylalanine by silver staining of polyacrylamide gels. *Anal. Biochem.* 194, 237-242.
- Wells, K.E. and Cordingley, J.S. (1991). *Schistosoma mansoni*: Eggshell formation is regulated by pH and calcium. *Exp. Parasitol.* 73, 295-310.
- Wells, K.E. and Cordingley, J.S. (1992). The Cell and Molecular Biology of Eggshell Formation in *Schistosoma mansoni*. In *Structure, Cellular Synthesis and Assembly of Biopolymers*. S.T. Case, ed. (New York: Springer-Verlag), pp. 97-113.