

7749-EN-09
DTIC.

**RUSSIAN INSECTS AND DISEASES
THAT COULD CONTROL US WATERWEEDS**

Final Technical Report

by

**V.F.Zaitzev, M.G.Volkovitch,
S.Ya.Reznik, S.A.Podlipaev, M.Yu.Dolgovskaya**

February 1996

United States Army
EUROPEAN RESEARCH OFFICE OF U.S. ARMY
London England
Contract Number: **N68171-95-C-9121**
Contractor: **Mr. Tracey Carr**

Approved for Public Release; distribution unlimited

19960430 042

DTIC QUALITY INSPECTED 1

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
<small>Please reading Bureau for this collection of information is estimated to average 1 hour per respondent, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this Bureau estimate of any other aspect of this collection of information, including suggestions for reducing this Bureau's reporting burden, Department for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.</small>				
1. AGENCY USE ONLY (Leave blank)		2. REPORT TYPE AND DATES COVERED		
		Final Report February 29, 1996 August 1995 - February 1996		
4. TITLE AND SUBTITLE			5. FUNDING NUMBERS	
Russian Insects and Diseases that could control US Waterweeds			C: N68171-95-C-9121 PR: WK2Q6C-7749-EN09	
6. AUTHOR(S)				
V.F.Zaitzev, M.G.Volkovitch, S.Ya.Reznik, S.A.Podlipaev, M.Yu.Dolgovskaya				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)			8. PERFORMING ORGANIZATION REPORT NUMBER	
Zoological Institute, Russian Academy of Sciences, Universitetskaya nab., 1. 199034, St.Petersburg, Russia			RF959001	
9. SPONSORING, MONITORING AGENCY NAME(S) AND ADDRESS(ES)			10. SPONSORING, MONITORING AGENCY REPORT NUMBER	
European Research Office of the U.S.Army, USARDSG-UK Environmental Sciences Branch Mr. Tracey Carr, Edison House, 223 Old Marylebone Road London NW1 5TH, United Kingdom				
11. SUPPLEMENTARY NOTES				
12. DISTRIBUTION/AVAILABILITY STATEMENT			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words)				
<p>A search for insects and diseases which attack Eurasian watermilfoil (<i>Myriophyllum spicatum</i> L.), waterchestnut (<i>Trapa natans</i> L.) and hydrilla (<i>Hydrilla verticillata</i> Royle) that are the noxious water weeds in the US was made from the literature, taxonomic collections and through consultations with specialists. Totally, 32 insect and 7 fungi species associated with the targeted weeds were found on the territory of the former Soviet Union. Most of Insecta are Coleoptera (14 species) and Lepidoptera (7 species). Besides, 5 Diptera, 4 Trichoptera, and 2 Homoptera species were revealed. Majority of above mentioned insects and all fungi are connected with <i>M. spicatum</i> (28 species). Particularly, three weevil species: <i>Phytobius leucogaster</i>, <i>Pelenomus canaliculatus</i> and <i>Bagous geniculatus</i>, which are reportedly oligophagous on <i>Myriophyllum</i> spp. may be considered as potential biocontrol agents.</p>				
14. SUBJECT TERMS			15. NUMBER OF PAGES	
water weeds, biocontrol, <i>Myriophyllum</i> , <i>Trapa</i> , <i>Hydrilla</i> , Eurasia, insects, fungi			22	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT	18. SECURITY CLASSIFICATION OF THIS PAGE	19. SECURITY CLASSIFICATION OF ABSTRACT	20. LIMITATION OF ABSTRACT	

**RUSSIAN INSECTS AND DISEASES
THAT COULD CONTROL US WATERWEEDS**

ABSTRACT

Eurasian watermilfoil (*Myriophyllum spicatum* L.), waterchestnut (*Trapa natans* L.) and hydrilla (*Hydrilla verticillata* Royle) that are the noxious water weeds in the US occur as native species in the Former Soviet Union (FSU). A search for the insects and diseases which attack these plants on the territory of the FSU was made from the literature, taxonomic collections and through consultations with specialists. Totally, 32 insect and 7 fungi species associated with the targeted weeds were found. Most of Insecta are Coleoptera (14 species) and Lepidoptera (7 species). Besides, 5 Diptera, 4 Trichoptera, and 2 Homoptera species were revealed. Majority of above mentioned insects and fungi are connected with *M. spicatum* (28 species). Particularly, three weevil species: *Phytobius leucogaster*, *Pelenomus canaliculatus* and *Bagous geniculatus*, which are reportedly oligophagous on *Myriophyllum spp.*, may be considered as potential biocontrol agents.

Keywords: water weeds, biocontrol, *Myriophyllum*, *Trapa*, *Hydrilla*, Eurasia, insects, fungi.

TABLE OF CONTENTS

1. Introduction	2
2. A short annotated list of organisms connected with the targeted weeds	4
3. Data on insect species connected with the targeted weeds	6
4. Data on pathogenic fungi strains isolated from the targeted weeds	16
5. Conclusions	17
6. Literature cited	18

INTRODUCTION


Three aquatic weeds: Eurasian watermilfoil (*Myriophyllum spicatum* L.), waterchestnut (*Trapa natans* L.) and hydrilla (*Hydrilla verticillata* Royle) that are the serious problems in the US occur as native species in Russia and other areas of the Former Soviet Union (FSU). This research has the goal of locating and employing insects and/or diseases to reduce the population densities of these plants in the US. For this purpose, a search for the insects and diseases which attack water plant species from the genera *Myriophyllum*, *Trapa*, and *Hydrilla* on the territory of the FSU was made from the literature, taxonomic collections and through consultations with specialists.

A short systematic list of the organisms connected with the targeted water weeds is enclosed. A comprehensive data on each of insect and fungi species is contained in the next two chapters. Each insect species is described by:

1. Synonyms.
2. Geographical distribution within and outside the FSU.
3. The targeted water weed association.
4. Host range, i.e. other plants association.
5. Kind of host association, particularly damaged part of the host plant.
6. References, i.e. the sources of information concerning given species.

Each of phytopathogenic fungi strain is characterized by the same way, but collection locality is described, instead of geographic distribution.

Totally, data from 78 references are included in the report, of which 49 references are in Russian. Besides, insect collection of Zoological Institute (St. Petersburg) was examined and assistance of 9 consulting specialists (taxonomists and ecologists from 3 research institutes) was used. Names and positions of all these scientists are recorded after the list of literature cited. In case when the contribution of a consulting specialist was particularly significant, his or her name is also included in 'References' in the end of a section presented information concerning given species.



Main geographic regions of the former Soviet Union and approximate northern borders of distribution of Eurasian watermilfoil (*Myriophyllum spicatum* L.) and waterchestnut (*Trapa natans* L.).

A SHORT ANNOTATED LIST
OF ORGANISMS CONNECTED WITH THE TARGETED WEEDS

Abbreviations:

- M - connected with *Myriophyllum spicatum*
T - connected with *Trapa natans*
* reportedly oligophagous on *Myriophyllum spp.*
? supposedly, feeding on *Myriophyllum spp.* is occasional.
! recorded as insect pest of agricultural plants.

I N S E C T A

H O M O P T E R A

A p h i d a e

- MT ! *Rhopalosiphum nymphaea* L.
M ! *Aphis nasturtii* Kaltentbach

C O L E O P T E R A

C h r y s o m e l i d a e

D o n a c i i n a e

- M *Macrolea appendiculata* Panz.
M *Macrolea mutica* Fabricius
M *Neohaemonia voronovae* L. Medvedev

G a l e r u c i n a e

- T ! *Galerucella nymphaeae* L.
T *Galerucella nipponensis* Laboissiere

A p i o n i d a e

N a n o p h y i n a e

- T *Nanophyes pubescens* Roelofs
T *Nanophyes japonicus* Roelofs

C u r c u l i o n i d a e

C e u t o r h y n c h i n a e

- M ? *Rhinoncus albicinctus* Gyllenhal
M * *Phytobius leucogaster* Marsham
M * *Pelenomus canaliculatus* Fahraeus
M *Eubrychius velutus* Beck

B a g o i n a e

- M ? *Bagous nodulosus* Gyllenhal
M * *Bagous geniculatus* Huchhuth

E r i r h i n i n a e

- M *Tanysphyrus lemnae* Paykull

T R I C H O P T E R A

L e p t o c e r i d a e

- M *Mystacides longicornis* L.

P h r y g a n e i d a e

- M *Phryganea grandis* L.
M *Phryganea bipunctata* Retzius

L i m n o p h i l l i d a e

- M *Limnophyllus stigma* Curtis.

L E P I D O P T E R A

P y r a l i d a e

N y m p h u l i n a e

- MT *Acentria ephemerella* Denis et Schiffermuller
M *Cataclysta lemnata* L.
MT ! *Nymphula nymphaeata* L.
M *Nymphula stagnata* Donovan
MT *Paraponyx stratiotatum* L.
M *Paraponyx nivalis* Denis et Schiffermuller
P y r a u s t i n a e
M ! *Nomophila noctuella* Denis et Schiffermuller

D I P T E R A

C h i r o n o m i d a e

O r t h o c l a d i i n a e

- MT ! *Cricotopus silvestris* Fabr.
C h i r o n o m i n a e
M *Endochironomus ex.gr. dispar* Meig.
MT *Endochironomus tendens* Thien (Fabricius)
M *Glyptotendipes gripekoveni* Kieff
M *Tanytarsus ex.gr. lauterborny* Kieff.

F U N G I

- M *Fusarium oxysporum*
M *Fusarium nivale*
M *Fusarium solani*
M *Fusarium sambucinum*
M *Fusarium sp1*
M *Fusarium sp2*
M *Rhizoctonia aderholdii*

DATA ON INSECT SPECIES CONNECTED WITH THE TARGETED WEEDS

I N S E C T A

H O M O P T E R A

A p h i d a e

Rhopalosiphum nymphaea L.

(=Siphonaphis nymphaea)

(=Rhopalosiphum najadum)

(=Rh. alicmea)

(=Aphis nymphaea)

(=A. butoni)

(=A. infuscata)

(=A. prunaria)

(=A. consona)

(=A. prunorum)

(=Hyadaphis sparganii)

Distribution: almost whole world.

Target weed species: *Myriophyllum spicatum*, *Trapa natans*.

Host range: feed also on other vascular water plants, at the first half of summer - on Rosaceae: cherry, plum, apricot, peach).

Kind of host association: feed on stem, may cause significant damage.

References: Mordvilko, 1929; Boerner, 1952; Bei-Bienko, 1964; Lekic, Mihajlovic, 1970; Kryzhanovskii, 1974; Spencer, Lekic, 1974; Heie, 1986; Ler, 1988.

Aphis nasturtii Kalténbach

Distribution: Europe, European part of FSU, Central Asia, N. America.

Target weed species: *Myriophyllum spicatum*.

Host range: polyphagous, was recorded on many plants including potato, tomato, pumpkin (insect pest and virus vector).

Kind of host association: feed on stem, may cause significant damage.

References: Tarbinskii, Plavilshchikov, 1948; Lekic, Mihajlovic, 1970; Spencer, Lekic, 1974, Kryzhanovskii, 1974.

COLEOPTERA

Chrysomelidae

Donaciinae

Macrolea appendiculata Panz.
(= *Haemonia appendiculata* Latr.)

Distribution: Europe, European part of FSU, Siberia, Kazakhstan, Central Asia.

Target weed species: *Myriophyllum spicatum*.

Host range: supposedly polyphagous on various vascular water plants. Was recorded on *Potamogeton*, *Stratiotes*, *Ruppia*.

Kind of host association: larvae and adults feed on leaves, stem and roots.

References: Ogloblin, 1948; Bei-Bienko, 1965; Ogloblin, Medvedev, 1971; Kutikova, Starobogatov, 1977; Medvedev, Zaitzev, 1978; Lopatin, 1986; Medvedev, Roginskaya, 1988; Dubeshko, Medvedev, 1989.

Macrolea mutica Fabricius, *Macrolea mutica japona* Jacoby

Distribution: Europe, European part of FSU, Siberia, Caucasus, Kazakhstan, Central Asia, Russian Far East, Mongolia, Japan.

Target weed species: *Myriophyllum spicatum*.

Host range: supposedly polyphagous on various vascular water plants. Was recorded on *Zostera*, *Ruppia*, *Zannichellia*, *Potamogeton*.

Kind of host association: feed on plant tissue.

References: Lopatin, 1960; Bei-Bienko, 1965; Lopatin, 1977; Medvedev, 1982; Lopatin, 1986; Medvedev, Roginskaya, 1988; Dubeshko, Medvedev, 1989; Medvedev, 1992; Medvedev, Dubeshko, 1992.

Neohaemonia voronovae L. Medvedev, 1977

Distribution: Mongolia, Siberia?.

Target weed species: *Myriophyllum spicatum*.

Host range: was recorded also on *Potamogeton*.

Kind of host association: larvae feed on roots, adults feed on leaves.

References: Medvedev, 1977; Medvedev, Voronova, 1979; Medvedev, Zaitzev, 1980; Medvedev, 1982, Korotyaev, pers.comm.

G a l e r u c i n a e

Galerucella nymphaeae L.
(= *Pyrrhalta nymphaeae* L.)

Distribution: Europe, Siberia, Kazakhstan, Central Asia, Mongolia, China, N. America.

Target weed species: *Trapa natans*.

Host range: polyphagous species, numerous host specific 'forms' with little morphological differences were described. Feeds mostly on water and hydrophyte plants. However, a form damaging to currant was reported from Siberia and a form damaging to strawberry was reported from Europe. In experiments *G. nymphaeae* is able to feed even on pod (*Phaseolus*), willow (*Salix*), and birch (*Betula*).

Kind of host association: larvae and adults feed on leaves, at high population density may cause serious damage.

References: Ogloblin, 1936; Isakov, 1954; Medvedev, 1955; Kozhanchikov, 1958; Gaevskaya, 1966; Kryzhanovskii, 1974; Lopatin, 1977; Medvedev, 1982; Lopatin, 1986; Medvedev, Roginskaya, 1988; Medvedev, Dubeshko, 1992.

Galerucella nipponensis Laboissiere.

Distribution: Russian Far East, Korea, Japan.

Target weed species: *Trapa japonica* (=natans?).

Host range: supposedly polyphagous, was recorded also on *Brasiena*, *Ludwigia*, *Lycopus*.

Kind of host association: feed on plant tissue.

References: Dubeshko, Medvedev, 1989

A p i o n i d a e

N a n o p h y i n a e

Nanophyes pubescens Roelofs

Distribution: Japan, Russian Far East.

Target weed species: *Trapa natans*.

Host range: unknown.

Kind of host association: unknown

References: Morimoto, 1964; Korotyaev, pers.comm.

Nanophyes japonicus Roelofs

Distribution: Japan, Russian Far East?.

Target weed species: *Trapa natans*.

Host range: unknown.

Kind of host association: Larvae live in the petiole, feed on plant tissue.

References: Morimoto, 1964; Korotyaev, pers.comm.

C u r c u l i o n i d a e

C e u t o r h y n c h i n a e

Rhinoncus albicinctus Gyllenhal

Distribution: Europe, European part of FSU, Siberia, China.

Target weed species: *Myriophyllum spicatum*.

Host range: supposedly oligophagous, mainly on *Polygonum amphibium* and related species, feeding on *M. spicatum* is occasional.

Kind of host association: larvae are steam-miners.

References: Bei-Bienko, 1965; Lekic, Mihajlovic, 1970; Spencer, Lekic, 1974; Korotyaev, pers.comm.

Phytobius leucogaster Marsham

(= *Litodactylus leucogaster* Redtenbacher)

(= *L. griseomicans* Schwarz)

(= *L. myriophylli* Gyllenhal)

(= *Lithodactylus leucogaster* Marsham)

(= *Amalus leucogaster* auct.)

(= *Hydaticus leucogaster* Schoenherr)

Distribution: Europe, European part of FSU, Siberia, Kazakhstan, Central Asia, Russian Far East, N. America.

Target weed species: *Myriophyllum spicatum*.

Host range: possibly oligophagous on *Myriophyllum* spp., was recorded also on *M. verticillatum* and *M. sibiricum*.

Kind of host association: larvae and adults feed on leaves, stem and buds.

References: Reichardt, Ogloblin, 1940; Wesenberg-Lund, 1943; Ogloblin, 1948; Pavlovskii, Lepneva, 1948; Bei-Bienko, 1965; Gaevskaya, 1966; Spencer, 1974; Spencer, Lekic, 1974; Egorov, 1988; Isaev, 1994; Greed, Sheldon, 1994b; Caldara, O'Brien, 1995; Korotyaev, pers.comm.

Pelenomus canaliculatus Fahraeus

(= *Phytobius canaliculatus*)

Distribution: Europe, European part of FSU, Siberia, Kazakhstan, Mongolia, Japan.

Target weed species: *Myriophyllum spicatum*.

Host range: was recorded also on *M. verticillatum* L.

Possibly, oligophagous on *Myriophyllum* spp.

Kind of host association: larvae feed on leaves.

References: Ogloblin, 1948; Isaev, 1994; Caldara, O'Brien, 1995; Korotyaev, pers.comm.

Eubrychius velutus Beck
(= *E. aquaticus* Thomson)
(= *Lithodactylus velutus* Beck)
(= *Rhynchaenus velutus* Beck)

Distribution: Europe, Siberia, Kazakhstan, Central Asia, Mongolia, Russian Far East, N. America.

Target weed species: *Myriophyllum spicatum*.

Host range: supposedly polyphagous, was recorded also on *M. verticillatum* and *Potamogeton* sp.

Kind of host association: adults and larvae feed on leaves, stem and buds under water.

References: Reichardt, Ogloblin, 1940; Wesenberg-Lund, 1943; Bei-Bienko, 1965; Gaevskaya, 1966; Spencer, Lekic, 1974; Egorov, 1988; Isaev, 1994; Caldara, O'Brien, 1995; Korotyaev, pers.comm.

B a g o i n a e

Bagous nodulosus Gyllenhal

Distribution: Europe, Siberia, Central Asia, Mongolia.

Target weed species: *Myriophyllum spicatum*.

Host range: feed mostly on *Butomus umbellatus*, but was also reported from *Caltha palustris*. Feeding on *M. spicatum* is supposedly occasional.

Kind of host association: causes damage to the stem.

References: Lekic, Mihajlovic, 1970; Spencer, Lekic, 1974; Isaev, 1994; Korotyaev, pers.comm.

Bagous geniculatus Huchhuth.

Distribution: South Europe, Caucasus, Central Asia, Pakistan.

Target weed species: *Myriophyllum spicatum*.

Host range: supposedly oligophagous on *Myriophyllum* spp. (was recorded also on *M. tuberculatum*, *M. indicum*).

Kind of host association: feed on plant tissue.

References: Khan and al., 1969; Spencer, Lekic, 1974, Korotyaev, pers.comm.

E r i r h i n i n a e

Tanysphyrus lemnae Paykull.

Distribution: Europe, European part of FSU, Siberia, Russian Far East, Japan, N. America.

Target weed species: *Myriophyllum spicatum*.

Host range: polyphagous, was recorded mostly on *Lemna*, *Spirodela*, *Caltha*.

Kind of host association: adults and larvae feed on leaves.

References: Bei-Bienko, 1965; Spencer, Lekic, 1974; Isaev, 1994; Caldara, O'Brien, 1995, Korotyaev, pers.comm.

TRICHOPTERA
Leptoceridae

Mystacides longicornis L.
(=Phryganea longicornis L.)

Distribution: Europe, European part of FSU, Siberia, Kazakhstan, N. America.

Target weed species: *Myriophyllum spicatum*.

Host range: was recorded on various water plants.

Kind of host association: feeds on fresh and decaying plant tissue of algae and vascular plants.

References: Lepneva, 1940; Ogloblin, 1948; Kolenkina, 1951; McGaha, 1952; Lepneva, 1964; Lepneva, 1966; Gaevskaya, 1966; Fischer, 1966.

Phryganeidae

Phryganea grandis L.

Distribution: Europe, European part of FSU, Siberia.

Target weed species: *Myriophyllum spicatum*.

Host range: supposedly polyphagous on algae and on vascular plants. Have a seasonal change of food specialization. In the end of summer feed on larvae of other insects.

Kind of host association: may feed on plant tissues.

References: Kolenkina, 1951; Kashkin, 1958; Kashkin, 1961; Smirnov, 1962; Fischer, 1964; Lepneva, 1964, 1966; Sozska, 1975a, Sozska, 1975b; Ivanov, pers. comm.

Phryganea bipunctata Retzius
(= *Ph. striata* auctorum)

Distribution: Europe, European part of FSU, Siberia, Russian Far East, Mongolia.

Target weed species: *Myriophyllum spicatum*.

Host range: supposedly polyphagous on algae and on vascular plants.

Kind of host association: larvae feed both on plant tissues and small invertebrates, *Myriophyllum* leaves are used to make a case for the first instar larva.

References: Pavlovskii, Lepneva, 1948; Kolenkina, 1951; Kachalova, 1960; Lepneva, 1964; Fischer, 1964; Lepneva, 1966; Ivanov, pers. comm.

L i m n o p h i l i d a e

Limnophilus stigma Curtis.

Distribution: Europe, European part of FSU, Caucasus, Siberia, Russian Far East.

Target weed species: *Myriophyllum spicatum*.

Host range: supposedly polyphagous on algae and on vascular plants.

Kind of host association: probably use *Myriophyllum* leaves to make a larval case.

References: Lepneva, 1964; Lepneva, 1966, Ivanov, pers. comm.

L E P I D O P T E R A

P y r a l i d a e

N y m p h u l i n a e

Acentria ephemerella Denis et Schiffermuller

(=*Acentropus niveus* Oliver)

(=*Acentria nivea* Oliver)

Distribution: Europe, European part of the FSU, Siberia, N.America.

Target weed species: *Myriophyllum spicatum*, *Trapa natans*.

Host range: polyphagous on various vascular water plants.

Kind of host association: first instar larvae mine the food plants, then caterpillars feed on plants, cutting stems and removing leaves.

References: Grunberg, 1909; Rimskii-Korsakov, 1940; Ogloblin, 1948; Pavlovskii, Lepneva, 1948; Kokocinski, 1963; Gaevskaya, 1966; Soszka, 1975a; Urban, 1975; Medvedev, 1986; Palm, 1986; Green and al., 1989; Ioganzen, Rostovtsev, 1994; Greed, Sheldon, 1994a; Newman, Maher, 1995.

Cataclysta lemnata L.

Distribution: Europe, European Part of FSU, Caucasus, Central Asia.

Target weed species: *Myriophyllum spicatum*.

Host range: polyphagous on various vascular water and hydrophyte plants.

Kind of host association: young larvae made mines, then feed on leaves and cut stems.

References: Tarbinskii, Plavilshchikov, 1948; Kashkin, 1959; Lekic, Mihajlovic, 1970; Spencer, Lekic, 1974; Medvedev, 1986.

Nymphula nymphaeata L.

(=*Elophila nymphaeata* L.)
(=*Hydrocampa nymphaeata* L.)
(=*Nausinoe nymphaeata* L.)

Distribution: Europe, European part of FSU, Caucasus, Siberia, Kazakhstan, Central Asia, Russian Far East, Mongolia, China, Japan.

Target weed species: *Myriophyllum spicatum*, *Trapa natans*.

Host range: polyphagous on various vascular water plants. Larvae may also cause damage to rice.

Kind of host association: larvae live inside the plant, feed on plant leaves, stem or flowers.

References: Berg 1950; Kashkin, 1959; Kashkin, 1961; Lekic, Mihajlovic, 1970; Spencer, Lekic, 1974; Lvovski, pers. comm.

Nymphula stagnata Donovan

(=*Parapoynx stagnata* Donovan)

Distribution: Europe, European part of FSU, Caucasus, Siberia, Central Asia.

Target weed species: *Myriophyllum spicatum*.

Host range: supposedly polyphagous: was recorded also on *Sparganium* and *Nuphar*.

Kind of host association: larvae boring in leaves and stem.

References: Spencer, Lekic, 1974; Lekic, Mihajlovic, 1970; Palm, 1986, Lvovski, pers. comm.

Parapoynx stratiotatum L.

(= *Parapoynx*)
(= *P. stratiotata* L. auct.)

Distribution: Europe, European part of FSU, Caucasus, Siberia, Kazakhstan, Russian Far East.

Target weed species: *Myriophyllum spicatum*, *Trapa natans*.

Host range: polyphagous on various vascular water plants.

Kind of host association: larvae feed on fresh leaves, more rarely on decaying plants

References: Karny, 1934; Ogloblin, 1948; Pavlovskii, Lepneva, 1948; Kashkin, 1959; Kashkin, 1961; Gaevskaya, 1966; Spencer, 1974; Spencer, Lekic, 1974; Urban, 1975; Soszka, 1975a; Soszka, 1975b; Medvedev, 1986; Lvovski, pers. comm.

Parapoynx nivalis Denis & Schiffermuller

(= *Parapoynx*)

Distribution: Europe, European part of FSU.

Target weed species: *Myriophyllum spicatum*.

Host range: supposedly polyphagous, feeding on *M. spicatum* is rare.

Kind of host association: feed on plant leaves.

References: Lekic, Mihajlovic, 1970; Spencer, Lekic, 1974; Medvedev, 1986.

P y r a u s t i n a e

Nomophila noctuella Denis et Schiffermuller

Distribution: Europe, European part of FSU, Central Asia, Pakistan, N. America.

Target weed species: *Myriophyllum spicatum*.

Host range: polyphagous, may cause damage to alfalfa and clover.

Kind of host association: larvae feed on leaves.

References: Hasenfuss, 1960; Spencer, Lekic, 1974; Medvedev, 1986, Lvovski, pers. comm.

D I P T E R A

C h i r o n o m i d a e

O r t h o c l a d i i n a e

Cricotopus silvestris Fabr.

Distribution: almost whole world, including FSU.

Target weed species: *Myriophyllum spicatum*, *Trapa natans*.

Host range: associated with various vascular water and hydrophyte plants, may cause damage to rice.

Kind of host association: ubiquitous species associated with macrophytes, larvae live in leaf mines, feed on fragments of living and dead plant tissue, periphytic algae, and small animals.

References: Chernovskii, 1949; Gosteva, 1950; Thienemann, 1954; Shilova, 1955; Konstantinov, 1958; Gaevskaya, 1966; Pankratova, 1970; Spencer, Lekic, 1974; Soszka, 1975a, Soszka, 1975b; Cranston, 1995; Prat, Rodrigues-Capitulo, 1995.

C h i r o n o m i n a e

Endochironomus ex.gr. dispar Meig.

Distribution: Europe, European part of FSU.

Target weed species: *Myriophyllum spicatum*.

Host range: associated with various vascular water plants.

Kind of host association: miner, often using tunnels made by other animals, mostly stems appeared to be infested. Larvae feed on detritus, small animal remains, also on fresh and dead plant tissue.

References: Chernovskii, 1949; Konstantinov, 1958; Soszka, 1975a; Soszka, 1975b; Urban, 1975; Starostin, 1992.

Endochironomus tendens Thien (Fabricius)

Distribution: Europe, European part of FSU, Siberia, Russian Far East, Central Asia, N. America.

Target weed species: *Myriophyllum spicatum*, *Trapa natans*.

Host range: associated with various vascular water plants.

Kind of host association: feed mostly on detritus, small contribution of animal remains and periphytic algae. The tissue of plants slightly contributed to the food.

References: Chernovskii, 1949; Thienemann, 1954; Konstantinov, 1958; Gaevskaya, 1966; Bei-Bienko, 1969; Soszka, 1975a; Soszka, 1975b; Kutikova, Starobogatov, 1977; Cranston, 1995; Prat, Rodrigues-Capitulo, 1995; Caspers, Heimbach, 1995.

Glyptotendipes gripekoveni Kieff

Distribution: Europe, European part of FSU, W. Siberia.

Target weed species: *Myriophyllum spicatum*.

Host range: associated with various vascular water plants.

Kind of host association: Larvae feed mainly on periphyton, boring mines mostly in stems, the tissue of plants slightly contributed to the food.

References: Chernovskii, 1949; Ioganzen, Rostovtsev, 1994; Soszka, 1975a; Soszka, 1975b; Urban, 1975.

Tanytarsus ex.gr. lauterborny Kieff.

Distribution: Europe, European part of FSU, Central Asia.

Target weed species: *Myriophyllum spicatum*.

Host range: associated with various vascular water plants.

Kind of host association: feed on fresh and dead tissue of vascular plants, algae, detritus, animals.

References: Chernovski, 1949; Konstantinov, 1958; Soszka, 1975a, Soszka, 1975b; Starostin, 1992,

DATA ON PATHOGENIC FUNGI STRAINS
ISOLATED FROM THE TARGETED WEEDS
(strains with experimentally proved pathogenicity)

Fusarium oxysporum

Collection place: Vorskla river, Belgorod district, European part of FSU.

Target weed species: *Myriophyllum spicatum*.

Host range: host specificity was not tested.

Kind of host association: medium virulence.

References: Dolgovskaya et al., 1994; Labutova, pers.comm.

Fusarium nivale

Collection place: brook in Pendzhikent, Tadzhikistan, Central Asia.

Target weed species: *Myriophyllum spicatum*.

Host range: host specificity was not tested.

Kind of host association: medium virulence.

References: Dolgovskaya et al., 1994; Labutova, pers.comm.

Fusarium solani

Collection place: shallow lake in Zeravshan reservation, Samarkand district, Uzbekistan, Central Asia.

Target weed species: *Myriophyllum spicatum*.

Host range: host specificity was not tested.

Kind of host association: low virulence.

References: Dolgovskaya et al., 1994; Labutova, pers.comm.

Fusarium sambucinum

Collection place: pond near Novgorod, European part of FSU.

Target weed species: *Myriophyllum spicatum*.

Host range: host specificity was not tested.

Kind of host association: low virulence.

References: Dolgovskaya et al., 1994; Labutova, pers.comm.

Fusarium sp1

Collection place: brook in Pendzhikent, Tadzhikistan, Central Asia.

Target weed species: *Myriophyllum spicatum*.

Host range: host specificity was not tested.

Kind of host association: low virulence.

References: Dolgovskaya et al., 1994; Labutova, pers.comm.

Fusarium sp2

Collection place: shoal-water in Zeravshan reservation, Samarkand district, Uzbekistan, Central Asia.

Target weed species: *Myriophyllum spicatum*.

Host range: host specificity was not tested.

Kind of host association: low virulence.

References: Dolgovskaya et al., 1994; Labutova, pers.comm.

Rhizoctonia aderholdii

Collection place: shallow pond in Zeravshan reservation, Samarkand district, Uzbekistan, Central Asia.

Target weed species: *Myriophyllum spicatum*.

Host range: host specificity was not tested.

Kind of host association: medium virulence.

References: Dolgovskaya et al., 1994; Labutova, pers.comm.

C O N C L U S I O N S

Totally, 32 insect and 7 fungi species associated with the targeted water weeds (*Myriophyllum spicatum*, *Trapa natans* and *Hydrilla verticillata*) were recorded from the territory of the Former Soviet Union (FSU). Most of Insecta are Coleoptera (14 species) and Lepidoptera (7 species). Besides, 5 Diptera, 4 Trichoptera, and 2 Homoptera species were found.

Majority of above mentioned insects and all fungi are connected with *M. spicatum* (35 species), 10 of insect species were recorded on *T. natans*, while we failed to find any reports on insects or fungi associated with *Hydrilla verticillata* in FSU. The reason supposedly is that *M. spicatum* and *T. natans* are widespread and rather common across the whole FSU territory excluding North (see map at p.3), while *Hydrilla* is represented by several disjunctive patches and rarely come into the view of insect collectors. It is believed that special field search will revealed a lot of insect and fungi species connected with all of three targeted weeds.

Concerning the possibility of using for biological control, 6 insect species are unsuitable (*Rhopalosiphum nymphaea*, *Aphis nasturtii*, *Galerucella nymphaeae*, *Nymphula nymphaeata*, *Nomophila noctuella*, *Cricotopus silvestris*) because these insects were recorded as causing damage to agricultural plants. Besides, all of Trichoptera, Lepidoptera and Diptera species connected with targeted water weeds are polyphagous and some of them are potentially dangerous to any culturing plant which may be partly covered with the water or even grows in damp soil. Moreover, all of Trichoptera and most of Diptera species connected with *M. spicatum* and *T. natans* are euryphagous: their larvae are able to feed both on fresh and decaying tissue of algae and vascular plants, and even on small animals, using water weeds rather as a case or as a shelter, than as a food.

On the contrary, some of Coleoptera, particularly Curculionidae species seems to be promising for weed biocontrol. *Phytobius leucogaster*, *Pelenomus canaliculatus* and *Bagous geniculatus*, reportedly oligophagous weevils fed on *Myriophyllum* species may be considered as potential biocontrol agents. *Ph. leucogaster* is a very wide spread species which was recorded also in North America. At present, *P. canaliculatus* and *B. geniculatus* are absent in US and may be proposed for the further investigations. Primarily, laboratory tests are necessary to check their specificity. Generally speaking, laboratory testing may significantly enlarge the host range of all insect species under study.

R E F E R E N C E S

- Bei-Bienko G.Ya. (Ed.). 1964. Key to determination of the insects of the European part of the USSR, I. Apterygota, Palaeoptera, Neoptera: Hemimetabola. "Nauka", Moscow-Leningrad: 936 pp. (In Russian).
- Bei-Bienko G.Ya. (Ed.). 1965. Key to determination of the insects of the European part of the USSR, II. Coleoptera, Strepsiptera. "Nauka", Moscow-Leningrad: 668 pp. (In Russian).
- Bei-Bienko G.Ya. (Ed.). 1969. Key to determination of the insects of the European part of the USSR, V. Diptera, Aphaniptera, Part 1. "Nauka", Leningrad: 807 pp. (In Russian).
- Berg C.O. 1950 Biology of certain aquatic caterpillars (Pyralididae: *Nymphula* spp.) which feed on Potamogeton. Transactions of the American Microscopical Society, 69(3): 254-266.
- Boerner C. 1952. Die Blattlaus Mitteleuropas. Schrift der Thuringischen Landesarbeitsgemeinschaft für Heilspflanzenkunde und Heilpflanzenbeschaffung in Weimar. Heft 4 und Mitteilungen der Thuringischen Botanischen Gesellschaft, 3. Weimar: 259 pp. (Lief 2): 250-284.
- Caldara R., O'Brien C.W. 1995. Curculionidae: Aquatic weevils of China (Coleoptera). In: Jach M.A. & L.Ji (Eds.) Water Beetles of China, 1. Wien: 389-408.
- Caspers N., Heimbach F. 1995. The Chironomids of an experimental pond system. In: Cranston P. (Ed.). Chironomids from genes to ecosystems. CSIRO, Australia: 273-275.
- Chernovskii A.A. 1949. Key to determination of the larvae of gnats of the family Tendipedidae. Keys to determination of the fauna of the USSR, published by Zoological Institute of Academy of Sciences of the USSR, 31. Izdatel'stvo AN SSSR, Moscow -Leningrad: 185 pp. (In Russian).
- Cranston P. (Ed.). 1995. Chironomids from genes to ecosystems. CSIRO, Australia: 482 pp.
- Dolgovskaya M.Yu, Podlipaev S.A., Reznik S.Ya., Volkovitch M.G. 1994. Preliminary results of the isolation of pathogenic fungi associated with Eurasian Water Milfoil, *Myriophyllum spicatum* L. in the former Soviet Union. Abstracts of 9th International Symposium of Aquatic Weeds. (Dublin, 12-16 September 1994) :13.
- Dubeshko L.N., Medvedev L.N. 1989. The ecology of leaf beetles of the Siberia and the Far East. Izdatel'stvo Irkutskogo Universiteta, Irkutsk: 224 pp. (In Russian).
- Egorov A.B. 1988. New data on the distribution and ecology nearwater weevils of subfamily Ceutorhynchinae (Coleoptera, Curculionidae) in the fauna of the Far East. In: Levanidova I.M., Makarchenko Ye.A. (Eds.). Fauna, systematics and biology of a freshwater invertebrates. Vladivostok: 60-66 (In Russian).
- Fischer F.C.J. 1964. Trichopterorum catalogus. V. Phryganeidae, Limnacentropodidae, Molannidae. Nederlandsche Entomologische Vereeniging, Amsterdam: 214 pp.
- Fischer F.C.J. 1966. Trichopterorum catalogus. VII. Leptoceridae. Pars 2. Nederlandsche Entomologische Vereeniging, Amstredam: 163 pp.
- Gaevskaya N.S. 1966. High aquatic plants in nutrition of freshwater animals. "Nauka", Moscow: 327 pp. (In Russian).
- Gosteva M.N. 1950. The gnat *Cricotopus silvestris* as a pest of fish spawn. Zoologicheskyy Zhurnal, 29(2): 187-188. (In Russian).

- Greed R.P., Sheldon S.P. 1994a. The effect of two herbivorous insect larvae on Eurasian Watermilfoil. *Journal of Aquatic Plant Management*, 32: 21-26.
- Greed R.P., Sheldon S.P. 1994b. Aquatic weevils (Coleoptera: Curculionidae) associated with northern watermilfoil (*Myriophyllum sibiricum*) in Alberta, Canada. *Entomological news* 105(2): 98-102.
- Green I.S., Smith D.L., Boylen C.W. 1989. Electron microscopic observations of the aquatic caterpillar *Acentropus niveus* (Olivier). *Proceedings of Annual Meeting of Electron Microscopic Society of America*. San Francisco Press, Inc. San Francisco, California: 906-907.
- Grunberg K. 1909. *Lepidoptera, Schmetterlinge. Die Susswasserfauna Deutschlands*. 8.
- Hasenfuss I. 1960. *Die larvalsystematic der Zunsler (Pyralidae)*. Akademie Verlag, Berlin: 263 pp.
- Heie O.E. 1986. The Aphidoidea (Homoptera) of Fennoscandia and Denmark. III. *Fauna Entomologica Scandinavia*, 17: 314 pp.
- Ioganzen A.A., Rostovtsev A.A. (Eds.). 1994. The lake Ubinskoe. Biological productivity and perspectives of piscicultural usage. St. Petersburg: 144 pp. (In Russian).
- Isaev A.Yu. 1994. Ecological-faunistic review of the weevils (Coleoptera: Apionidae, Rhynchophoridae, Curculionidae) of Ul'yanovsk district. *Priroda Ul'yanovskoi oblasti*, 4. Ul'yanovsk: 77 pp. (In Russian).
- Isakov Yu.A. 1954. To the methodology of investigation of nutrition of the waterfowl birds. *Bulleten' Moskovskogo Obshchestva Ispytatelei Prirody [MOIP]. Otdelenie biologii*, 59(1): 33-36 (In Russian).
- Kachalova O.L. 1960. Nutrition of the larvae of some water insects - fodder objects of fishes. *Izvestiya AN Latvii SSR*, 6(155): 147-152 (In Russian).
- Karny H.H. 1934. *Biologie der Wasserinsekten*. Wien: 311 pp.
- Kashkin N.I. 1958. Consumption of higher water vegetation by the larvae of *Phryganea grandis* L. *Trudy Moskovskogo Technicheskogo Instituta Rybnoi Promyshlennosti i Khoziaistva*, 9: 146-160 (In Russian).
- Kashkin N.I. 1959. Nutrition of caterpillars of *Paraponyx stratiotata* L., *Nymphula nymphaeata* L. and *Cataclysta lemnata* L. (Pyralidae, Lepidoptera). *Trudy Moskovskogo Technicheskogo Instituta Rybnoi Promyshlennosti i Khoziaistva*, 10: 162-176 (In Russian).
- Kashkin N.I. 1961. To the extent of utilization of higher water plants by some phytophagous invertebrates (on an example of Iakhromskoe storage pool of the channel Moscow - Volga). *Trudy Murmanskogo Morskogo biologicheskogo Instituta*, 3(7): 170-184 (In Russian).
- Khan A.G., Rahim A., Ahmad F. 1969. Studies on the natural enemies of *Myriophyllum* spp. (for the U.S.A. and New Zealand). *Commonwealth Agricultural Bureaus, Commonwealth Institute of biological control*: 68-69.
- Kokocinski W. 1963. Remarks on biology of *Acentropus niveus* Oliv. (Lepidoptera, Pyralidae). *Polskie Pismo entomologiczne*, 33 (2): 233-243.
- Kolenkina L.V. 1951. Nutrition of the larvae of some Trichoptera. *Trudy Vsesoyuznogo Gidrobiologicheskogo obshchestva*, III: 44-47 (In Russian).

- Konstantinov A.S. 1958. Biology of Chironomids and their cultivation. Izdatel'stvo "Kommunist", Saratov: 362 pp. (In Russian).
- Kozhanchikov I.V. 1958. Biological peculiarities of the european species of the genus Galerucella and differentiation of biological forms in Galerucella lineola F. Trudy Zoologicheskogo Instituta AN SSSR, 24. Izdatel'stvo AN SSSR, Moscow-Leningrad: 271-332 (In Russian).
- Kryzhanovskii O.L. (Ed.). 1974. Insects and ticks - pests of agricultural plants, 2. Coleoptera. "Nauka", Leningrad: 235 pp. (In Russian).
- Kutikova L.A., Starobogatov Ya.I. (Eds.). 1977. Key to determination of the freshwater invertebrates of the European part of the USSR. (Plancton and bentos). "Gidrometeoizdat", Leningrad: 510 pp. (In Russian).
- Lekic M, Mihajlovic L. 1970. Entomofauna of Myriophyllum spicatum L. (Haloragidaceae) as an aquatic weed in Jugoslavia. Arhiv za Poljoprivedne Nauke, 23. Beograd: 63-76.
- Lepneva S.G. 1940. Trichoptera. In: Pavlovskii Ye.N., Zhadin V.I. (Eds.). Life of fresh waters. 1. Moskva-Leningrad: 191-223 (In Russian).
- Lepnevã S.G. 1964. Trichoptera. Fauna of the USSR, 2, Part 1. Izdatel'stvo AN SSSR, Moskva-Leningrad: 562 pp. (In Russian).
- Lepneva S.G. 1966. Trichoptera. Fauna of the USSR, 2, Part 2. Izdatel'stvo AN SSSR, Moskva-Leningrad: 562 pp. (In Russian).
- Ler P.A. (Ed.). 1988. Key to determination of the insects of the Far East of the USSR, II. Homoptera and Hemiptera. "Nauka", Leningrad: 972 pp. (In Russian).
- Ler P.A. (Ed.). 1992. Key to determination of the insects of the Far East of the USSR, III. Coleoptera, Part 2. "Nauka", St.Petersburg: 704 pp. (In Russian)
- Lopatin I.K. 1960. Materials on the fauna and ecology of leaf beetles (Coleoptera, Chrysomelidae) of the Southern bank of the Dnieper River. Entomologicheskoe Obozrenie, 39(3): 629-642 (In Russian).
- Lopatin I.K. 1977. The leaf beetles (Chrysomelidae) of Central Asia and Kazakhstan. Keys to determination of the fauna of the USSR, published by Zoological Institute of Academy of Sciences of the USSR, 113. "Nauka", Leningrad: 270 pp. (In Russian).
- Lopatin I.K. 1986. The leaf beetles in Byelorussian and Baltic faunas (Key to determination). Vysshaya shkola, Minsk: 131 pp. (In Russian).
- McGaha Y.J. 1952. The limnological relations of insects to certain aquatic flowering plants. Transactions of the American Microscopical Society, 71: 355-382.
- Medvedev G.S. (Ed.). 1986. Key to determination of the insects of the European part of the USSR, IV. Lepidoptera, Part 3. "Nauka", Leningrad: 504 pp. (In Russian).
- Medvedev L.N. 1955. The leaf beetles. In: Vrediteli lesa, 2. Moskva-Leningrad: 546-578 (In Russian).
- Medvedev L.N. 1977. Finding of the member the American genus Neochaemonia Szek. (Coleoptera, Chrysomelidae) in Mongolia. Doklady AN SSSR, 236(2): 488-490 (In Russian).
- Medvedev L.N. 1982. The leaf beetles of Mongolian People's Republic. Key to determination. "Nauka", Moscow: 303 pp. (In Russian).

- Medvedev L.N. 1992. In: Ler P.A. (Ed.). Key to determination of the insects of the Far East of the USSR, 3. Coleoptera, Part 2. "Nauka", S. Petersburg: 533-602 (In Russian).
- Medvedev L.N., Dubeshko L.N. 1992. Key to determination of the leaf beetles of the Siberia. Isdatel'stvo Irkutskogo Universiteta, Irkutsk: 224 pp. (In Russian).
- Medvedev L.N., Roginskaya E.Ya. 1988. Catalogue of host-plants of the leaf beetles of the USSR. Moscow: 192 pp. (In Russian).
- Medvedev L.N., Voronova N.V. 1979. On the chrysomelid-beetle fauna of Mongolia (Coleoptera, Chrysomelidae), 3. Nasekomye Mongolii, 6. "Nauka", Leningrad: 97-134 (In Russian).
- Medvedev L.N., Zaitzev Yu.M. 1978. The larvae of the leaf beetles of the Siberia and Far-East. "Nauka", Moscow: 182 pp. (In Russian).
- Medvedev L.N., Zaitzev Yu.M. 1980. New data on chrysomelid beetles larvae from Mongolia (Coleoptera, Chrysomelidae). Nasekomye Mongolii, 7. "Nauka", Leningrad: 97-106 (In Russian).
- Mordvilko A.K. 1929. Food plant of the Aphididae of the USSR. Trudy po prikladnoi entomologii, 14(1). Bureau of Applied Entomology, State Institute of Experimental Agronomy. Leningrad: 101 pp. (In Russian).
- Morimoto K. 1964. Key and illustrations for the identification of the curculionoid-beetles of Japan and the Ryukyus. Kontyu, 32(1): 81-90.
- Newman R.M., Maher L.M. 1995. New records and distribution of Aquatic insect herbivores of watermilfoils (Haloragidaceae: Myriophyllum spp.) in Minnesota. Entomological News, 106(1): 6-18.
- Ogloblin D.A. 1936. The leaf beetles, Galerucinae. Fauna SSSR, Nasekomye zhestkokrylye, 26(1). Isdatel'stvo AN SSSR, Moscow-Leningrad: 455 pp. (In Russian).
- Ogloblin D.A. 1948. Chrysomelidae. The leaf beetles. pp. 472-507. In: Tarbinskii S.P., Plavilshchikov N.N. (Eds.). Key to determination of the insects of the European part of the USSR. "Nauka", Moscow-Leningrad: 1128 pp. (In Russian).
- Ogloblin D.A., Medvedev L.N. 1971. The larvae of the leaf beetles (Coleoptera, Chrysomelidae) of the European part of the USSR. Keys to determination of the fauna of the USSR, published by Zoological Institute of Academy of Sciences of the USSR, 106. "Nauka", Leningrad: 123 pp. (In Russian).
- Palm E. 1986. Nordeuropas Pyralider - med saerlight henblik pa den danske fauna (Lepidoptera: Pyralidae). Denmarks Dyreliv, 3. Fauna Boger Kobenhaven: 287pp.
- Pankratova V. Ya. 1970. Larvae and pupae of Chironomids of subfamily Orthocladiinae of the USSR (Diptera - Chironomidae = Tendipedidae). Keys to determination of the fauna of the USSR, published by Zoological Institute of Academy of Sciences of the USSR, 102. Izdatel'stvo AN SSSR, Leningrad: 341 pp. (In Russian).
- Pavlovskii E.N., Lepneva S.G. 1948. Sketches on the life of freshwater animals. "Sovetskaya Nauka", Leningrad: 458 pp. (In Russian).
- Prat N., Rodrigues-Capitulo A. 1995. Population dynamics of Chironomidae associated with *Nymphaea alba* leaves in springs in the Ebro Delta (N.E. Spain). pp. 297-304. In: Cranston P. (Ed.). Chironomids from genes to ecosystems. CSIRO, Australia: 482 pp.

- Reichardt A.N., Ogloblin D.A. 1940. Beetles (Coleoptera). In: Pavlovskii Ye.N., Zhadin V.I. (Eds.). Life of fresh waters, 1. Moscow-Leningrad: 158-188 (In Russian).
- Rimskii-Korsakov M.N. 1940. Lepidopterous or butterflies (Lepidoptera). In: Pavlovskii Ye.N., Zhadin V.I. (Eds.). Life of fresh waters, 1. Moscow-Leningrad: 224-226 (In Russian).
- Shilova A.I. 1955. To the nutrition of the larva of *Cricotopus silvestris* F. in connection with the character of structure of mouthparts. Doklady AN SSSR, 100(6): 1191-1193 (In Russian).
- Smirnov N.N. 1962. On nutrition of Caddis Worms *Phryganea grandis* L. Hydrobiologia, 19(3): 252-261.
- Soszka G.J. 1975a. The invertebrates on submerged Macrophytes in three Masurian Lakes. Ecologia Polska, 23: 371-391.
- Soszka G.J. 1975b. Ecological relations between invertebrates and submerged Macrophytes in the lake littoral. Ecologia Polska, 23: 393-415.
- Spencer N.R. 1974. Insect enemies of aquatic weeds. PANS, 20(4): 444-450.
- Spencer N.R., Lekic M. 1974. Prospects for biological control of Eurasian Watermilfoil. Weed Science, 22(4): 401-404.
- Starostin I.V. 1992. Fauna of the inner water pools of the Turkmenistan. "Ylym", Ashgabat: 256 pp. (In Russian).
- Tarbinskii S.P., Plavilshchikov N.N. (Eds.). 1948. Key to determination of the insects of the European part of the USSR. "Sel'khozgiz", Moscow-Leningrad: 1127 pp. (In Russian).
- Thienemann A. 1954. Chironomus. Leben Verbreitung und wirtschaftliche Bedeutung der Chironomiden. In: Die Bunnengewasser, 20. Stuttgart: 834 pp.
- Urban E. 1975. The mining fauna in four macrophyte species in Mikolajskie Lake. Ecologia Polska, 23(3): 417-435.
- Wesenberg-Lund C. 1943. Biologie der Susswasserinsekten. J.Springer, Berlin - Wien: 682 pp.

A LIST OF CONSULTING SPECIALISTS

- Dr. **E.V.Balushkina**, Laboratory of Fresh Water Hydrobiology, Zoological Institute, St.Petersburg, Russia (Diptera: Chironomidae).
- Dr. **A.F.Emeljanov**, Laboratory of Insect Taxonomy, Zoological Institute, St.Petersburg, Russia (Homoptera).
- Dr. **M.I.Falkovich**, Laboratory of Insect Taxonomy, Zoological Institute, St.Petersburg, Russia (Lepidoptera).
- Dr. **V.D.Ivanov**, Entomology Department, Faculty of Biology, St. Petersburg State University (Trichoptera).
- Dr. **A.B.Korotyaev**, Laboratory of Insect Taxonomy, Zoological Institute, St.Petersburg, Russia (Coleoptera).
- Dr. **N.M.Labutova**, Institute of Agricultural Microbiology, St.Petersburg, Russia (Fungi).
- Dr. **S.Yu.Sinev**, Laboratory of Insect Taxonomy, Zoological Institute, St.Petersburg, Russia (Lepidoptera).
- Dr. **Ya.I.Starobogatov**, Laboratory of Fresh Water Hydrobiology, Zoological Institute, St.Petersburg, Russia (Freshwater invertebrates).
- Dr. **A.B.Stekol'schikov**, Laboratory of Insect Taxonomy, Zoological Institute, St.Petersburg, Russia (Homoptera: Aphidae).