


AL/AO-TR-1996-0065



**A  
R  
M  
S  
T  
R  
O  
N  
G  
  
L  
A  
B  
O  
R  
A  
T  
O  
R  
Y**

**1995 ANNUAL US AIR FORCE TUBERCULOSIS REPORT**

**Joy M. Miller  
Martin A. Puckett**

**AEROSPACE MEDICINE DIRECTORATE  
EPIDEMIOLOGIC RESEARCH DIVISION  
EPIDEMIOLOGY SERVICES BRANCH  
2601 West Gate Road, Suite 114  
Brooks Air Force Base, TX 78235-5241**

**May 1996**

**Final Technical Report for 1995**

Approved for public release; distribution is unlimited.

**19960624 244**

**DTIC QUALITY INSPECTED 1**

**AIR FORCE MATERIEL COMMAND  
BROOKS AIR FORCE BASE, TEXAS**

## NOTICES

When Government drawings, specifications, or other data are used for any purpose other than in connection with a definitely Government-related procurement, the United States Government incurs no responsibility or any obligation whatsoever. The fact that the Government may have formulated or in any way supplied the said drawings, specifications, or other data, is not to be regarded by implication, or otherwise in any manner construed, as licensing the holder, or any other person or corporation; or as conveying any rights or permission to manufacture, use, or sell any patented invention that may in any way be related thereto.

The mention of trade names or commercial products in this publication is for illustration purposes and does not constitute endorsement or recommendation for use by the United States Air Force.


The Office of Public Affairs has reviewed this report, and it is releasable to the National Technical Information Service, where it will be available to the general public, including foreign nationals.

Government agencies and their contractors registered with Defense Technical Information Center (DTIC) should direct requests for copies to: Defense Technical Information Center (DTIC), 8725 John J. Kingman Road, Suite 0944, Ft. Belvoir, VA 22060-6218.

Non-Government agencies may purchase copies of this report from: National Technical Information Services (NTIS), 5285 Port Royal Road, Springfield, VA 22161.

This technical report has been reviewed and is approved for publication.

  
JOY M. MILLER, Captain, USAF, BSC  
Project Scientist

  
JAMES A. WRIGHT, Colonel, USAF, MC  
Chief, Epidemiologic Research Division

# REPORT DOCUMENTATION PAGE

Form Approved  
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188) Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE May 1996	3. REPORT TYPE AND DATES COVERED Final Report 1995	
4. TITLE AND SUBTITLE  1995 Annual US Air Force Tuberculosis Report		5. FUNDING NUMBERS	
6. AUTHOR(S) Joy M. Miller Martin A. Puckett		8. PERFORMING ORGANIZATION REPORT NUMBER  AL/AO-TR-1996-0065	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Armstrong Laboratory Aerospace Medicine Directorate Epidemiologic Research Div/Epidemiology Services Branch 2601 West Gate Road, Suite 114 Brooks AFB TX 78235-5241		10. SPONSORING / MONITORING AGENCY REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)		11. SUPPLEMENTARY NOTES	
12a. DISTRIBUTION / AVAILABILITY STATEMENT		12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words)  This report evaluates the 1995 US Air Force (USAF) Tuberculosis (TB) Prevention and Control Program. The report analyzes TB mortality, morbidity, and tuberculosis screening data from 88 medical treatment facilities worldwide. The data are compared to US tuberculosis incidence data and to USAF morbidity and screening data for 1992-1994. In 1995, there were no reported deaths attributed to tuberculosis. Among USAF active duty personnel, the tuberculosis incidence rate continued to decline (1995 incidence per 100,000 personnel was 0.49 compared to 1994 incidence of 0.71). Out of the 131,883 TB skin tests (TSTs) administered to active duty personnel, 2,940 were eligible for isoniazid (INH) therapy. Among the nonactive duty population, there were 93,201 TSTs administered. A total of 2,467 nonactive duty persons were positive reactors under 35 years of age or recent converters (71.2% placed on INH).			
14. SUBJECT TERMS Tuberculosis, recent reactors, converter, Tuberculosis Skin Test (TST)			15. NUMBER OF PAGES 20
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED			16. PRICE CODE
18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED	19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED	20. LIMITATION OF ABSTRACT UL	

# Table of Contents

## 1995 Annual US Air Force Tuberculosis Report

International TB Profile Summary.....	1
National TB Profile Summary.....	1
USAF TB Profile Summary.....	2
Tuberculosis Mortality in the USAF.....	2
Tuberculosis Morbidity in the USAF.....	2
USAF Tuberculosis Screening and Chemoprophylaxis Program.....	2
Limitations of the Data.....	3
Table 1.....	4
Table 2.....	4
Figure 1.....	5
Figure 2.....	6
References.....	7
Additional Reading.....	9
Appendix 1.....	11

# 1995 Annual US Air Force Tuberculosis Report

## Introduction

United States (US) military missions, including humanitarian support and other peace-keeping activities, in areas with endemic tuberculosis (TB) represent a health risk for United States Air Force (USAF) personnel. TB is transmitted as airborne droplets that may stay suspended in the air for long periods. Persons living or working in communities with endemic tuberculosis may become infected even without direct, face-to-face contact with native populations. Protection of USAF personnel and their dependents requires vigilant surveillance using approved methods of administering tuberculin skin tests and prompt, complete treatment of those individuals identified as newly positive or recent converters.

This report presents information on the TB prevention and control program within the USAF. The narrative section of the report describes the 1995 USAF data on tuberculin skin test results, the number of active cases of TB, and the epidemiologic trends. To put the USAF TB experience in perspective, a summary of the international and national TB profile is provided. The second section provides the substantiating data and includes the tables and graphs showing the USAF data compared to US data.

## International TB Profile Summary<sup>1</sup>

An estimated 1.7 billion people (one-third of the world population) are currently infected with TB, including nearly half of the world's refugees. Approximately eight million new cases of tuberculosis occur each year. In 1990, 4% of the new cases of TB were attributable to human immunodeficiency virus (HIV) co-infection. By the year 2000, this number is expected to increase to 14%. Worldwide, an estimated 30 million persons are infected with drug-resistant TB. The majority of persons infected with active TB will be young parents and workers, often in their most productive years.

In 1995, nearly 3 million people worldwide died from TB. Tuberculosis now kills more adults than any other infectious disease, including Acquired Immune Deficiency Syndrome (AIDS), malaria, cholera, and other tropical diseases combined. Over 95% of these deaths will be in the developing world where TB accounts for 25% of avoidable adult mortality.

## National TB Profile Summary<sup>2,3</sup>

From 1985 through 1992, the number of TB cases reported annually in the US surged 20%, from 22,201 to 26,673. Factors associated with the increase of TB during this time period include the HIV/AIDS epidemic; immigration of persons from countries where TB incidence rates are 10-30

<sup>1</sup> World Health Organization Report on the Tuberculosis Epidemic, 1995  
([http://www.who.ch/programmes/gtb/GTB\\_Homepage.html](http://www.who.ch/programmes/gtb/GTB_Homepage.html))

<sup>2</sup> CDC. Tuberculosis Morbidity, United States, 1994. MMWR 1995; 44:387-95

<sup>3</sup> CDC. Press release. World TB Day News Conference, March 22, 1996

times higher than in the United States; transmission of TB among persons residing in congregate settings such as hospitals, prisons, and homeless shelters; and declines in resources for TB control. In 1992, increased funding for TB programs allowed improved management of TB cases and thus intensified efforts against the disease. In 1993, the Division of TB Elimination (DTBE), Centers for Disease Control and Prevention (CDC), implemented an expanded TB surveillance system to monitor and target groups at risk for TB, assess drug-susceptibility results for initial and final *M. tuberculosis* isolates for each culture-positive TB patient, and evaluate outcome of TB cases.

Recently the CDC reported there was a 6.4% decrease in the number of reported TB cases in the US, from 24,361 new cases reported in 1994 to 22,812 new cases in 1995 (case rates decreased from 9.4 per 100,000 in 1994 to 8.7 per 100,000 in 1995). This represents the third consecutive year that reported cases of TB have declined. Several factors contributed to the decline in reports of new TB cases, including increased program funding and diligent application of directly observed therapy (DOT). Unfortunately, in 1995, 40% of the states reported either no change or an increase in TB cases from the previous year. TB rates increased 28% in Arizona, 11% in Minnesota, 9-10% in Iowa, Louisiana, and Pennsylvania, and 7% in Wisconsin.

## **Air Force TB Profile Summary**

### **Tuberculosis Mortality in the USAF**

Analysis of the Retrospective Case Mix Analysis Systems (RCMAS) data base showed no deaths associated with tuberculosis either among USAF active duty or dependent/retired personnel who were admitted to Department of Defense hospitals. The data base records disposition of USAF personnel, their dependents, and retirees who were hospitalized. The search included primary diagnosis of tuberculosis (ICD code 010-018.9) as cause of death.

### **Tuberculosis Morbidity in the USAF**

**Table 1, USAF TB Detection and Control Program by MAJCOM, 1994**, summarizes the reports from each of the medical treatment facilities (MTF) within the major commands (MAJCOM). Divided into two sections, Table 1 contains data on tuberculin skin test results and number of active cases of TB identified among active duty personnel (AD) and among dependents and retirees (nonactive duty [NAD]).

Two cases of tuberculosis were reported among the AD population and 30 cases among NAD duty personnel. In 1995, the incidence rate of TB among AD personnel was 0.49 per 100,000. **Figure 1** shows the trend of TB incidence rate since 1982 and compares the USAF experience with that of the US. Historically, the AD USAF TB incidence rate has been considerably lower than the US rate. The trend for the AD USAF population shows a decline of tuberculosis overall, with two peaks occurring in 1983 and 1987. The rate has declined since 1987.

### **USAF Tuberculosis Screening and Chemoprophylaxis Program**

In 1995, a total of 225,084 tuberculin skin tests (TST) were administered. This represents an increase for AD of 13.6% and a decrease of 2.6% for NAD from the number of tests administered

in 1994. The increase in TST administered to AD personnel may reflect added screening due to deployments and peace-keeping missions in endemic areas.

In 1995, there were 5,407 persons (2,940 AD and 2,467 NAD) testing positive on the tuberculin skin test. A total of 4,451 people were placed on isoniazid (INH) as a result of the screening program (**Table 1**). The percentage of TST positive AD personnel placed on INH has risen from a low of 71% in 1991 to 91.7% in 1995 (**Figure 2**). Among NAD, the percentage has remained relatively constant (71.2% for 1995), probably reflecting the fact that this population is not mandated to report for follow-up therapy.

MTFs reported a total of 1,123 people were either not placed on, or were removed from, INH therapy (**Table 2**). Of this group, those removed due to pregnancy or permanent change of station (PCS) typically receive follow-up after delivery or upon arrival at their next duty station. **Table 1** shows the number of persons diagnosed at a previous duty station who are receiving INH therapy at the current duty station.


#### **Limitations of the Data**


The terms "active duty" and "nonactive duty" have not been defined in AFI 48-115, *TB Detection and Control Program*. The active duty group may encompass active duty USAF, Air National Guard, and Reserves and active duty members of other services who obtain care in USAF MTFs. The nonactive duty population comprises those other than active duty (retirees, dependents, or other beneficiaries) who utilize the MTF. MTFs reporting by major command are listed in **Appendix 1**.

The incidence rate of TB among AD personnel was determined by using the mid-year population data obtained from the *1995 Annual US USAF Sexually Transmitted Disease Report*. Incidence rates of newly diagnosed AD positive reactors < 35 years of age and those of AD recent converters could not be determined because the denominator data were not stratified by age or TST result category. Incidence rates for the NAD duty population cannot be determined since mid-year population data are not available for this group.

  
JOY M. MILLER, Capt, USAF, BSC  
Chief, Disease Surveillance

Approved

  
JAMES A. WRIGHT, Colonel, USAF, MC, CFS  
Chief, Epidemiologic Research Division

  
MARTIN A. PUCKETT, SSgt, USAF  
NCOIC, Epidemiology Services Branch

**TABLE 1. USAF TB DETECTION AND CONTROL PROGRAM  
BY MAJCOM, 1995**

MAJCOM											
ACTIVE DUTY	ACC	AFSPC	USAFE	AFMC	USAFA	ADW	AMC	AETC	PACAF	AFSOC	TOTAL
<b>Newly Diagnosed Positive Under Age 35</b>	132	37	59	70	4	2	253	686	158	5	1,406
Placed on INH	105	28	30	56	3	2	224	613	137	5	1,203
Percentage on INH	80%	76%	51%	80%	75%	100%	89%	89%	87%	100%	86%
New Positive - Previous Base	21	11	4	12	0	2	19	20	27	4	120
Placed on INH	21	10	4	9	0	2	15	17	24	3	105
Percentage Placed on INH	100%	91%	100%	75%	N/A	100%	79%	85%	89%	75%	88%
<b>Recent Converters Any Age</b>	227	72	145	151	8	10	378	348	189	6	1,534
Placed on INH	191	63	91	122	8	10	348	295	120	5	1,253
Percentage on INH	84%	88%	63%	81%	100%	100%	92%	85%	63%	83%	82%
Recent Converters - Previous Base	43	14	9	9	2	6	21	17	25	2	148
Placed on INH	41	12	9	9	2	5	17	15	22	2	134
Percentage Placed on INH	95%	86%	100%	100%	100%	83%	81%	88%	88%	100%	91%
<b>Total Newly Diagnosed Positive or Converter</b>	359	109	204	221	12	12	631	1034	347	11	2940
Total on INH	358	113	134	196	13	19	604	940	303	15	2695
Percent on INH	99.7%	103.7%	65.7%	88.7%	108.3%	158.3%	95.7%	90.9%	87.3%	136.4%	91.7%
<b>Total TB Skin Tests Given and Read</b>	14,872	2,944	11,929	8,955	672	1,509	24,636	49,290	16,665	411	131,883
Number of active TB Cases	0	0	0	0	0	0	2	0	0	0	2
<b>MIDYEAR POPULATION</b>	105,800	23,499	36,866	57,373	6,774	2,744	54,429	71,076	41,528	7,000	407,089
<b>RATE OF ACTIVE TB PER 100,000</b>											0.49
NONACTIVE DUTY											
<b>Newly Diagnosed Positive Under Age 35</b>	230	52	153	124	8	4	175	77	487	11	1,321
Placed on INH	151	39	68	80	3	4	131	49	310	7	842
Percentage Placed on INH	66%	75%	44%	65%	38%	100%	75%	64%	64%	64%	64%
New Positive - Previous Base	21	1	6	11	0	1	11	8	17	0	76
Placed on INH	19	0	5	8	0	1	9	7	13	0	62
Percentage Placed on INH	90%	0%	83%	73%	N/A	100%	82%	88%	76%	N/A	82%
<b>Recent Converters Any Age</b>	200	56	165	110	2	6	125	210	265	7	1,146
Placed on INH	147	50	80	62	2	6	95	161	188	6	797
Percentage Placed on INH	74%	89%	48%	56%	100%	100%	76%	77%	71%	86%	70%
Recent Converters - Previous Base	20	4	5	4	0	1	4	16	5	2	61
Placed on INH	17	3	5	2	0	1	4	16	5	2	55
Percentage Placed on INH	85%	75%	100%	50%	N/A	100%	100%	100%	100%	100%	90%
<b>Total Newly Diagnosed Positive or Converter</b>	430	108	318	234	10	10	300	287	752	18	2467
Total on INH	334	92	158	152	5	12	239	233	516	15	1756
Percent on INH	77.7%	85.2%	49.7%	65.0%	50.0%	120.0%	79.7%	81.2%	68.6%	83.3%	71.2%
<b>Total TB Skin Tests</b>	17,508	5,272	14,339	12,049	481	1,042	11,710	11,628	18,287	885	93,201
Number of active TB Cases	6	2	0	0	0	0	10	6	6	0	30

**Table 2. SUMMARY OF REASONS WHY PATIENTS WERE NOT PLACED ON INH PROPHYLATIC THERAPY**

	ACC	AFSPC	USAFE	AFMC	USAFA	ADW	AMC	AETC	PACAF	AFSOC	TOTAL
Pregnancy	82	16	54	32	3	1	25	33	64	5	315
Liver Disease	4	1	1	0	0	0	0	2	1	0	9
Patient Refusal	5	5	16	19	1	0	16	4	41	0	107
PCS	37	4	150	22	1	0	11	52	142	0	419
Other Medical	74	14	27	79	1	0	72	103	105	0	475
<b>TOTAL NOT PLACED INH</b>	<b>202</b>	<b>40</b>	<b>248</b>	<b>152</b>	<b>6</b>	<b>1</b>	<b>124</b>	<b>194</b>	<b>353</b>	<b>5</b>	<b>1123</b>

Figure 1

ACTIVE DUTY USAF & US POPULATION  
ACTIVE TUBERCULOSIS CASE RATES, 1982-1995





Figure 2

AIR FORCE PERSONNEL  
PERCENT TST POSITIVE PLACED ON INH PREVENTIVE THERAPY, 1990-1995



### References

1. American Thoracic Society/CDC. Treatment of tuberculosis and tuberculosis infection in adults and children. *Am J Respir Crit Care Med* 1994;149:1359-74.
2. CDC. Tuberculosis Morbidity, United States, 1994. *MMWR* 1995;44:387-95.
3. CDC. Proportionate mortality from pulmonary tuberculosis associated with occupations--28 States, 1979-1990. *MMWR* 1995; 44 (1):14-19.

### Additional Reading

1. CDC. Summary of notifiable diseases, United States, 1993. MMWR 1994; 42 (53):3-12, 58-61.
2. CDC. Expanded tuberculosis surveillance and tuberculosis morbidity--United States, 1993. MMWR 1994; 43 (20):361-6.
3. CDC. Guidelines for preventing the transmission of *Mycobacterium tuberculosis* in health care facilities, 1994. MMWR 1994; 43 (RR-13):1-312.
4. CDC. Tuberculosis control laws - United States, 1993, recommendations of the Advisory Council for the Elimination of Tuberculosis (ACET). MMWR 1993; 42(RR-15).
5. CDC. Approaches to improving adherence to antituberculosis therapy - South Carolina and New York, 1986-1991. MMWR 1993; 42( 4):75-81.
6. CDC. Initial therapy for tuberculosis in the era of multidrug resistance, recommendations of the Advisory Council for the Elimination of Tuberculosis. MMWR 1993; 42( RR-7).
7. CDC. Tuberculosis Morbidity - United States, 1992. MMWR 1993; 42:696-7, 703-4.
8. CDC. Exposure of passengers and flight crew to *Mycobacterium tuberculosis* on commercial aircraft, 1992-1995. MMWR 1995; 44(8):137-140.
9. Clark RA. OSHA enforcement policy and procedures for occupational exposure to tuberculosis. Infect Control Hosp Epidemiol 1993; 14:694-8.
10. Decker MD. OSHA enforcement policy for occupational exposure to tuberculosis. Infect Control Hosp Epidemiol 1993; 14:689-93.
11. Ellner JJ, Hinman AR, Dooley SW, et al. Tuberculosis symposium: emerging problems and promise. J Infect Dis 1993; 168:537-51.
12. Fraser VJ, Kilo CM, Bailey TC, Medoff G, Dunagan WC. Screening of physicians for tuberculosis. Infect Control Hosp Epidemiol 1994; 15:95-100.
13. Gerberding JL. Occupational infectious diseases or infectious occupational diseases? Bridging the views on tuberculosis control. Infect Control Hosp Epidemiol 1993; 14:686-8.
14. Gostin LO. Controlling the resurgent tuberculosis epidemic. A 50-state survey of TB statutes and proposals for reform. JAMA 1993; 269:255-61.

15. Iseman MD. Treatment of multidrug-resistant tuberculosis. *N Engl J Med*. 1993; 329:784-91.
16. Jarvis WR, et al. Respirators, recommendations, and regulations: The controversy surrounding protection of health care workers from tuberculosis. *Ann Int Med* 1995;122-146.
17. Muder RR, Brennen C, Yu KT. Choosing appropriate criteria for tuberculin positivity and conversion in a long-term care facility. *Infect Control Hosp Epidemiol* 1993; 14:523-6.
18. Menzies D, et al. Tuberculosis among health care workers. *N Engl J Med* 1995; 332:92-98.
19. Sepkowitz KA. Tuberculosis and the health care worker: A historical perspective. *Ann of Intern Med* 1994; 120:71-9.
20. Shearer BG. MDR-TB - another challenge from the microbial world. *JADA* 1994; 125:43-9.
21. Smith BR. OSHA's TB rulemaking. *Occupational Health and Safety*. April 1995:48-51.
22. Tuberculosis. *Med. Clin. North Am.* 1993; 77.
23. Voelker R. New federal stances on TB control may be confusing to health care facilities. *JAMA* 1993; 270:1903-4.
24. Weis SE, Slocum PC, Blais FX, King B, Nunn M, Matney B, Gomez E, Foresman BH. The effect of directly observed therapy on the rates of drug resistance and relapse in tuberculosis. *NEJM* 1994; 330:1179-84.

## Appendix 1

### Reporting Installation Medical Treatment Facilities by MAJCOM

ACC - Barksdale, Beale, Cannon, Castle, Davis-Monthan, Dyess, Ellsworth, Griffiss, Holloman, Howard, K.I. Sawyer, Lajes, Langley, Little Rock, MacDill, Minot, Moody, Mt. Home, Nellis, Offutt, Pope, Seymour Johnson, Shaw, Whiteman

AETC - Altus, Columbus, Goodfellow, Keesler, Lackland, Laughlin, Luke, Maxwell, Randolph, Reese, Sheppard, Tyndall, Vance

AFMC - Brooks, Edwards, Eglin, Hanscom, Hill, Kelly, Kirtland, Los Angeles, McClellan, Robins, Tinker, Wright-Patterson

AFSPC - Peterson, Vandenberg, Malmstrom, Patrick, Onizuka, FE Warren

AMC - Andrews, Charleston, Dover, Fairchild, Grand Forks, March, McChord, McConnell, McGuire, Plattsburgh, Scott, Travis

PACAF - Andersen, Elmendorf, Eielson, Hickam, Kadena, Kunsan, Misawa, Osan, Yokota

ADW - Bolling

USAFE - RAF Alconbury, Aviano, RAF Chicksands, Incirlik, Izmir, Lakenheath, Ramstein, Spangdahlem/Bitburg, Sembach, Rhein-Main

USAF Academy