

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE	3. REPORT TYPE AND DATES COVERED FINAL REPORT 01 Mar 93 - 31 Jul 96	
4. TITLE AND SUBTITLE OPTICAL METROLOGY OF MAGNETICALLY TRAPPED HYDROGEN			5. FUNDING NUMBERS 61102F 2301/DS	
6. AUTHOR(S) Dr Kleppner			AFOSR-TR-96 97 0010	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) MIT, Room 26-237 Cambridge, MA 02139			10. SPONSORING / MONITORING AGENCY REPORT NUMBER F49620-93-1-0215	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) AFOSR/NE 110 Duncan Avenue Suite B115 Bolling AFB DC 20332-8080			11. SUPPLEMENTARY NOTES	
12a. DISTRIBUTION / AVAILABILITY STATEMENT APPROVED FOR PUBLIC RELEASE: DISTRIBUTION UNLIMITED			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) A new technique for spectroscopy of atomic hydrogen has been developed, opening the way to a thousand-fold increase in spectroscopic resolution of hydrogen or any neutral atomic species, and perhaps making possible a new type of UV optical atomic clock. Two-photon Doppler-free spectroscopy of atoms confined in a magnetic trap has been carried out to observe the 1S to 2S two-photon (243nm) transition to a metastable state with an extremely long lifetime, 1/8 sec. The trap is highly non-perturbative, permitting observation times limited ultimately by the natural lifetime, yielding a natural linewidth of about 1 Hz. Initially a resolution of about three kHz was achieved, higher than the best then achieved by other techniques and apparently limited only by laser jitter. Signal rates as high as 3000 counts per second were observed. The collection efficiency in the initial version of the apparatus is less than 10 ⁻⁵ , and the actual signal rate is greater than 10 ⁸ counts per second, almost astronomically high compared to previous methods. A direct measurement of the life time of the metastable 2S atoms was made and found to be close to the theoretical lifetime of 0.12 sec. This is the first time the 2S lifetime has been measured.				
14. SUBJECT TERMS			15. NUMBER OF PAGES	
			16. PRICE CODE	
			20. LIMITATION OF ABSTRACT	
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED	18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED	19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED		

19970109 056

FAX COVER SHEET

To: Dr. Ralph Kelley

From: Daniel Kleppner

M.I.T., room 26-237

Cambridge, MA. 02139

phone: (617) 253-6811

FAX : (617) 253-4876

internet: DK@kleppner.mit.edu

Date Wed Dec 18 10:17:52 EST 1996

Pages (including cover): 3

MASSACHUSETTS INSTITUTE OF TECHNOLOGY
DEPARTMENT OF PHYSICS

DANIEL KLEPPNER
Lester Wolfe Professor of Physics
Associate Director, Research Laboratory of Electronics

address: MIT
room 26-237
Cambridge, MA 02139

phone: 617/253-6811
fax: 617/253-4876
dk@amo.mit.edu

December 18, 1996

Dr. Ralph Kelley
Air Force Office of Scientific Research
Bolling Air Force Base, Building 410
Washington, DC 20332

Dear Dr. Kelley,

This is the Final Report for AFOSR Grant F49620-93-1-0215.

Under this grant we have developed a new technique for spectroscopy of atomic hydrogen which opens the way to a thousand-fold increase in spectroscopic resolution of hydrogen—or indeed any neutral atomic species—and may make possible a new type of optical atomic clock operating in the UV region.

We have succeeded in carrying out two-photon Doppler-free spectroscopy of atoms that are confined in a magnetic trap. The transition we observe, the $1S \rightarrow 2S$ two-photon (243nm) transition, is to a metastable state with an extremely long lifetime, 1/8 sec. The trap is highly non-perturbative, permitting observation times limited ultimately by the natural lifetime, yielding a natural linewidth of about 1 Hz. In our initial experiments we achieved a resolution of about three kHz, higher than the best then achieved by other techniques and apparently limited only by laser jitter, which we are now trying to improve. We have observed signal rates as high as 3000 counts per second. The collection efficiency in the initial version of our apparatus is less than 10^{-5} , and so the actual signal rate is greater than 10^8 counts per second, which compared to previous methods, is almost astronomically high. In addition, we have directly measured the life time of the metastable $2S$ atoms, and have observed it to be close to the theoretical lifetime of 0.12 sec. This is the first time the $2S$ lifetime has been measured.

The performance of our system gives promise that a new type of optical clock, based on the $1S \rightarrow 2S$ two-photon transition in hydrogen, may be feasible. We have started an analysis of this system with a view to creating such a clock. In addition, the spectroscopic techniques developed in this work provide a powerful tool for studying Bose-Einstein condensation in hydrogen, and the possible creation of a coherent atomic beam of hydrogen- a hydrogen atom laser.

PUBLICATIONS AND THESES

Two-photon spectroscopy of trapped atomic hydrogen, Claudio L. Cesar, Dale G. Fried, Thomas C. Killian, Adam D. Polcyn, Jon C. Sandberg, Ite A. Yu, Thomas J. Greytak, Daniel Kleppner, and John Doyle, Phys. Rev. Lett. **77**, 255 (1996)

Two-Photon Spectroscopy of Trapped Atomic Hydrogen, C. L. Cesar, D. G. Fried, T. C. Killian, A. D. Polcyn, J. C. Sandberg, J. M. Doyle, I. A. Yu, T. J. Greytak and D. Kleppner, Proceedings of the Fifth Symposium on Frequency Standards and Metrology, J. C. Bergquist, ed. (World Scientific, Singapore) 1996, p 365.

Doppler-Free Spectroscopy of Trapped Atomic Hydrogen, T.C. Killian, D.G. Fried, C.L. Cesar, A.D. Polcyn, T.J. Greytak and Daniel Kleppner, Proceedings of the Fifteenth International Conference of Atomic Physics, to be published

(In preparation) Two-photon Doppler-Free Spectroscopy of Trapped Atoms, Claudio L. Cesar and Daniel Kleppner

THESES

Claudio L. Cesar, Ph.D. thesis Two-Photon Spectroscopy of Trapped Atomic Hydrogen, M.I.T., December 1995

Sincerely,



Daniel Kleppner
Lester Wolfe Professor of Physics