

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188


Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE 4/21/97	3. REPORT TYPE AND DATES COVERED FINAL REPORT 4/1/92 - 3/31/94	
4. TITLE AND SUBTITLE The Use of Combinatorial Heavy and Light Chain Libraries and Site Specific Mutagenesis to Create Antibody Biosensors for Metal Ions			5. FUNDING NUMBERS G N00014-91-J-1593	
6. AUTHOR(S) Stephen J. Benkovic			8. PERFORMING ORGANIZATION REPORT NUMBER N/A	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) The Pennsylvania State University Department of Chemistry 414 Wartik Laboratory University Park, PA 16802				
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Harold Bright Office of Naval Research 800 N. Quincey Street Arlington, VA 22217-5000			10. SPONSORING/MONITORING AGENCY REPORT NUMBER N/A	
11. SUPPLEMENTARY NOTES N/A				
12a. DISTRIBUTION/AVAILABILITY STATEMENT No known limitations			12b. DISTRIBUTION CODE	
DISTRIBUTION STATEMENT A Approved for public release Distribution Unlimited				
13. ABSTRACT (Maximum 200 words) We investigated the potential catalytic activity of the Zn(II) containing antibody against a variety of ester, and phosphomonoester and diester substrates. They were designed on the basis of their structural resemblance to the 43C9 hapten. None of the materials proved to be substrates for the Zn(II) containing antibody and the project was terminated.				
19970528 023				
14. SUBJECT TERMS 43C9 antibody, p-nitrophenol, F _{ab} fragments			15. NUMBER OF PAGES 3	
			16. PRICE CODE N/A	
17. SECURITY CLASSIFICATION OF REPORT unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT unclassified	20. LIMITATION OF ABSTRACT UL	

Grant# N00014-91-J-1593

April 21, 1997

I wish to make the attached Annual Progress Report serve as part of my Final Report.



Date: 4-22-97

Stephen J. Benkovic
Evan Pugh Professor and
Eberly Chair in Chemistry

DTIC QUALITY INSPECTED 3

ANNUAL PROGRESS REPORT

GRANT#: N00014-91-J-1593

PRINCIPAL INVESTIGATOR: Stephen J. Benkovic

INSTITUTION: The Pennsylvania State University

GRANT TITLE: The Use of Combinatorial Heavy & Light Chain Libraries and Site Specific Mutagenesis to Create Antibody Biosensors for Metal Ions

REPORTING PERIOD: June 1, 1993 - March 31, 1994

AWARD PERIOD: April 1, 1992 - March 31, 1994

OBJECTIVE: To construct F_{ab} fragments that possess metal ion binding ligands in juxtaposition to the antigen combining site, so that the binding of both antigen and metal produces observable chemical or spectral changes in the antigen.

APPROACH: A recursive protocol will be used to create the required F_{ab} fragments. i) Antibodies with the desired binding functions will be induced with the appropriate immunogen and F_{abs} obtained from recombinant libraries, ii) F_{abs} will be screened for antigen binding and those with appropriate affinities isolated, sequenced and overexpressed in *E.coli*; iii) The structures of selected F_{abs} will be deduced from modeling (in collaboration with Drs. Getzoff and Roberts, Scripps Clinic) and amino acid motifs required for metal ion binding will be introduced by site specific mutagenesis; iv) Purified F_{abs} will be tested for their metal ion affinity and response to a transducing substrate; v) improvements in the binding and catalytic properties of the F_{ab} will be sought by additional rounds of mutagenesis and/or chain shuffling with the original light or heavy chain libraries.

ACCOMPLISHMENTS (last 12 months): We have selected the antibody, 43C9, which efficiently catalyzes the hydrolysis of aromatic ester and anilides as a monoclonal antibody, and have converted it to a single chain form, 43C9SCA. The expression vector, pJS118 (documented in our previous annual progress report) permits secretion of the SCA from *E.coli* and serves as a vehicle for site specific mutagenesis.

A Zn(II) bind site has been introduced into catalytic antibody 43C9, which accelerates the hydrolysis of an aromatic amide by a factor of 10^6 over the rate of the uncatalyzed reaction. The binding site was constructed using a computer model of the 43C9 Fv fragment and the ligand field was based on that found in carbonic anhydrase. In 43C9, the site consisted of three histidine residues: an existing histidine at position H35 as well as two site-directed mutations (N-H33-H and Y-H95-H). The mutant 43C9 antibody binds Zn(II) with a K_D value of $1.54 \pm 0.09 \mu M$ and the binding of Zn(II) quenches the intrinsic antibody fluorescence by 85%, allowing this antibody to function

as a biosensor for Zn(II). The large extent of fluorescence quenching by the diamagnetic Zn(II) ion may result from changes in hydrogen bonding between one of the putative ligands (His H35) and Trp H47. This site was highly selective for Zn(II) and the binding affinities for Mg(II), Ca(II), Mn(II), Fe(III), Co(II), Ni(II), Cu(II) and Cd(II) were at least two orders of magnitude lower. By measuring the absorbance spectrum of *p*-nitrophenol bound to the mutant antibody in the presence of increasing concentrations of Zn(II), we demonstrated that the antibody can simultaneously accommodate both Zn(II) and *p*-nitrophenol and that they are in sufficiently close proximity to interact electronically.

SIGNIFICANCE: We have our first evidence that we have constructed a metal binding antibody with high metal ion specificity but also capable of binding an external ligand.