


^C
~~SP~~-96-4041

The Isomer State of HF-178(16+) Studing:
Theoretical Investigation.

Taras M. Zajac

Uzhgorod State University, Department of Physics,
294000, Voloshin st., 32/142, Uzhgorod,
UKRAINE



19970814 071

REPORT DOCUMENTATION PAGE

Form Approved OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE 1997	3. REPORT TYPE AND DATES COVERED Final Report	
4. TITLE AND SUBTITLE The Isomer State of HF-178(16+): Theoretical Investigation		5. FUNDING NUMBERS F6170896W0251	
6. AUTHOR(S) Dr. Taras Zajac			
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Uzhgorod State University Voloshina St. 32/142 Uzhgorod 294000 Ukraine		8. PERFORMING ORGANIZATION REPORT NUMBER N/A	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) EOARD PSC 802 BOX 14 FPO 09499-0200		10. SPONSORING/MONITORING AGENCY REPORT NUMBER SPC 96-4041	
11. SUPPLEMENTARY NOTES			
12a. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution is unlimited.		12b. DISTRIBUTION CODE A	
13. ABSTRACT (Maximum 200 words) This report results from a contract tasking Uzhgorod State University as follows: The contractor will investigate the nuclear levels of the 178Hf isomer and the transition probabilities between the levels as described in his proposals.			
14. SUBJECT TERMS		15. NUMBER OF PAGES 24	
		16. PRICE CODE N/A	
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED	18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED	19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED	20. LIMITATION OF ABSTRACT UL

NSN 7540-01-280-5500

Standard Form 298 (Rev. 2-89)
Prescribed by ANSI Std. Z39-18
298-102

Report Special Project SPS-96-4041 of Contract
F61708-96-W0251, Notice of Contract Award

Contents. p.

The Isomer State of HF-178(16+) Studing: Theoretical
Investigation.

Abstract 3

Introduction

4

1.Vibrative-rotative modell and total hamiltonian for
deformed nuclei

5

2.Coriolisis interaction for deformed nuclei and noneadiabatic
effects 9

3.Nillson shell modell and probability ratio calculating

12

4. The results of calculation ratio $B(E\lambda)$ for transitions in
178-Hf 15

Abstract.

In this paper total experimental and theoretical data about Hf-178 levels are presented: levels of Hf-178 and decay, electromagnetic momentum of Hf-178, ratio $B(E2)$, $B(E3)$, $B(E4)$ and others ratio and parameters has been calculated in Nilsson model. The results of calculation are presented, and compared with experimental data and others calculations.

Introduction.

The scheme of long-lifetime isomer γ -laser seen to be the most natural and easy for experimental layout. Such may was already proposed in the first application by L.A.Rivlin [1]. Using long-lifetime nuclei as initial object for γ -laser creation was also suggested by other authors [2,3]. Excited radioactive nuclei in long-lifetime isomer state are radiochemically separated from nuclei in ground state.

Then during a period, much lower than the lifetime, a crystal is grown of radioactive isomer by one of the known techniques, e.g. Bridgman technique of crystallization from melt. In such crystal the inversion condition holds practically always. Then it is necessary to cool the crystal to low temperatures to obtain the conditions for Mossbauer effect observation, and the crystal begins to emit. But since one fails to preserve the line of natural width, the stimulated emission will not occur.

And therefore all activity on creation of long-lifetime isomer γ -laser was directed to search of methods for Mossbauer radiation linewidth reduction to its natural value.

Line broadening may be uniform and non-uniform. The uniform line broadening arises e.g. at vibrations of atoms in the crystal lattice, or electron spin fluctuations in paramagnetic crystals. Similar reasons broadening can be eliminated or minimized by the process of freeing e.g. by deep cooling of the sample to the temperature much lower than 1 K. It is quite possible to realise such conditions at modern level of physics and engineering. But there is a lot of technical obstacles on this way, and the problem seems to become purely experimental. However, theoretical studies, especially, in our opinion, elucidation of the mechanism of interaction of gamma-quanta with the nucleus for each separate specific level, studying of the degree of collectivity of various levels excitation processes and

studies of their structure, should localize and reduce the number of the most urgent problems.

The longest-lifetime isomer, known at present is ^{178}Hf , having been found in [4] with the energy of 2446.0 (16+, K=16). In the same paper the scheme of this isomeric state decay was proposed [6]. The structure of ^{178}Hf level was studied by considering decay of two ^{178}Ta isomers and in (n,g) reaction. ^{178}Hf nucleus is a very interesting, because it is an even-even nucleus. There is a wide variety of data available on the studies of the levels of this nucleus. The studies of this nucleus seem to be very interesting, since the interest to the problem of long-lifetime isomer g-laser is still strong, what was demonstrated at Garlas-95 school [5-9].

Our view point on this problem consists in studying the nature of the level, the transitions between which are important for creating inverse population of $^{178m2}\text{Hf}$ isomer by calculating probabilities of the transitions between the levels, using various models. It is the level nature which determines the model choice. Since, as noted above, the higher levels are a mixture of a great amount of states, as a rule, of all the states of the band, it seems important to construct realistic bases for the studies of these states and the choice of the corresponding interaction potentials.

1. VIBRATIONAL-ROTATIONAL MODEL AND TOTAL HAMILTONIAN FOR DEFORMED NUCLEI.

It is known, that in the region of axial-symmetric deformed nuclei for lowly situated excitation states the projection of spine I into the axis of symmetry of nucleus K is being a good quantum number [10]. Together with this position, it is interesting to find out in what conditions the non-maintainance of the number K is begin and how this transition is going through.

With the increase of energy excitation the density of nucleus levels grows greatly and some levels are changing into the combinations of different configurations with the varied numbers of quasi-partical [11]. However, if in such exiations it were possible to neglect the powers of Coriolis the levels would form the rotation bands being characterized by a quantum number within the systems of axial symmetry [10].

From the results of the analysis on data of neutrons capture by nuclei it is possible to conclude, that for the neutron resonance states connected with the energy excitations of nuclea, near to the energy of connection of neuron B_n . The possible dimensions of the projection K by the given meaning are practically equally possible [10-11]. The possible mechanism causing the non-maintainance of the K is the Coriolis's mixing.

The Corriolis's interaction leads to the connection of inner and collective rotation of deformed nucleus and mixing states with different K and within strong Coriolis's combination the resulting state of the nucleus J^π will present the superposition of states $I^\pi K$,

with all possible at the given meanings of the K. In the frameworks of the abovementioned issues certain interest presents the study of K-isomer stimulation of depended nuclei in the reactions of inelastic scattering gamma-quanta.

For the nuclei with constant deformation of interaction is:

$$\mathbb{H}_{int} = \sum_i f(r_i) \sum_{\mu} \alpha_{2\mu}^* Y_{2\mu}^{(i)} \dots \quad (1.1)$$

between individual nucleons and collective vibrations is rather tight and the use of perturbation theory is not positive any more. Therefore the connection with the deformed core of nucleus must be considered precisely. This leads to the model of films for deformed nuclei, that is to the monopartitive films in the deformed potential. The full Hamiltonian has a general structure:

$$\mathbb{H} = \mathbb{H}_{coll} + \mathbb{H}_{sp} + \mathbb{H}_{int} \quad (1.2),$$

where

$$\mathbb{H}_{sp} = \sum_i \left\{ \mathbb{T}_i + V(r_i, \mathbb{P}_i, \mathbb{S}_i) \right\} \quad (1.3)$$

$\mathbb{H}_{sp}$ – Hamiltonian of shell model, $\mathbb{T}_i$ – kinetic energy, V_i – potential of shell model for i particle.

However, the collective Hamiltonian now presents the correspondent collective Hamiltonian of the deformed nucleus and namely vibrative-rotative Hamiltonian. The used projected model is illustrated on figure 1, which shows how two nucleons move around

the deformed core while the core itself according to the vibrative-rotative model can vibrate and rotate.

It is understandable that the task of such kind may be solved with the use of other models as well being based on the model of asymmetric rotator. For the low situated collective excitations the both models give almost the same results, we prefer the dynamic approach of the vibrative-rotative model.

The Hamiltonian of the vibrative-rotative model consists of rotational moment of the core $\hat{M}$. The total angular momentum of the system $\hat{P}$ consists of the moment of the core and the moment of appeared particles $\hat{J}$, $\hat{J} = \sum_{v=1,2,\dots} \hat{J}_v$, where $\hat{J}_v$ - angular momenta of individual particles.

Substituting $\hat{M} = \hat{P} - \hat{J}$ into the vibrative-rotative Hamiltonian (1.2) it is easy to get H_{coll} for deformed nuclei, which preserve some additional parts over core. The result has the following form:

$$\hat{H} = \hat{H}_{\text{rot}} + \hat{H}_{\text{vib}} + \hat{H}_{\text{vibrot}} \quad (1.4)$$

where

$$\hat{H}_{\text{rot}} = \frac{(\hat{P} - \hat{J})^2 \hbar^2 - (\hat{P}_3 - \hat{J}_3) \hbar^2}{2\mathcal{J}_0} + \frac{(\hat{P} - \hat{J})^2 \hbar^2}{16B\eta^2}; \quad (1.4 \text{ a})$$

$$\hat{H}_{\text{vib}} = \frac{\hbar^2}{2B} \left[\frac{\partial^2}{\partial \xi^2} + \frac{1}{2} \frac{\partial^2}{\partial \eta^2} \right] + \frac{1}{2} C_{\sigma} \xi^2 + C_{\sigma} \eta^2 - \frac{\hbar^2}{16B\eta^2}; \quad (1.4 \text{ b})$$

$$\begin{aligned}
H_{vibrot} = & \frac{(\mathbb{P} - \mathcal{J})^2 h^2 - (\mathbb{P}_3 - J_3) h^2}{2 \mathcal{J}_0} \left[2 \frac{\eta^2}{\beta_0^2} - 2 \frac{\xi}{\beta_0} + 3 \frac{\xi^2}{\beta_0^2} \right] + \\
& + \frac{(\mathbb{P}_+ - \mathcal{J}_+)^2 h^2 - (\mathbb{P}_- - J_-) h^2}{4 \mathcal{J}_0} \left[2\sqrt{6} \frac{\xi\eta}{\beta_0^2} - \frac{2}{3} \sqrt{6} \frac{\eta}{\beta_0} \right] \quad (1.4 c)
\end{aligned}$$

and the element of volume in space of collective coordinates $d\tau = d\Omega d\xi d\eta$.

It is interesting to mention that the prediction of film model for deformed nuclei concerning different monopartitive levels are almost true. Most of the predicted stimulated levels are found in experiment even if their energy is in compatible with the theory. Spins of the ground states are well explained in the frames of this model. However, the theoretical predictions are based on the precise meaning of nucleus deformation which must be formed of the data of meanings $B(E2)$ for transition within the limits of the main band. Deformation being obtained this way is always connected with experimental mistakes and one-partical spectrum may be approximated only within the limits of these.

If the Nilsson's levels are set up one can try to count the other features of deformed nuclei using knowledge of the correspondent monopartitive wave functions. For example, one can try to predict the possibilities $E1$, $M1$, $E2$ - transition.

2. CORIOLIS INTERACTION FOR DEFORMED NUCLEI.

Let us consider the hamiltonian of perturbation H' in the form:

$$\begin{aligned}
 H' = & \frac{\hbar^2}{2\mathcal{I}_0} \mathcal{J}^2 - \frac{\hbar^2}{2\mathcal{I}_0} [\mathcal{P}_+ \mathcal{J}_- + \mathcal{P}_- \mathcal{J}_+ + 2\mathcal{P}_3 \mathcal{J}_3] + \\
 & \frac{\hbar^2}{2\mathcal{I}_0} [\mathcal{P}_+^2 - \mathcal{P}_3^2 + \mathcal{J}_+^2 - \mathcal{J}_3^2 + \mathcal{P}_+ \mathcal{J}_- - \mathcal{P}_- \mathcal{J}_+] \left[\frac{2\eta^2}{\beta_0^2} - 2\frac{\xi}{\beta_0} + 3\frac{\xi^2}{\beta_0^2} \right] + \\
 & + \frac{\hbar^2}{4\mathcal{I}_0} [\mathcal{P}_+^2 + \mathcal{P}_-^2 + \mathcal{J}_+^2 - \mathcal{J}_-^2 - 2(\mathcal{P}_+ \mathcal{J}_+ - \mathcal{P}_- \mathcal{J}_-)] \left[2\sqrt{6}\frac{\xi\eta}{\beta_0^2} - \frac{2}{3}\sqrt{6}\frac{\eta}{\beta_0} \right] - \\
 & - M\omega^2 r^2 [\xi Y_{20} + \eta(Y_{22} + Y_{2-2})]
 \end{aligned} \quad (2.1)$$

One of the most important members in H is Coriolis interaction :

$$H_{\text{coriolis}} = \hbar^2 / 2\mathcal{I}_0 (\mathcal{P}_+ \mathcal{J}_- + \mathcal{P}_- \mathcal{J}_+ + 2\mathcal{P}_3 \mathcal{J}_3) = -(\hbar^2 / \mathcal{I}_0) (\mathcal{P} \cdot \mathcal{J}) \quad (2.2)$$

Matrix elements of this expression are easily calculated with the help of wavefunctions:

$$\begin{aligned}
 \langle IMK'\Omega' n n_f \alpha | H_{\text{coriolis}} | IMK\Omega n n_f \alpha \rangle = \\
 = \frac{\hbar^2}{2\mathcal{I}_0} \sum_j C_{\Omega'}^{(a)*} C_{\Omega}^{(a)} \left\{ \left[\delta_{K'K-j} \delta_{\Omega'\Omega-1} + (-1)^{I'+1/2} (-1)^{j-1/2} \delta_{K'-(K-1)} \delta_{\Omega'-(\Omega-1)} \right] [(I+K)(I-K+1) \times \right. \\
 \times (j+\Omega)(j-\Omega+1)]^{1/2} \} + \left[\delta_{K'K+j} \delta_{\Omega'\Omega+1} + (-1)^{I'+1/2} (-1)^{j-1/2} \delta_{K'-(K+1)} \delta_{\Omega'-(\Omega+1)} \right] \times \\
 \times [(I-K)(I+K+1)(j-\Omega)(j+\Omega+1)]^{1/2} \} - \frac{\hbar^2}{2\mathcal{I}_0} 2K\Omega \delta_{K'KB} \delta_{\Omega'\Omega}
 \end{aligned} \quad (2.3)$$

$$\mathcal{P}_\chi \mathcal{P}_\lambda - \mathcal{P}_\lambda \mathcal{P}_\chi = -i\mathcal{P}_{\chi \times \lambda} \quad (2.4 a)$$

$$J_x J'_\lambda - J'_\lambda J_x = J_{x \times \lambda} \quad (2.4 \text{ b})$$

$$J_x I'_\lambda - I'_\lambda J_x = 0 \quad (2.4 \text{ c})$$

The commutational proportion was used here for operators I_ν , J_ν , which lead to matrix elements:

$$\langle IMK | J_\pm | IMK \pm 1 \rangle = \sqrt{(I \mp K)(I \pm K + 1)} \quad (2.5)$$

$$\langle j\Omega | J_m | j\Omega \pm 1 \rangle = \sqrt{(j \mp \Omega)(j \pm \Omega + 1)} \quad (2.6)$$

The dependence on corner moment I in the example is found only in that part which is proportional to the member in brackets. It contributes to diagonal matrix elements with $K' = K$ and $\Omega' = \Omega$ only for band with $K = 1/2$. This matrix element is usually written like :

$$\begin{aligned} \langle IM \frac{1}{2} \frac{1}{2} n_2 n_0 \alpha | \hat{H}_{\text{coriolis}} | IM \frac{1}{2} \frac{1}{2} n_2 n_0 \alpha \rangle &= -\frac{\hbar^2}{2J_0} \sum_j C_{j\Omega}^{(\alpha)*} C_{j\Omega}^{(\alpha)} \left\{ (-1)^{(I+\frac{1}{2})} (-1)^{(j-\frac{1}{2})} (I + \frac{1}{2})(j + \frac{1}{2}) \right\} - \\ \frac{\hbar^2}{4J_0} &= \frac{\hbar^2}{2J_0} (-1)^{(I+\frac{1}{2})} (I + \frac{1}{2}) a - \frac{\hbar^2}{4J_0}, \end{aligned} \quad (2.7)$$

where

$$a = \sum_j (-1)^{j-\frac{1}{2}} (j + \frac{1}{2}) |C_{j1/2}^{(\alpha)}|^2 \quad (2.8)$$

The so called parameter of solution – a .

This name it got because of the correspondence to partitive break of connection of monopartitive and rotative movements.

Considering the Coriolis's member in the first range of perturbation theory is the following expression:

$$E_{IK\Omega n_2 \alpha} = \mathcal{E}_{\alpha\Omega} + (I(I+1) - (K-\Omega)^2) \frac{1}{2} \varepsilon + \left(\frac{1}{2} |K-\Omega| + 1 + 2n_2 \right) E_\gamma + \\ + \left(n_0 + \frac{1}{2} \right) E_\beta - a \left[(-1)^{I+\frac{1}{2}} \left(I + \frac{1}{2} \right) \right] \frac{1}{2} \varepsilon \delta_{K\frac{1}{2}} - 2K\Omega \frac{\varepsilon}{2} \quad (2.9)$$

The influence of Coriolis's forces on the bands with $K' = 1/2$ may be very strong. In the considerable meanings of parameter the order of levels in the rotative band may be changed.

For $a=0$ levels are situated according to the rule $I(I+1)$. In the considerable positive or negative meanings a order of levels is becoming quite different.

In general sense the hamiltonian $\mathcal{H}$ mixes 2 rotative bands for which $\Delta K = \pm 1$. It is obvious, that this effect will be small (maximum) if nonperturbed bands are greatly divided an energy. However, if the connection of the part with ostov is weak, these lands are overlapping greatly and the correction from the side of Coriolis's forces is important. In this case it is possible to get the analytical expression which describes the influence of Coriolis's powers on the own energy of too bands, by means of simple diagonalisation of matrix 2×2 the corresponding solution are equation:

$$\begin{vmatrix} E_{IK\Omega n_2 n_0 \alpha} - E \langle IK\Omega n_2 n_0 \alpha | \mathcal{H}_{coriolis} | IK + 1 \Omega + 1 n_2 n_0 \alpha \rangle \\ \langle IK + 1 \Omega + 1 n_2 n_0 \alpha | \mathcal{H}_{coriolis} | IK\Omega n_2 n_0 \alpha \rangle E_{IK+1\Omega+1 n_2 n_0 \alpha} - E \end{vmatrix} = 0 \quad (2.10)$$

where

$$E_{\pm}(IK\Omega n_2 n_{\theta} \alpha) = \frac{1}{2} \left\{ E_{IK\Omega n_2 n_{\theta} \alpha} + E_{IK+1\Omega+1n_2 n_{\theta} \alpha} \pm \right. \\ \left. \pm \Delta E \left[1 + 4 \left| \frac{IK+1\Omega+1n_2 n_{\theta} \alpha | \hat{H}_{corolis} | IK\Omega n_2 n_{\theta} \alpha}{\Delta E} \right|^2 \right]^{1/2} \right\} \quad (2.11)$$

the matrix elements being involved into this formulae are solved from the expression (1) and may be written in more simple form:

$$\langle IMK+1\Omega+1n_2 n_{\theta} \alpha | \hat{H}_{corolis} | IMK\Omega n_2 n_{\theta} \alpha \rangle = A_K [(I-K)(I+K+1)]^{1/2} \quad (2.12)$$

where

$$A_K = -\frac{1}{2} \varepsilon \sum_j C_{jK+1}^{(\alpha)*} C_{jK}^{(\alpha)} [(j-K)(j+K+1)]^{1/2} \quad (2.13)$$

It is apparent that K' is not a good quantitative number any more even if the ostov still possesses aximal symmetry. Only in case when the part is connected considerably with the ostov, number ΔB is becoming increased and K will be approximated by good quantitative number.

3. NILSON SHELL-MODELL AND PROBABILITY RATIO CALCULATING.

In considerably deformed nuclei one can differentiate monopartitative passages which are connected with the change of inner wave function X_n and collective passages which leave the inner monopartitative structure untouched. Within the collective passages

which do not touch φ_{coll} but only change the rotative state of the system D_α .

The differentiation of movement in nucleus on collective and inner correspond to prediction that the wave function being the solution of wave equation for the nucleus has the following form of expression:

$$\Psi = X \varphi_{\text{coll}} D_\alpha \quad (3.1)$$

where X – corresponds to inner movement of nucleons, which may be expressed using the notion of independent movement of parts in fixed non-special field. Ψ describes the rotation of nucleus concerning its equilibrium form, and D_α – presents the collective rotative movement of system as a whole [11].

For calculating $B(\lambda, I \rightarrow I')$ we are using following expression:

$$B(\lambda, I \rightarrow I') = \sum_{\mu M'} \left| \langle \Omega', I' K' M' | \mathfrak{M}''(\lambda, \mu) | \Omega, I K M \rangle \right|^2 \quad (3.2)$$

$$T(\lambda) = \frac{8\pi(\lambda+1)}{\lambda[(2\lambda+1)!!]^2} \frac{1}{h} \left(\frac{\omega}{C} \right)^{2\lambda+1} B(\lambda) \quad (3.3)$$

$$\mathfrak{M}''(\lambda, \mu) = \sum_v \mathcal{D}_{\mu\nu}^\lambda(\theta_i) \mathfrak{M}'(\lambda, \nu) \quad (3.4)$$

$$B(\lambda, I \rightarrow I') = |\langle I \lambda K K' - K | I \lambda I' K' \rangle \int \chi_{\Omega'}^{I+} \mathfrak{M}(\lambda, K' - K) \chi_{\Omega} d\tau + \langle I \lambda K - K' - K | I \lambda I' - K' \rangle \int [(-)^{I'-I'} \chi_{-\Omega'}]^* \times \mathfrak{M}(\lambda, -K' - K) \chi_{\Omega} d\tau|^2 \quad (3.5)$$

$$\int \mathcal{D}_{M'K'}^{I'+} \mathcal{D}_{\mu\lambda}^\lambda \mathcal{D}_{MK}^I d\Omega^3 = \frac{8\pi^2}{2I'+1} \langle I \lambda M \mu | I \lambda I' M' \rangle \langle I \lambda K \nu | I \lambda I' K' \rangle \quad (3.6)$$

$$\int Y_{I\Lambda}^+ Y_{\lambda\nu} Y_{I\Lambda} d\Omega^2 = \sqrt{\frac{(2I+1)(2\lambda+1)}{4\pi(2I+1)}} \langle I \lambda I \Lambda \nu | I \lambda I \Lambda' \rangle \langle I \lambda 0 0 | I \lambda I 0 \rangle \quad (3.7)$$

$$\begin{aligned} 1 \cdot (\nabla \mathbf{r}^\lambda \gamma_{\lambda\nu}) = & \sqrt{\frac{2\lambda+1}{2\lambda-1}} \left[\sqrt{\lambda^2 - \nu^2} l_z r^{\lambda-1} \gamma_{\lambda-\nu} + \frac{1}{2} \sqrt{(\lambda-\nu)(\lambda-\nu-1)} \times \right. \\ & \left. \times l_+ r^{\lambda-1} \gamma_{\lambda-\nu+1} - \frac{1}{2} \sqrt{(\lambda+\nu)(\lambda+\nu-1)} l_- r^{\lambda-1} \gamma_{\lambda-\nu-1} \right] \end{aligned} \quad (3.8)$$

$$l_+ = l_x + i l_y, \quad l_- = l_x - i l_y \quad (3.8 a)$$

$$\begin{aligned} B(E\lambda, I \rightarrow I') = & e^2 \left(1 + (-)^{\lambda} \frac{Z}{A^{\frac{1}{2}}} \right)^2 \left(\frac{\hbar}{M\omega_0} \right)^{\lambda} \frac{2\lambda+1}{4\pi} \times \\ & \times |\langle I\lambda K K' - K | I\lambda I' K' \rangle + b_{E\lambda} (-)^{I'+K'} \langle I\lambda K - K' - K | I\lambda I' - K' \rangle|^2 G_{E\lambda}^2 \end{aligned} \quad (3.9)$$

$$\begin{aligned} b_{E\lambda} = & \frac{(-)^{K'+1/2+I}}{G_{E\lambda}} \left\{ \sum_{I'} \langle N' I' | r^{\lambda} | N I \rangle \sqrt{\frac{2I+1}{2I'+1}} \langle I\lambda 00 | I\lambda I' 0 \rangle \times \right. \\ & \times \sum_{\Lambda' \Lambda \Sigma' \Sigma} \delta_{-\Sigma', \Sigma} a_{I\Lambda'} a_{I\Lambda} \langle I\lambda \Lambda - K' - K | I\lambda I' - \Lambda' \rangle, \end{aligned} \quad (3.9 a)$$

$$G_{E\lambda} = \sum_{I'} \langle N' I' | r^{\lambda} | N I \rangle \sqrt{\frac{2I+1}{2I'+1}} \langle I\lambda 00 | I\lambda I' 0 \rangle \times \sum_{\Lambda' \Lambda \Sigma' \Sigma} \delta_{-\Sigma', \Sigma} a_{I\Lambda'} a_{I\Lambda} \langle I\lambda \Lambda K' - K | I\lambda I' \Lambda' \rangle. \quad (3.9 b)$$

$$\begin{aligned} B(M\lambda, I \rightarrow I') = & \left(\frac{\hbar}{2Mc} \right) \left(\frac{\hbar}{M\omega_0} \right)^{\lambda-1} \frac{1}{4} \frac{2\lambda+1}{4\pi} |\langle I\lambda K K' - K | I\lambda I' K' \rangle + \\ & b_{M\lambda} (-)^{I'+K'} \langle I\lambda K - K' - K | I\lambda I' - K' \rangle|^2 G_{M\lambda}^2 \end{aligned} \quad (3.10)$$

$$\begin{aligned}
b_{M\lambda} = & \frac{(-)^{K'+1/2+I}}{G_{M\lambda}} \sum_{II'} \langle N'I | r^{\lambda-1} | N\lambda \rangle \langle \lambda - 100 | \lambda - 1I' 0 \rangle \sqrt{\frac{2I+1}{2I+3}} \times \\
& \times \sum_{\Lambda'\Lambda\Sigma\Sigma'} a_{\Lambda'} a_{\Lambda} \left\{ g_3 \left[A(q) \delta_{-\Sigma',E} (-)^{\Sigma-1/2} \langle \lambda - 1\Lambda q | \lambda - 1I' -\Lambda' \rangle + \right. \right. \\
& + B(q) \delta_{\Sigma',1/2} \delta_{\Sigma,1/2} \langle \lambda - 1\Lambda q + 1 | \lambda - 1I' -\Lambda' \rangle - \\
& - C(q) \delta_{\Sigma',-1/2} \delta_{\Sigma,-1/2} \langle \lambda - 1\Lambda q - 1 | \lambda - 1I' -\Lambda' \rangle + \\
& + \frac{2}{\lambda+1} g \delta_{-\Sigma',\Sigma} \left[A(q) (-2\Lambda') \langle \lambda - 1\Lambda q | \lambda - 1I' -\Lambda' \rangle + \right. \\
& + B(q) \sqrt{(I+\Lambda')(I-\Lambda'+1)} \langle \lambda - 1\Lambda q + 1 | \lambda - 1I' -\Lambda' + 1 \rangle - \\
& \left. \left. - C(q) \sqrt{(I-\Lambda')(I+\Lambda'+1)} \langle \lambda - 1\Lambda q - 1 | \lambda - 1I' -\Lambda' + 1 \rangle \right] \right\} \quad (3.10 a)
\end{aligned}$$

$$\begin{aligned}
G_{M\lambda} = & \sum_{II'} \langle N'I | r^{\lambda-1} | N\lambda \rangle \langle \lambda - 100 | \lambda - 1I' 0 \rangle \times \\
& \times \sqrt{\frac{2I+1}{2I+3}} \sum_{\Lambda'\Lambda\Sigma\Sigma'} a_{\Lambda'} a_{\Lambda} \left\{ g_3 \left[A(k) \delta_{\Sigma',E} (-)^{\Sigma-1/2} \langle \lambda - 1\Lambda k | \lambda - 1I' \Lambda' \rangle + \right. \right. \\
& + B(k) \delta_{\Sigma',1/2} \delta_{\Sigma,1/2} \langle \lambda - 1\Lambda k + 1 | \lambda - 1I' \Lambda' \rangle - \\
& - C(k) \delta_{\Sigma',-1/2} \delta_{\Sigma,-1/2} \langle \lambda - 1\Lambda k - 1 | \lambda - 1I' \Lambda' \rangle + \\
& + \frac{2}{\lambda+1} g \delta_{\Sigma',\Sigma} \left[A(k) 2\Lambda' \langle \lambda - 1\Lambda k | \lambda - 1I' \Lambda' \rangle + \right. \\
& + B(k) \sqrt{(I-\Lambda')(I+\Lambda'+1)} \langle \lambda - 1\Lambda k + 1 | \lambda - 1I' \Lambda' + 1 \rangle - \\
& \left. \left. - C(k) \sqrt{(I+\Lambda')(I-\Lambda'+1)} \langle \lambda - 1\Lambda k - 1 | \lambda - 1I' \Lambda' - 1 \rangle \right] \right\}, \quad (3.10 b)
\end{aligned}$$

$$A(\nu) = \sqrt{\lambda^2 - \nu^2}, \quad (3.11 a)$$

$$B(\nu) = \sqrt{(\lambda - \nu)(\lambda - \nu - 1)}, \quad (3.11 b)$$

$$C(\nu) = \sqrt{(\lambda + \nu)(\lambda + \nu - 1)}, \quad (3.11 c)$$

$$k = K' - K, \quad (3.11 d)$$

$$q = -K' - K. \quad (3.11 e)$$

$$G_{M1} = (g_{\Omega} - g_R) 2\Omega \quad (3.11 f)$$

$$b_0 = -\frac{(-)^I}{g_{\Omega} - g_R} \left\{ (g_s - g_R) \sum_I a_{10}^2 + 2(g_s - g_R) \sum_I \sqrt{I(I+1)} a_{10} a_{11} \right\} \quad (3.12)$$

$$B_0(M1) = \frac{(3)}{64\pi} \left(\frac{1h}{2Mc} \right)^2 \frac{2I' + 1}{I' + 1} C_0^2 \left| 1 + b_0(-)^{I'-1/2} \right|^2 \quad (3.13)$$

$$2b_0 G_0 = G_0 - 2(g_I - g_R)a + g_s - 2g_I + g_R \quad (3.14)$$

$$G_0 = 3\mu - a(g_I - g_R) + \frac{1}{2}g_s + g_I - 2g_R \quad (3.15)$$

and

$$b_0 = -\frac{1}{2G_0} \left[3\mu + a(g_I - g_R) + \frac{1}{2}g_s - g_I - g_R \right] \quad (3.16)$$

The matrix element:

$$\langle N' I | r^2 | N I \rangle = \left[\frac{\Gamma(n + \lambda) \Gamma(n + \lambda)}{\Gamma(n + t - \nu + \lambda) \Gamma(n + t - \nu' + \lambda)} \right]^{1/2} \nu'! \nu! \times \quad (3.17)$$

$$\sum_{\sigma} \frac{\Gamma(t + \sigma + \lambda)}{\sigma! (n - \sigma)! (n - \sigma)! (\sigma + \nu - n)! (\sigma + \nu' - n)!}$$

where

$$n = \frac{1}{2}(N - \lambda), \quad (3.17 a)$$

$$n' = \frac{1}{2}(N' - \lambda), \quad (3.17 b)$$

$$\nu = \frac{1}{2}(I - t + \lambda), \quad (3.17 c)$$

$$\nu' = \frac{1}{2}(I - t' + \lambda), \quad (3.17 d)$$

$$t = \frac{1}{2}(I + t' + \lambda + 1) \quad (3.17 e)$$

and where

$$n \geq \sigma \geq n - \nu \quad (3.18 a)$$

$$n' \geq \sigma \geq n' - \nu' \quad (3.18 b)$$

$$I + \lambda \geq t \geq I - \lambda, \quad (3.18 c)$$

$$N + \lambda \geq N' \geq N \quad (3.18 d)$$

4. THE RESULTS OF CALCULATION RATIO $\hat{A}(E\lambda)$.

For example, few ratio $B(E\lambda)$ and few $B(E\lambda)$ are presented:

exp.	theor.
$\frac{2+0(1276.7) \rightarrow 2+0(93.2)}{0+0(1199.4) \rightarrow 2+0(93.2)} > 2.44 [33]$	3.2
$\frac{2+0(1276.7) \rightarrow 2+0(93.2)}{2+0(1276.7) \rightarrow 0+0(0.0)} = 10.0(25) [33]$	12.5
$\frac{2+0(1276.7) \rightarrow 4+0(306.6)}{\quad\quad\quad} = 1.13[33]$	1.5
$2+0(1199.4) \rightarrow 2+0(93.2)$	
$\frac{4+0(1450.4) \rightarrow 4+0(306.6)}{4+0(450.4) \rightarrow 2+0(93.2)} > 42.8[33]$	45.0

 $B(E2)$ in Hf-178

$0+0(0) \rightarrow 2+0(1276.7) = 0.002(1)$	[21]	0.03
	0.024(2)	[28]
	0.018(7)	[16]
$0+0(0) \rightarrow 2+2(1276.7) = 0.100(8)$	[21]	0.05
	0.113(12)	[32]
	0.017(3)	[28]
	0.015(4)	[16]
$0 + 0(0) \rightarrow 2+0(1496.4) = 0.01$	[21]	0.02
	0.017(10)	[28]
	0.013(2)	[16]

 $X(E0/E2)$ in Hf-178

$0+0(1199.4) \rightarrow 0+0(0) = 0.02(6)$	[33]	0.18
--	------	------

0.155(11) [50]
0.160(9) [32]

$$2+0(1276.7) \rightarrow 2+0(93.2) = 0.026(4) \quad [33] \quad 1.4$$

$$0.13 < X < 0.23 \quad [50]$$

$$1.35(18) \quad [50]$$

$$1.56(15) \quad [32]$$

$$4+0(1450.4) \rightarrow 4+0(306.6) = 0.15(7) \quad [33]$$

0.19

References

1. Rivlin L.A. // Voprosi radioelektroniky. v.1. 1963. ¹ 6. p.42.
2. Vali W., Vali V. // Proc.IEEE.1963.v.182.p.1248.
3. Baldwin G.C., Neissel J.P., Tonks I. // Proc.IEEE. 1963. v.128. p.1247.
4. Helmer K.G., Reich C.W. Decay of an isomeric state in 178-Hf with
K > 16. Nucl.Phys. 1968.- v.A114.- p.649-662.
5. Helmer K.G., Reich C.W. Half-life of 178m2-Hf and its neutron capture
production.- Nucl.Phys.1973.- v.-A211. -p.1-6.
6. Collins C.B., Carrol J.J. Process in the pumping of a gamma-ray
laser//
First International Gamma_Ray Laser Workshop, GARALAS-95.
Technical Digest of Abstracts, 19-23 August, 1995, p.2.; Collins
C.B.,Carrol J.J. K mixing in 178-Hf // First International
Gamma_Ray Laser Workshop, GARALAS-95.Technical Digest of
Abstracts, 19-23 August, 1995,p.11;

7. Oganessian Yu.Ts., Karamian S.A. K mixing in nuclear reactions
// First International Gamma-Ray Laser Workshop, GARALAS-95.
Technical Digest of Abstracts, 19-23 August, 1995, p.3.
8. Coussemont R., Neyens G. Gamma-ray lasers with long-lived
isomers // First International Gamma-Ray Laser Workshop,
GARALAS-95. Technical Digest of Abstracts, 19-23 August, 1995,
p.2.
9. Constantinescu O. Production, chemical and isotopic separation
of the high-spin isomer $^{178m2}\text{Hf}$ // First International
Gamma-Ray Laser Workshop, GARALAS-95. Technical Digest of
Abstracts, 19-23 August, 1995, p.2.
10. Bohr A., Mottelson B.R. Structure of atomic nuclei // 1977,
v.1,2.
11. Nilsson S.G. Kgl.Danske Viden sk. Selsk., Mat.-Fys. Medd.,
1955., v.29, 116, p.1-68.
12. Hamilton J.H., Varnell L., Ronninger L.L. et al.
Bull.Amer.Phys.Soc.
1974., v.19., p.579.
13. Fossan D.B., Hersking B. Rotational-state lifetimes // Phys.Lett.
1962., v.2., p.155-157.
14. Fossan D.B., Hersking B. Half-lives of first excited $2+$ state
($150 < A < 190$) // Nucl.Phys. 1963. v.40., p.24-33.
15. Bodendstedt E., Korner H.J., Gerdau E. et al. Das
gyromagnetische
verhältnis des $2+$ - rotationsniveaus von ^{178}Hf und weitere
untersuchungen zum Zerfall von ^{178}W // Z.Phys. 1962., v.168.,
p.103-114.
16. Karlsson E., Matthiss E., Ogaza S. The gyromagnetic ratio of the
first rotational state in ^{178}Hf // Arkiv Fysik. 1962., v.22., p.257-
266.
17. Ronningen R.M., Hamilton J.H., Ramayya A.V. et al. Reduced
transition probabilities of vibrational states in $^{154-160}\text{Gd}$ and
 $^{176-180}\text{Hf}$ // Phys.Rev.1977., v.C15., p.1671-1678.
18. Hamilton J.H., Ramayya A.V., Ronningen R.M. et al. Isomeric
trapping
following Coulomb excitation of high-spin states in ^{178}Hf //
Phys.Lett. 1982., v.112B., p.327-330.

19. Alexander K.R., Brinnckmann H.F. Aktivierungsquerschnitte von ^{178m}Hf (4.3 sec.) und ^{179m}Hf (18.6 sec.) für thermischeneutronen // Nucl. Phys. 1962., v.32., p.482-489.
20. Button E.J., Smith T.M. Half-life measurements of some neutron induced isomers // Thesis. Naval Scholl. 1965., AD-620384.
21. Ronningen R.M., Hamilton J.H., Varnell L. et al. E2 and E4 reduced matrix elements of $^{154}, ^{156}, ^{158}, ^{160}\text{-Gd}$ and $^{176}, ^{178}, ^{180}\text{-Hf}$ // Phys.Rev. 1977., v.C16., p.2208-2217.
22. Varnell L., Hamilton., Robinson R.L. Collective states in $^{178}\text{-Hf}$ via Coulomb excitation // Phys.Rev. 1971., v.C3., p.1265-1268.
23. Walker P.M., Ward D., Hauser et al. g-Factors for $^{172}, ^{173}, ^{174}, ^{178}\text{-Hf}$ K-isomers // Nucl.Phys. 1980., v. A349., p.1-9.
24. Kostov L.K., Rotter H., Heiser C. et al. Absolute E1, K=1 transition probabilities between multi-quasiparticle high-spin states in $^{176}\text{-Hf}$ and $^{178}\text{-Hf}$ // Nucl.Phys. 1982. v. A376. p.451-462.
25. Khoo T.L., Lovhiden G. Structural changes in the yrast states of $^{178}\text{-Hf}$ // Phys.Lett. 1977., v. 67B., p.271-275.
26. Abou-Leila H. Determination des moments d'inertie et des deformations a partir des mesures des periodes courtes dans les noyaux pair.pair deforms // Ann.Phys.(Paris). 1967. v.2., p.181-195.
27. Namenson A., Jackson H.E., Smither R.K. Highe Energy g-spectra Resulting from Neutron Capture in Hafnium Isotope* // Phys.Rev., 1966., v.146, 13, p.844-851.
28. Stelson P.H., Girodzins L. // Nucl.Data. 1965., v.A1., p.26
29. Hamilton J.G. New works on programm // Izv.Akad.Nauk SSSR., ser,Fiz. 1976., v.740., p.18-37.
30. R.K.Smither Gamma-Ray Spectrum from Thermal-Neutron Capture in $^{177}\text{-Hf}$ and Associated Energy Levels in $^{178}\text{-Hf}$ // Phys.Rev. 1962. v.129., N4., p.1691-1699.
31. Namesson A.I., Bolotin H.H. Thermal-Neutron-Capture Gamma-Ray

Studies of the Excited State of Odd-A Hafnium Isotopes // Phys.Rev. v.158.,N4.,p.1206-1124.

32. Hamilton J.H., Varnell L., Ronninger L.L. et al. E2 and E4 moments in 176,178,180-Hf // Bull.Amer.Phys.Soc. 1974.,v.19., p.579.JH11.
33. Hamilton J.H.,Kumar K.,Varnell L. et al. E0/E2 transition strengths and interpretations of 0+, 2+ states in 178-Hf // Phys.Rev. 1974. v.C10. p.2540-2544.
34. Fogelberg B.,Baclin A. Levels in 178-Hf studied in the (n,g) reaction // Nucl.Phys. 1971., v.A171., p.353-383.
35. Haque A.M.I., Richter R., Gelberg A. et al. A study of the low-lying states in 178-Hf through the neutron capture reaction// Proc. Intern. Symp. Capture Gamma-ray Spectroscopy and Related Topics. 1984. Knoxville. Tenn. AIP Conf.Proc. 1985, N125,p.423-426.
36. Hamilton J.H.,Ramayya A.V., Little P.E. New band-mixing anomalies in 178-Hf // Phys.Rev.Lett. 1970. v.25., p.946-949.
37. Chou Y.W.,Howes R.H.,Swerdlow P.H. Mosbauer measurements of quadrupole-moment ratios of the first excited 2+ states of 176, 178, 180-Hf // Phys.Lett. 1970. v.32B. p.682-684.
38. Van Klinken J., Venema W.Z., Janssens R.V.E. et al. K-forbidden decays in 178-Hf; M4 decay of an yrast state // Nucl.Phys. 1980., v.A339. p.189-204.
39. Little P.E.,Hamilton J.H.,Ramayya A.V. Level structure of 178-Hf // Phys.Rev. 1972., v.C5., p.252-270.
40. Stelson P.H.,Girodzins L. // Nucl.Data. 1965., v.A1.,p.21.
41. Casten R.F., von Brentano P.,Haque A.M.I. Evidence for an underlying SU(3) structure near neutron number N=104 // Phys.Rev. 1985., v.C31., p.1991-1994.
42. Novoselsky A. Strongly deformed even-even nuclei in the IBA-2 model // Nucl.Phys. 1988., v.A483., p.282-294.
43. Bogdzely A.A., Boneva C.T., Vasiljeva O.V. et al. Studing cascade g-transition in 178-Hf nuclei// Izv. Akad. Nauk. SSSR., ser.Fiz., 1987., v.51, p.1882-1888.

44. Bonatson D. Parameter systematics of variable moment of inertia models in the NpN scheme // Phys.Lett. 1987., v.187B., p.1-6.