

ARMY RESEARCH LABORATORY



Elastic Scattering from a Spherical Body Eccentrically Located Within a Sphere: Theoretical Derivation

by Gordon Videen and Eric Sandvol

ARL-TR-1394

August 1997

19970821 030

DTIC QUALITY INSPECTED 3

Approved for public release; distribution unlimited.

The findings in this report are not to be construed as an official Department of the Army position unless so designated by other authorized documents.

Citation of manufacturer's or trade names does not constitute an official endorsement or approval of the use thereof.

Destroy this report when it is no longer needed. Do not return it to the originator.

Abstract

The scattered fields from a spherical body eccentrically located within a host sphere are found by a derivation that satisfies the boundary conditions at both interfaces. The source, which may be composed of any linear combination of S and P waves, is also arbitrarily located within the host sphere. The scattering system has applications in seismic scattering, since the scattering interaction between a scatterer and the Earth's surface is significant when the scatterer is located near the surface. This interaction can affect subsequent seismograms.

Contents

1	Introduction	1
2	Theory	2
2.1	Scattering Sphere	4
2.2	Host Sphere	7
2.3	Fields Interior to the Host Sphere (Earth)	9
2.4	Incident Field	11
3	Implications	13
A	Boundary Coefficients	15
B	Vector Translations	19
	References	23
	Distribution	26
	Report Documentation Page	28

Figures

- 1 Geometry of scattering system showing coordinate systems of host (x_1, y_1, z_1) , scatterer (x_2, y_2, z_2) , and source (x_3, y_3, z_3) . . . 2

1. Introduction

Scattering of elastic waves by spherical bodies is often studied in the context of seismic theory. The simplest system for which an analytical solution can readily be obtained for the entire scattered elastic wave field is that of a sphere in an infinite elastic medium (Ying and Truell, 1956; Einspruch, Witterholt, and Truell, 1960; Dubrovskiy and Morozhnik, 1989). More complicated techniques must be employed to take into consideration the effects of multiple scattering (Wu, 1985; Wu and Aki, 1985) and irregularities in the geometry of scatterers (Waterman, 1969, 1976; Varatharajulu and Pao, 1976). When an inhomogeneity is located near an interface, the interaction between the inhomogeneity and the interface is not insignificant compared to the incident field, and must be considered when the scattered fields are calculated.

In this report, we derive the equations describing the scattered field from a spherical scattering body eccentrically located within a host sphere. The interface, in this case, is the host sphere (free) surface, which can be made large compared with the size of the scatterer. Such a system is of interest in seismology, since it can be used to describe the scattered fields due to inhomogeneities located near the surface of the Earth (host sphere). In order to solve this problem, we expand the relevant wave fields in terms of vector spherical harmonics and use the interface boundary conditions at both interfaces. In order to solve both sets of boundary conditions simultaneously, we translate the spherical vector harmonics representing the individual fields between the coordinate systems of the host and scattering spheres. This work is similar to previous works in which eccentric sources are embedded within a spherical host (Thompson, 1973; Glenn *et al*, 1985; Rial and Moran, 1986; Zhao and Harkrider, 1992); however, in addition to an eccentric source, we also include an eccentrically located scatterer. The notation used in this report is similar to that of a previous paper on light scattering from a host sphere containing an eccentric inclusion (Videen *et al*, 1995).

2. Theory

The geometry of the scattering system is shown in figure 1. A host sphere of radius a and elastic moduli λ and μ is centered on the x_1, y_1, z_1 coordinate system. A scattering sphere of radius a' and elastic moduli λ' and μ' is centered on the x_2, y_2, z_2 coordinate system, located at $(x_1 = 0, y_1 = 0, z_1 = d)$. A source is centered on the x_3, y_3, z_3 coordinate system, located at $(r_1 = r_s, \theta_1 = \theta_s, \varphi_1 = 0)$ such that \hat{z}_3 is parallel to \hat{r}_s , and \hat{y}_1 is parallel to \hat{y}_3 . A time dependence of $\exp(-i\omega t)$ is implicit. Real seismic sources have a more complicated time dependence. For linear, homogeneous systems, we can solve the equations for a general time-dependent source using integral Fourier transformations:

$$f(x_j) = \int_{-\infty}^{\infty} F(k_j) \exp(-ix_j k_j) dk_j, \quad (1)$$

$$F(k_j) = \frac{1}{2\pi} \int_{-\infty}^{\infty} f(x_j) \exp(ix_j k_j) dx_j. \quad (2)$$

In this report we solve for the amplitudes of the scattered fields at specific frequencies $F(k_j)$, which we can transform to find the time-dependent amplitudes using equation (1).




Figure 1. Geometry of scattering system showing coordinate systems of host (x_1, y_1, z_1) , scatterer (x_2, y_2, z_2) , and source (x_3, y_3, z_3) .

We reach the general solution by satisfying the boundary conditions on both spherical interfaces simultaneously. In doing so, we consider the fields incident on the interfaces of the two subsystems (host and scattering spheres) separately. We use the vector spherical harmonics (Hansen vectors), which have the following form in this derivation:

$$\begin{aligned} \mathbf{L}_{nm,j}^{(\rho)} = & \hat{r}_j \left[z_n^{(\rho)'}(kr_j) \tilde{P}_n^m(\cos \theta_j) e^{im\varphi_j} \right] + \\ & \hat{\theta}_j \left[\frac{1}{kr_j} z_n^{(\rho)}(kr_j) \frac{d}{d\theta_j} \tilde{P}_n^m(\cos \theta_j) e^{im\varphi_j} \right] + \\ & \hat{\varphi}_j \left[\frac{1}{kr_j} z_n^{(\rho)}(kr_j) \frac{im}{\sin \theta_j} \tilde{P}_n^m(\cos \theta_j) e^{im\varphi_j} \right], \end{aligned} \quad (3)$$

$$\begin{aligned} \mathbf{M}_{nm,j}^{(\rho)} = & \hat{\theta}_j \left[\frac{im}{\sin \theta_j} z_n^{(\rho)}(kr_j) \tilde{P}_n^m(\cos \theta_j) e^{im\varphi_j} \right] - \\ & \hat{\varphi}_j \left[z_n^{(\rho)}(kr_j) \frac{d}{d\theta_j} \tilde{P}_n^m(\cos \theta_j) e^{im\varphi_j} \right], \end{aligned} \quad (4)$$

$$\begin{aligned} \mathbf{N}_{nm,j}^{(\rho)} = & \hat{r}_j \left[\frac{1}{kr_j} z_n^{(\rho)}(kr_j) n(n+1) \tilde{P}_n^m(\cos \theta_j) e^{im\varphi_j} \right] + \\ & \hat{\theta}_j \left[\frac{1}{kr_j} \frac{d}{dr_j} (r_j z_n^{(\rho)}(kr_j)) \frac{d}{d\theta_j} \tilde{P}_n^m(\cos \theta_j) e^{im\varphi_j} \right] + \\ & \hat{\varphi}_j \left[\frac{1}{kr_j} \frac{d}{dr_j} (r_j z_n^{(\rho)}(kr_j)) \frac{im}{\sin \theta_j} \tilde{P}_n^m(\cos \theta_j) e^{im\varphi_j} \right], \end{aligned} \quad (5)$$

where the prime denotes a derivative with respect to the argument, the index j corresponds to the coordinate system used ($j = 1, 2$), and $z_n^{(\rho)}(kr_j)$ are the spherical Bessel functions of the first, second, third, or fourth kind ($\rho = 1, 2, 3, 4$) (Ben-Menahem and Singh, 1981), and

$$\tilde{P}_n^m(\cos \theta_j) = \sqrt{\frac{(2n+1)(n-m)!}{2(n+m)!}} P_n^m(\cos \theta_j), \quad (6)$$

where $P_n^m(\cos \theta_j)$ are the associated Legendre polynomials.

It is convenient to consider the vector surface harmonics, \mathbf{P}_{nm} , \mathbf{B}_{nm} , and \mathbf{C}_{nm} when solving the boundary conditions. The vector surface harmonics

are defined in terms of the vector spherical harmonics:

$$\mathbf{L}_{nm,j}^{(\rho)} = \mathbf{P}_{nm,j} z_n^{(\rho)'}(kr_j) + \mathbf{B}_{nm,j} \frac{1}{kr_j} z_n^{(\rho)}(kr_j), \quad (7)$$

$$\mathbf{M}_{nm,j}^{(\rho)} = \mathbf{C}_{nm,j} z_n^{(\rho)}(kr_j), \quad (8)$$

$$\mathbf{N}_{nm,j}^{(\rho)} = \mathbf{P}_{nm,j} \frac{1}{kr_j} z_n^{(\rho)}(kr_j) n(n+1) + \mathbf{B}_{nm,j} \left[\frac{1}{kr_j} \frac{d}{dr_j} (r_j z_n^{(\rho)}(kr_j)) \right]. \quad (9)$$

We start the derivation by expanding the fields in terms of the vector spherical harmonics and satisfying the boundary conditions on the interface of the scattering sphere.

2.1 Scattering Sphere

First we examine the fields that strike the outer surface of the scattering sphere. We consider an arbitrary field incident on the system that can be expanded with the spherical Bessel functions of the first kind, $j_n(kr_2)$:

$$\begin{aligned} \mathbf{u}_{sou}^2 &= \sum_{n=0}^{\infty} \sum_{m=-n}^n a_{nm}^{(1)} \mathbf{L}_{nm,2}^{(1)} + a_{nm}^{(2)} \mathbf{M}_{nm,2}^{(1)} + a_{nm}^{(3)} \mathbf{N}_{nm,2}^{(1)} \quad (10) \\ &= \sum_{n=0}^{\infty} \sum_{m=-n}^n \mathbf{P}_{nm,2} \left[a_{nm}^{(1)} z_n^{(1)'}(k_\alpha r_2) + a_{nm}^{(3)} n(n+1) \frac{z_n^{(1)}(k_\beta r_2)}{k_\beta r_2} \right] \\ &\quad + \mathbf{B}_{nm,2} \left(a_{nm}^{(1)} \frac{z_n^{(1)}(k_\alpha r_2)}{k_\alpha r_2} + a_{nm}^{(3)} \left[\frac{z_n^{(1)}(k_\beta r_2)}{k_\beta r_2} + z_n^{(1)'}(k_\beta r_2) \right] \right) \\ &\quad + \mathbf{C}_{nm,2} a_{nm}^{(2)} z_n^{(1)}(k_\beta r_2) \\ &= \sum_{n=0}^{\infty} \sum_{m=-n}^n A_{nm}^{(1)}(k_\alpha r_2, k_\beta r_2) \mathbf{P}_{nm,2} + A_{nm}^{(2)}(k_\alpha r_2, k_\beta r_2) \mathbf{B}_{nm,2} + A_{nm}^{(3)}(k_\beta r_2) \mathbf{C}_{nm,2}, \end{aligned}$$

where k_α and k_β are the compression and shear wavenumbers external to the scattering sphere, respectively. Similarly, k'_α and k'_β are the compression and shear wavenumbers internal to the scattering sphere, respectively. The coefficients $a_{nm}^{(1)}$, $a_{nm}^{(2)}$, and $a_{nm}^{(3)}$ represent the field from the source. Similarly, the scattered field from the scattering sphere may be expanded with the spherical Bessel functions of the third kind, $h_n^{(1)}(kr_2)$:

$$\begin{aligned}
\mathbf{u}_{sca}^2 &= \sum_{n=0}^{\infty} \sum_{m=-n}^n b_{nm}^{(1)} \mathbf{L}_{nm,2}^{(3)} + b_{nm}^{(2)} \mathbf{M}_{nm,2}^{(3)} + b_{nm}^{(3)} \mathbf{N}_{nm,2}^{(3)} \\
&= \sum_{n=0}^{\infty} \sum_{m=-n}^n B_{nm}^{(1)}(k_{\alpha}r_2, k_{\beta}r_2) \mathbf{P}_{nm,2} + B_{nm}^{(2)}(k_{\alpha}r_2, k_{\beta}r_2) \mathbf{B}_{nm,2} + B_{nm}^{(3)}(k_{\beta}r_2) \mathbf{C}_{nm,2} .
\end{aligned} \tag{11}$$

This scattered field interacts with the host sphere (free) surface, which reflects incoming spherical waves onto the scattering sphere. These may be expanded with the spherical Bessel functions of the fourth kind, $h_n^{(2)}(kr_2)$:

$$\begin{aligned}
\mathbf{u}_{ref}^2 &= \sum_{n=0}^{\infty} \sum_{m=-n}^n c_{nm}^{(1)} \mathbf{L}_{nm,2}^{(4)} + c_{nm}^{(2)} \mathbf{M}_{nm,2}^{(4)} + c_{nm}^{(3)} \mathbf{N}_{nm,2}^{(4)} \\
&= \sum_{n=0}^{\infty} \sum_{m=-n}^n C_{nm}^{(1)}(k_{\alpha}r_2, k_{\beta}r_2) \mathbf{P}_{nm,2} + C_{nm}^{(2)}(k_{\alpha}r_2, k_{\beta}r_2) \mathbf{B}_{nm,2} + C_{nm}^{(3)}(k_{\beta}r_2) \mathbf{C}_{nm,2} .
\end{aligned} \tag{12}$$

The fields internal to the scattering sphere may be expanded into standing waves with spherical Bessel functions of the first kind:

$$\begin{aligned}
\mathbf{u}_{int}^2 &= \sum_{n=0}^{\infty} \sum_{m=-n}^n d_{nm}^{(1)} \mathbf{L}_{nm,2}^{(1)} + d_{nm}^{(2)} \mathbf{M}_{nm,2}^{(1)} + d_{nm}^{(3)} \mathbf{N}_{nm,2}^{(1)} \\
&= \sum_{n=0}^{\infty} \sum_{m=-n}^n D_{nm}^{(1)}(k'_{\alpha}r_2, k'_{\beta}r_2) \mathbf{P}_{nm,2} + D_{nm}^{(2)}(k'_{\alpha}r_2, k'_{\beta}r_2) \mathbf{B}_{nm,2} + D_{nm}^{(3)}(k'_{\beta}r_2) \mathbf{C}_{nm,2} .
\end{aligned} \tag{13}$$

With the expansions of the fields defined, we can now satisfy the boundary conditions at the interface of the scattering sphere. The stress vector across a spherical surface for a displacement field given by

$$\mathbf{u} = U(r)\mathbf{P}_{nm} + V(r)\mathbf{B}_{nm} + W(r)\mathbf{C}_{nm} \tag{14}$$

can be calculated as (Ben-Menahem and Singh, 1981)

$$\begin{aligned}
\mathbf{T}(\hat{r}) = & \hat{r} \left[(\lambda + 2\mu) \frac{dU}{dr} + 2\lambda \frac{U}{r} - n(n+1)\lambda \frac{V}{r} \right] \tilde{P}_n^m(\cos\theta) e^{im\varphi} \\
& + \hat{\theta} \mu \left[\left(\frac{dV}{dr} + \frac{U-V}{r} \right) \frac{\partial}{\partial\theta} + \left(\frac{dW}{dr} - \frac{W}{r} \right) \frac{1}{\sin\theta} \frac{\partial}{\partial\varphi} \right] \tilde{P}_n^m(\cos\theta) e^{im\varphi} \\
& + \hat{\varphi} \mu \left[\left(\frac{dV}{dr} + \frac{U-V}{r} \right) \frac{1}{\sin\theta} \frac{\partial}{\partial\theta} - \left(\frac{dW}{dr} - \frac{W}{r} \right) \frac{\partial}{\partial\varphi} \right] \tilde{P}_n^m(\cos\theta) e^{im\varphi}.
\end{aligned} \tag{15}$$

In order for the displacement field and the stress tensor to be continuous, the following six boundary conditions must be satisfied at $r_2 = a'$:

$$A_{nm}^{(1)}(k_\alpha r_2, k_\beta r_2) + B_{nm}^{(1)}(k_\alpha r_2, k_\beta r_2) + C_{nm}^{(1)}(k_\alpha r_2, k_\beta r_2) = D_{nm}^{(1)}(k'_\alpha r_2, k'_\beta r_2), \tag{16}$$

$$A_{nm}^{(2)}(k_\alpha r_2, k_\beta r_2) + B_{nm}^{(2)}(k_\alpha r_2, k_\beta r_2) + C_{nm}^{(2)}(k_\alpha r_2, k_\beta r_2) = D_{nm}^{(2)}(k'_\alpha r_2, k'_\beta r_2), \tag{17}$$

$$A_{nm}^{(3)}(k_\beta r_2) + B_{nm}^{(3)}(k_\beta r_2) + C_{nm}^{(3)}(k_\beta r_2) = D_{nm}^{(3)}(k'_\beta r_2), \tag{18}$$

$$\begin{aligned}
& \left[(\lambda + 2\mu) \frac{d}{dr_2} + \frac{2\lambda}{a'} \right] \left[A_{nm}^{(1)}(k_\alpha r_2, k_\beta r_2) + B_{nm}^{(1)}(k_\alpha r_2, k_\beta r_2) + C_{nm}^{(1)}(k_\alpha r_2, k_\beta r_2) \right] - \\
& n(n+1) \frac{\lambda}{a'} \left[A_{nm}^{(2)}(k_\alpha r_2, k_\beta r_2) + B_{nm}^{(2)}(k_\alpha r_2, k_\beta r_2) + C_{nm}^{(2)}(k_\alpha r_2, k_\beta r_2) \right] = \\
& \left[(\lambda' + 2\mu') \frac{d}{dr_2} + \frac{2\lambda'}{a'} \right] D_{nm}^{(1)}(k'_\alpha r_2, k'_\beta r_2) - n(n+1) \frac{\lambda'}{a'} D_{nm}^{(2)}(k'_\alpha r_2, k'_\beta r_2),
\end{aligned} \tag{19}$$

$$\begin{aligned}
& a' \mu \frac{d}{dr_2} \left[A_{nm}^{(2)}(k_\alpha r_2, k_\beta r_2) + B_{nm}^{(2)}(k_\alpha r_2, k_\beta r_2) + C_{nm}^{(2)}(k_\alpha r_2, k_\beta r_2) \right] + \\
& \mu \left[A_{nm}^{(1)}(k_\alpha r_2, k_\beta r_2) + B_{nm}^{(1)}(k_\alpha r_2, k_\beta r_2) + C_{nm}^{(1)}(k_\alpha r_2, k_\beta r_2) \right] - \\
& \mu \left[A_{nm}^{(2)}(k_\alpha r_2, k_\beta r_2) + B_{nm}^{(2)}(k_\alpha r_2, k_\beta r_2) + C_{nm}^{(2)}(k_\alpha r_2, k_\beta r_2) \right] \\
& = \mu' \left\{ a' \frac{d}{dr_2} D_{nm}^{(2)}(k'_\alpha r_2, k'_\beta r_2) + \left[D_{nm}^{(1)}(k'_\alpha r_2, k'_\beta r_2) - D_{nm}^{(2)}(k'_\alpha r_2, k'_\beta r_2) \right] \right\},
\end{aligned} \tag{20}$$

$$\mu \left[a' \frac{d}{dr_2} - 1 \right] \left[A_{nm}^{(3)}(k_\beta r_2) + B_{nm}^{(3)}(k_\beta r_2) + C_{nm}^{(3)}(k_\beta r_2) \right] = \mu' \left[a' \frac{d}{dr_2} - 1 \right] D_{nm}^{(3)}(k'_\beta r_2). \quad (21)$$

We are interested primarily in the external fields of the scattering sphere. By substituting equations (16) to (18) into equations (20) to (21), we can eliminate the internal field coefficients and express the scattered external field coefficients in terms of the incoming field coefficients. Then, by expanding the vector surface harmonics in terms of the vector spherical harmonics (given in eq (10)), we can derive the following expressions for the vector spherical harmonic coefficients:

$$b_{nm}^{(1)} = a_{nm}^{(1)} Q_{nm}^{11} + a_{nm}^{(3)} Q_{nm}^{13} + c_{nm}^{(1)} R_{nm}^{11} + c_{nm}^{(3)} R_{nm}^{13}, \quad (22)$$

$$b_{nm}^{(2)} = a_{nm}^{(2)} Q_{nm}^{22} + c_{nm}^{(2)} R_{nm}^{22}, \quad (23)$$

and

$$b_{nm}^{(3)} = a_{nm}^{(1)} Q_{nm}^{31} + a_{nm}^{(3)} Q_{nm}^{33} + c_{nm}^{(1)} R_{nm}^{31} + c_{nm}^{(3)} R_{nm}^{33}. \quad (24)$$

Explicit expressions for Q_{nm}^{11} , Q_{nm}^{13} , R_{nm}^{11} , R_{nm}^{13} , Q_{nm}^{22} , R_{nm}^{22} , Q_{nm}^{31} , Q_{nm}^{33} , R_{nm}^{31} , and R_{nm}^{33} are given in appendix A.

2.2 Host Sphere

We next examine the fields in the host (Earth-centered) coordinate system. First we consider a source that can be expanded with spherical Bessel functions of the third kind, $h_n^{(1)}(kr_1)$:

$$\begin{aligned} \mathbf{u}_{sou}^1 &= \sum_{n=0}^{\infty} \sum_{m=-n}^n e_{nm}^{(1)} \mathbf{L}_{nm,1}^{(3)} + e_{nm}^{(2)} \mathbf{M}_{nm,1}^{(3)} + e_{nm}^{(3)} \mathbf{N}_{nm,1}^{(3)} \\ &= \sum_{n=0}^{\infty} \sum_{m=-n}^n E_{nm}^{(1)}(r_1) \mathbf{P}_{nm,1} + E_{nm}^{(2)}(r_1) \mathbf{B}_{nm,1} + E_{nm}^{(3)}(r_1) \mathbf{C}_{nm,1}. \end{aligned} \quad (25)$$

The scattered and incident fields from the scattering sphere may be expanded with the spherical Bessel functions of the third kind, $h_n^{(1)}(kr_1)$,

$$\begin{aligned} \mathbf{u}_{sca}^1 &= \sum_{n=0}^{\infty} \sum_{m=-n}^n f_{nm}^{(1)} \mathbf{L}_{nm,1}^{(3)} + f_{nm}^{(2)} \mathbf{M}_{nm,1}^{(3)} + f_{nm}^{(3)} \mathbf{N}_{nm,1}^{(3)} \\ &= \sum_{n=0}^{\infty} \sum_{m=-n}^n F_{nm}^{(1)}(r_1) \mathbf{P}_{nm,1} + F_{nm}^{(2)}(r_1) \mathbf{B}_{nm,1} + F_{nm}^{(3)}(r_1) \mathbf{C}_{nm,1}, \end{aligned} \quad (26)$$

and the spherical Bessel functions of the fourth kind, $h_n^{(2)}(kr_1)$,

$$\begin{aligned} \mathbf{u}_{ref}^1 &= \sum_{n=0}^{\infty} \sum_{m=-n}^n g_{nm}^{(1)} \mathbf{L}_{nm,1}^{(4)} + g_{nm}^{(2)} \mathbf{M}_{nm,1}^{(4)} + g_{nm}^{(3)} \mathbf{N}_{nm,1}^{(4)} \\ &= \sum_{n=0}^{\infty} \sum_{m=-n}^n G_{nm}^{(1)}(r_1) \mathbf{P}_{nm,1} + G_{nm}^{(2)}(r_1) \mathbf{B}_{nm,1} + G_{nm}^{(3)}(r_1) \mathbf{C}_{nm,1} . \end{aligned} \quad (27)$$

In appendix B, we provide expansions whereby relationships can be derived between the $a_{nm}^{(p)}$ and $e_{nm}^{(p)}$, $b_{nm}^{(p)}$ and $f_{nm}^{(p)}$, and $c_{nm}^{(p)}$ and $g_{nm}^{(p)}$ coefficients. The expansions (26) and (27) are valid in the region where $r_1 > d$: i.e., near the host sphere (free) surface. Hence, we need not be concerned with the pole that occurs at $r_1 = 0$ for the scattered and incident fields, since the expansion is not valid in this region. Since the stress vector across a free surface must be zero, the following boundary conditions apply at $r_1 = a$:

$$\begin{aligned} \left[(\lambda + 2\mu) \frac{d}{dr_1} + \frac{2\lambda}{a} \right] \left(E_{nm}^{(1)}(r_1) + F_{nm}^{(1)}(r_1) + G_{nm}^{(1)}(r_1) \right) = \\ n(n+1) \frac{\lambda}{a} \left(E_{nm}^{(2)}(r_1) + F_{nm}^{(2)}(r_1) + G_{nm}^{(2)}(r_1) \right), \end{aligned} \quad (28)$$

$$a \frac{d}{dr_1} \left(E_{nm}^{(2)}(r_1) + F_{nm}^{(2)}(r_1) + G_{nm}^{(2)}(r_1) \right) + \quad (29)$$

$$\left(E_{nm}^{(1)}(r_1) + F_{nm}^{(1)}(r_1) + G_{nm}^{(1)}(r_1) \right) - \left(E_{nm}^{(2)}(r_1) + F_{nm}^{(2)}(r_1) + G_{nm}^{(2)}(r_1) \right) = 0 ,$$

$$a \frac{d}{dr_1} \left(E_{nm}^{(3)}(r_1) + F_{nm}^{(3)}(r_1) + G_{nm}^{(3)}(r_1) \right) - E_{nm}^{(3)}(r_1) + F_{nm}^{(3)}(r_1) + G_{nm}^{(3)}(r_1) = 0 . \quad (30)$$

As with the scattering-sphere coefficients, we express the coefficients in the host-sphere coordinate system in the following form:

$$g_{nm}^{(1)} = \left(e_{nm}^{(1)} + f_{nm}^{(1)} \right) S_{nm}^{11} + \left(e_{nm}^{(3)} + f_{nm}^{(3)} \right) S_{nm}^{13} , \quad (31)$$

$$g_{nm}^{(2)} = \left(e_{nm}^{(2)} + f_{nm}^{(2)} \right) S_{nm}^{22} , \quad (32)$$

and

$$g_{nm}^{(3)} = \left(e_{nm}^{(1)} + f_{nm}^{(1)} \right) S_{nm}^{31} + \left(e_{nm}^{(3)} + f_{nm}^{(3)} \right) S_{nm}^{33} . \quad (33)$$

In appendix A, we give explicit expressions for S_{nm}^{11} , S_{nm}^{13} , S_{nm}^{22} , S_{nm}^{31} , and S_{nm}^{33} .

2.3 Fields Interior to the Host Sphere (Earth)

The displacement fields given by equations (10) to (12) are equivalent to the displacement fields given by equations (25) to (27). By performing a translation of the spherical vector harmonics from the host coordinate system to the scattering-sphere coordinate system (using the results of app B), we can put equations (31) to (33) into the following form:

$$e_{nm}^{(1)} S_{nm}^{11} + e_{nm}^{(3)} S_{nm}^{13} = \sum_{n'=0}^{\infty} c_{n'm}^{(1)} C_{n',n}^{(m,1)} - b_{n'm}^{(1)} S_{nm}^{11} C_{n',n}^{(m,1)} \quad (34)$$

$$- b_{n'm}^{(3)} S_{nm}^{13} A_{n',n}^{(m,1)} - b_{n'm}^{(2)} S_{nm}^{13} B_{n',n}^{(m,1)},$$

$$S_{nm}^{22} e_{nm}^{(2)} = \sum_{n'=0}^{\infty} c_{n'm}^{(2)} A_{n',n}^{(m,1)} + c_{n'm}^{(3)} B_{n',n}^{(m,1)} - b_{n'm}^{(2)} S_{nm}^{22} A_{n',n}^{(m,1)} - b_{n'm}^{(3)} S_{nm}^{22} B_{n',n}^{(m,1)}, \quad (35)$$

and

$$e_{nm}^{(1)} S_{nm}^{31} + e_{nm}^{(3)} S_{nm}^{33} = \sum_{n'=0}^{\infty} c_{n'm}^{(3)} A_{n',n}^{(m,1)} + c_{n'm}^{(2)} B_{n',n}^{(m,1)} - b_{n'm}^{(1)} S_{nm}^{31} C_{n',n}^{(m,1)} \quad (36)$$

$$- b_{n'm}^{(3)} S_{nm}^{33} A_{n',n}^{(m,1)} - b_{n'm}^{(2)} S_{nm}^{33} B_{n',n}^{(m,1)}.$$

Substituting equations (22) to (24) into these expressions yields

$$e_{nm}^{(1)} S_{nm}^{11} + e_{nm}^{(3)} S_{nm}^{13} = - \sum_{n'=0}^{\infty} a_{n'm}^{(1)} \left(Q_{n'm}^{11} S_{nm}^{11} C_{n',n}^{(m,1)} + Q_{n'm}^{31} S_{nm}^{13} A_{n',n}^{(m,1)} \right) + \quad (37)$$

$$a_{n'm}^{(2)} Q_{n'm}^{22} S_{nm}^{13} B_{n',n}^{(m,1)} +$$

$$a_{n'm}^{(3)} \left(Q_{n'm}^{13} S_{nm}^{11} C_{n',n}^{(m,1)} + Q_{n'm}^{33} S_{nm}^{13} A_{n',n}^{(m,1)} \right) +$$

$$c_{n'm}^{(1)} \left(R_{n'm}^{11} S_{nm}^{11} C_{n',n}^{(m,1)} + R_{n'm}^{31} S_{nm}^{13} A_{n',n}^{(m,1)} - C_{n',n}^{(m,1)} \right) +$$

$$c_{n'm}^{(2)} R_{n'm}^{22} S_{nm}^{13} B_{n',n}^{(m,1)} +$$

$$c_{n'm}^{(3)} \left(R_{n'm}^{13} S_{nm}^{11} C_{n',n}^{(m,1)} + R_{n'm}^{33} S_{nm}^{13} A_{n',n}^{(m,1)} \right),$$

$$\begin{aligned}
e_{nm}^{(2)} S_{nm}^{22} = & - \sum_{n'=0}^{\infty} a_{n'm}^{(1)} \left(Q_{n'm}^{31} S_{nm}^{22} B_{n',n}^{(m,1)} \right) + c_{n'm}^{(1)} \left(R_{n'm}^{31} S_{nm}^{22} B_{n',n}^{(m,1)} \right) + \\
& a_{n'm}^{(2)} \left(Q_{n'm}^{22} S_{nm}^{22} A_{n',n}^{(m,1)} \right) + c_{n'm}^{(2)} \left(R_{n'm}^{22} S_{nm}^{22} A_{n',n}^{(m,1)} - A_{n',n}^{(m,1)} \right) + \\
& a_{n'm}^{(3)} \left(Q_{n'm}^{33} S_{nm}^{22} B_{n',n}^{(m,1)} \right) + c_{n'm}^{(3)} \left(R_{n'm}^{33} S_{nm}^{22} B_{n',n}^{(m,1)} - B_{n',n}^{(m,1)} \right),
\end{aligned} \tag{38}$$

and

$$\begin{aligned}
e_{nm}^{(1)} S_{nm}^{31} + e_{nm}^{(3)} S_{nm}^{33} = & - \sum_{n'=0}^{\infty} a_{n'm}^{(1)} \left(Q_{n'm}^{11} S_{nm}^{31} C_{n',n}^{(m,1)} + Q_{n'm}^{31} S_{nm}^{33} A_{n',n}^{(m,1)} \right) + \\
& a_{n'm}^{(2)} \left(Q_{n'm}^{22} S_{nm}^{33} B_{n',n}^{(m,1)} \right) + \\
& a_{n'm}^{(3)} \left(Q_{n'm}^{13} S_{nm}^{31} C_{n',n}^{(m,1)} + Q_{n'm}^{33} S_{nm}^{33} A_{n',n}^{(m,1)} \right) + \\
& c_{n'm}^{(1)} \left(R_{n'm}^{11} S_{nm}^{31} C_{n',n}^{(m,1)} + R_{n'm}^{31} S_{nm}^{33} A_{n',n}^{(m,1)} \right) + \\
& c_{n'm}^{(2)} \left(R_{n'm}^{22} S_{nm}^{33} B_{n',n}^{(m,1)} - B_{n',n}^{(m,1)} \right) + \\
& c_{n'm}^{(3)} \left(R_{n'm}^{13} S_{nm}^{31} C_{n',n}^{(m,1)} + R_{n'm}^{33} S_{nm}^{33} A_{n',n}^{(m,1)} - A_{n',n}^{(m,1)} \right).
\end{aligned} \tag{39}$$

The $a_{nm}^{(p)}$ and $e_{nm}^{(p)}$ coefficients are related by equations (B-14) to (B-16) (app B). Since the $e_{nm}^{(p)}$ coefficients are the coefficients for the incident field, which is assumed to be known, the $a_{nm}^{(p)}$ can be easily determined, and equations (37) to (39) represent three sets of equations containing three sets of unknowns $c_{nm}^{(p)}$. Numerical results based on this analytic solution can be acquired following the same algorithms used in light scattering (Videen *et al*, 1995). Equations relating the various fields are given by equations (22) to (24) and (31) to (33); the coefficients used in these equations can be calculated from the equations given in appendix A. Computational results obtained from similar expressions in the field of light scattering have shown that the coefficients of the scattering harmonic terms tend to have a negligible contribution to the scattered fields for $n > ka + 4(ka)^{1/3} + 2$ (Wiscombe, 1980; Bohren and Huffman, 1983). It is likely that a similar truncation criterion holds for the elastic scatter. With the truncation of the summations given by equations (37) to (39), a solution can be found through matrix inversion. With the $a_{nm}^{(p)}$ and $c_{nm}^{(p)}$ known, $b_{nm}^{(p)}$ can be determined by equations (22) to (24). Hence, the entire displacement field within the host sphere can be found.

2.4 Incident Field

Although the solution derived in section 2.3 is completely general, in many applications we are interested in a specific case where a point source is located at some position within the host sphere ($r_1 = r_s$, $\theta_1 = \theta_s$, $\varphi_1 = 0$). We specify a source coordinate system (x_3, y_3, z_3) such that \hat{z}_3 is parallel to \hat{r}_s and \hat{y}_1 is parallel to \hat{y}_3 , as shown in figure 1. The derivation is valid only when the source is closer to the center of the host sphere than is the scattering-sphere center ($r_s < d$). The limitation of the validity is a result of the expansion that we use in translating the vector spherical harmonics. If the source is farther from the center of the host sphere than the scattering sphere ($r_s > d$), then we would need to use a different set of translation coefficients, and the derivation would be slightly different (Stratton, 1941). The source can be expanded as

$$\mathbf{u}_{\text{sou}}^3 = \sum_{n=0}^{\infty} \sum_{m=-n}^n h_{nm}^{(1)} \mathbf{L}_{nm,3}^{(3)} + h_{nm}^{(2)} \mathbf{M}_{nm,3}^{(3)} + h_{nm}^{(3)} \mathbf{N}_{nm,3}^{(3)}. \quad (40)$$

This field can be expressed in a coordinate system whose origin is the same as the host (Earth) coordinate system, but whose axes are parallel to the source coordinate system by a translation of a distance r_s in the $-\hat{z}_3$ direction. The source in this (x_4, y_4, z_4) is

$$\mathbf{u}_{\text{sou}}^4 = \sum_{n=0}^{\infty} \sum_{m=-n}^n i_{nm}^{(1)} \mathbf{L}_{nm,4}^{(3)} + i_{nm}^{(2)} \mathbf{M}_{nm,4}^{(3)} + i_{nm}^{(3)} \mathbf{N}_{nm,4}^{(3)}. \quad (41)$$

We can rotate this vector field using the results of appendix B to give the coefficients of equation (25).

We provide formulae by which the incident field coefficients can be derived for two cases. These represent simple, but solvable sources. The incident field coefficients from more complicated sources can be substituted for the coefficients given in this section. In the first case, we consider the field due to an explosion. In this case, the source pressure wave is not angularly dependent, and we can express it as $-p_o \mathbf{L}_{00,3}^{(3)}$, or as

$$h_{nm}^{(1)} = h_{nm}^{(1)p} = -p_o \delta_{n,0} \delta_{m,0}, \quad (42)$$

$$h_{nm}^{(2)} = h_{nm}^{(2)p} = 0, \quad (43)$$

and

$$h_{nm}^{(3)} = h_{nm}^{(3)p} = 0. \quad (44)$$

In the second case, we consider a source that emits shear waves only, whose shearing stress $\tau_{r\varphi} = \tau_o \sin \theta_3$ is applied at the inner surface of a spherical cavity centered on the source. In this case the source coefficients are of the form

$$h_{nm}^{(1)} = h_{nm}^{(1)s} = 0 , \quad (45)$$

$$h_{nm}^{(2)} = h_{nm}^{(2)s} = \tau_o \delta_{n,1} \delta_{m,0} , \quad (46)$$

and

$$h_{nm}^{(3)} = h_{nm}^{(3)s} = 0 . \quad (47)$$

If the shear is in another direction, we can use the vector rotation results given in appendix B. The final expressions for the source coefficients in the host sphere coordinate system are given by equations (B-31) to (B-33).

3. Implications

In this report, we derive equations describing the scattered fields from a spherical scatterer located eccentrically within a host sphere. The source is also arbitrarily located within the host sphere, and can be chosen to simulate a P wave, S wave, or any linear combination of the two. We can use this theory in seismology to predict the scattering interaction between a scattering body and the Earth's surface.

The scattered fields are complete, and exact solutions for the elastic wave-field produced by the spherical scatterer are expressed. Because the solution presented here is almost entirely analytical, it avoids the need for much of the computationally intensive algorithms, such as finite-difference schemes, that would normally be required to solve this problem. Also, because we have made very few assumptions concerning the boundary conditions, the analytical solution given here should be well behaved.

This theory provides a basis that should prove useful in understanding and predicting seismic scattering. Using techniques developed in electromagnetic scattering (Videen *et al*, 1995), it should be possible to extend this theory to describe the scattering from an arbitrary body rather than a spherical one. Hence, this theory is an important first step in demonstrating a method for analytically deriving scattered elastic wave fields from anomalies such as magma chambers or subducting slabs near the Earth's surface.

Appendix A. Boundary Coefficients

We can find the coefficients that satisfy the boundary conditions at the scattering sphere outer surface ($r_2 = a'$) by first eliminating the dependence on the internal field coefficients. Performing some algebraic manipulation and taking advantage of the differential equation defining spherical Bessel functions,

$$r^2 z_n''(r) + 2r z_n'(r) + [r^2 - n(n+1)] z_n(r) = 0, \quad (\text{A-1})$$

we can put the coefficients that satisfy the boundary conditions on the scattering sphere into the following form:

$$Q_{nm}^{11} = -\frac{A_1^1 B_2^3 - A_2^1 B_1^3}{B_1^1 B_2^3 - B_2^1 B_1^3}, \quad (\text{A-2})$$

$$Q_{nm}^{13} = -\frac{A_1^3 B_2^3 - A_2^3 B_1^3}{B_1^1 B_2^3 - B_2^1 B_1^3}, \quad (\text{A-3})$$

$$Q_{nm}^{22} = -\frac{J_n(1, 1)}{J_n(3, 1)}, \quad (\text{A-4})$$

$$Q_{nm}^{31} = \frac{A_1^1 B_2^1 - A_2^1 B_1^1}{B_1^1 B_2^3 - B_2^1 B_1^3}, \quad (\text{A-5})$$

$$Q_{nm}^{33} = \frac{A_1^3 B_2^1 - A_2^3 B_1^1}{B_1^1 B_2^3 - B_2^1 B_1^3}, \quad (\text{A-6})$$

and

$$R_{nm}^{11} = -\frac{C_1^1 B_2^3 - C_2^1 B_1^3}{B_1^1 B_2^3 - B_2^1 B_1^3}, \quad (\text{A-7})$$

$$R_{nm}^{13} = -\frac{C_1^3 B_2^3 - C_2^3 B_1^3}{B_1^1 B_2^3 - B_2^1 B_1^3}, \quad (\text{A-8})$$

$$R_{nm}^{22} = -\frac{J_n(4, 1)}{J_n(3, 1)}, \quad (\text{A-9})$$

$$R_{nm}^{31} = \frac{C_1^1 B_2^1 - C_2^1 B_1^1}{B_1^1 B_2^3 - B_2^1 B_1^3}, \quad (\text{A-10})$$

$$R_{nm}^{33} = \frac{C_1^3 B_2^1 - C_2^3 B_1^1}{B_1^1 B_2^3 - B_2^1 B_1^3}, \quad (\text{A-11})$$

where

$$\begin{aligned} \mathcal{A}_1^n &= (A_1^n D_2^3 - A_2^n D_1^3) (D_3^1 D_4^3 - D_4^1 D_3^3) \\ &\quad - (A_3^n D_4^3 - A_4^n D_3^3) (D_1^1 D_2^3 - D_2^1 D_1^3) \end{aligned} \quad (\text{A-12})$$

$$\begin{aligned} \mathcal{A}_2^n &= (A_1^n D_2^3 - A_2^n D_1^3) (D_2^1 D_3^3 - D_3^1 D_2^3) \\ &\quad - (A_2^n D_3^3 - A_3^n D_2^3) (D_1^1 D_2^3 - D_2^1 D_1^3) \end{aligned} \quad (\text{A-13})$$

$$\begin{aligned} \mathcal{B}_1^n &= (B_1^n D_2^3 - B_2^n D_1^3) (D_3^1 D_4^3 - D_4^1 D_3^3) \\ &\quad - (B_3^n D_4^3 - B_4^n D_3^3) (D_1^1 D_2^3 - D_2^1 D_1^3) \end{aligned} \quad (\text{A-14})$$

$$\begin{aligned} \mathcal{B}_2^n &= (B_1^n D_2^3 - B_2^n D_1^3) (D_2^1 D_3^3 - D_3^1 D_2^3) \\ &\quad - (B_2^n D_3^3 - B_3^n D_2^3) (D_1^1 D_2^3 - D_2^1 D_1^3) \end{aligned} \quad (\text{A-15})$$

$$\begin{aligned} \mathcal{C}_1^n &= (C_1^n D_2^3 - C_2^n D_1^3) (D_3^1 D_4^3 - D_4^1 D_3^3) \\ &\quad - (C_3^n D_4^3 - C_4^n D_3^3) (D_1^1 D_2^3 - D_2^1 D_1^3) \end{aligned} \quad (\text{A-16})$$

$$\begin{aligned} \mathcal{C}_2^n &= (C_1^n D_2^3 - C_2^n D_1^3) (D_2^1 D_3^3 - D_3^1 D_2^3) \\ &\quad - (C_2^n D_3^3 - C_3^n D_2^3) (D_1^1 D_2^3 - D_2^1 D_1^3) \end{aligned} \quad (\text{A-17})$$

$$A_j^1 = H_{2j-1}(1, k_\alpha, \lambda, \mu, a'), \quad (\text{A-18})$$

$$A_j^3 = H_{2j}(1, k_\beta, \lambda, \mu, a'), \quad (\text{A-19})$$

$$B_j^1 = H_{2j-1}(3, k_\alpha, \lambda, \mu, a'), \quad (\text{A-20})$$

$$B_j^3 = H_{2j}(3, k_\beta, \lambda, \mu, a'), \quad (\text{A-21})$$

$$C_j^1 = H_{2j-1}(4, k_\alpha, \lambda, \mu, a'), \quad (\text{A-22})$$

$$C_j^3 = H_{2j}(4, k_\beta, \lambda, \mu, a'), \quad (\text{A-23})$$

$$D_j^1 = H_{2j-1}(1, k'_\alpha, \lambda', \mu', a'), \quad (\text{A-24})$$

$$D_j^3 = H_{2j}(1, k'_\beta, \lambda', \mu', a'), \quad (\text{A-25})$$

$$J_n(i, j) = H_n(i, k_\beta, \lambda, \mu, a') - \frac{z_n^{(i)}}{z_n^{(j)}} H_n(j, k'_\beta, \lambda', \mu', a'),$$

and

$$H_1(p, k, \lambda, \mu, a) = 2\mu \left[z_n^{(p)'}(ka) - \frac{z_n^{(p)}(ka)}{ka} \right], \quad (\text{A-26})$$

$$H_2(p, k, \lambda, \mu, a) = 2\mu(n^2 + n - 1) \frac{z_n^{(p)}(ka)}{ka} - 2\mu z_n^{(p)'}(ka) - \mu k a z_n^{(p)}(ka), \quad (\text{A-27})$$

$$H_3(p, k, \lambda, \mu, a) = \left[\frac{2\mu n(n+1)}{ka} - ka(\lambda + 2\mu) \right] z_n^{(p)}(ka) - 4\mu z_n^{(p)'}(ka),$$

$$H_4(p, k, \lambda, \mu, a) = \mu n(n+1)H_1(p, k, a). \quad (\text{A-28})$$

The coefficients that satisfy the boundary conditions on the host-sphere outer surface ($r_1 = a$) are

$$S_{nm}^{11} = \frac{H_3(3, k_\alpha, \lambda, \mu, a)H_2(4, k_\beta, \lambda, \mu, a) - H_1(3, k_\alpha, \lambda, \mu, a)H_4(4, k_\beta, \lambda, \mu, a)}{H_1(4, k_\alpha, \lambda, \mu, a)H_4(4, k_\beta, \lambda, \mu, a) - H_3(4, k_\alpha, \lambda, \mu, a)H_2(4, k_\beta, \lambda, \mu, a)} \quad (\text{A-29})$$

$$S_{nm}^{13} = \frac{H_4(3, k_\beta, \lambda, \mu, a)H_2(4, k_\beta, \lambda, \mu, a) - H_2(3, k_\beta, \lambda, \mu, a)H_4(4, k_\beta, \lambda, \mu, a)}{H_1(4, k_\alpha, \lambda, \mu, a)H_4(4, k_\beta, \lambda, \mu, a) - H_3(4, k_\alpha, \lambda, \mu, a)H_2(4, k_\beta, \lambda, \mu, a)} \quad (\text{A-30})$$

$$S_{nm}^{22} = -\frac{H_1(3, k_\beta, \lambda, \mu, a)}{H_1(4, k_\beta, \lambda, \mu, a)} \quad (\text{A-31})$$

$$S_{nm}^{31} = \frac{H_3(3, k_\alpha, \lambda, \mu, a)H_1(4, k_\alpha, \lambda, \mu, a) - H_1(3, k_\alpha, \lambda, \mu, a)H_3(4, k_\alpha, \lambda, \mu, a)}{H_3(4, k_\alpha, \lambda, \mu, a)H_2(4, k_\beta, \lambda, \mu, a) - H_1(4, k_\alpha, \lambda, \mu, a)H_4(4, k_\beta, \lambda, \mu, a)} \quad (\text{A-32})$$

$$S_{nm}^{33} = \frac{H_4(3, k_\beta, \lambda, \mu, a)H_1(4, k_\alpha, \lambda, \mu, a) - H_2(3, k_\beta, \lambda, \mu, a)H_3(4, k_\alpha, \lambda, \mu, a)}{H_3(4, k_\alpha, \lambda, \mu, a)H_2(4, k_\beta, \lambda, \mu, a) - H_1(4, k_\alpha, \lambda, \mu, a)H_4(4, k_\beta, \lambda, \mu, a)} \quad (\text{A-33})$$

Appendix B. Vector Translations

Stein (1961) and Cruzan (1962) derived translation addition theorems for vector spherical wave functions that can be used to express the displacement fields in one coordinate system in terms of displacement fields in another coordinate system. These results are also given by Ben-Menahem and Singh (1981). We use their results to find relationships between the coefficients $a_{nm}^{(p)}$ and $e_{nm}^{(p)}$, $b_{nm}^{(p)}$ and $f_{nm}^{(p)}$, and $c_{nm}^{(p)}$ and $g_{nm}^{(p)}$. For a translation along the z -axis with no rotation, the vector spherical harmonics are related by

$$\mathbf{L}_{nm,2}^{(p)} = \sum_{n'=0}^{\infty} C_{n,n'}^{(m,i)} \mathbf{L}_{n'm,1}^{(q)}, \quad (\text{B-1})$$

$$\mathbf{M}_{nm,2}^{(p)} = \sum_{n'=0}^{\infty} A_{n,n'}^{(m,i)} \mathbf{M}_{n'm,1}^{(q)} + B_{n,n'}^{(m,i)} \mathbf{N}_{n'm,1}^{(q)}, \quad (\text{B-2})$$

$$\mathbf{N}_{nm,2}^{(p)} = \sum_{n'=0}^{\infty} B_{n,n'}^{(m,i)} \mathbf{M}_{n'm,1}^{(q)} + A_{n,n'}^{(m,i)} \mathbf{N}_{n'm,1}^{(q)}. \quad (\text{B-3})$$

If $p = 3$ or 4 , then $i = 1$ in the region where $r_1 < |d|$, and $i = p$ in the region where $r_i > |d|$. Recurrence relations for the scalar translation coefficients $C_{n,n'}^{(m,i)}$ were derived by Bobbert and Vlieger (1986), which greatly facilitate calculations. The results are

$$C_{0,n'}^{(0,i)} = \sqrt{2n'+1} z_{n'}^{(i)}(kd), \quad (\text{B-4})$$

$$C_{-1,n'}^{(0,i)} = -\sqrt{2n'+1} z_{n'}^{(i)}(kd), \quad (\text{B-5})$$

$$C_{n+1,n'}^{(0,i)} = \frac{1}{(n+1)} \sqrt{\frac{2n+3}{2n'+1}} n' \sqrt{\frac{2n+1}{2n'-1}} C_{n,n'-1}^{(0,i)} + n \sqrt{\frac{2n'+1}{2n-1}} C_{n-1,n'}^{(0,i)} - (n'+1) \sqrt{\frac{2n+1}{2n'+3}} C_{n,n'+1}^{(0,i)}, \quad (\text{B-6})$$

$$\sqrt{(n-m+1)(n+m)(2n'+1)} C_{n,n'}^{(m,i)} = \sqrt{(n'-m+1)(n'+m)(2n'+1)} C_{n,n'}^{(m-1,i)} - \quad (\text{B-7})$$

$$kd \sqrt{\frac{(n'-m+2)(n'-m+1)}{(2n'+3)}} C_{n,n'+1}^{(m-1,i)} - kd \sqrt{\frac{(n'+m)(n'+m-1)}{(2n'-1)}} C_{n,n'-1}^{(m-1,i)}.$$

From the scalar translation coefficients, the vector translation coefficients can be easily derived:

$$A_{n,n'}^{(m,i)} = C_{n,n'}^{(m,i)} - \frac{kd}{n'+1} \sqrt{\frac{(n'-m+1)(n'+m+1)}{(2n'+1)(2n'+3)}} C_{n,n'+1}^{(m,i)} \quad (\text{B-8})$$

$$- \frac{kd}{n'} \sqrt{\frac{(n'-m)(n'+m)}{(2n'+1)(2n'-1)}} C_{n,n'-1}^{(m,i)},$$

$$B_{n,n'}^{(m,i)} = \frac{-imkd}{n'(n'+1)} C_{n,n'}^{(m,i)}. \quad (\text{B-9})$$

From these equations, we see that

$$A_{n,n'}^{(m,p,q)} = A_{n,n'}^{(-m,p,q)}, \quad (\text{B-10})$$

$$B_{n,n'}^{(m,p,q)} = B_{n,n'}^{(-m,p,q)}, \quad (\text{B-11})$$

$$C_{n,n'}^{(m,p,q)} = C_{n,n'}^{(-m,p,q)}. \quad (\text{B-12})$$

Since $\mathbf{u}_{sou}^1 = \mathbf{u}_{sou}^2$, $\mathbf{u}_{sca}^1 = \mathbf{u}_{sca}^2$, and $\mathbf{u}_{int}^1 = \mathbf{u}_{int}^2$, we can derive expressions relating the displacement coefficients using equations (B-1) to (B-3):

$$a_{nm}^{(1)} = \sum_{n'=0}^{\infty} e_{n'm}^{(1)} C_{n',n}^{(m,3)}(-k_{\beta}d), \quad (\text{B-13})$$

$$a_{nm}^{(2)} = \sum_{n'=0}^{\infty} e_{n'm}^{(2)} A_{n',n}^{(m,3)}(-k_{\beta}d) + e_{n'm}^{(3)} B_{n',n}^{(m,3)}(-k_{\beta}d), \quad (\text{B-14})$$

$$a_{nm}^{(3)} = \sum_{n'=0}^{\infty} e_{n'm}^{(3)} A_{n',n}^{(m,3)}(-k_{\beta}d) + e_{n'm}^{(2)} B_{n',n}^{(m,3)}(-k_{\beta}d) \quad (\text{B-15})$$

$$f_{nm}^{(1)} = \sum_{n'=0}^{\infty} b_{n'm}^{(1)} C_{n',n}^{(m,1)}(k_{\alpha}d), \quad (\text{B-16})$$

$$f_{nm}^{(2)} = \sum_{n'=0}^{\infty} b_{n'm}^{(2)} A_{n',n}^{(m,1)}(k_{\beta}d) + b_{n'm}^{(3)} B_{n',n}^{(m,1)}(k_{\beta}d), \quad (\text{B-17})$$

$$f_{nm}^{(3)} = \sum_{n'=0}^{\infty} b_{n'm}^{(3)} A_{n',n}^{(m,1)}(k_{\beta}d) + b_{n'm}^{(2)} B_{n',n}^{(m,1)}(k_{\beta}d), \quad (\text{B-18})$$

$$g_{nm}^{(1)} = \sum_{n'=0}^{\infty} c_{n'm}^{(1)} C_{n',n}^{(m,1)}(k_{\alpha}d), \quad (\text{B-19})$$

$$g_{nm}^{(2)} = \sum_{n'=0}^{\infty} c_{n'm}^{(2)} A_{n',n}^{(m,1)}(k_{\beta}d) + c_{n'm}^{(3)} B_{n',n}^{(m,1)}(k_{\beta}d), \quad (\text{B-20})$$

$$g_{nm}^{(3)} = \sum_{n'=0}^{\infty} c_{n'm}^{(3)} A_{n',n}^{(m,1)}(k_{\beta}d) + c_{n'm}^{(2)} B_{n',n}^{(m,1)}(k_{\beta}d). \quad (\text{B-21})$$

Finally, we note that the scalar translation coefficients $C_{n,n'}^{(m,i)}$ used in deriving the vector translation coefficients $A_{n,n'}^{(m,i)}$ and $B_{n,n'}^{(m,i)}$ are, in general, different from the vector translation coefficients $C_{n,n'}^{(m,i)}$ used in the translation of the P waves ($\mathbf{L}_{mn}^{(p)}$), since in general the P and S waves have different spatial frequencies. In addition, the translations for the source coefficients $a_{nm}^{(p)}$ and $e_{nm}^{(p)}$ are in opposite directions.

The translation from the source coordinate system to the rotated host coordinate system is similar:

$$i_{nm}^{(1)} = \sum_{n'=0}^{\infty} h_{n'm}^{(1)} C_{n',n}^{(m,1)}(k_{\alpha}r_s), \quad (\text{B-22})$$

$$i_{nm}^{(2)} = \sum_{n'=0}^{\infty} h_{n'm}^{(2)} A_{n',n}^{(m,1)}(k_{\beta}r_s) + h_{n'm}^{(3)} B_{n',n}^{(m,1)}(k_{\beta}r_s), \quad (\text{B-23})$$

$$i_{nm}^{(3)} = \sum_{n'=0}^{\infty} h_{n'm}^{(3)} A_{n',n}^{(m,1)}(k_{\beta}r_s) + h_{n'm}^{(2)} B_{n',n}^{(m,1)}(k_{\beta}r_s). \quad (\text{B-24})$$

Using the results of Stein (1960), we present the coefficients in a rotated coordinate system. The vector harmonics can be expanded as

$$\mathbf{L}_{nm,j}^{(p)} = \sum_{m'=0}^{\infty} D_{m'}^{(n,m)} \mathbf{L}_{nm',i}^{(p)}, \quad (\text{B-25})$$

$$\mathbf{M}_{nm,j}^{(p)} = \sum_{m'=0}^{\infty} D_{m'}^{(n,m)} \mathbf{M}_{nm',i}^{(q)}, \quad (\text{B-26})$$

$$\mathbf{N}_{nm,j}^{(p)} = \sum_{m'=0}^{\infty} D_{m'}^{(n,m)} \mathbf{N}_{nm',i}^{(q)}, \quad (\text{B-27})$$

where

$$D_{m'}^{(n,m)} = \exp [i(m'\alpha + m\gamma)] \left[\frac{(n+m')!(n-m')!}{(n+m)!(n-m)!} \right]^{1/2} \quad (\text{B-28})$$

$$\times \sum_{\sigma} \binom{n+m}{n-m'-\sigma} \binom{n-m}{\sigma} (-1)^{n+m-\sigma} [\cos(\beta/2)]^{2\sigma+m'+m} \quad (\text{B-29})$$

$$[\sin(\beta/2)]^{2n-2\sigma-m'-m}$$

and α , β , and γ are Euler angles using the convention of Edmonds (1957). The coefficients in the rotated coordinate system, $a_{nm,i}$, can be expressed in the unrotated coordinate system, $a_{nm,j}$, as

$$a_{nm,j} = \sum_{m'=0}^{\infty} D_{m'}^{(n,m)} a_{nm',i} . \quad (\text{B-30})$$

Therefore, the source coefficients in the host coordinate system for the point source can be expressed as

$$e_{nm}^{(1)} = \sum_{m'=0}^{\infty} D_{m'}^{(n,m)} \sum_{n'=0}^{\infty} h_{n'm'}^{(1)} C_{n',n}^{(m',1)}(k_{\alpha} r_s) , \quad (\text{B-31})$$

$$e_{nm}^{(2)} = \sum_{m'=0}^{\infty} D_{m'}^{(n,m)} \sum_{n'=0}^{\infty} h_{n'm'}^{(2)} A_{n',n}^{(m',1)}(k_{\beta} r_s) + h_{n'm'}^{(3)} B_{n',n}^{(m',1)}(k_{\beta} r_s) , \quad (\text{B-32})$$

$$e_{nm}^{(3)} = \sum_{m'=0}^{\infty} D_{m'}^{(n,m)} \sum_{n'=0}^{\infty} h_{n'm'}^{(3)} A_{n',n}^{(m',1)}(k_{\beta} r_s) + h_{n'm'}^{(2)} B_{n',n}^{(m',1)}(k_{\beta} r_s) . \quad (\text{B-33})$$

References

1. Ben-Menahem, A., and S. J. Singh, 1981. *Seismic Waves and Sources*, Springer-Verlag, New York.
2. Bobbert, P. A., and J. Vlieger, 1986. "Light scattering by a sphere on a substrate," *Physica* **137A**, 209–241.
3. Bohren, C. F., and D. R. Huffman, 1983. *Absorption and Scattering of Light by Small Particles*, Wiley, New York.
4. Cruzan, O. R., 1962. "Translational addition theorems for spherical vector wave functions," *Q. Appl. Math.* **20**, 33–40.
5. Dubrovskiy, V. A., and V. S. Morozhnik, 1989. "Nonstationary scattering of elastic waves by spherical inclusion," *Izvestiya, Earth Physics* **25**, 679–686.
6. Edmonds, A. R., 1957. *Angular Momentum in Quantum Mechanics*, Princeton University Press, Princeton, New Jersey, 45–51.
7. Einspruch, N. G, E. J. Witterholt, and R. Truell, 1960. "Scattering of a plane transverse wave by a spherical obstacle in an elastic medium," *J. Appl. Phys.* **31**, 806–818.
8. Glenn, L. A., A.J.C. Ladd, B. Moran, and K. A. Wilson, 1985. "Elastic radiation from explosively-loaded ellipsoidal cavities in an unbounded medium," *Geophys. J. R. Astron. Soc.* **86**, 231–242.
9. Rial, J. A., and B. Moran, 1986. "Radiation patterns for explosively-loaded axisymmetric cavities in an elastic medium: analytic approximations and numerical results," *Geophys. J. R. Astron. Soc.* **86**, 855–862.
10. Stein, S., 1961. "Addition theorems for spherical wave functions," *Q. Appl. Math.* **19**, 15–24.
11. Stratton, J. A., 1941. *Electromagnetic Theory*, McGraw-Hill, New York.

12. Thompson, W. Jr., 1973. "Acoustic radiation from a spherical source embedded eccentrically within a fluid sphere," *J. Acoust. Soc. Am.* **54**, 1694-1707.
13. Varatharajulu, V., and Y.-H. Pao, 1976. "Scattering matrix for elastic waves. I. Theory," *J. Acoust. Soc. Am.* **60**, 556-566.
14. Videen, G., D. Ngo, P. Chýlek, and R. G. Pinnick, 1995. "Light scattering from a sphere with an irregular inclusion," *J. Opt. Soc. Am. A* **12**, 922-928.
15. Waterman, P. C., 1969. "New formulation of acoustic scattering," *J. Acoust. Soc. Am.* **45**, 1417-1429.
16. Waterman, P. C., 1976. "Matrix theory of elastic waves scattering," *J. Acoust. Soc. Am.* **60**, 567-580.
17. Wiscombe, W. J., 1980. "Improved Mie scattering algorithms," *Appl. Opt.* **19**, 1505-1509.
18. Wu, R.-S., 1985. "Multiple scattering and energy transfer of seismic waves—separation of scattering effect from intrinsic attenuation—I. Theoretical modelling," *Geophys. J. R. Astron. Soc.* **82**, 57-80.
19. Wu, R.-S., and K. Aki, 1985. "Scattering characteristics of elastic waves by an elastic heterogeneity," *Geophysics* **50**, 4.
20. Ying, C. F., and R. Truell, 1955. "Scattering of a plane longitudinal wave by a spherical obstacle in an isotropically elastic solid," *J. Appl. Phys.* **27**, 1086-1097.
21. Zhao, L.-S., and D. G. Harkrider, 1992. "Wave fields from an off-center explosion in an embedded solid sphere," *Bull. Seismol. Soc. Am.* **82**, 1927-1955.

Distribution

Admnstr
Defns Techl Info Ctr
Attn DTIC-OCF
8725 John J Kingman Rd Ste 0944
FT Belvoir VA 22060-6218

Central Intellignc Agency Dir DB Standard
Attn GE 47 QB
Washington DC 20505

Chairman Joint Chiefs of Staff
Attn J5 R&D Div
Washington DC 20301

Defns Intllgnc Acgy
Attn DT 2 Wpns & Sys Div
Washington DC 20301

Dir of Defns Rsrch & Engrg
Attn DD TWP
Attn Engrg
Washintgon DC 20301

Ofc of the Secy of Defns
Attn ODDRE (R&AT) G Singley
Attn ODDRE (R&AT) S Gontarek
The Pentagon
Washington DC 20301-3080

OIR CSB CRB
Attn A M Jones
RB 1413 OHM
Washington DC 20505

US Dept of Energy
Attn KK 22 K Sisson
Attn Techl Lib
Washington DC 20585

Commanding Officer
Attn NMCB23
6205 Stuart Rd Ste 101
FT Belvoir VA 22060-5275

CECOM
Attn PM GPS COL S Young
FT Monmouth NJ 07703

CECOM RDEC Electronic Systems Div Dir
Attn J Niemela
FT Monmouth NJ 07703

CECOM
Sp & Terrestrial Commctn Div
Attn AMSEL-RD-ST-MC-M H Soicher
FT Monmouth NJ 07703-5203

DARPA
Attn B Kaspar
Attn J Pennella
Attn L Stotts
Attn Techl Lib
3701 N Fairfax Dr
Arlington VA 22203-1714

Dir of Chem & Nuc Ops DA DCSOPS
Attn Techl Lib
Washington DC 20310

Dpty Assist Secy for Rsrch & Techl
Attn SARD-TR R Chait
Attn SARD-TT D Chait
Attn SARD-TT F Milton Rm 3E479
Attn SARD-TT K Kominos
Attn SARD-TT R Reisman
Attn SARD-TT T Killion
Attn SARD-TT C Nash Rm 3E479
The Pentagon Rm 3E476
Washington DC 20310-0103

DUSD Space
Attn 1E765 J G McNeff
3900 Defense Pentagon
Washington DC 20301-3900

Hdqtrs Dept of the Army
Attn DAMO-FDQ D Schmidt
400 Army Pentagon
Washington DC 20310-0460

OSD
Attn OUSD(A&T)/ODDDR&E(R) J Lupo
The Pentagon
Washington DC 20301-7100

US Army Engrg Div
Attn HNDED FD
PO Box 1500
Huntsville AL 35807

Distribution

US Army Matl Cmnd
Dpty CG for RDE Hdqtrs
Attn AMCRD BG Beauchamp
5001 Eisenhower Ave
Alexandria VA 22333-0001

US Army Matl Cmnd
Prin Dpty for Acquisition Hdqtrs
Attn AMCDCG-A D Adams
5001 Eieenhower Ave
Alexandria VA 22333-0001

US Army Matl Cmnd
Prin Dpty for Techlgy Hdqtrs
Attn AMCDCG-T M Fisetete
5001 Eisenhower Ave
Alexandria VA 22333-0001

US Army Mis & Spc Intllgnc Ctr
Attn AIAMS YDL
Redstone Arsenal AL 35898-5500

US Army NGIC
Attn Rsrch & Data Branch
220 7th Stret NE
Charlottesville VA 22901-5396

US Army Nuc & Cheml Agency
7150 Heller Loop Ste 101
Springfield VA 22150-3198

US Army Rsrch Lab
Attn SLCRO-D
PO Box 12211
Research Triangle Park NC 27709-2211

US Army Strtgc Defns Cmnd
Attn CSSD H MPL Techl Lib
Attn CSSD H XM Dr Davies
PO Box 1500
Huntsville AL 35807

US Military Academy
Dept of Mathematical Sci
Attn MAJ D Engen
West Point NY 10996

USAASA
Attn MOAS-AI W Parron
9325 Gunston Rd Ste N319
FT Belvoir VA 22060-5582

Chief of Nav OPS Dept of the Navy
Attn OP 03EG
Washington DC 20350

GPS Joint Prog Ofc Dir
Attn COL J Clay
2435 Vela Way Ste 1613
Los Angeles AFB CA 90245-5500

Ofc of the Dir Rsrch and Engrg
Attn R Menz
Pentagon Rm 3E1089
Washington DC 20301-3080

Special Assist to the Wing Cmndr
Attn 50SW/CCX CAPT P H Bernstein
300 O'Malley Ave Ste 20
Falcon AFB CO 80912-3020

USAF SMC/CED
Attn DMA/JPO M Ison
2435 Vela Way Ste 1613
Los Angeles AFB CA 90245-5500

ARL Electromag Group
Attn Campus Mail Code F0250 A Tucker
University of TX
Austin TX 78712

US Army Rsrch Lab
Attn AMSRL-CI-LL Tech Lib (3 copies)
Attn AMSRL-CS-AL-TA Mail & Records
Mgmt
Attn AMSRL-CS-AL-TP Techl Pub (3 copies)
Attn AMSRL-IS-EE G Videen (15 copies)
Adelphi MD 20783-1197

REPORT DOCUMENTATION PAGE			<i>Form Approved OMB No. 0704-0188</i>	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE August 1997	3. REPORT TYPE AND DATES COVERED Progress, 1 Oct 1996 to 1 Oct 1997	
4. TITLE AND SUBTITLE Elastic Scattering from a Spherical Body Eccentrically Located Within a Sphere: Theoretical Derivation			5. FUNDING NUMBERS PE: 61102A DA: B53A	
6. AUTHOR(S) Gorden Videen (ARL) and Eric Sandvol (Cornell University)				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) U.S. Army Research Laboratory Attn: AMSRL-IS-EE 2800 Powder Mill Road Adelphi, MD 20783-1197			8. PERFORMING ORGANIZATION REPORT NUMBER ARL-TR-1394	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Research Laboratory 2800 Powder Mill Road Adelphi, MD 20783-1197			10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES AMS code: 611102.53A11 ARL PR: 7FEJ60				
12a. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited.			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) <p>The scattered fields from a spherical body eccentrically located within a host sphere are found by a derivation that satisfies the boundary conditions at both interfaces. The source, which may be composed of any linear combination of <i>S</i> and <i>P</i> waves, is also arbitrarily located within the host sphere. The scattering system has applications in seismic scattering, since the scattering interaction between a scatterer and the Earth's surface is significant when the scatterer is located near the surface. This interaction can affect subsequent seismograms.</p>				
14. SUBJECT TERMS Elastic scattering, magma chambers, sphere scattering, seismic scatter			15. NUMBER OF PAGES 32	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT UL	