

AD _____

CONTRACT NUMBER DAMD17-93-C-3084

TITLE: An *In Vitro* Assay for Toxic Particulates

PRINCIPAL INVESTIGATOR: Richard H. C. San, Ph.D.

CONTRACTING ORGANIZATION: Microbiological Associates, Inc.
Rockville, Maryland 20850

REPORT DATE: 16 October 1997

TYPE OF REPORT: Final

PREPARED FOR: Commander
U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for public release;
distribution unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

19971114 048

DTIC QUALITY INSPECTED 2

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

| | | | |
|---|--|---|---|
| 1. AGENCY USE ONLY - (Leave blank) | 2. REPORT DATE 16 October 1997 | 3. REPORT TYPE AND DATES COVERED Final (15 Feb 93 - 15 May 94) | |
| 4. TITLE AND SUBTITLE An In Vitro Assay for Toxic Particulates | | 5. FUNDING NUMBERS DAMD17-93-C-3084 | |
| 6. AUTHOR(S) San, Richard H.C., Ph.D. | | | |
| 7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Microbiological Associates, Inc. Rockville, Maryland 20850 | | 8. PERFORMING ORGANIZATION REPORT NUMBER | |
| 9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012 | | 10. SPONSORING / MONITORING AGENCY REPORT NUMBER | |
| 11. SUPPLEMENTARY NOTES | | | |
| 12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited | | 12b. DISTRIBUTION CODE | |
| 13. ABSTRACT (Maximum 200 words) The purpose of this study is to evaluate the use of cells from a mouse macrophage cell line RAW 264.7 as an <i>in vitro</i> test for determining the cytotoxicity and mutagenicity of particulates potentially hazardous to the respiratory system. Mutagenicity assays were conducted on the hypoxanthine guanine phosphoribosyl transferase (HGPRT) gene locus. The mutant frequency in the untreated control in some experiments was substantially higher than the spontaneous frequency of between 0 and 10 mutants per 10 ⁶ cells reported by Hitchins <i>et al.</i> (1992). It remains to be determined as to whether the variability in spontaneous mutant frequency is an inherent feature of this cell line. The inability to demonstrate mutagenicity with low fire or high fire beryllium oxide indicates that the assay procedure still requires further refinement. No mutagenic activity was demonstrated with red dye mix. A dose-related increase in mutant frequency was observed with 10% trinitrotoluene (TNT)-contaminated compost in only one experiment. Since the TNT-contaminated compost was not tested at concentrations higher than 100 µg/mL using the 18-hour extended treatment regimen, additional experiments are warranted prior to making comments on the potential mutagenicity of soils contaminated with munitions. | | | |
| 14. SUBJECT TERMS Toxins, Cytotoxicity, Mutagenicity, Berillium Oxide, Ethyl Methanesulfonate, Trinitrotoluene | | 15. NUMBER OF PAGES 22 | |
| | | 16. PRICE CODE | |
| 17. SECURITY CLASSIFICATION OF REPORT Unclassified | 18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified | 19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified | 20. LIMITATION OF ABSTRACT Unlimited |

FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the US Army.

_____ Where copyrighted material is quoted, permission has been obtained to use such material.

_____ Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

PS ✓ Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.

_____ In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and Use of Laboratory Animals of the Institute of Laboratory Resources, National Research Council (NIH Publication No. 86-23, Revised 1985).

_____ For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.

_____ In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

_____ In the conduct of research utilizing recombinant DNA, the investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules.

_____ In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories.

Richard X October 16, 1997

PI - Signature

Date

TABLE OF CONTENTS

| | Page |
|---|------|
| Introduction | 1 |
| Materials and Methods | 2 |
| Test Materials | 2 |
| Cell Line and Culture Conditions | 2 |
| Cytotoxicity Assay | 3 |
| Mutagenicity Assay | 4 |
| Results | 5 |
| Low and High Fire Berillium Oxide (BeO) | 5 |
| Ethyl Methanesulfonate (EMS) | 6 |
| Comparison of RAW 264.7 from Two Different Stock Cultures | 6 |
| Trinitrotoluene (TNT)-Contaminated Compost and Uncontaminated Soil Sample | 6 |
| Red Dye Mix | 6 |
| Conclusions | 6 |
| References | 7 |
| Data Tables | 9 |

INTRODUCTION

Toxic assault through the respiratory tract is a major threat in both working and domestic environments. Most of these toxic materials are small, soluble molecules which can be easily tested for cytotoxicity and genotoxicity by current short term tests. However, around 10% of airborne pollution is made up of particulates (Amdur, 1986). These may be smokes, fly ash, mineral dusts or other particles of 5 microns or less and therefore able to pass into the lungs (Menzel and Amdur, 1986). Since there is no direct assay for particulates, extracts of particulate materials are often tested in the established assays (Brusick, 1982; Zelikoff, 1985). This approach assumes that the active portion of the particulate is extractable and that the non-extractable solid portion of the particulate is inactive. However, there is evidence that certain solids are directly genotoxic (asbestos, glass fibers and certain metals) and that the form and size of the particle is critical in determining its toxicity (Sincook *et al.*, 1982; Hesterberg *et al.*, 1983). These properties would be missed in an assay based on extraction of particulate material. Thus, there is a need for an *in vitro* assay system utilizing cells which can actively phagocytize the particles to be tested and allowing assessment of their effects through several endpoints.

Attempts to use traditional short term tests to assess cytotoxicity and genotoxicity of particulates directly have met with varying success. Zelikoff *et al.* (1985) used V79 Chinese hamster lung fibroblasts to study the cytotoxicity of silica, alumina, wood charcoal and coal fly ash. They were able to demonstrate phagocytosis of these particles but it is not clear what percentage of the particles were absorbed. Costa and Heck (1982) studied the ability of Chinese hamster ovary cells to phagocytize specific crystalline and amorphous nickel, cobalt, copper and cadmium compounds. Between 14% and 49% of the cells ingested the crystalline salts while only 3% to 6% of the cells phagocytized the amorphous salts. These results were consistent with results of the Syrian hamster embryo transformation assay which showed greater transformation by the crystalline salts. The percentage of cells which actually phagocytized the particles was low. Rossman *et al.* (1987) reviewed the genetic toxicity of metal compounds in *in vitro* systems. They found that bacterial systems give mixed results in comparison to *in vivo* systems. Genotoxic effects of soluble salts were detected in these systems, while the effects of insoluble salts were not. Cultured, primary rat liver cells were a better model for determining cytotoxicity than bacteria. However, these cells were nondividing and needed to be obtained directly from donor animals and could not be easily used in mutagenicity assays.

Macrophages play an important role in the lung in trapping, concentrating, and processing, through both superoxide and lysosomal enzyme systems, potentially genotoxic particles (Drath *et al.*, 1976). Macrophages ingest most of the particles which pass into the bronchiolar and alveolar regions of the lungs. Thus, macrophages would be ideal target cells in an assay for assessment of cytotoxicity of particulates. Freshly isolated alveolar macrophages have been used as targets for this purpose. Metals (Castranova *et al.*, 1980), fly ash (Hill *et al.*, 1982), tars and other combustion products (Hill *et al.*, 1983), and volcanic dust (McLemore *et al.*, 1984) have been studied with a number of endpoints. However, as these are primary, nondividing cells, they cannot be used for study of many genotoxic endpoints. Mutation assays require cell division to allow fixation and expression of genotoxic lesions. Therefore, a continuous, dividing cell line, with the phagocytic and metabolic properties of a normal macrophage, would combine the desired characteristics for an *in vitro* assay to measure the genotoxic potential of particulates.

The purpose of the study presented in this report is the development and validation of an inexpensive and rapid *in vitro* test for determining the cytotoxicity and genotoxicity of particulates potentially hazardous to the respiratory system. The short term tests currently available were designed for the testing of soluble test materials and were adapted for the study of particulates with target cells possessing limited phagocytic ability. The assay system evaluated in this report used a mouse macrophage cell line, RAW 264.7, which is avidly phagocytic and will undergo a respiratory burst. This cell line grows readily in culture, a feature which will allow both cytotoxic and genotoxic endpoints to be monitored. Preliminary studies have been conducted to establish the basic parameters for both a cytotoxicity assay and an HGPRT gene based forward mutation assay. These studies showed that RAW 264.7 cells exhibited a dose dependent but differential cytotoxic response to high versus low fire beryllium oxide particles and dose dependent mutagenic responses to two soluble mutagens, ultraviolet radiation (UVC) and ethyl methanesulfonate. The goals for the present study are to complete the work on establishing optimal conditions for the assays, validate the cytotoxicity assay with known toxic particulates and the mutation assay with several known classes of mutagenic particulates.

MATERIALS AND METHODS

Test Materials

Low Fire Beryllium Oxide and High Fire Beryllium Oxide were obtained as a gift from Dr. John H. Harbell, Institute for *In Vitro* Sciences, Gaithersburg, Maryland. The following test materials were provided by the project officer: trinitrotoluene (TNT)-contaminated compost, 10% TNT-contaminated compost, uncontaminated soil sample, and red dye mix (a mixture of Solvent Red 1 and Disperse Red 11).

Cell Line and Culture Conditions

The RAW 264.7 macrophage cell line was derived from an Abelson leukemia virus infected BAB/14 mouse by Raschke and co-workers (1978). The cell line is avidly phagocytic towards both antibody coated and uncoated particles, has Fc and complement receptors, and produces lysozyme. These cells have been shown to actively phagocytose and kill bacteria, as well as undergo the "respiratory burst" and limited release of superoxide characteristic of alveolar macrophages (McGown *et al.*, 1985a). The symosan induced respiratory burst was measured after poisoning mitochondrial oxygen consumption with KCN and was found to consume between 2 and 3 nmoles per 10^6 cells per minute. Thus, this cell line combines the avid phagocytic activity and important metabolic properties of the normal macrophage with the ability to grow and divide in culture. The use of this functional macrophage cell line allows the study of genetic endpoints, as well as cytotoxicity, without the need to sacrifice animal donors for each assay. In addition, a number of physiologically relevant studies will also be possible with this assay system. The study of combined soluble mutagen and particulate, as might occur with smokes, would be greatly facilitated by this target cell which is sensitive to both. Similarly, the lung cells may be subject to the combined effects of the respiratory burst and toxic challenge and this system would allow the study of such a synergy.

Several pertinent preliminary studies on the RAW 264.7 cells and culture system have been completed. These include media and serum requirements, growth in suspension culture,

phagocytic activity, population doubling time, clonability, propensity to form colonies on plastic, and chromosome number.

The RAW 264.7 cell line was originally isolated in minimal essential medium (MEM) with 10% fetal bovine serum. It grows well in RPMI 1640 and Ham's F12 media without hypoxanthine. The cells attach to wettable surfaces (polystyrene or glass) but not to polypropylene or siliconized glass. Thus, they can be cultured either in attached or suspension culture. Phagocytic activity is greater and more uniform in suspension culture. After a 2-hour challenge, over 95% of the cells were observed to have ingested carbon or 2-micron latex particles (J. Harbell, personal communication). The population doubling time, in suspension, is between 18 and 20 hours, and the population continues to divide after ingesting particles. The cells can be cloned and they form discrete colonies on plastic, two characteristics essential to a clonal mutation assay system. The cell line is pseudodiploid (modal count of 40). An HGPRT^{-/-} subclone was produced for use in reconstruction experiments. Transmission electron micrographs of cell clusters grown in suspension or attached culture did not show gap junctions. Therefore, cross feeding between mutants and nonmutants during the 6-thioguanine mutant selection phase of the HGPRT assay would seem unlikely. For, example, cross feeding between mutant and nonmutant CHO cells limits the cell density to only 2×10^5 cells on each mutant selection plate.

The standard culture medium used in this study was RPMI 1640 supplemented with 10% fetal bovine serum. This culture medium was used for both the cytotoxicity and mutagenicity assay, and for both attached and suspension cultures. Unless specified otherwise, cells were exposed to the test materials in suspension culture. To remove uningested test material, if appropriate, the cells were allowed to attach to flasks and the monolayer rinsed to remove the free particles. Cells were removed from the plastic, for counting or passaging, by treatment with Ca⁺⁺ and Mg⁺⁺ free Hanks' balanced salt solution (HBSS). Ethylenediaminetetraacetic acid (EDTA; 0.02%) might also be added to facilitate cell release. The use of trypsin was avoided as it tended to activate the membrane of the RAW 264.7 cells.

Cytotoxicity Assay

Cytotoxicity can be manifested in many forms including loss of differentiated function, cessation of cell replication, and/or cell death. The RAW 264.7 cell line is appropriate for the study of all of these endpoints. McGown *et al.* (1985b) have shown that trivalent arsenicals can inhibit respiratory burst and bacterial cell killing (differentiated function), reduce cell replication, and kill cells outright depending on the dose. Preliminary studies demonstrated that changes in population number determined by cloning cells can be used for measuring cytotoxicity. Toxicity could also be determined by assessment of active uptake of particulates. The loss of differentiated function can be determined through measurement of changes in the respiratory burst induced by symosan particles.

For the cytotoxicity assay, 10 ml suspension cultures at 5×10^4 cells per ml were prepared in 15 ml polypropylene tubes. Serial dilutions of the test material were added to the appropriately labelled tubes. Untreated and carrier/solvent (if appropriate) control cultures were also prepared. The cells were exposed, in suspension culture on a roller drum, for 5 hours (unless specified otherwise) and then counted. For assessment of toxicity based on determination of uptake of **particulates**, an aliquot of the cells was withdrawn, and the cells were fixed with methanol,

stained with 10% Giemsa and the number of cells which ingested particles determined by examination under a microscope. The remainder of the cells was carried forward to determine cloning efficiency. In cases where separation of the uningested test material from the cells was desired, the cells would be plated into 25 cm² flasks and allowed to attach for 30 minutes. The monolayer was then rinsed with culture medium to remove the uningested test material. The cultures were fed with fresh medium and allowed to grow for 18 to 24 hours. Cells were then detached, counted and reseeded at approximately 3 cells per well in 96-well microtiter plates or at 100 cells per 60 mm petri plate. After 8 to 10 days, the colonies were fixed with methanol, stained with 10% Giemsa, counted and evaluated for cytotoxic effects. Data were presented as the relative cloning efficiency of the treated populations as compared to that in the untreated control population.

Mutagenicity Assay

The hypoxanthine guanine phosphoribosyl transferase (HGPRT) gene locus on the X chromosome has been used as a target for forward mutation studies because mammalian cells are naturally hemizygous at this locus. The RAW 264.7 cell line is hemizygous with a spontaneous frequency of between 0 and 10 mutants per 10⁶ cells (Hitchins *et al.*, 1992). The cells will form colonies, do not appear to form gap junctions, and are not inhibited by rat liver S9 activation. These characteristics are necessary characteristics for a target cell line in a forward mutation assay.

In preliminary experiments, the HGPRT^{-/-} subclone was developed and used in reconstruction experiments to determine parameters for the assay using the same approach as that used by Hsie *et al.* (1974) in the development of the HGPRT assay in CHO cells. These experiments allowed the determination of the optimal mutant expression period required to achieve maximum mutant frequency yields, the mutant selection period, the amount of selective agent (6-thioguanine) required, and the maximum cell plating density during the mutant selection phase. Two mutagens, ultraviolet radiation (UVC; Hitchins *et al.*, 1992) and ethyl methanesulfonate (EMS; Bigger, unpublished data), were tested in the assay and dose dependent increases comparable to those achieved in the CHO/HGPRT assay were achieved in the RAW 264.7 cell assay.

For the mutagenicity assay, cells were treated with test article and a portion of the treated cells were used to determine cytotoxicity as described above for the cytotoxicity assay. For mutagenic evaluation, the remainder of the cells were returned to culture for five to seven days to allow expression of the mutation at the HGPRT locus. At the time of selection, cells were seeded at 2 x 10⁵ cells per 60 mm petri dish (or, at 5 x 10³ cells/well in a 96-well microtiter plate) in the presence of 4 x 10⁻⁵M 6-thioguanine; for cloning efficiency, cells were seeded at 100 cells per 60 mm petri dish (or, at 3 cells/well in a 96-well microtiter plate).

For seeding in 60 mm petri dishes, the computation of cloning efficiency and mutant frequency was as follows:

$$\text{Cloning efficiency (\%)} = \frac{\text{Total colonies counted}}{\text{Dishes counted} \times 100 \text{ cells}} \times 100$$

$$\text{Mutants}/10^6 \text{ clonable cells} = \frac{\text{Total mutant colonies} \times 10^6}{\text{Number selection dishes} \times \text{cloning efficiency} \times 2 \times 10^5 \text{ cells}}$$

For seeding in 96-well microtiter plates, the computation of cloning efficiency and mutant frequency was as follows:

$$\text{Cloning efficiency} = \frac{-\text{Natural log (number of empty wells/total number of wells)} \times 100}{\text{Number of cells seeded per well}}$$

$$\text{Mutants}/10^6 \text{ clonable cells} = \frac{\text{Cloning efficiency on selection plates}}{\text{Cloning efficiency on non-selection plates}}$$

RESULTS

Low and High Fire Berillium Oxide (BeO). Preliminary studies were conducted with low and high fire BeO particles (Bigger, unpublished data). RAW 264.7 cells and Chinese hamster ovary (CHO) cells were compared with respect to their susceptibility to the toxic effects of these particles. Cells from both of these lines showed a dose dependent cytotoxic response to the low fire particles but the response of the RAW 264.7 cells (1% relative cytotoxicity at 50 $\mu\text{g}/\text{mL}$) showed significantly more sensitivity than that of the CHO cells (60%). CHO cells showed no cytotoxic response to the high fire particles while the RAW 264.7 cells showed a high response (24%). The differential response to the two types of particles is the result of the inherent physical characteristics of these two particles. The processing temperatures used to produce these metal oxides yield significantly different particle structures. The high fire particles tend to have a much smoother surface and therefore less surface area than the more porous low fire particles.

Treatment of RAW 264.7 cells with low fire BeO did not result in a dose-related increase in mutant frequency (i.e., mutants per 10^6 clonable cells) over a dose range of 5 to 200 $\mu\text{g}/\text{mL}$, regardless of whether mutant selection was conducted at 5 days or 7 days post-treatment (Table 1). The increase in mutant frequency at 25 $\mu\text{g}/\text{mL}$ (day 7 selection) was less than 40 mutants per 10^6 clonable cells above that in the untreated control and was not considered to have any biological significance. It was noted that the mutant frequency in the untreated controls (for both day 5 and day 7 selections) was substantially higher than the spontaneous frequency of between 0 and 10 mutants per 10^6 cells reported by Hitchins *et al.* (1992).

Treatment of RAW 264.7 cells with high fire BeO did not result in a dose-related increase in mutant frequency over a dose range of 100 to 500 $\mu\text{g}/\text{mL}$ (mutant selection conducted at 5 days post-treatment, although an increase in mutant frequency (at 100 $\mu\text{g}/\text{mL}$) of 51 mutants per 10^6 clonable cells above that in the untreated control was noted (Table 2). The mutant frequency in the untreated controls was substantially higher than the spontaneous frequency of between 0 and 10 mutants per 10^6 cells reported by Hitchins *et al.* (1992).

The preceding experiments were conducted with treatment of cells in suspension and passage of the cells as suspension cultures until the time of plating for mutant selection. Since the cell generation time was about 18 hours for attached cultures and about 24 hours for suspension cultures, the shorter generation time of attached cultures may decrease the time needed for fixation and expression of mutations. On the other hand, the use of suspension cultures may help to minimize the difficulty of cell removal from flasks with attached cultures. In light of these considerations, an experiment was conducted to compare the effect of passaging cells as

suspension cultures and that of passaging cells as attached cultures following treatment. Low fire and high fire BeO, along with EMS (as positive control) and untreated and solvent controls, were included in this comparative study. As presented in Table 3 (Trial 1) and Table 4 (Trial 2), there was essentially no difference in the results between the use of suspension cultures and attached cultures following treatment in the yield of mutants.

Ethyl Methanesulfonate (EMS). For use as a positive control in the RAW 264.7 mutagenicity assay system, EMS was evaluated in two independent experiments. A dose-related increase in mutant frequency was observed in both experiments (Table 5).

Comparison of RAW 264.7 from Two Different Stock Cultures. To ensure that the stock culture in this laboratory (Stock Culture 2) was functioning in a manner similar to that of cells from another laboratory (Stock Culture 1), the cells from both stock cultures were compared with respect to their response to the cytotoxic and mutagenic effects of EMS and low fire BeO. The mutagen EMS elicited a dose-related increase in toxicity (Table 6) and mutagenicity (Table 7) in the cells from both stock cultures, while low fire BeO was essentially non-toxic (Table 6) and non-mutagenic (Table 7) to cells from both stock cultures.

Trinitrotoluene (TNT)-Contaminated Compost and Uncontaminated Soil Sample. In the initial mutagenicity assay using a 5-hour treatment time, essentially no cytotoxicity and no mutagenic activity were elicited by TNT-contaminated compost, 10% TNT-contaminated compost and uncontaminated soil sample (Table 8, Trial 1). Similar results were obtained with TNT-contaminated compost and uncontaminated soil sample in an independent experiment (Table 8, Trial 2). Using an 18-hour extended treatment time, no mutagenic activity was observed with TNT-contaminated compost at concentrations ranging from 1 to 100 $\mu\text{g/mL}$ and uncontaminated soil sample at concentrations of 1 and 10 $\mu\text{g/mL}$ (Table 9, Trial 1). Upon retesting at higher concentrations (10 to 500 $\mu\text{g/mL}$), a dose-related increase in mutant frequency was observed with 10% TNT-contaminated compost at concentrations of 100 and 500 $\mu\text{g/mL}$ (Table 9, Trial 2). No mutagenic activity was observed with uncontaminated soil sample in the retest at concentrations of 1 and 10 $\mu\text{g/mL}$ (Table 9, Trial 2). However, the mutant frequency in the untreated controls in this experiment was substantially higher than the spontaneous frequency of between 0 and 10 mutants per 10^6 cells reported by Hitchins *et al.* (1992).

Red Dye Mix. No mutagenic activity was elicited by red dye mix at concentrations ranging from 5 to 100 $\mu\text{g/mL}$ (Table 10).

CONCLUSIONS

The inability to demonstrate mutagenicity with low fire BeO or high fire BeO indicates that the assay procedure still requires further refinement. The mutant frequency in the untreated control in some experiments was substantially higher than the spontaneous frequency of between 0 and 10 mutants per 10^6 cells reported by Hitchins *et al.* (1992). It remains to be determined as to whether the variability in spontaneous mutant frequency is an inherent feature of this cell line, as in the case of Chinese hamster ovary cells in the CHO/HGPRT assay system. No mutagenic activity was demonstrated with red dye mix. A dose-related increase in mutant frequency was observed with 10% TNT-contaminated compost in only one experiment. Since the TNT-contaminated compost was not tested at concentrations higher than 100 $\mu\text{g/mL}$ using the 18-hour

extended treatment regimen, additional experiments are warranted prior to making comments on the potential mutagenicity of soils contaminated with munitions.

REFERENCES

- Amdur, M.O. (1986) Air Pollutants. In: Toxicology, The Basic Science of Poisons, Klaasen, C.D., Amdur M.O. and Doull, J. (eds.), MacMillan Publishing Co., New York NY, pp. 801-824.
- Brusick, D. (1982) Genetic toxicology. In: Principles and Methods of Toxicology, Hayes, A.W. (ed.), Raven Press, New York.
- Castranova, V., Bowman, L., Miles, P.R. and Reasor, M.J. (1980) Toxicity of metal ions to alveolar macrophages. *Am. J. Int. Med.* 1:349-357.
- Costa, M. and Heck, J.D. (1982) Specific nickel compounds as carcinogens. *Trends in Pharmacological Sciences* 3:408-410.
- Drath, D.B., Harper, A.A. and Karnoysky, M.L. (1976) Biochemical characteristics of alveolar macrophages. In: Lung Cells in Disease, Nouhuys, A. (ed.), North-Holland Publishing Co., New York, NY, pp. 153-161.
- Hesterberg, T.W., Tsutsui, T. and Barrett, J.C. (1983) Neoplastic transformation of Syrian hamster embryo (SHE) cells by asbestos and fiberglass: The importance of fiber dimension. *Proc. Amer. Assoc. Cancer Res.* 24:96.
- Hill, J.O., Gray, R.H., DeNee, P.B. and Newton, G.J. (1982) Comparative damage to alveolar macrophages after phagocytosis of respirable particles. *Environmental Research* 27:95-109.
- Hill, J.O., Royer, E., Mitchell, C.E. and Durcher, J.S. (1983) *In vitro* cytotoxicity to alveolar macrophages of tar from a low btu coal gasifier. *Environmental Research* 31:484-492.
- Hitchins, V.M., Lytle, C.D., Withrow, T.J., Thomas, D.P., Sigler, C.I., Stahl, R.W. and Harbell, J.W. (1992) Response of RAW 264.7 murine macrophage cells to UVC radiation. *In Vitro Toxicology* 5:39-49.
- Hsie, A.W., Brimer, P.A., Mitchell, T.J. and Gosslee, D.G. (1974) The dose-response relationship for ethyl methanesulfonate-induced mutations at the hypoxanthine-guanine phosphoribosyl transferase locus in Chinese hamster ovary cells. *Somatic Cell Genet.* 1:247-261.
- McGown, E.L., Orencole, S.F., Rusnak, M.G., Hillman, J.R. and Schwartz, B.D. (1985a) Bactericidal capability and respiratory burst characteristics of the murine macrophage cell line RAW 264.7. Letterman Army Institute of Research, Institute Report o. 195.
- McGown, E.L., Harbell, J.W., Cumlao, C.R. and O'Connor, R.J. (1985b) Use of *in vitro* cell systems to study cytotoxic effects of vesicant arsenicals and efficacy of antidotes. *Proceedings of the Fifth Annual Chemical Defense Bioscience Review:* 292-301.

McLemore, T.L., Mauldin, J.E., Marshall, M.V., Teague, R., Ford, G., Busbee, D.L., Wray, N.P., Griffin, A.C. and Greenberg, S.D. (1984) Biological effects of Mount Saint Helens volcanic ash on cultured human alveolar macrophages. *J. Amer. Coll. of Toxicol.* 3:25-35.

Menzel, D.B. and Amdur, M.O. (1986) Toxic responses of the respiratory system. In: *Toxicology, The Basic Science of Poisons*, Klaasen, C.D., Amdur, M.O. and Doull, J. (eds.), MacMillan Publishing Co., New York NY, pp. 330-358.

Raschke, W.C., Baird, S., Ralph, P. and Nakoinz, I. (1978) Functional macrophage cell lines transformed by Abelson leukemia virus. *Cell* 15:261-267.

Rossmann, T.G., Zelikoff, J.T., Agarwal, S. and Kneip, T.J. (1987) Genetic toxicology of metal compounds: An examination of appropriate cellular models. *Toxicological and Environmental Chemistry* 14:251-262.

Sincock, A.M., Delhanty, J.D.A. and Casey, G. (1982) A comparison of the cytogenetic response to asbestos and glass fibre in Chinese hamster and human cell lines: Demonstration of growth inhibition in primary human fibroblasts. *Mutation Res.* 101:257-268.

Zelikoff, J.T., Atkins, Jr., N.M., Rossmann, T.G. and Daisey, J.M. (1985) Cytotoxicity of fine particles with and without adsorbed polycyclic aromatic hydrocarbons using Chinese hamster lung cells. *Environmental International* 11:331-339.

Table 1
 Mutagenicity Assay of Low Fire Beryllium Oxide
 Using RAW 264.7 Macrophage Cells

| Treatment | Colonies per Dish | | Total # Colonies | Cloning Efficiency (%) | Mutant Colonies per Selection Dish | | | | | Total Mutant Colonies | Mutants per 10 ⁶ Clonable Cells | |
|--------------------------------|-------------------|-----|------------------|------------------------|------------------------------------|-----|-----|-----|-----|-----------------------|--|-------|
| | | | | | 1 | 2 | 3 | 4 | 5 | | | |
| <u>Day 5 Selection</u> | | | | | | | | | | | | |
| Untreated | 99 | 112 | 122 | 333 | 111 | 59 | 42 | 44 | 49 | 64 | 258 | 93.0 |
| Low Fire Berrilium Oxide | | | | | | | | | | | | |
| 5 µg/ml | 208 | 118 | 102 | 428 | 143 | 58 | 57 | 51 | 86 | 54 | 306 | 85.8 |
| 10 µg/ml | 104 | 216 | 139 | 459 | 153 | 50 | 49 | 54 | 55 | 42 | 250 | 65.4 |
| 25 µg/ml | 106 | 105 | 158 | 369 | 123 | 48 | 65 | 63 | 64 | 65 | 305 | 99.2 |
| 50 µg/ml | 101 | 126 | 228 | 455 | 152 | 39 | 47 | 47 | 33 | 53 | 219 | 57.8 |
| 100 µg/ml | 108 | 83 | 106 | 297 | 99 | 47 | 38 | 39 | 49 | 42 | 215 | 86.9 |
| 200 µg/ml | 106 | 110 | 121 | 337 | 112 | 18 | 22 | 27 | 27 | 20 | 114 | 40.6 |
| EMS (positive control) | | | | | | | | | | | | |
| 0.6 µl/ml | 116 | 182 | 95 | 393 | 131 | 185 | 141 | 184 | 136 | 128 | 774 | 236.3 |
| DMSO (solvent control for EMS) | | | | | | | | | | | | |
| 10 µl/ml | 122 | 116 | 124 | 362 | 121 | 71 | 48 | 53 | 53 | 57 | 282 | 93.5 |
| <u>Day 7 Selection</u> | | | | | | | | | | | | |
| Untreated | 148 | 134 | 168 | 450 | 150 | 45 | 51 | 49 | 44 | 60 | 249 | 66.4 |
| Low Fire Berrilium Oxide | | | | | | | | | | | | |
| 5 µg/ml | 135 | 174 | 159 | 468 | 156 | 110 | 94 | 85 | 99 | 98 | 486 | 124.6 |
| 10 µg/ml (not plated) | | | | | | | | | | | | |
| 25 µg/ml | 145 | 145 | 162 | 452 | 151 | 68 | 84 | 81 | 85 | 77 | 395 | 104.9 |
| 50 µg/ml | 166 | 126 | 139 | 431 | 144 | 66 | 63 | 63 | 65 | 51 | 308 | 85.8 |
| 100 µg/ml | 150 | 120 | 138 | 408 | 136 | 34 | 36 | 22 | 45 | 28 | 165 | 48.5 |
| 200 µg/ml | 170 | 167 | 151 | 488 | 163 | 27 | 38 | 43 | 42 | 30 | 180 | 44.3 |
| EMS (positive control) | | | | | | | | | | | | |
| 0.6 µl/ml | 118 | 128 | 136 | 382 | 127 | 202 | 173 | 189 | 206 | 219 | 989 | 310.7 |
| DMSO (solvent control for EMS) | | | | | | | | | | | | |
| 10 µl/ml | 127 | 150 | 142 | 419 | 140 | 58 | 61 | 69 | 74 | 53 | 315 | 90.2 |

Cloning efficiency = cloning efficiency at time of plating for mutant selection.
 Untreated Control = Hanks' balanced salt solution (HBSS); EMS = ethyl methanesulfonate.

Table 2
 Mutagenicity Assay of High Fire Beryllium Oxide
 Using RAW 264.7 Macrophage Cells

| Treatment | Colonies per Dish | | Total # Colonies | Cloning Efficiency (%) | Mutant Colonies per Selection Dish | | | | | Total Mutant Colonies | Mutants per 10 ⁶ Clonable Cells | |
|--------------------------------|-------------------|-----|------------------|------------------------|------------------------------------|----|----|----|----|-----------------------|--|-------|
| | | | | | 1 | 2 | 3 | 4 | 5 | | | |
| <u>Day 5 Selection</u> | | | | | | | | | | | | |
| Untreated | 91 | 153 | 133 | 377 | 126 | 50 | 54 | 36 | 42 | 52 | 234 | 74.5 |
| High Fire Berrilium Oxide | | | | | | | | | | | | |
| 100 µg/ml | 68 | 87 | 69 | 224 | 75 | 41 | 55 | 42 | 46 | 51 | 235 | 125.9 |
| 200 µg/ml | 88 | 94 | 77 | 259 | 86 | 17 | 17 | 19 | 22 | 18 | 93 | 43.1 |
| 300 µg/ml | 83 | 94 | 69 | 246 | 82 | 38 | 42 | 36 | 34 | 30 | 180 | 87.8 |
| 400 µg/ml | 72 | 73 | 84 | 229 | 76 | 13 | 7 | 7 | 12 | 11 | 50 | 26.2 |
| 500 µg/ml | 47 | 69 | 76 | 192 | 64 | 7 | 22 | 11 | 10 | ND | 50 | 39.1 |
| EMS (positive control) | | | | | | | | | | | | |
| 0.6 µl/ml | 52 | 49 | 54 | 155 | 52 | 64 | 52 | 43 | 43 | 66 | 268 | 207.5 |
| DMSO (solvent control for EMS) | | | | | | | | | | | | |
| 10 µl/ml | 71 | 90 | 78 | 239 | 80 | 46 | 39 | 41 | 39 | 46 | 211 | 105.9 |

Cloning efficiency = cloning efficiency at time of plating for mutant selection.
 Untreated Control = Hanks' balanced salt solution (HBSS); EMS = ethyl methanesulfonate;
 ND = no data.

Table 3

Mutagenicity Assay of High Fire and Low Fire Beryllium Oxide
Using RAW 264.7 Macrophage Cells in 96-Microwell Plates (Trial 1)

| Treatment | Viability Assessment Plates | | | Mutant Selection Plates | | | |
|--|-----------------------------|-----------------------|------------------------|-------------------------|-----------------------|------------------------|--|
| | Empty Wells Counted | Total # Wells Counted | Cloning Efficiency (%) | Empty Wells Counted | Total # Wells Counted | Cloning Efficiency (%) | Mutants per 10 ⁶ Clonable Cells |
| <u>Using Cells Passaged as Suspension Cultures</u> | | | | | | | |
| Untreated | 19 | 96 | 54.0 | 184 | 192 | 0.0009 | 15.8 |
| High Fire Berrilium Oxide | | | | | | | |
| 5 µg/ml | 18 | 96 | 55.8 | 168 | 192 | 0.0027 | 47.9 |
| 50 µg/ml | 21 | 96 | 50.7 | 173 | 192 | 0.0021 | 41.1 |
| Low Fire Berrilium Oxide | | | | | | | |
| 5 µg/ml | 21 | 96 | 50.7 | 186 | 192 | 0.0006 | 12.5 |
| 50 µg/ml | 31 | 96 | 37.7 | 174 | 192 | 0.0020 | 52.3 |
| EMS (positive control) | | | | | | | |
| 0.3 µl/ml | 20 | 96 | 52.3 | 155 | 192 | 0.0043 | 81.9 |
| 1.0 µl/ml | 48 | 96 | 23.1 | 129 | 192 | 0.0080 | 344.2 |
| DMSO (solvent control for EMS) | | | | | | | |
| 10 µl/ml | 19 | 96 | 54.0 | 172 | 192 | 0.0022 | 40.7 |
| <u>Using Cells Passaged as Attached Cultures</u> | | | | | | | |
| Untreated | 49 | 192 | 45.5 | 373 | 384 | 0.0006 | 12.8 |
| High Fire Berrilium Oxide | | | | | | | |
| 5 µg/ml | 28 | 192 | 64.2 | 355 | 384 | 0.0016 | 24.5 |
| 50 µg/ml | 36 | 192 | 55.8 | 351 | 384 | 0.0018 | 32.2 |
| Low Fire Berrilium Oxide | | | | | | | |
| 5 µg/ml | 36 | 192 | 55.8 | 163 | 192 | 0.0033 | 58.7 |
| 50 µg/ml | 48 | 192 | 46.2 | 372 | 384 | 0.0006 | 13.7 |
| EMS (positive control) | | | | | | | |
| 0.3 µl/ml | 23 | 192 | 70.7 | 304 | 384 | 0.0047 | 66.1 |
| 1.0 µl/ml | 48 | 192 | 23.1 | 149 | 192 | 0.0051 | 219.5 |
| DMSO (solvent control for EMS) | | | | | | | |
| 10 µl/ml | 46 | 192 | 47.6 | 354 | 384 | 0.0016 | 34.2 |

Cloning efficiency = cloning efficiency at time of plating for mutant selection.
Untreated Control = Hanks' balanced salt solution (HBSS); EMS = ethyl methanesulfonate;
- = not tested; ND = no data.

Table 4
 Mutagenicity Assay of High Fire and Low Fire Beryllium Oxide
 Using RAW 264.7 Macrophage Cells in 96-Microwell Plates (Trial 2)

| Treatment | <u>Viability Assessment Plates</u> | | | <u>Mutant Selection Plates</u> | | | |
|--|------------------------------------|-----------------------|------------------------|--------------------------------|-----------------------|------------------------|--|
| | Empty Wells Counted | Total # Wells Counted | Cloning Efficiency (%) | Empty Wells Counted | Total # Wells Counted | Cloning Efficiency (%) | Mutants per 10 ⁶ Clonable Cells |
| <u>Using Cells Passaged as Suspension Cultures</u> | | | | | | | |
| Untreated | 4 | 96 | 105.9 | 165 | 192 | 0.0030 | 28.6 |
| High Fire Berrilium Oxide | | | | | | | |
| 5 µg/ml | 1 | 96 | 152.1 | 150 | 192 | 0.0049 | 32.5 |
| 20 µg/ml | 4 | 96 | 105.9 | 142 | 192 | 0.0060 | 57.0 |
| 50 µg/ml | 1 | 96 | 152.1 | 165 | 192 | 0.0030 | 19.9 |
| Low Fire Berrilium Oxide | | | | | | | |
| 5 µg/ml | 4 | 96 | 105.9 | 98 | 192 | 0.0135 | 127.0 |
| 20 µg/ml | 10 | 96 | 75.4 | 149 | 191 | 0.0050 | 65.9 |
| 50 µg/ml | 3 | 96 | 115.5 | 144 | 192 | 0.0058 | 49.8 |
| EMS (positive control) | | | | | | | |
| 0.3 µl/ml | 12 | 96 | 69.3 | 129 | 192 | 0.0080 | 114.8 |
| 1.0 µl/ml | 9 | 96 | 78.9 | 4 | 192 | 0.0774 | 981.2 |
| DMSO (solvent control for EMS) | | | | | | | |
| 10 µl/ml | 6 | 96 | 92.4 | 156 | 192 | 0.0042 | 44.9 |
| <u>Using Cells Passaged as Attached Cultures</u> | | | | | | | |
| Untreated | 8 | 96 | 82.8 | 188 | 192 | 0.0004 | 5.1 |
| High Fire Berrilium Oxide | | | | | | | |
| 5 µg/ml | 15 | 96 | 61.9 | 175 | 186 | 0.0012 | 19.7 |
| 20 µg/ml | 15 | 96 | 61.9 | 150 | 188 | 0.0045 | 73.0 |
| 50 µg/ml | 11 | 96 | 72.2 | 132 | 146 | 0.0020 | 27.9 |
| Low Fire Berrilium Oxide | | | | | | | |
| 5 µg/ml | 11 | 96 | 72.2 | 189 | 192 | 0.0003 | 4.4 |
| 20 µg/ml | 11 | 96 | 72.2 | 168 | 173 | 0.0006 | 8.1 |
| 50 µg/ml | 24 | 96 | 46.2 | 155 | 161 | 0.0008 | 16.4 |
| EMS (positive control) | | | | | | | |
| 0.3 µl/ml | 12 | 96 | 69.3 | 147 | 174 | 0.0034 | 48.7 |
| DMSO (solvent control for EMS) | | | | | | | |
| 10 µl/ml | 9 | 95 | 78.6 | 165 | 192 | 0.0030 | 38.6 |

Cloning efficiency = cloning efficiency at time of plating for mutant selection.
 Untreated Control = Hanks' balanced salt solution (HBSS); EMS = ethyl methanesulfonate;
 - = not tested; ND = no data.

Table 5
 Mutagenicity Assay of Ethyl Methanesulfonate (EMS)
 Using RAW 264.7 Macrophage Cells

| Treatment | Colonies per Dish | | Total # Colonies | Cloning Efficiency (%) | Mutant Colonies per Selection Dish | | | | | Total Mutant Colonies | Mutants per 10 ⁵ Clonable Cells | |
|---------------------------------|-------------------|-----|------------------|------------------------|------------------------------------|----|----|----|----|-----------------------|--|-------|
| | | | | | 1 | 2 | 3 | 4 | 5 | | | |
| <u>Trial 1: Day 7 Selection</u> | | | | | | | | | | | | |
| Untreated | 134 | 99 | 98 | 331 | 110 | 16 | 12 | 17 | 10 | 16 | 71 | 64.4 |
| EMS | | | | | | | | | | | | |
| 0.2 µl/ml | 112 | ND | 100 | 212 | 71 | 10 | 10 | 8 | 12 | 6 | 46 | 65.1 |
| 0.4 µl/ml | 102 | 95 | 112 | 309 | 103 | 9 | 24 | 17 | 6 | 13 | 69 | 67.0 |
| 0.6 µl/ml | 71 | 72 | 81 | 224 | 75 | 37 | 37 | 55 | 47 | 47 | 223 | 298.7 |
| 0.8 µl/ml | 70 | 101 | 72 | 243 | 81 | 16 | 11 | 34 | 26 | 21 | 108 | 133.3 |
| 1.0 µl/ml | 91 | 81 | 78 | 250 | 83 | 13 | 11 | 17 | 9 | 16 | 66 | 79.2 |
| DMSO (solvent control) | | | | | | | | | | | | |
| 10 µl/ml | 100 | 126 | 98 | 324 | 108 | 6 | 5 | 4 | 0 | 8 | 23 | 21.3 |
| <u>Trial 6: Day 6 Selection</u> | | | | | | | | | | | | |
| Untreated | 81 | 71 | 96 | 248 | 83 | 16 | 5 | 6 | 7 | 7 | 41 | 49.6 |
| EMS | | | | | | | | | | | | |
| 0.2 µl/ml | 71 | 92 | 69 | 232 | 77 | 7 | 4 | 5 | 12 | 5 | 33 | 42.7 |
| 0.4 µl/ml | 50 | 70 | 52 | 172 | 57 | 11 | 11 | 6 | 11 | 13 | 52 | 90.7 |
| 0.6 µl/ml | 72 | 73 | 63 | 208 | 69 | 19 | 20 | 16 | 28 | 28 | 111 | 160.1 |
| 0.8 µl/ml | 58 | 71 | 71 | 200 | 67 | 15 | 18 | 14 | 17 | 20 | 84 | 126.0 |
| 1.0 µl/ml | 60 | 79 | 60 | 199 | 66 | 25 | 22 | 20 | 22 | 22 | 111 | 167.3 |
| DMSO (solvent control) | | | | | | | | | | | | |
| 10 µl/ml | 58 | 64 | 50 | 172 | 57 | 6 | 5 | 5 | 8 | 9 | 33 | 57.6 |

Cloning efficiency = cloning efficiency at time of plating for mutant selection.
 Untreated Control = Hanks' balanced salt solution (HBSS).

Table 6

Cytotoxicity of Ethyl Methanesulfonate (EMS) and Low Fire Berrilium Oxide
in RAW 264.7 Macrophage Cells from Two Different Stock Cultures

| | Empty Wells Counted per Dish | | Cloning Efficiency (%) | Relative Cloning Efficiency (%) |
|--------------------------|------------------------------------|----|------------------------------|--|
| <u>Stock Culture 1</u> | | | | |
| Untreated | 37 | 30 | 35.1 | 122 |
| EMS | | | | |
| 0.3 μ l/ml | 35 | 20 | 41.7 | 145 |
| 0.6 μ l/ml | 29 | 35 | 36.6 | 127 |
| 1.0 μ l/ml | 66 | 53 | 16.0 | 55 |
| Low Fire Berillium Oxide | | | | |
| 5 μ g/ml | 25 | 33 | 39.0 | 139 |
| 20 μ g/ml | 30 | 30 | 38.8 | 135 |
| 50 μ g/ml | 32 | 34 | 35.6 | 124 |
| DMSO (solvent control) | | | | |
| 10 μ l/ml | 41 | 40 | 28.8 | 100 |
| <u>Stock Culture 2</u> | | | | |
| Untreated | 31 | 33 | 36.6 | 96 |
| EMS | | | | |
| 0.3 μ l/ml | 33 | 26 | 39.3 | 103 |
| 0.6 μ l/ml | 43 | 47 | 25.3 | 66 |
| 1.0 μ l/ml | 78 | 75 | 7.6 | 20 |
| Low Fire Berillium Oxide | | | | |
| 5 μ g/ml | 38 | 34 | 32.7 | 86 |
| 20 μ g/ml | 25 | 20 | 48.4 | 127 |
| 50 μ g/ml | 28 | 29 | 40.5 | 106 |
| DMSO (solvent control) | | | | |
| 10 μ l/ml | 36 | 25 | 38.2 | 100 |

Cloning efficiency based on total of 192 wells counted.
Untreated Control = Hanks' balanced salt solution (HBSS).

Table 7

Mutagenicity Assay of Ethyl Methanesulfonate (EMS) and Low Fire Berrilium Oxide
in RAW 264.7 Macrophage Cells from Two Different Stock Cultures

| Treatment | <u>Viability Assessment Plates</u> | | | <u>Mutant Selection Plates</u> | | | |
|--------------------------|------------------------------------|-----------------------------|------------------------------|--------------------------------|-----------------------------|------------------------------|---|
| | Empty Wells Counted | Total # Wells Counted | Cloning Efficiency (%) | Empty Wells Counted | Total # Wells Counted | Cloning Efficiency (%) | Mutants per 10 ⁶ Clonable Cells |
| <u>Stock Culture 1</u> | | | | | | | |
| Untreated | 41 | 192 | 51.5 | 166 | 192 | 0.0029 | 56.6 |
| EMS | | | | | | | |
| 0.3 μ l/ml | 30 | 192 | 61.9 | 186 | 192 | 0.0006 | 10.3 |
| 0.6 μ l/ml | 26 | 192 | 66.7 | 165 | 192 | 0.0030 | 45.5 |
| 1.0 μ l/ml | 60 | 132 | 38.8 | 120 | 176 | 0.0077 | 197.6 |
| Low Fire Berillium Oxide | | | | | | | |
| 5 μ g/ml | 34 | 158 | 57.7 | 174 | 192 | 0.0020 | 34.1 |
| 20 μ g/ml | 45 | 147 | 48.4 | 192 | 192 | 0.0000 | 0 |
| 50 μ g/ml | 85 | 192 | 27.2 | 168 | 192 | 0.0027 | 98.3 |
| DMSO (solvent control) | | | | | | | |
| 10 μ l/ml | 58 | 192 | 39.9 | 185 | 192 | 0.0007 | 18.6 |
| <u>Stock Culture 2</u> | | | | | | | |
| Untreated | 30 | 192 | 61.9 | 161 | 192 | 0.0035 | 56.9 |
| EMS | | | | | | | |
| 0.3 μ l/ml | 98 | 192 | 22.4 | 164 | 168 | 0.0005 | 21.5 |
| 0.6 μ l/ml | 58 | 192 | 39.9 | 166 | 192 | 0.0029 | 72.9 |
| 1.0 μ l/ml | 18 | 192 | 78.9 | 50 | 192 | 0.0269 | 341.0 |
| Low Fire Berillium Oxide | | | | | | | |
| 5 μ g/ml | 30 | 192 | 61.9 | 182 | 192 | 0.0011 | 17.3 |
| 20 μ g/ml | 35 | 192 | 56.7 | 169 | 192 | 0.0026 | 45.0 |
| 50 μ g/ml | 41 | 192 | 51.5 | 173 | 192 | 0.0021 | 40.5 |
| DMSO (solvent control) | | | | | | | |
| 10 μ l/ml | 36 | 192 | 55.8 | 163 | 184 | 0.0024 | 43.4 |

Cloning efficiency = cloning efficiency at time of plating for mutant selection.
Untreated Control = Hanks' balanced salt solution (HBSS).

Table 8
Mutagenicity Assay of Trinitrotoluene (TNT)-Contaminated Compost
and Uncontaminated Soil Sample Using RAW 264.7 Macrophage Cells

| Treatment | Colonies | | Total # Colonies | Cloning Efficiency (%) | Mutant Colonies per Selection Dish | | | | | Total Mutant Colonies | Mutants per 10 ⁶ Clonable Cells | |
|-----------------------------------|----------|-----|---------------------|------------------------------|--|----|----|----|----|-----------------------------|---|-------|
| | per Dish | | | | 1 | 2 | 3 | 4 | 5 | | | |
| <u>5-Hour Treatment (Trial 1)</u> | | | | | | | | | | | | |
| Untreated | 137 | 136 | 179 | 452 | 151 | 16 | 11 | 18 | 11 | 12 | 68 | 18.1 |
| TNT-Contaminated Compost | | | | | | | | | | | | |
| 10 µg/ml | 157 | 93 | 207 | 457 | 152 | 19 | 20 | 6 | 17 | ND | 62 | 20.4 |
| 50 µg/ml | 169 | 151 | 153 | 483 | 161 | 15 | 10 | 7 | 10 | 13 | 55 | 13.7 |
| 100 µg/ml | 146 | 148 | ND | 294 | 147 | 14 | 21 | 17 | 14 | 18 | 84 | 22.9 |
| 500 µg/ml | 98 | 83 | 107 | 288 | 96 | 5 | 15 | 7 | 19 | 9 | 55 | 22.9 |
| 10% TNT-Contaminated Compost | | | | | | | | | | | | |
| 10 µg/ml | 151 | 157 | 131 | 439 | 146 | 5 | 10 | 11 | 8 | 10 | 44 | 12.0 |
| Uncontaminated Soil Sample | | | | | | | | | | | | |
| 10 µg/ml | 141 | 111 | 162 | 414 | 138 | 5 | 7 | 7 | 4 | 7 | 30 | 8.7 |
| 50 µg/ml | 82 | 110 | 102 | 294 | 98 | 32 | 29 | 33 | 26 | 22 | 142 | 58.0 |
| 100 µg/ml | 105 | 105 | 106 | 316 | 105 | 7 | 10 | 10 | 16 | ND | 43 | 20.4 |
| EMS (positive control) | | | | | | | | | | | | |
| 0.6 µl/ml | 113 | 131 | 121 | 365 | 122 | 78 | 59 | 94 | 63 | 70 | 364 | 119.7 |
| DMSO (solvent control) | | | | | | | | | | | | |
| 10 µl/ml | 188 | 195 | 200 | 583 | 194 | 15 | 28 | 22 | 28 | 38 | 131 | 27.0 |
| <u>5-Hour Treatment (Trial 2)</u> | | | | | | | | | | | | |
| TNT-Contaminated Compost | | | | | | | | | | | | |
| 10 µg/ml | 60 | 63 | 83 | 206 | 69 | 3 | 6 | 2 | 1 | 2 | 14 | 8.2 |
| 50 µg/ml | 39 | 84 | 98 | 221 | 74 | 10 | 7 | 9 | 7 | 4 | 37 | 20.1 |
| 100 µg/ml | 101 | 104 | 113 | 318 | 106 | 8 | 6 | 6 | 5 | 10 | 35 | 13.2 |
| 500 µg/ml | 71 | 81 | 97 | 249 | 83 | 2 | 7 | 5 | 9 | 3 | 26 | 12.5 |
| Uncontaminated Soil Sample | | | | | | | | | | | | |
| 10 µg/ml | 83 | 70 | 64 | 217 | 72 | 15 | 17 | 20 | 12 | 18 | 82 | 45.3 |
| EMS (positive control) | | | | | | | | | | | | |
| 0.6 µl/ml | 40 | 57 | 39 | 136 | 45 | 22 | 28 | 35 | 23 | 20 | 128 | 112.9 |
| DMSO (solvent control) | | | | | | | | | | | | |
| 10 µl/ml | 82 | 94 | 82 | 258 | 86 | 4 | 2 | 1 | 1 | 3 | 11 | 5.1 |

Cloning efficiency = cloning efficiency at time of plating for mutant selection.
Untreated Control = Hanks' balanced salt solution (HBSS); ND = no data.

Table 9
 Mutagenicity Assay of Trinitrotoluene (TNT)-Contaminated Compost
 and Uncontaminated Soil Sample Using RAW 264.7 Macrophage Cells

| Treatment | Colonies per Dish | | Total # Colonies | Cloning Efficiency (%) | Mutant Colonies per Selection Dish | | | | | Total Mutant Colonies | Mutants per 10 ⁶ Clonable Cells | |
|------------------------------------|-------------------|-----|------------------|------------------------|------------------------------------|-----|-----|-----|-----|-----------------------|--|-------|
| | | | | | 1 | 2 | 3 | 4 | 5 | | | |
| <u>18-Hour Treatment (Trial 1)</u> | | | | | | | | | | | | |
| Untreated | 72 | 74 | 97 | 243 | 81 | 4 | 7 | 2 | 5 | 4 | 22 | 10.9 |
| TNT-Contaminated Compost | | | | | | | | | | | | |
| 1 µg/ml | 81 | 73 | 88 | 242 | 81 | 1 | 7 | 1 | 1 | 4 | 14 | 6.9 |
| 10 µg/ml | 80 | 95 | 77 | 252 | 84 | 7 | 8 | 14 | 12 | 8 | 49 | 23.3 |
| 50 µg/ml | 66 | 70 | 62 | 198 | 66 | 5 | 3 | 2 | 4 | 2 | 16 | 9.7 |
| 100 µg/ml | 53 | 53 | 42 | 148 | 49 | 3 | 3 | 4 | 6 | 7 | 23 | 18.6 |
| Uncontaminated Soil Sample | | | | | | | | | | | | |
| 1 µg/ml | 69 | 86 | 66 | 221 | 74 | 6 | 7 | 5 | 7 | 10 | 35 | 19.0 |
| 10 µg/ml | 80 | 77 | 83 | 240 | 80 | 9 | 12 | 12 | 7 | 6 | 46 | 23.0 |
| EMS (positive control) | | | | | | | | | | | | |
| 0.1 µl/ml | 82 | 98 | 93 | 273 | 91 | 31 | 34 | 27 | 21 | 20 | 133 | 58.5 |
| 0.3 µl/ml | 36 | 46 | 52 | 134 | 45 | 121 | 108 | 111 | 124 | 118 | 582 | 521.2 |
| DMSO (solvent control) | | | | | | | | | | | | |
| 10 µl/ml | 77 | 69 | 74 | 220 | 73 | 9 | 4 | 3 | 12 | 7 | 35 | 19.1 |
| <u>18-Hour Treatment (Trial 2)</u> | | | | | | | | | | | | |
| Untreated | 118 | 129 | 122 | 369 | 123 | 36 | 47 | 36 | 52 | 36 | 207 | 67.3 |
| 10% TNT-Contaminated Compost | | | | | | | | | | | | |
| 100 µg/ml | 106 | 117 | 111 | 334 | 111 | 35 | 70 | 58 | 50 | 62 | 275 | 98.8 |
| 500 µg/ml | 87 | 93 | 129 | 309 | 103 | 111 | 93 | 131 | 128 | 125 | 592 | 229.9 |
| Uncontaminated Soil Sample | | | | | | | | | | | | |
| 1 µg/ml | 69 | 86 | 66 | 221 | 74 | 6 | 7 | 5 | 7 | 10 | 35 | 19.0 |
| 10 µg/ml | 80 | 77 | 83 | 240 | 80 | 9 | 12 | 12 | 7 | 6 | 46 | 23.0 |
| EMS (positive control) | | | | | | | | | | | | |
| 0.4 µl/ml | 22 | 49 | 41 | 112 | 37 | 17 | 34 | 20 | 31 | ND | 102 | 136.6 |
| DMSO (solvent control) | | | | | | | | | | | | |
| 10 µl/ml | 134 | 145 | 134 | 413 | 138 | 37 | 46 | 57 | 62 | 69 | 271 | 78.7 |

Cloning efficiency = cloning efficiency at time of plating for mutant selection.
 Untreated Control = Hanks' balanced salt solution (HBSS); ND = no data.

Table 10

Mutagenicity Assay of Red Dye Mix Using RAW 264.7 Macrophage Cells

| Treatment | Colonies per Dish | | Total # Colonies | Cloning Efficiency (%) | Mutant Colonies per Selection Dish | | | | | Total Mutant Colonies | Mutants per 10 ⁶ Clonable Cells | |
|------------------------|-------------------|-----|------------------|------------------------|------------------------------------|----|----|----|----|-----------------------|--|------|
| | 1 | 2 | | | 3 | 4 | 5 | | | | | |
| Untreated | 121 | 130 | 101 | 352 | 117 | 13 | 16 | 14 | 19 | 9 | 71 | 24.2 |
| Red Dye Mix | | | | | | | | | | | | |
| 5 µg/ml | 141 | 140 | 145 | 426 | 142 | 10 | 15 | 6 | 8 | 5 | 44 | 12.4 |
| 10 µg/ml | 106 | 99 | 138 | 343 | 114 | 8 | 5 | 5 | 9 | 10 | 37 | 12.9 |
| 50 µg/ml | 161 | 127 | 132 | 420 | 140 | 5 | 8 | 4 | 7 | 8 | 32 | 9.1 |
| 100 µg/ml | 180 | 220 | 239 | 639 | 213 | 10 | 6 | 8 | 14 | 13 | 51 | 9.6 |
| EMS (positive control) | | | | | | | | | | | | |
| 0.6 µl/ml | 172 | 175 | 131 | 478 | 159 | 51 | 54 | 55 | 56 | 64 | 280 | 70.3 |
| DMSO (solvent control) | | | | | | | | | | | | |
| 10 µl/ml | 165 | 185 | 193 | 543 | 181 | 17 | 12 | 15 | 10 | 13 | 67 | 14.8 |

Cloning efficiency = cloning efficiency at time of plating for mutant selection.
 Untreated Control = Hanks' balanced salt solution (HBSS).