

REPORT DOCUMENTATION PAGE

Form Approved
OMB NO. 0704-0188

Public Reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comment regarding this burden estimates or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave Blank)		2. REPORT DATE February 5, 1998		3. REPORT TYPE AND DATES COVERED Final Technical Report 1994-1997	
4. TITLE AND SUBTITLE Evolution of Regulatory Genes Governing Catabolic Pathways in <i>Acinetobacter</i>				5. FUNDING NUMBERS DAAH04-94-G-0406	
6. AUTHOR(S) L. Nicholas Ornston					
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Department of Molecular, Cellular and Developmental Biology, Yale University P.O. Box 208103, New Haven, CT 06520-8103				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U. S. Army Research Office P.O. Box 12211 Research Triangle Park, NC 27709-2211				10. SPONSORING / MONITORING AGENCY REPORT NUMBER ARO 33124.11-LS	
11. SUPPLEMENTARY NOTES The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision, unless so designated by the documentation.					
12 a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited.				12 b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) Microbial control of biodegradative enzymes generally is exercised at the level of transcription. Thus an understanding of existing pathways for biodegradation demands knowledge, not only of the enzymes, but also of the mechanisms that control their synthesis. In this research program we have explored the properties of an evolutionarily related set of genes that control transcription of catabolic enzymes. Despite their similarities, the regulatory genes exert highly specific controls at the level of DNA. Using newly developed procedures based on random polymerase chain reaction (PCR) mutagenesis, we have been able to select mutants in which one of the transcriptional activators has been inactivated by a wide range of amino acid substitutions. This advance has established a basis for analysis of how the primary structure of the activator protein influences its activity. Furthermore, we have shown that relatively minor genetic modification is sufficient to alter one regulatory gene so that it can assume the function of an evolutionarily related regulatory gene that had been deleted from the chromosome.					
14. SUBJECT TERMS Biodegradation, Bacteria, Regulation, Transcription, Mutation, Polymerase Chain Reaction (PCR), Mutagenesis				15. NUMBER OF PAGES	
				16. PRICE CODE	
17. SECURITY CLASSIFICATION OR REPORT UNCLASSIFIED	18. SECURITY CLASSIFICATION ON THIS PAGE UNCLASSIFIED	19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED		20. LIMITATION OF ABSTRACT UL	

NSN 7540-01-280-5500

Standard Form 298 (Rev.2-89)
Prescribed by ANSI Std. Z39-18
298-102

19980520 025

REPORT DOCUMENTATION PAGE (SF298)
(Continuation Sheet)

Final Progress Report

Evolution of Regulatory Genes Governing Catabolic Pathways in *Acinetobacter*

L. N. Ornston

Department of Molecular, Cellular and Developmental Biology, Yale University

Biodegradation of complex chemicals requires the coordinated activities of numerous enzymes. Microbial control of such enzymes generally is exercised at the level of transcription. Thus an understanding of existing pathways for biodegradation demands knowledge, not only of the enzymes, but also of the mechanisms that control their synthesis. In this research program we have explored the properties of an evolutionarily related set of genes that control transcription of catabolic enzymes. Despite their similarities, the regulatory genes exert highly specific controls at the level of DNA. Nevertheless, relatively minor genetic modification is sufficient to alter one regulatory gene so that it can assume the function of another regulatory gene that has been deleted from the chromosome.

The first of the regulatory genes to be elucidated was *pobR* from an *Acinetobacter* strain that is unique among bacteria in its extraordinary competence for natural transformation. This unusual genetic property has greatly facilitated both analysis and manipulation of genes governing biodegradation. Mutations inactivating *pobR* prevent the inducible expression of *pobA*, structural gene for the hydroxylase that converts *p*-hydroxybenzoate to protocatechuate. Organisms containing such mutations are readily selected in strains containing a mutation that blocks protocatechuate catabolism at the level of a toxic metabolite.

Genetic and biochemical analysis demonstrated that *pobR* and *pobA* are divergently transcribed and physically separated by a 134 bp intergenic region containing an operator to which *PobR* binds. This DNA is characterized by a 40 bp segment containing direct repetitions of a 9 bp signature sequence (TGTCCGATG) followed by 9 bp of apparently unrelated sequence directly preceding an inversion of the 9 bp signature sequence. As we have learned through this investigation and the work of others, variants of this operator sequence govern transcription of different genes associated with catabolism of aromatic acids.

Spontaneous mutations blocking *pobR* are unusual in that they tend to be caused by disruption introduced by the newly discovered insertion sequence *IS1236*. These mutations do not occur at a single hotspot but rather are distributed throughout the gene. Thus it appears that the insertions are not determined so much by sequence as by another property of *pobR* DNA, perhaps its topology.

REPORT DOCUMENTATION PAGE (SF298)
(Continuation Sheet)

DNA altered by IS1236 is increased in length by 1.2 kb and therefore is easily identified by Polymerase Chain Reaction (PCR). Therefore the background of spontaneous mutation in *pobR* offered a system in which a novel technique for recovery of mutants containing random mutations generated within a targeted region of DNA by PCR replication errors during amplification. Primers on either side of the target (in this case *pobR*) are used to amplify it, and the amplified DNA serves as donors for recipient strains in which the function of the targeted DNA prevents growth. Transformation with *Taq*-amplified DNA *pobR* DNA increased the mutations frequency 240-fold and reduced the level of IS1236 inserts to undetectable levels. Sequence analysis of 89 of the mutant *pobR* alleles showed that the mutations were predominantly single-nucleotide substitutions broadly distributed within *pobR*. Promoter mutations were recovered, as were two mutations likely to block *pobR* translation. One third of the recovered mutations conferred a leaky or temperature-sensitive phenotype. Therefore the general application of this procedure is likely to afford a sensitive assessment of mutations causing partially impaired function that can be detected at the level of phenotype.

A separate study led to characterization of PcaU, the transcriptional activator of the *pca* operon which encodes enzymes for dissimilation of protocatechuate. The *pcaU* regulatory system was discovered by sequencing of DNA flanking a mutation designated *pcaP1* now known to be a regulatory mutation that blocks transcription of the *pca* operon of structural genes. This operon is separated from the divergently transcribed *pcaU* by 282 bp of DNA containing both *pcaP1* and a 40 bp DNA segment that closely resembles the *pobR* operator. PcaU and PobR share common ancestry as indicated by 54% identity of their amino acid sequences. Characterization of *pcaU* was delayed by the fact that knockout mutations in this gene prevent neither expression of the *pca* operon nor growth of the bacteria with protocatechuate at 37°, the temperature at which this strain of *Acinetobacter* is conventionally grown in the laboratory. Exposure of *pcaU* knockout strains to 22° does prevent their growth with protocatechuate, so this temperature was used for genetic characterization of the *pca* regulatory system. This work is still in progress, but it is notable that single nucleotide substitutions in either *pcaU* or *pobR* were sufficient to alter the product of the one gene so that it could complement knockout mutations in the other gene. Thus very little genetic information is required to achieve the extraordinary specificity that allows cells to distinguish between *p*-hydroxybenzoate (4-hydroxybenzoate) and protocatechuate (3,4-dihydroxybenzoate) as modulators effecting transcriptional control.

Important components of regulatory systems are the proteins that govern transport of potential growth substrates into the cell. Sequencing has revealed genes for multiple transport systems, and the available evidence indicates that these systems have overlapping substrate specificities. The physiological functions of the transport systems are being established by characterization of the phenotypes of strains in which the transport genes have been inactivated by mutation.

REPORT DOCUMENTATION PAGE (SF298)
(Continuation Sheet)

Publications

Gerischer, U., D'Argenio, D. A., and L. N. Ornston. 1996. IS1236, a newly discovered member of the IS3 family, exhibits varied patterns of insertion into the *Acinetobacter calcoaceticus* chromosome. Microbiology 142:1825-1831.

Kok, R. G., D. A. D'Argenio, and L. N. Ornston. 1997. Combining localized PCR mutagenesis and natural transformation in direct genetic analysis of a transcriptional regulator gene, *pobR*. J. Bacteriol. 179:4270-4276.

Gerischer, U., A. Segura, and L. N. Ornston. 1998. PcaU, transcriptional activator of genes for protocatechuate utilization in *Acinetobacter*. J. Bacteriol. In press.

Participating Scientific Personnel

L. N. Ornston, Principal Investigator

Ulrike Gerischer, Ruben Kok and Ana Segura, Postdoctoral associates

David A. D'Argenio, graduate student