


BOSTON UNIVERSITY  
SCHOOL OF MEDICINE

Thesis

REGULATION OF COL $\alpha$ 1(I), LO AND COX-1 mRNA EXPRESSION BY  
PROSTAGLANDIN E<sub>2</sub> IN HUMAN LUNG EMBRYONIC FIBROBLASTS, IMR-90



by

JAMES K. CHOUNG

B.S., United States Military Academy, 1989

19980721 084

Submitted in partial fulfillment of the  
Requirements for the degree of  
Master of Arts  
1998

DTIC QUALITY INSPECTED 5

BOSTON UNIVERSITY  
SCHOOL OF MEDICINE

Thesis

REGULATION OF COL $\alpha$ 1(I), LO AND COX-1 mRNA EXPRESSION BY  
PROSTAGLANDIN E<sub>2</sub> IN HUMAN LUNG EMBRYONIC FIBROBLASTS, IMR-90

by

JAMES K. CHOUNG

B.S., United States Military Academy, 1989

19980721 084

Submitted in partial fulfillment of the

Requirements for the degree of

Master of Arts

1998

**Personally Identifiable  
Information Redacted**

DTIC QUALITY INSPECTED 5


CPT JAMES K. CHOUNG  
3102C Paterson Loop  
West Point, NY 10996  
6 July 1998

Defense Technical Information Center  
ATTN: DTIC-OCC  
8725 John K. Kingman Rd  
Ft. Belvoir  
Alexandria, VA 22060-6218

Dear Whom It May Concern:

My social security number is [REDACTED] I have received my Master of Arts degree in biochemistry from Boston University School of Medicine, and I have enclosed the thesis that was submitted for this degree completion. I am sending this document as a requirement in the United States Army Student Detachment Handbook.

Sincerely,



James K. Choung  
CPT, USA

Approved by

First Reader Linda Taylor

Linda Taylor, Ph.D.

Assistant Professor of Biochemistry

Second Reader Peter Polgar

Peter Polgar, Ph.D.

Professor of Biochemistry

## ACKNOWLEDGEMENTS

The completion of this thesis and my education at Boston University could not have been possible without the help of two great people, Dr Linda Taylor and Dr Peter Polgar. They became a part of my life ever since I began a new life in academia and since my return to the United States from an overseas military tour as an U. S. Army officer. I am greatly indebted to them. Despite my lack of knowledge and experience in this science field, Dr Polgar took me under his wings and gave me a chance to explore and learn the art of science. On the other hand, Dr. Taylor was a great teacher from whom I was able to learn the laboratory techniques that were crucial in completing this thesis. From time to time, they were also my family counselors. For this, my family and I are very grateful.

My gratitude also extends to all the members of the laboratory, the enviable bachelor with great knowledge in science, Dr Greg Prado; the father of a newborn girl to whom I wish much happiness and success, Mike Yang; and the master of Northern blot with magical hands for good experimental results, Charles Zhou. They all gave me an unlimited assistance in the laboratory and their friendship made my days at the laboratory very enjoyable. Especially, I wish to thank Charles for taking his time to study with me and to teach me the many aspects of the laboratory work as well as the course work.

Lastly, I am grateful to my lovely wife, Ji and charming daughter, Elizabeth. They were most patient with my late hours, weekend studies and

many broken dinner dates. Ji has always given me a full support in all my endeavors and for this I am very thankful. And to my darling Elizabeth, I want to thank her for giving me many laughs and inspiration to succeed.

REGULATION OF COL $\alpha$ 1(I), LO AND COX-1 mRNA EXPRESSION BY  
PROSTAGLANDIN E<sub>2</sub> IN HUMAN LUNG EMBRYONIC FIBROBLASTS, IMR-90

JAMES K. CHOUNG

Boston University, School of Medicine, 1998

Major Professor: Linda Taylor, Ph.D., Assistant Professor of Biochemistry

ABSTRACT

PGE<sub>2</sub> decreases the synthesis of collagen and lysyl oxidase, key factors involved in fibrosis and tissue scarring. The mechanism by which PGE<sub>2</sub> induces its inhibitory effect has been explored in human embryonic lung fibroblasts, IMR-90, and Rat-1 cells. In IMR-90, PGE<sub>2</sub> limited COL $\alpha$ 1(I) and LO mRNA expressions, but increased mRNA level of COX-1 which catalyzes PGE<sub>2</sub> synthesis. Through Northern blot analysis, EP2, EP3 and EP4 receptor mRNAs were detected. Among these, EP2 mRNA expression was predominant, while EP3 and EP4 mRNAs were expressed in lesser degree with EP3 expression showing the least amount. EP1 receptor was not detected. Agonists, 11-deoxy PGE<sub>1</sub> for EP2/EP4 and sulprostone for EP3/EP1, were used to determine the receptor type and possible signaling mechanism involved in mediating the PGE<sub>2</sub> effect. 11-deoxy PGE<sub>1</sub> demonstrated similar potency in limiting not only the basal but also TGF- $\beta$  and IL-1 $\beta$  stimulated COL $\alpha$ 1(I) and LO mRNA levels, respectively. Sulprostone showed no effect. Since EP2 and EP4 are known to generate cAMP, forskolin

was used to examine this possibility. Forskolin showed similar results as PGE<sub>2</sub> and 11-deoxy PGE<sub>1</sub>. However, forskolin with PGE<sub>2</sub> showed greater inhibition of COL $\alpha$ 1(I) mRNA level than either effector, alone. Rat-1 cells were used to further examine the importance of EP2 receptor. In contrast to IMR-90, PGE<sub>2</sub> had no effect on COL $\alpha$ 1(I) mRNA level, but forskolin increased it by almost 10-fold. Interestingly, only EP4 mRNA was detected in Rat-1 cells. These results suggest that EP2 is crucial in suppressing COL $\alpha$ 1(I) and LO mRNA levels, subsequently limiting collagen deposition.

## TABLE OF CONTENTS

|                               |     |
|-------------------------------|-----|
| TITLE PAGE                    | i   |
| READER'S APPROVAL PAGE        | ii  |
| ACKNOWLEDGEMENTS              | iii |
| ABSTRACT                      | v   |
| TABLE OF CONTENTS             | vii |
| LIST OF FIGURES               | ix  |
| INTRODUCTION                  |     |
| Overview                      | 1   |
| Collagen and Lysyl Oxidase    | 3   |
| TGF- $\beta$ and IL-1 $\beta$ | 7   |
| COX and PGE <sub>2</sub>      | 13  |
| PGE <sub>2</sub> Receptors    | 15  |
| Cyclic AMP                    | 17  |
| MATERIALS AND METHODS         |     |
| Materials                     | 21  |
| Cell Culture                  | 21  |
| cDNA Probe Preparation        | 22  |
| Northern Blot Analysis        | 24  |
| Cyclic AMP Assay              | 27  |

## TABLE OF CONTENTS - CONTINUED

### RESULTS

|                                                                                                                                           |    |
|-------------------------------------------------------------------------------------------------------------------------------------------|----|
| Effect of PGE <sub>2</sub> on COL $\alpha$ 1(I) mRNA in IMR-90                                                                            | 28 |
| EP Receptor Profile in IMR-90                                                                                                             | 28 |
| Effect of 11-deoxy PGE <sub>1</sub> and Sulprostone on COL $\alpha$ 1(I), LO and COX-1<br>mRNA in IMR-90                                  | 33 |
| Effect of Forskolin on COL $\alpha$ 1(I), LO and COX-1 mRNA in IMR-90                                                                     | 40 |
| Cyclic AMP Assay for IMR-90 with Forskolin, PGE <sub>2</sub> and 11-deoxy PGE <sub>1</sub>                                                | 49 |
| Cyclic AMP Assay for IMR-90 with PGE <sub>2</sub> Concentrations                                                                          | 52 |
| Effect of 11-deoxy PGE <sub>1</sub> on TGF- $\beta$ and IL-1 $\beta$ stimulated COL $\alpha$ 1(I)<br>and LO mRNA, respectively, in IMR-90 | 52 |
| EP Receptor Profile in Rat-1 Cells                                                                                                        | 59 |
| Effect of PGE <sub>2</sub> on COL $\alpha$ 1(I) in Rat-1 Cells                                                                            | 59 |
| Cyclic AMP Assay for Rat-1 Cells with Forskolin and PGE <sub>2</sub>                                                                      | 64 |
| DISCUSSION                                                                                                                                | 67 |
| REFERENCES                                                                                                                                | 73 |

## LIST OF FIGURES

|                                                                                                                                 |    |
|---------------------------------------------------------------------------------------------------------------------------------|----|
| FIGURE 1: Effect of PGE <sub>2</sub> on COL $\alpha$ 1(I) mRNA in IMR-90 .....                                                  | 29 |
| FIGURE 2: Northern Blot Analysis of EP Receptors in IMR-90 .....                                                                | 31 |
| FIGURE 3: Effect of 11-deoxy PGE <sub>1</sub> on COL $\alpha$ 1(I) mRNA in IMR-90 .....                                         | 34 |
| FIGURE 4: Effect of 11-deoxy PGE <sub>1</sub> on LO mRNA in IMR-90 .....                                                        | 36 |
| FIGURE 5: Effect of 11-deoxy PGE <sub>1</sub> on COX-1 mRNA in IMR-90 .....                                                     | 38 |
| FIGURE 6: Effect of Sulprostone on COL $\alpha$ 1(I), LO<br>and COX-1 mRNA in IMR-90 .....                                      | 41 |
| FIGURE 7: Effect of PGE <sub>2</sub> and Forskolin on COL $\alpha$ 1(I) mRNA in IMR-90 .....                                    | 43 |
| FIGURE 8: Effect of PGE <sub>2</sub> and Forskolin on LO mRNA in IMR-90 .....                                                   | 45 |
| FIGURE 9: Effect of PGE <sub>2</sub> and Forskolin on COX-1 mRNA in IMR-90 .....                                                | 47 |
| FIGURE 10: Effect of Forskolin, PGE <sub>2</sub> and 11-deoxy PGE <sub>1</sub> on<br>cAMP Production in IMR-90 .....            | 50 |
| FIGURE 11: Effect of PGE <sub>2</sub> on cAMP Production in IMR-90 .....                                                        | 53 |
| FIGURE 12: Effect of TGF- $\beta$ , 11-deoxy PGE <sub>1</sub> and PGE <sub>2</sub> on<br>COL $\alpha$ 1(I) mRNA in IMR-90 ..... | 55 |
| FIGURE 13: Effect of IL-1 $\beta$ , 11-deoxy PGE <sub>1</sub> and PGE <sub>2</sub> on<br>LO mRNA in IMR-90 .....                | 57 |
| FIGURE 14: Northern Blot Analysis of EP Receptors in Rat-1 Cells .....                                                          | 60 |

LIST OF FIGURES - CONTINUED

|                                                                               |    |
|-------------------------------------------------------------------------------|----|
| FIGURE 15: Effect of PGE <sub>2</sub> and Forskolin on COL $\alpha$ 1(I) mRNA |    |
| in Rat-1 Cells -----                                                          | 62 |
| FIGURE 16: Effect of Forskolin and PGE <sub>2</sub> on cAMP Production        |    |
| in Rat-1 Cells -----                                                          | 65 |

## **INTRODUCTION**

### **Overview**

Deposition of large amounts of collagen is one of the prominent attributes of fibrosis [Zhang et al., 1994; McDonald et al., 1986] and tissue scarring [Whitby and Ferguson, 1991]. The regulation of collagen deposition involves a number of effectors, including prostaglandin E<sub>2</sub> (PGE<sub>2</sub>) [Goldstein and Polgar, 1982; Raisz et al., 1993], an enzyme, lysyl oxidase (LO) [Siegel et al., 1970; Rucker and Murray, 1978], and cytokines such as transforming growth factor-beta (TGF- $\beta$ ) [Fine et al., 1990; Diaz et al., 1989; Krupsky et al., 1996] and interleukin-1 (IL-1) [Bhatnagar et al., 1986; Kahari et al., 1987; Pasternak et al., 1986; El Maradny et al., 1996]. The effect of PGE<sub>2</sub> on the regulation of the type I collagen  $\alpha$ 1 [COL $\alpha$ 1(I)] synthesis and deposition at the receptor and transcriptional levels is the focus of this dissertation.

In response to tissue injury, macrophages release a plethora of membrane-perturbing stimuli including IL-1 $\beta$  [Cannon et al., 1992; Zhang et al., 1993] and TGF- $\beta$  in wound cleft [Broadly et al., 1988; Grotendorst et al., 1988]. One of the initial effects of these cytokines seems to result in an overall increase in collagen synthesis. TGF- $\beta$  is widely known to cause an increase in collagen synthesis by cells such as fibroblasts [Reed et al., 1994; Gallivan et al., 1997; Krupsky et al., 1996; Fine et al., 1990; Diaz et al., 1989]. On the other hand, IL-1 $\beta$  has been found to increase the synthesis of LO [Roy et al., 1996] which is an active participant during collagen deposition [Pinnell and Martin, 1968]. In

addition, these two cytokines actively induce PGE<sub>2</sub> synthesis, apparently through arachidonate metabolism catalyzed by cyclooxygenase-1 (COX-1) and COX-2 [Jackson et al., 1993]. Of the many responses PGE<sub>2</sub> elicits during the immune response, it decreases collagen levels, both steady state [Brilla et al., 1995; Goldstein and Polgar, 1982; Fine et al., 1992] and TGF- $\beta$  stimulated levels [Fine et al., 1990; Diaz et al., 1989; Krupsky et al., 1996]. Also, it decreases both steady state and IL-1 $\beta$  stimulated LO mRNA levels [Roy et al., 1996].

This evidence suggests that PGE<sub>2</sub> and its receptor(s) are important factors, if not the central factors, in collagen deposition. PGE<sub>2</sub> induces its effect in an autocrine fashion through four receptor subtypes, EP1, EP2, EP3 and EP4 [Coleman et al., 1994; Ichikawa et al., 1996]. It is not known which EP receptor subtype(s) is/are involved in effecting collagen deposition. The first objective of this research is to determine the EP receptor subtype profile in two *in vitro* models, mortal human embryonic lung fibroblasts (IMR90) and immortal rat embryonic fibroblasts (Rat-1). The second objective is to pinpoint the receptor subtype(s) involved in mediating the PGE<sub>2</sub> effect on collagen deposition. The third objective is to determine the secondary signal messenger involved in the receptor subtype(s) signal transduction. By achieving these objectives, the investigator hopes to provide some insight into critical aspects of collagen deposition that may contribute to therapeutic ramifications in fibrosis and tissue scarring.

### **Collagen and Lysyl Oxidase**

Fibrosis develops from an excessive accumulation of collagen in the extracellular matrix and demonstrates many characteristics of wound healing following tissue injury [Siegel, 1979]. In fibrotic lung, such as in idiopathic chronic fibrosis, the increase in the total amount of the type I collagen synthesis and its deposition in the interstitium and alveolar spaces are the most prominent features [Seyer et al., 1976].

Collagen molecules consist of three helical polypeptides, called  $\alpha$  chains, wound about each other, forming a triple helix. Type I collagen [Birk and Silver, 1984] usually consists two  $\alpha 1(I)$  chains and one  $\alpha 2(I)$  chain [Piez et al., 1963], or homotrimers of  $\alpha 1(I)$  chains, if no  $\alpha 2(I)$  is produced [Miller and Matukas, 1969]. Type I collagen is fibril forming collagen and is one of the main types found in the connective tissues [Bornstein, 1974]. It is not only the principal collagen of skin and bone, but also the major interstitial collagen [Bornstein, 1974].

The most commonly held view of collagen molecule folding begins with synthesis of procollagen from the three pro $\alpha$ -chains [Bornstein, 1974; Bienkowski and Gotkin, 1995]. The pro $\alpha$ -chains are rich in proline and glycine which appear to play important roles in triple helix formation. Each chain consists of about 1000 amino acid residues [Kang et al., 1967]. The chain is arranged in a left-handed helix with three amino acid residues per turn and contains glycine as every third residue [Bornstein, 1974]. Its sequence is widely viewed as Gly-X-Y, where X is commonly proline and Y is proline or hydroxyproline. According to

Bornstein (1974), the synthesis of a pro $\alpha$ -chain begins on the membrane-bound ribosomes, and it is injected into the lumen of the endoplasmic reticulum. The synthesized pro $\alpha$ -chain becomes hydroxylated at selected proline and lysine residues with some glycosylation at selected hydroxylysine residues.

Subsequently, the three pro $\alpha$ -chains wind about each other with extension peptides at the amino (N-) and carboxy (C-) termini, forming a triple helix. The helical formation in each chain is stabilized in part by the ring structure of proline, while glycine, with its small size, allows the tight packing of the three helical chains to form the ultimate triple helix. The triple helix is further stabilized with hydrogen bonds between the chains.

The triple helix is secreted into the extracellular matrix [Lenaers et al., 1971] as procollagen via secretory vesicles formed in the Golgi apparatus. The extension peptides of procollagen are then cleaved by the procollagen peptidases, producing collagen molecules [Lapiere et al., 1971]. The collagen molecules then associate laterally and head-to-end, ultimately forming fibrils [Robins and Duncan, 1983].

Thus far, almost 20 genetically distinct collagens have been found [Bienkowski and Gotkin, 1995] while about 25  $\alpha$  chains have been identified. The type I collagen accounts for 90% of body collagen and it is found in many tissues, including bone, skin, tendon, ligaments, cornea, and internal organs. The genes coding for the pro $\alpha$ 1(I) and pro $\alpha$ 2(I) chains, COL1A1 and COL1A2, are located on Chromosomes 17 and 7, respectively [Sandell and Boyd, 1990].

The tensile strength of collagen fibrils is achieved through covalent cross-links between lysine residues of collagen molecules within the fibril. Such cross-linking is initiated through LO in the extracellular space. LO is an amine oxidase which contains a copper ion and an aromatic carbonyl (pyridoxal 5'-phosphate or pyrroloquinoline quinone) as cofactors [Myers et al., 1985; Kagan, 1986]. Its molecular weight in various human tissues, determined through Western blotting of urea extracts, is approximately 30 kilo Daltons (kDa) [Kuivaniemi, 1985].

The LO oxidatively deaminates  $\epsilon$ -amino groups of lysyl or hydroxylysyl residues in collagen fibrils to peptidyl  $\alpha$ -aminoadipic- $\delta$ -semialdehyde [Pinnell and Martin, 1968]. This process yields highly reactive aldehyde groups, allysine or hydroxyallysine, respectively, which spontaneously condense with the other vicinal aldehydes to form aldol cross-links, or with the other  $\epsilon$ -amino groups of lysyl or hydroxylysyl residues to form Schiff base cross-links [Siegel, 1979].

The intramolecular covalent bonds, or the cross-links, in the triple helix of pro $\alpha$ -chains occur at two allysine residues located at the short non-helical, terminal region of the chains [Robins and Duncan, 1983]. These reactive residues combine in aldol-type condensation, known as allysine aldol. On the other hand, intermolecular cross-links results from one of the two pathways, one based on lysine aldehyde pathway and the other on hydroxylysine aldehyde pathway [Ricard-Blum and Ville, 1989]. These cross-links allow formation of covalent bonds between different  $\alpha$  chains of the same collagen type or different

collagen types. The selection of the crosslinking pathway appears to be dictated by the type of tissue in which collagen is produced [Eyre et al., 1984].

To date, no one seems to refute the suggestion of Rucker and Murray (1978) that the enzymatic activity of LO may be the sole initiator of cross-link formation for collagen. As early as 1968, Peacock et al., had shown that inhibition of the cross-linking process with lathyrogens, an irreversible lysyl oxidase inhibitor, apparently limited scar formation and retarded fibrosis. Since then, other possible therapeutic approaches have been explored to reduce collagen accumulation. Some of these approaches have targeted the regulation of tissue inhibitors of metalloproteinases (TIMPs), an inhibitor of collagenase [Shapiro et al., 1992; Pardo et al., 1992], or the production and activation of intracellular collagenases such as matrix metalloproteinases (MMPs) [Horwitz et al., 1977].

Recently, cytokines such as TGF- $\beta$  and IL-1 $\beta$  have been shown to have prominent effects in the regulation of collagen synthesis. Of these two, TGF- $\beta$  appears to be the key factor, contributing to the increase in collagen mRNA synthesis [Ritzenthaler et al., 1993; Fine et al., 1990; Diaz et al., 1989]. On the other hand, IL-1 $\beta$  has been shown to both enhance [Elias et al., 1990] and inhibit [Diaz et al., 1993; Armendariz-Borunda et al., 1992] collagen synthesis. Nonetheless, IL-1 $\beta$  seems to have a great potential to facilitate collagen accumulation, for it has been shown to increase LO synthesis [Roy et al., 1996].

### **TGF- $\beta$ and IL-1 $\beta$**

During the wound healing process, the blood-born monocytes recruit the macrophages to the wound cleft [Leibovich and Ross, 1975]. Subsequently, macrophages release a battery of growth factors as well as the cytokines, such as TGF- $\beta$  and IL-1 $\beta$ , at the wound cleft. These cytokines are also synthesized and released by surrounding connective tissue cells [Bienkowski and Gotkin, 1995; Gauldie et al., 1993] and endothelial cells [Phan et al., 1991].

The cellular response during wound healing appears to parallel the fibrotic response. In rats, the intratracheal administration of bleomycin elicits an acute inflammatory response in the lung followed by the development of pulmonary fibrosis. However, the mechanism by which bleomycin causes fibrosis is not understood. It has been proposed to involve the pulmonary macrophage which is considered a central cell in the cytokine network of the lung [Scheule et al., 1992]. In any case, in a bleomycin-induced pulmonary fibrosis, Breen et al. (1992) showed an increase in the TGF- $\beta$  mRNA levels in rat lung fibroblasts in culture, while Phan et al. (1991) demonstrated a similar increase of TGF- $\beta$  mRNA levels in rat lung endothelial cells. Further, Scheule et al. (1992) found using an *in vitro* model that bleomycin also increased IL-1 $\beta$  synthesis and secretion by human alveolar macrophages. Overall, these results seem to suggest that the cytokines, such as TGF- $\beta$  and IL-1 $\beta$ , could be the key factors in the development of fibrosis.

The TGF- $\beta$  superfamily constitutes local mediating factors which are classified into three subgroups, and are involved in many cellular processes in vertebrates [Brand and Schneider, 1995]. The three subgroups consist of the five isoforms of TGF- $\beta$ s (TGF- $\beta$ 1 -  $\beta$ 5), activins and bone morphogenetic proteins (BMPs) [Massague, 1990]. All the members of the TGF- $\beta$  superfamily are first synthesized as large latent complexes which are inactive [Derynck et al., 1985; Purchio et al., 1988]. In the latent form, TGF- $\beta$  is noncovalently bound to the latency associated peptide (LAP), a homodimer formed from the propeptide region of TGF- $\beta$  [Miyazono et al., 1988]. The LAP is disulfide linked to latent TGF- $\beta$  binding protein which has features in common with extracellular matrix proteins, and it targets latent TGF- $\beta$  to the matrix [Derynck et al., 1985; Miyazono et al., 1988; Munger et al., 1997]. Although the physiologic mechanism for activation of TGF- $\beta$  latent form is not clear, it has been activated through a cleavage at a conserved RXXR motif in the mature C-terminal dimer and the N-terminal propeptide region [Brand and Schneider, 1995].

There are three known TGF- $\beta$  receptors, types I, II and III. In most mammalian cells, these receptors have approximate molecular weights of 50 kDa, 80 kDa and 280 kDa, respectively [Brand and Schneider, 1995]. In mutant mink lung epithelial cells, it has been shown that types I and II receptors primarily mediate the signaling of TGF- $\beta$  [Laiho et al., 1990; Brand and Schneider, 1995]. On the other hand, the type III, binds TGF- $\beta$  with high affinity, but has no

apparent signaling function [Brand and Schneider, 1995]. The type I and II receptors, both have N-terminal cysteine-rich extracellular domains followed by hydrophobic single-pass transmembrane domain and serine-threonine kinase intracellular domains at the C-terminus [Wrana and Pawson, 1997; Padgett et al., 1997]. However, unlike the type II receptor, the type I contains a glycine-serine-rich region immediately preceding the kinase domain. Phosphorylation of this region has been implicated in the signaling activity [Padgett et al., 1997].

One TGF- $\beta$  signaling model consists of TGF- $\beta$  activation by release of the mature growth factor from its LAP. The mature TGF- $\beta$  then binds to the type II receptor which phosphorylates and activates the type I receptor, which subsequently activates an intracellular signaling cascade [Wrana and Pawson, 1997; Padgett et al., 1997]. Currently, the signaling mechanisms involving other combinations of the receptor types are still being explored. Recently, the discovery of a TGF- $\beta$  downstream signaling protein, a mammalian Mad (*Mothers against decapentaplegic*) related protein called smad, gave a glimpse of how this cytokine induces its signal [de Caestecker et al., 1997; Heldin et al., 1997]. However, there has been no indication of such protein being involved in the TGF- $\beta$  signaling mechanism responsible for collagen synthesis.

IL-1 has been known as "endogenous pyrogen" as early as the 1940's for its ability to induce fever [Dinarello, 1988]. The IL-1 gene family has three members, IL-1 $\alpha$  [Lomedico et al., 1984], IL-1 $\beta$  [Auron et al., 1984] and IL-1 receptor antagonist (IL-1Ra) [Seckinger et al., 1987; Hannum et al., 1990].

Although IL-1 $\alpha$  and IL-1 $\beta$  in humans have shown only about 26% amino acid sequence homology, both bind to the same receptors with similar affinities [Kuno and Matsushima, 1994]. The two are also similar in their translational and post-translational modifications. Through immunoelectron microscopy, Stevenson et al. (1992) detected that the IL-1 $\alpha$  and IL-1 $\beta$ , 31kDa precursor molecules, were translated on cytoskeletal-associated free polyribosomes in human monocytes. Also, both lack leader sequences and undergo myristoylation on lysine residues in their N-terminal propieces [Stevenson et al., 1993].

The precursor molecule of IL-1 $\alpha$  is fully active, and has been known to remain intracellularly [Dinarello, 1996]. It has also been found to translocate into the nucleus and bind to DNA [Maier et al., 1994]. Such nuclear targeting was due to the basic cluster, KVLKKRR, at the N-terminus of IL-1 $\alpha$  precursor. However, IL-1 $\alpha$  precursor can be cleaved to become 17kDa molecule either by the activation of the calcium dependent, membrane associated cysteine-rich proteases called calpains [Kobayashi et al., 1990] or by the extracellular proteases [Kobayashi et al., 1991]. In contrast, IL-1 $\beta$  precursor must be enzymatically cleaved by IL-1 $\beta$  cleaving enzyme (ICE) in the cytosol before it can become active [Black et al., 1988; Walker et al., 1994]. After cleavage, the 17kDa, mature IL-1 $\beta$  is secreted into the extracellular space through a putative membrane [Dinarello, 1996].

While IL-1 $\alpha$  and IL-1 $\beta$  are agonists for IL-1 receptors [Alheim et al., 1997], the IL-1Ra is an antagonist [Arend et al., 1990]. Unlike IL-1 $\alpha$  and IL-1 $\beta$ , IL-1Ra is

translated in the endoplasmic reticulum with a leader sequence and transported to the Golgi apparatus [Dinarello, 1996]. Subsequently, it is released into the extracellular space upon cleavage of the leader sequence. Using a murine recombinant IL-1Ra in human synovial cells and rabbit articular chondrocytes, Arend et al. (1990) showed that IL-1Ra inhibited IL-1 $\alpha$  and IL-1 $\beta$  induced PGE<sub>2</sub> production in a dose dependent manner. Further, the same experiment showed that the collagenase production by the synovial cells was also inhibited.

Although IL-1Ra is an antagonist, its binding affinity to the IL-1 receptors is similar to the two agonists. Using a murine thymoma cell line, Dripps et al. (1991) showed that IL-1Ra bound to the 80 kDa IL-1 receptor with an affinity (KD = 150 pM) similar to that of IL-1 $\alpha$  and IL-1 $\beta$ . The amino acid sequence homology of the IL-1Ra to the mature, or activated, IL-1 $\beta$  is about 26% [Dinarello, 1996].

There are two types, type I (80kDa) and II (60-68kDa), of IL-1 receptors that bind to IL-1 $\alpha$  and IL-1 $\beta$  [Kuno and Matsushima, 1994]. The two receptors have single-membrane spanning segments and the ligand binding domains which possess three immunoglobulin-like domains [Dinarello, 1996]. These receptors also exist in soluble forms and have been found in inflammatory synovial fluid and in other pathologic body fluids [Arend et al., 1994].

The type I receptor is considered the main receptor involved in IL-1 signal transduction while the type II receptor is known as a "decoy receptor". Binding of IL-1 to the type II receptor does not generate signal [Colotta et al., 1993]. One of the main reasons for the difference in their signaling abilities seems to be in their

cytoplasmic regions. While the cytoplasmic domain of the type I receptor in human contains about 215 amino acids, the type II cytoplasmic domain only consists of 29 amino acids [Kuno and Matsushima, 1994]. Nevertheless, both receptor genes share the same chromosomal locations, 2q12-22 [Kuno and Matsushima, 1994].

The signal transduction by IL-1 receptor appears to require the binding of IL-1 to the type I receptor and subsequent binding of this complex to the IL-1 receptor-accessory protein (IL-1RAcP) [Greenfeder et al., 1995]. The IL-1RAcP is a single-membrane spanning segment as the type I receptor, and the cytoplasmic domains of both IL-1RAcP and type I receptor contain the same amino acid domains commonly found in the GTPase superfamily [Hopp, 1995]. Although their signaling mechanism is still being explored, the heterodimer complex, formed by IL-1 receptor and IL-1RAcP, is likely to initiate the IL-1 signaling cascade through the association of its cytoplasmic domain with GTPase activating proteins [Dinarello, 1996]. The IL-1RAcP does not form a complex with either the ligand-bound type II receptor or IL-1Ra bound to the type I receptor [Dinarello, 1996].

In a number of experiments, IL-1 has been shown to effect PGE<sub>2</sub> synthesis and cAMP production. In the human lung fibroblast cell line, TIG-1, the IL-1 $\alpha$  or IL-1 $\beta$  increased PGE<sub>2</sub> synthesis and secondary cAMP production [Takii et al., 1992]. In the same experiment, indomethacin, a COX inhibitor, inhibited the IL-1 induced PGE<sub>2</sub> synthesis and adenosine 3',5'-cyclic monophosphate

(cAMP) production. Kol et al. (1997) also showed IL-1 $\beta$  increased arachidonic acid release, prostanoid accumulation and cytosolic phospholipase A<sub>2</sub> (PLA<sub>2</sub>) transcripts as well as PLA<sub>2</sub> activities. These effects all contribute to the PGE<sub>2</sub> production. Similarly, Roy et al. (1996) showed IL-1 $\beta$  treated IMR90 cells increased PGE<sub>2</sub> synthesis by more than 5-fold above the control level. Additionally, IL-1 $\beta$  increased COX-1 and COX-2 mRNA expressions by 3-fold and by more than 5-fold, respectively [Roy et al., 1996].

### **COX and PGE<sub>2</sub>**

Prostaglandins and related fatty acid derivatives are one of the naturally occurring autocooids that seem to induce a variety of cellular responses. Under physiological conditions, prostaglandin synthesis takes place following the liberation of precursor fatty acid arachidonate from the membrane lipids and other lipid esters by phospholipases [Lands and Samuelsson, 1968]. Subsequently, the arachidonate is transformed into prostaglandin H<sub>2</sub> by the enzyme prostaglandin endoperoxide H synthase (PGHS) and PGE<sub>2</sub> is formed from PGH<sub>2</sub> by prostaglandin E synthase [Samuelsson et al., 1975].

The PGHS is a single protein with two catalytic sites, cyclooxygenase and peroxidase active sites. The cyclooxygenase converts arachidonate into unstable PGG<sub>2</sub> through oxygenation and cyclization of a pentane ring [Samuelsson, 1975]. Then the peroxidase reduces PGG<sub>2</sub> to PGH<sub>2</sub>. There are two isoforms of PGHS, PGHS-1 and -2 which are also referred to COX-1 and COX-2, respectively [Smith and Dewitt, 1996]. These two isozymes exist as

homodimers, heme-containing, glycosylated proteins which are membrane bound. However, both are encoded by separate genes located on different chromosomes, Chromosomes 9 and 1 for COX-1 and COX-2, respectively [Crofford, 1997].

The COX-1 lacks a TATA box which is typical of a housekeeping gene. Also, it is constitutively expressed in most cells [Kraemer et al., 1992]. On the other hand, COX-2 is undetectable in most mammalian tissues, but it can be induced during the state of acute inflammation [Seibert et al., 1994; Vane et al., 1994]. The COX-2 promoter contains a TATA box and other regulatory elements which are common in highly regulated genes [Appleby et al., 1994]. These elements are known to respond during inflammation to nuclear factor- $\kappa$ B, cAMP response element binding protein and the CAAT enhancer binding proteins [Appleby et al., 1994].

Jackson et al. (1993) demonstrated that IL-1 $\beta$  alone increased PGE<sub>2</sub> synthesis in IMR90 by 8-fold, while IL-1 $\beta$  and TGF- $\beta$  together increased PGE<sub>2</sub> synthesis by 25-fold. In addition, IL-1 $\beta$  and TGF- $\beta$  increased PLA<sub>2</sub> and COX-1 mRNA, respectively, while IL-1 $\beta$  rapidly induced COX-2 mRNA in fibroblasts [Jackson et al., 1993; DeWitt and Meade, 1993]. In addition, Roy et al. (1996) showed that the half-life of COX-2 mRNA ( $t_{1/2}$  = 30 min) is shorter than COX-1 mRNA ( $t_{1/2}$  = 4 h). Also, PGE<sub>2</sub> induced COX-1 synthesis [Roy et al., 1996]. These results suggested that COX-2 may provide the initial catalytic activity required for the increased synthesis of PGE<sub>2</sub> at the beginning of an inflammatory

response. Subsequently, PGE<sub>2</sub> could increase the synthesis of COX-1 which may participate in maintaining the elevated level of PGE<sub>2</sub> [Roy et al., 1996]. Overall, the results from Jackson et al. (1993) and Roy et al. (1996) suggest that perhaps the increase in PGE<sub>2</sub> synthesis in response to cytokines, IL-1 $\beta$  and TGF- $\beta$ , could very well be due to the coordinated regulation of the COX-1 and COX-2.

Although the functional importance of COX isozymes are still being elucidated, their importance has been noted in idiopathic pulmonary fibrosis. According to Wilborn et al. (1995), lung fibroblasts isolated from the patients with idiopathic pulmonary fibrosis showed decreased level of COX-2 expression compared to the normal lung fibroblasts. Consequently, the diseased fibroblasts' capacity to synthesize PGE<sub>2</sub> was diminished.

### **PGE<sub>2</sub> Receptors**

PGE<sub>2</sub> exerts its diversified biological action on cells in an autocrine fashion utilizing four PGE<sub>2</sub> receptor (EP) subtypes, EP1 [Funk et al., 1993], EP2 [Regan et al., 1994], EP3 [Regan et al., 1994], and EP4 [An et al., 1993; Bastien et al., 1994]. These receptors are G-protein coupled, seven transmembrane receptors [Coleman et al., 1994; Ichikawa et al., 1996]. EP1 receptors have been found to be coupled to increases in intracellular Ca<sup>2+</sup> in Chinese hamster ovary (CHO) cells [Kato et al., 1995] and COS cells [Funk et al., 1993]. On the other hand, both EP2 and EP4 receptors have been known to stimulate adenylate cyclase, thus increasing cAMP [Coleman et al., 1994; Regan et al., 1994; Bastien et al., 1994; Mori et al., 1996; Nishigaki et al., 1996].

However, there are some differences between the EP2 and EP4 receptors, both structurally and functionally. For example, in the cloned mouse EP4 receptor, there are about 3.5 fold more serine and threonine residues in the C-terminal than the cloned mouse EP2 receptor [Nishigaki et al., 1996]. Also, EP4 receptor contains a potential PKA phosphorylation sites whereas none was found on EP2 receptor [Nishigaki et al., 1996]. Such phosphorylation sites may be critical in receptor desensitization [Liggett et al., 1993] and subsequently its signal transduction.

Unlike EP1, EP2 and EP4 receptors, EP3 receptors exist in a number of isoforms. Thus far, six isoforms of EP3 (EP3A, EP3D, EP3E, EP3F, EP3e, EP3f) have been found in human [Pierce et al., 1995]. All cDNAs from human EP3 isoforms showed identical amino acid sequences from their N-terminal to the 11 residues after the putative seventh transmembrane domain [Pierce et al., 1995]. The most obvious difference between the isoforms was the length of the C-terminal, resulting from alternative mRNA splicing [Woodward et al., 1997]. This variance in length may cause the receptor isoforms to function differently. For example, EP3E isoform has the shortest C-terminal region, consisting of only six amino acids following the splice site, and there are no serine or threonine residues. In contrast, EP3D isoform has the longest C-terminal region with 29 amino acids following the splice site and has seven serine and threonine residues [Pierce et al., 1995]. Lefkowitz (1993) noted that the phosphorylation of serine and threonine residues appeared to be involved in G-protein coupled

receptor desensitization. Further, An et al. (1994) reported that PGE<sub>2</sub> induced EP3E isoform desensitization was rapid and transient while EP3D isoform demonstrated slow and persistent desensitization.

The functions of EP3 isoforms are not only diversified, but also contradicting, sometimes. All human EP3 isoforms expressed in baby hamster kidney (BHK) cells inhibited adenylate cyclase activity upon stimulation with PGE<sub>2</sub> [Schmid et al., 1995]. Of all the isoforms, EP3A isoform was most efficient in inhibiting adenylate cyclase. However, overexpression of EP3A isoform also increased cAMP levels in response to PGE<sub>2</sub>. Kotani et al. (1995) also showed that some human EP3 isoforms in COS-7 and CHO cells raised cAMP levels when incubated with EP3 agonist, M&B-28767. In addition, the human EP3A, B and C isoforms expressed in BHK cells increased intracellular Ca<sup>2+</sup> in response to PGE<sub>2</sub> [Schmid et al., 1995]. Moreover, both human [An et al., 1994] and mouse [Irie et al., 1994] EP3 isoforms in CHO cells, increased intracellular Ca<sup>2+</sup> in varying degrees and decreased cAMP levels.

### **Cyclic AMP**

Cyclic AMP is an intracellular mediator that is synthesized from ATP by the enzyme adenylate cyclase. To date, 11 isoforms of adenylate cyclase have been found in mammalian cells [Houslay and Milligan, 1997]. This enzyme is activated by the Gs $\alpha$  subunit of trimeric G-protein.

The cyclic AMP plays a vital role in signal transduction of many G protein coupled, seven transmembrane receptors. Increases in intracellular cAMP levels

results in various cellular processes such as protein metabolism, secretory, pro-inflammatory cytokine production, apoptosis and growth control [Houslay and Milligan, 1997]. For example, the increase in cAMP level by parathyroid hormone in rat osteoblastic osteosarcoma cells, decreases collagen mRNA synthesis and mitogenesis of the cells [Pun, 1989]. Such losses may contribute to the loss of bone in hyperparathyroidism.

The effects derived from increased levels of cAMP production usually involve the cAMP-dependent protein kinase, known as protein kinase A (PKA) [Krebs, 1989]. The PKA consists of tetramers containing two regulatory (R) subunits and two catalytic (C) subunits [Nesterova et al., 1996]. Upon binding of cAMP, the two C subunits dissociate from the R subunits and become active, phosphorylating target proteins. There are a number of mammalian isoforms for both subunits: RI $\alpha$ , RI $\beta$ , RII $\alpha$ , RII $\beta$ , C $\alpha$ , C $\beta$ , C $\gamma$  [Nesterova et al., 1996; Dell'Acqua and Scott, 1997]. Each isoform of R subunit can associate with any of the C subunits while different R subunits can heterodimerize together [Doskeland et al., 1993; Nesterova et al., 1996; Dell'Acqua and Scott, 1997]. While C subunit isoforms show no distinguishable difference in substrate specificity and interaction with either R subunits, the dimeric R subunits show distinct cAMP binding affinities and different localization in the cells [Taylor et al., 1990].

Additionally, the effects of cAMP are further modulated through the control of cAMP phosphodiesterase (PDE) activity [Houslay and Milligan, 1997]. The

cAMP PDE hydrolyzes cAMP to 5'AMP, thus inactivating cAMP activities. To date, over 30 isoforms of PDE have been reported for mammalian cells [Houslay and Milligan, 1997].

A number of hormones mediate their signals through cAMP. How these hormones activate their respective signaling pathways by inducing the correct pool of enzymes, which includes adenylate cyclase, PKA and/or cAMP PDE, is still not clear. Recently, a "targeting hypothesis" has been proposed in response to this phenomenon [Dell'Acqua and Scott, 1997]. This hypothesis proposes that the activation of a specific enzyme is determined by its association with targeting subunits, now known as anchoring protein [Johnson et al., 1997; Visconti et al., 1997; Feliciello et al., 1997; Pawson and Scott, 1997; Dell'Acqua and Scott, 1997].

Interestingly, one of the signaling mediators reported to influence collagen synthesis is cAMP. Saltzman et al. (1982) and Baum et al. (1978) showed increased levels of cAMP suppressed collagen synthesis. In contrast, Fine et al (1992) reported that the inhibitory effect of PGE<sub>2</sub> in collagen synthesis was not mediated through cAMP.

Overall, the pathologic hallmark of tissue scarring and fibrosis is collagen accumulation. TGF- $\beta$  and IL-1 $\beta$  increase collagen and LO mRNAs, respectively. In combination or alone, these cytokines also increase COX-1 and COX-2 expressions, and PGE<sub>2</sub> synthesis. PGE<sub>2</sub> has shown to abrogate both collagen and LO mRNA expressions, even in the presence of the two cytokines. The

effect of PGE<sub>2</sub> is clearly inhibitory in both collagen synthesis and deposition. PGE<sub>2</sub> in essence may play the central role in this process. However, the mechanism involved in such inhibition is not yet clear. The interaction between PGE<sub>2</sub> and its receptor subtypes activate signal transduction pathways that are independent and sometimes contradictory in nature. Nonetheless, the elucidation of this complicated signaling mechanism could provide valuable information leading to limitation or inhibition of tissue scarring and/or fibrosis.

## **MATERIALS AND METHODS**

### **Materials**

PGE<sub>2</sub> and forskolin were purchased from Sigma (St Louis, MO); IL-1 $\beta$  was purchased from Genzyme (Cambridge, MA); sulprostone and TGF- $\beta$  were purchased from R & D Systems Inc. (Minneapolis, MN); and 11-deoxy PGE<sub>1</sub> was purchased from Cayman Chemical Co (Ann Arbor, MI). EP1 cDNA was a gift of Dr. C.D. Funk (University of Pennsylvania); EP2, and EP3 were gifts of Dr. J. W. Regan (University of Arizona); EP4 cDNA, and COL $\alpha$ 1 cDNA were the gifts of Dr. E. J. Goetzl (University of California Medical Center at San Francisco) and Dr. RH Goldstein of Boston University School of Medicine, respectively. Human COX-1 cDNA was a gift from Dr. T. Hla, Holland Laboratory, American Red Cross; and human LO cDNA was a gift from Dr. H. Kagan of Boston University School of Medicine.

### **Methods**

**Cell culture** IMR90 cells were obtained from the Coriell Institute for Medical Research (Camden, NJ). The cells were grown to confluence in 100x20 mm sterile Falcon tissue culture plates for Northern blot experiments, and in a 12-well plates for cAMP assay. These cells were cultured at 37°C in Minimum Essential Medium Eagle (MEM; Mediatech, Inc., Herndon, VA) containing 10% fetal bovine serum (FBS) supplemented with 50units/ml penicillin and 50 $\mu$ g/ml streptomycin. Cells were used at population doubling level of 18 to 30. Before initiation of experiments, the cells were maintained in MEM containing 0.5% FBS for 24 h,

and incubated with or without (control) test agents for 18 h for Northern blot experiments and 30 min for cAMP assays.

Rat-1 cells were obtained from Dr. Robert Weinberg (Whitehead institute, MIT). Cells were grown to confluence in 100x20 mm sterile Falcon tissue culture plates at 37°C in Dulbecco's Modified Eagle Medium (DMEM; Life Technologies Grand Island, NY) containing 5% FBS supplemented with 50units/ml penicillin and 50µg/ml streptomycin. The cells were incubated with or without (control) test agents for 18 h for Northern blot experiments and 30 min for cAMP assays.

#### **cDNA Probe preparation**

**EP1 cDNA isolation from pcDNA-1** For transformation of EP1 cDNA in pcDNA-1 (Invitrogen, Carlsbad, CA), 100µl of competent cells, MC 1061/p3 (Invitrogen), was thawed on ice. Then, 5µl of 0.5M β-mercaptoethanol was added and cells were mixed gently. Subsequently, it was further mixed with 0.5µl of the EP1 cDNA in pcDNA-1. The mixture was incubated on ice for 30 min, and at 42°C for 30 s, then back on ice for 2 min. 900µl of SOC media (Invitrogen) was added to the mixture and it was incubated at the 37°C for 1 h. The mixture was centrifuged for 1 min at 12800x g, and all but 50µl of the supernatant was discarded. The pellet was resuspended in remaining 50µl of supernatant. The transformed cells were plated on 95x15 mm plastic petri dish from Fisher Scientific Co. containing Luria-Bertani (LB) agar supplemented with 10mg/ml tetracyclin (TET) and 5 mg/ml ampicillin (AMP), and were stored overnight at 37°C. A colony was then selected and was grown overnight at 37°C in LB with

TET and AMP. The plasmid was prepared by a miniprep procedure. Briefly, 1.2ml of the overnight prep was centrifuged at 12800x g for 2 min. Then the pellet was resuspended using 300 $\mu$ l of TENS (10mM Tris pH 8.0, 1mM EDTA, 0.1N NaOH and 0.4% SDS) and stored in ice for 5 min. Then, 150 $\mu$ l of 3M NaOAc (pH 5.2) was added and the mixture was stored in ice for 8 min prior to centrifuging for 8 min at 12800x g. The supernatant (approximately 450 $\mu$ l) was transferred to a clean tube and precipitated with 2x volume (approximately 900 $\mu$ l) of 100% EtOH. Then, the mixture was centrifuged at 12800x g for 10 min and the pellet was washed two times with 70% EtOH. The pellet was air dried and resuspended in 40 $\mu$ l of TE (10mM Tris Cl pH 8.0, and 1mM EDTA pH 8.0) containing 10 $\mu$ g/ml RNase. Purified EP1 cDNA in pcDNA-1 was then digested with EcoRI from Stratagene (La Jolla, CA) and electrophoresed on 1x Tris-acetate (TAE) DNA gel (1.2% agarose) along with Lambda DNA-Bst E II digest from New England Biolabs, Inc (Beverly, MA) as a molecular weight standard (1x TAE consists of 0.04M Tris-acetate and 0.001M EDTA). The 1.4-kb band representing EP1 cDNA in the agarose gel was excised and EP1 cDNA was extracted from the gel using the Compass Kit from American Bioanalytical (Natick, MA).

#### **EP2, EP3, EP4, COL, LO and COX-1 cDNA isolation from their vectors**

Samples of each cDNA already transformed in JM109 bacterial cells and stored in 50% glycerol at -80°C were amplified by growing them overnight at 37°C in LB supplemented with AMP (50 $\mu$ g/ml). Then, the plasmids were purified using

miniprep procedure as described for EP1 cDNA isolation. The cDNAs were excised from their vectors using specific restriction endonucleases (EP2 – Sac I and Hind III; EP3 – Hind III and Xba I; EP4 – Bam H I and Xba I; COL – Pst I; LO – Eco RI; and COX-1 – Hind III and Xba I). All the restriction endonucleases were obtained from New England Biolabs, Inc.. The digested plasmids were electrophoresed on 1x TAE gel (1.2% agarose) along with Lambda DNA-Bst E II digest as a molecular weight standard. The bands representing respective cDNAs in the agarose gel were excised and extracted from the gel using Compass Kit.

**Quantitative analysis** 4µl of each purified cDNA was electrophoresed on 1x TAE, DNA gel (1.2% agarose) alongside known concentrations of bradykinin (BK) receptor cDNA to determine the concentration of the newly purified cDNAs. 40ng, 120ng, 200ng, 400ng and 800ng of BK receptor cDNA was run. The intensity of each digested cDNA insert observed through ethidium bromide staining was then compared to the BK receptor cDNA concentrations. From this, the respective stock concentrations for each cDNA from the miniprep were approximated. 30ng of cDNA was used for all probe preparations.

**Northern blot analysis** Total RNA was extracted from cells by the guanidinium thiocyanate-phenol-chloroform procedure according to the methods described previously [Taylor et al., 1992]. To maximize the RNA concentration, two plates (100x20 mm) of cells were used for each type of incubation for IMR-90 and Rat-1 cells. Each plate of cells was treated with 2ml of Solution D (2M guanidinium-

thiocyanate, 0.04M Na Citrate pH7.0, 207 $\mu$ M  $\beta$ -mercapto ethanol and 0.6% Sarcosyl). Cells were scraped into a sterile Fisher test tube. Subsequently, 240 $\mu$ l of 2M NaOAc, 2ml of phenol (pH 4.5) and 960 $\mu$ l of chloroform were added, vortexing after each addition. The mixture was incubated in ice for 20 min and centrifuged at 9500 rpm (10000x g) for 20 min at 4°C using SS-34 Sorvall centrifuge. Then the top layers (approximately 2ml) were carefully transferred to new tubes using 1ml pipette and the duplicates were pooled in one tube. An equal volume of isopropanol was added to each tube and the mixture was stored at -20°C overnight. On the following day, the mixture was centrifuged at 9750 rpm (10780x g) for 20 min at 4°C. The supernatant was carefully removed with glass pipette to prevent any disturbance of the pellet. The pellet was resuspended with 300 $\mu$ l of Solution D with  $\beta$ -mercaptoethanol and transferred to 1.5ml microfuge tubes. 300 $\mu$ l of isopropanol was added and the tubes vortexed. The mixture was left at -20°C overnight. On the following day, the mixture was centrifuged at 12000 rpm (13500x g) for 25 min at 4°C. The supernatant was discarded and the pellet was washed with 1ml of cold 75% EtOH. The mixture was centrifuged for 5 min at 12000 rpm at 4°C and washed again with another 1ml of 75% EtOH. The mixture was centrifuged for 10 min at 12000 rpm (13500x g) at 4°C. EtOH was discarded and the pellet was air dried for approximately 2 min. About 35-40 $\mu$ l of DEPC treated H<sub>2</sub>O was added to each tube and the pellets were left to dissolve on ice for about 2 h before storing in -80°C. For quantitation, 4 $\mu$ l of RNA was added to 996 $\mu$ l of H<sub>2</sub>O and ultraviolet absorbance

was read at 260nm. For RNA gel, 10 $\mu$ g of total RNA was fractionated by electrophoresis on a 1% agarose and 8% formaldehyde denaturing gel. The fractionated RNA was transferred to nylon membranes from NEN Life Science Products (Boston, MA) and fixed by ultraviolet crosslinking. Approximately 30ng of cDNAs for EP receptors (EP1, EP2, EP3 and EP4), COL $\alpha$ 1, LO, and COX-1 were labeled with  $^{32}$ P using Ready-to-Go DNA Labeling beads (-dCTP) from Pharmacia Biotech (Piscataway, NJ). The RNA blots were probed with the  $^{32}$ P-labeled cDNA probes at 65°C in a Rapid-hybridization buffer from Amersham Life Science (Amersham, UK.) and washed with 2x SSC buffer (0.3M NaCl, 0.03M Na Citrate, 1% SDS) at 28°C for low stringency then with 0.2x SSC buffer (0.03M NaCl, 0.003M Na Citrate, 0.1% SDS) at 50°C for high stringency. Kodak Scientific Imaging Film from Eastman Kodak Company (Rochester NY) was exposed to the  $^{32}$ P-labeled nylon membrane in -80°C for autoradiography. Radioactivity associated with each band was quantitated with an Instantimager, an electronic autoradiography program (Packard, Meriden, CT). The relative intensity of hybridization with COX-1 probe in Figure 9 was measured using Sigma Scan/Image Program (Jandel Scientific, San Rafael, CA). The cpm values were corrected to eliminate loading errors as follows. The ethidium bromide stained 18s rRNAs were scanned through a video snapshot program (Snappy, Play Inc., Rancho Cordova, CA). The intensities of the ethidium bromide stained 18s rRNAs were measured using a Sigma Scan/Image Program (Jandel Scientific, San Rafael, CA). The cpm values were then divided by the 18s rRNA

values. To normalize the results, values were expressed as either % of control or fold increase.

**cAMP assay** The cells were labeled with  $^3\text{H}$ -adenine for 24 h. After labeling, cells were washed with DMEM buffered with 20mM HEPES at pH 7.5. The cells were then incubated in 20mM HEPES buffered DMEM containing 100mM 3-isobutyl-1-methylxanthine (IBMX) and with or without test agents for 30 min. Reactions were stopped by aspirating the medium and adding 1ml ice cold 5% TCA containing 1mM ATP and 1mM cAMP. After 30 min incubation on ice, the TCA extract was collected and applied to Dowex columns containing analytical grade resins, AG 1-x8 Resin (100-200 mesh formate form) from Bio-Rad Laboratories (Hercules, CA). The columns were washed with 3ml of water. A total volume of 4ml was collected in scintillation vials which represented the  $^3\text{H}$ -ATP and  $^3\text{H}$ -ADP fractions. Cyclic AMP was then eluted from the Dowex columns directly onto Alumina columns (Sigma Chemical Co., St. Louis, MO) with 10ml water. cAMP was eluted from the Alumina with 6ml 0.1M imidazole, pH 7.5. The radioactivity of the cAMP fraction was quantified by liquid scintillation spectrometry (Wallac, Inc., Gaithersburg, MD).

## **RESULTS**

### **Effect of PGE<sub>2</sub> on COL $\alpha$ 1(I) mRNA in IMR-90**



In a concentration dependent manner, PGE<sub>2</sub> reduced COL $\alpha$ 1(I) mRNA levels in IMR-90 cells. This is illustrated in Figure 1. The COL $\alpha$ 1(I) mRNA level was reduced at 1nM, and further decreased in a stepwise manner up to 1 $\mu$ M. Then, from 1 $\mu$ M to 100 $\mu$ M, the COL $\alpha$ 1(I) mRNA level appeared to show no further change. Therefore, PGE<sub>2</sub> induced the maximal inhibition on COL $\alpha$ 1(I) at 1 $\mu$ M.

### **EP Receptor Profile in IMR-90**

PGE<sub>2</sub> effects cellular response using the four EP receptors, EP1, EP2, EP3 and EP4 [Coleman et al., 1994; Ichikawa et al., 1996]. Thus, the EP receptor profile in IMR-90 cells was examined (Figure 2). The total RNA was hybridized with EP1, EP2, EP3 and EP4 receptor probes. The expected sizes were approximately 1.6-kb for EP1 [Funk et al., 1993], 3.1-kb for EP2 [Regan et al., 1994], 2.4-kb for EP3 [Regan et al., 1994] and 3.8-kb for EP4 [An et al., 1993]. As shown in Figure 2, the bands for approximate sizes of EP2, EP3 and EP4 were detected. Among these, the EP2 receptor expression appeared the most prominent while the EP3 receptor expression was the least. EP1 receptor expression was not detected. Narumiya (1994) reported similar results in mouse lung in which the expression of EP2 mRNA was predominant compared to the other EP receptors.

Figure 1

Effect of  $\text{PGE}_2$  on  $\text{COL}\alpha 1(\text{I})$  mRNA in IMR-90




## Legend for Figure 1

IMR-90 cells were made quiescent as described in the methods section. The cells were then untreated (control) or treated with the respective concentrations of PGE<sub>2</sub> for 18 h. Total RNA isolated from 2 separately treated 100x20mm culture plates under the same conditions was combined. 10µg of total RNA was fractionated by electrophoresis and transferred to nylon membrane as described in the methods section. 18s rRNA represents the loading. Then, RNA was hybridized with COLα1(I) cDNA. PGE<sub>2</sub> concentrations are expressed as log of molar concentration. C = control.

Figure 2

Northern Blot Analysis of EP receptors  
in IMR-90



## Legend for Figure 2

Northern blot analysis was performed on 10 $\mu$ g of total RNA isolated from confluent, quiescent, untreated IMR-90 cells. Each lane was probed with specific receptor cDNA. The positions of the 28s and 18s rRNAs are marked with solid arrow heads. The hybridizing transcripts of EP2, EP3 and EP4 are marked with solid-lined arrows (→). The dotted-lined arrow (---→) represents the expected 1.6-kb hybridizing transcript of EP1 receptor which was reported by Funk et al (1993) and Fedyk and Phipps (1996) in human erythroleukemia cells. Autoradiography was performed at -80°C. The radiogram is a representative of results obtained from three separate experiments.

### **Effect of 11-deoxy PGE<sub>1</sub> and Sulprostone on COL $\alpha$ 1(I), LO and COX-1**


#### **mRNA in IMR-90**

In order to identify the type of EP receptor that was mediating PGE<sub>2</sub> response in IMR-90 cells, the EP2/EP4 agonist, 11-deoxy PGE<sub>1</sub>, and EP3/EP1 agonist, sulprostone, were used. The effects of 11-deoxy PGE<sub>1</sub> and sulprostone were examined with respect to COL $\alpha$ 1(I) as well as LO and COX-1 mRNA levels.

The Figures 3 - 5 show the effect of varying concentrations (10nM-10 $\mu$ M) of 11-deoxy PGE<sub>1</sub> on COL $\alpha$ 1(I) (Figure 3), LO (Figure 4) and COX-1 (Figure 5) gene expression, respectively. 11-deoxy PGE<sub>1</sub> at 10 $\mu$ M decreased the COL $\alpha$ 1(I) mRNA level the most, decreasing it by more than 50% from the control. 1 $\mu$ M of 11-deoxy PGE<sub>1</sub> also decreased COL $\alpha$ 1(I) mRNA level, but to a lesser degree. However, 100nM of 11-deoxy PGE<sub>1</sub> had no apparent effect while 10nM increased COL $\alpha$ 1(I) mRNA level by approximately 30%. In Figure 4, 11-deoxy PGE<sub>1</sub> at 10 $\mu$ M again had the strongest effect in decreasing LO mRNA level. 10 $\mu$ M of 11-deoxy PGE<sub>1</sub> decreased LO mRNA level by 50%. The 11-deoxy PGE<sub>1</sub> concentrations (1 $\mu$ M - 10nM) also decreased LO mRNA levels in a concentration dependent manner and did not show the biphasic effect observed with COL $\alpha$ 1(I) shown in Figure 3. From 1 $\mu$ M to 10nM, there was a stepwise decrease in the 11-deoxy PGE<sub>1</sub>'s inhibitory effect on LO mRNA levels. The effect of 11-deoxy PGE<sub>1</sub> concentrations (1 $\mu$ M - 10nM) on COX-1 mRNA level was opposite from its effect on LO (Figure 5). This figure shows 11-deoxy PGE<sub>1</sub>

Figure 3

Effect of 11-deoxy PGE<sub>1</sub> on COL $\alpha$ 1(I) mRNA in IMR-90




## Legend for Figure 3

IMR-90 cells were untreated (control) or treated with 11-deoxy PGE<sub>1</sub> (10μM, 1μM, 100nM and 10nM) for 18 h in MEM containing 0.5% FBS. Total RNA was then isolated as described in the methods section. RNA from two separate incubations of the same condition was then combined. 10μg of total RNA was fractionated by electrophoresis and transferred to nylon membrane as described in the methods section. Then, RNA was hybridized with COLα1(I) cDNA. 18s rRNA represents the loading. Control #1 media is treated with 0.1% ethanol while Control #2 is treated with 0.01% ethanol. The controls were treated with ethanol, for 11-deoxy PGE<sub>1</sub> stock was dissolved in ethanol. There are two controls, for the amount of ethanol in 10μM was different than the amounts for 1μM - 1nM.

Figure 4

Effect of 11-deoxy PGE<sub>1</sub> on LO mRNA in IMR-90





## Legend for Figure 4

IMR-90 cells were untreated (control) or treated with 11-deoxy PGE<sub>1</sub> (10 $\mu$ M, 1 $\mu$ M, 100nM and 10nM) for 18 h in MEM containing 0.5% FBS. Total RNA was then isolated as described in methods. RNA from two separate incubations of the same condition was then combined. 10 $\mu$ g of total RNA was fractionated by electrophoresis and transferred to nylon membrane as described in the methods section. Then, RNA was hybridized with LO cDNA. 18s rRNA represents the loading control. Control #1 media is treated with 0.1% ethanol while Control #2 is treated with 0.01% ethanol. The controls were treated with ethanol, for 11-deoxy PGE<sub>1</sub> stock was dissolved in ethanol. There are two controls, for the amount of ethanol in 10 $\mu$ M was different than the amounts for 1 $\mu$ M - 1nM.

Figure 5

Effect of 11-deoxy PGE<sub>1</sub> on COX-1 mRNA in IMR-90



## Legend for Figure 5

IMR-90 cells were untreated (control) or treated with 11-deoxy PGE<sub>1</sub> (10μM, 1μM, 100nM and 10nM) for 18 h in MEM containing 0.5% FBS. Total RNA was then isolated as described in methods. RNA from two separate incubations of the same condition was then combined 10μg of total RNA was fractionated by electrophoresis and transferred to nylon membrane as described in the methods section. Then, RNA was hybridized with COX-1 cDNA. 18s rRNA represents loading control. Control #1 media is treated with 0.1% ethanol while Control #2 is treated with 0.01% ethanol. The controls were treated with ethanol, for 11-deoxy PGE<sub>1</sub> stock was dissolved in ethanol. There are two controls, for the amount of ethanol in 10μM was different than the amounts for 1μM - 1nM.


increased COX-1 mRNA levels in a concentration dependent manner in which the effect was maximal at 10 $\mu$ M and minimal at 10nM. This figure also did not show the biphasic effect seen in Figure 3. Two different controls were used for Figures 3 - 5. Control #1 was for 10 $\mu$ M 11-deoxy PGE<sub>1</sub> while Control #2 was for 11-deoxy PGE<sub>1</sub> concentrations of 1 $\mu$ M - 10nM. The only difference in the composition of these control vehicles was that there was more ethanol in Control #1 than Control #2. There was some difference between Control #1 and Control #2 in their basal mRNA levels for all three genes. Figure 6 shows the effect of sulprostone on COL $\alpha$ 1(I), LO and COX-1 mRNA levels, respectively. In contrast to the effects seen with 11-deoxy PGE<sub>1</sub>, sulprostone (1nM-1 $\mu$ M) appeared to have little or no effect on the three genes.

#### **Effect of Forskolin on COL $\alpha$ 1(I), LO and COX-1 mRNA in IMR-90**

The EP2 and EP4 receptors have been reported to be associated with the cAMP signaling path [Scutt et al., 1995]. Therefore, the effect of forskolin, an adenylyl cyclase activator, was compared to that of PGE<sub>2</sub> on the three genes using IMR-90 cells. These results are shown in Figures 7 - 9. In Figure 7, PGE<sub>2</sub> (1 $\mu$ M) or forskolin (25 $\mu$ M), showed a comparable effect in limiting COL $\alpha$ 1(I) mRNA levels. PGE<sub>2</sub> or forskolin decreased COL $\alpha$ 1(I) mRNA levels by 50%. Interestingly, when 1 $\mu$ M PGE<sub>2</sub> was co-incubated with 25 $\mu$ M forskolin, the decreasing effect on the COL $\alpha$ 1(I) mRNA level was much greater than the effect with either PGE<sub>2</sub> or forskolin, alone. Similarly, as shown in Figure 8, forskolin

Figure 6

Effect of Sulprostone on COL $\alpha$ 1(I), LO  
and COX-1 mRNA in IMR-90





## Legend for Figure 6

IMR-90 cells were untreated (control) or treated with sulprostone ( $1\mu\text{M}$ ,  $100\text{nM}$  and  $10\text{nM}$ ) for 18 h in MEM containing 0.5% FBS. Total RNA was then isolated as described in methods. RNA from two separate incubations of the same condition was then combined.  $10\mu\text{g}$  of total RNA was fractionated by electrophoresis and transferred to nylon membrane as described in the methods section. Then, RNA was hybridized with COL $\alpha$ 1(I), LO and COX-1 cDNA. 18s rRNA represents the loading control. CONTL = control.  $1\mu\text{M}$  -  $1\text{nM}$  = sulprostone concentrations.

Figure 7

Effect of  $\text{PGE}_2$  and Forskolin on  
 $\text{COL}\alpha 1(\text{I})$  mRNA in IMR-90




## Legend for Figure 7

IMR-90 cells were made quiescent as described in the methods section. The cells were untreated (control) or treated with PGE<sub>2</sub> (1μM) and/or forskolin (25μM) for 18 h. Total RNA was isolated as described in the methods section. 10μg of total RNA was fractionated by electrophoresis and transferred to nylon membrane as described in the methods section. Then, RNA was hybridized with COLα1(I) cDNA. Northern blot analysis is a representative of three separate experiments. 18s rRNA represents the loading control.

Figure 8

Effect of PGE<sub>2</sub> and Forskolin on  
LO mRNA in IMR-90





## Legend for Figure 8

IMR-90 cells were made quiescent as described in the methods section. The cells were untreated (control) or treated with PGE<sub>2</sub> (1μM) and/or forskolin (25μM) for 18 h. Total RNA was isolated. 10μg of total RNA was fractionated by electrophoresis and transferred to nylon membrane as described in the methods section. Then, RNA was probed with LO cDNA. Northern blot analysis is a representative of three separate experiments. 18s rRNA represents the loading control.

Figure 9

Effect of PGE<sub>2</sub> and Forskolin on  
COX-1 mRNA in IMR-90



## Legend for Figure 9

IMR-90 cells were made quiescent as described in the methods section. The cells were untreated (control) or treated with PGE<sub>2</sub> (1μM) and/or forskolin (25μM) for 18 h. Total RNA was isolated. 10μg of total RNA was fractionated by electrophoresis and transferred to nylon membrane as described in the methods section. Then, RNA was probed with COX-1 cDNA. Northern blot analysis is a representative of three separate experiments. 18s rRNA represents the loading control.


(25 $\mu$ M) also decreased LO mRNA levels by approximately 25%, while PGE<sub>2</sub> (1 $\mu$ M) limited LO mRNA slightly more than forskolin. However, no additive effect was observed when the two were combined. In Figure 9, forskolin (25 $\mu$ M) increased COX-1 mRNA levels by more than 50% from the control level. PGE<sub>2</sub> (1 $\mu$ M) increased COX-1 mRNA level by approximately 25%. The co-incubation of PGE<sub>2</sub> and forskolin increased COX-1 mRNA level that was slightly greater than the effect of forskolin. The co-incubation of PGE<sub>2</sub> and forskolin increased COX-1 mRNA levels almost 2-fold above the control.

#### **Cyclic AMP Assay for IMR-90 with Forskolin, PGE<sub>2</sub> and 11-deoxy PGE<sub>1</sub>**

One explanation for the greater inhibition by PGE<sub>2</sub> in the presence of forskolin was that the dose of either effector alone did not maximize the cAMP levels. This hypothesis was tested with cAMP assay (Figure 10). As shown, forskolin at 2.5 $\mu$ M increased cAMP levels by approximately 5-fold. The increase in cAMP levels paralleled the increase in forskolin concentration. Although forskolin at 250 $\mu$ M increased the cAMP levels the most, it was not much different from the effect at 100 $\mu$ M. In any case, PGE<sub>2</sub> at 1 $\mu$ M also produced sub-optimal increases in cAMP levels. As expected 1 $\mu$ M of PGE<sub>2</sub>, co-incubated with 25 $\mu$ M of forskolin, generated the maximal cAMP levels that was comparable to forskolin at 250 $\mu$ M. Finally, since 11-deoxy PGE<sub>1</sub> demonstrated similar effects as PGE<sub>2</sub> on the three genes of interest, 1 $\mu$ M of 11-deoxy PGE<sub>1</sub> was tested to determine its

Figure 10

Effect of Forskolin, PGE<sub>2</sub> and 11-deoxy PGE<sub>1</sub>  
on cAMP Production in IMR-90



## Legend for Figure 10

IMR-90 cells were made quiescent and labeled with  $^3\text{H}$ -adenine as described in the methods section. The cells were treated with IBMX and with or without (control) varying concentrations of forskolin (2.5 $\mu\text{M}$  - 250 $\mu\text{M}$ ), PGE<sub>2</sub> (1 $\mu\text{M}$ ) with or without forskolin (25 $\mu\text{M}$ ), or 11-deoxy PGE1 (1 $\mu\text{M}$ ) for 30 min.  $^3\text{H}$ -cAMP was eluted and quantitated using liquid scintillation spectrometry. Values for the experiment are the means  $\pm$  standard error of three experiments.

CONTL = control.

effect on cAMP. 11-deoxy PGE<sub>1</sub> increased cAMP levels, but its effect was slightly less than PGE<sub>2</sub> (1μM) or forskolin (25μM) alone.

#### **Cyclic AMP Assay for IMR-90 with PGE<sub>2</sub> Concentrations**


Since the effect of 1μM of PGE<sub>2</sub> was sub-optimal as shown in Figure 10, the investigator attempted to determine the maximal PGE<sub>2</sub> concentration required for optimal cAMP production in IMR-90 cells (Figure 11). The cAMP levels began to increase even at 1nM of PGE<sub>2</sub>. Then, the levels began to plateau at 100nM and were maximized at 1μM. The half-maximal effective dose was  $3.32 \times 10^{-8}$  nM.

#### **Effect of 11-deoxy PGE<sub>1</sub> on TGF-β and IL-1β stimulated COLα1(I) and LO mRNA levels, respectively, in IMR-90**

Since 11-deoxy PGE<sub>1</sub> demonstrated similar effects as PGE<sub>2</sub> on the three genes of interest and on cAMP levels, 11-deoxy PGE<sub>1</sub> was examined to see if it can also limit the cytokine induced increases in COLα1(I) and LO mRNA levels (Figure 12 and 13, respectively). As shown in Figure 12, TGF-β increased the COLα1(I) mRNA levels by about 50%. 11-deoxy PGE<sub>1</sub> decreased the basal and the TGF-β stimulated COLα1(I) mRNA levels by about 50% and more than 60%, respectively. PGE<sub>2</sub> showed almost identical inhibitory effects on the basal and TGF-β stimulated COLα1(I) mRNA levels as 11-deoxy PGE<sub>1</sub>. In Figure 13, IL-1β stimulated LO mRNA levels by more than 2-fold from the control. 11-deoxy PGE<sub>1</sub> or PGE<sub>2</sub> decreased the basal LO mRNA levels by about 50%. In addition, 11-deoxy PGE<sub>1</sub> decreased the IL-1β stimulated increase in LO mRNA levels by

Figure 11

Effect of  $\text{PGE}_2$  on cAMP Production in IMR-90





## Legend for Figure 11

IMR-90 cells were made quiescent and labeled with  $^3\text{H}$ -adenine as described in the methods section. The cells were treated with IBMX and with or without (control) varying concentrations of  $\text{PGE}_2$  (1nM - 100 $\mu\text{M}$ ) for 30 min.  $^3\text{H}$ -cAMP was eluted as described in the methods section and quantitated using liquid scintillation spectrometry. The values for the experiment are the means  $\pm$  standard error of three experiments. C = control. 11-DP = 11-deoxy  $\text{PGE}_1$ .

Figure 12

Effect of TGF- $\beta$ , 11-deoxy PGE<sub>1</sub> and PGE<sub>2</sub> on COL $\alpha$ 1(I) mRNA in IMR-90





## Legend for Figure 12

IMR-90 cells were grown to confluence and made quiescent as described in the methods section. The cells were untreated (control) or treated with PGE<sub>2</sub> (1μM) or 11-deoxy PGE<sub>1</sub> (1μM) with or without TGFβ (100ng/ml) for 18 h. Total RNA isolated from 2 separately treated 100x20mm culture plates was combined. 10μg of total RNA was electrophoresed and transferred to nylon membrane as described in the methods section. Then, the RNA was hybridized with COLα1(I) cDNA. 18s rRNA represents the loading control.

Figure 13

Effect of IL-1 $\beta$ , 11-deoxy PGE<sub>1</sub> and PGE<sub>2</sub> on  
LO mRNA in IMR-90



## Legend for Figure 13

IMR-90 cells were grown to confluence and made quiescent as described in the methods section. The cells were untreated (control) or treated with PGE<sub>2</sub> (1μM) or 11-deoxy PGE<sub>1</sub> (1μM) with or without IL-1β (100pg/ml) for 18 h. Total RNA isolated from 2 separately treated 100x20mm culture plates was combined. 10μg of total RNA was fractionated by electrophoresis and transferred to nylon membrane as described in the methods section. Then, the RNA was hybridized with LO cDNA. 18s rRNA represents loading control.

more than 50%. PGE<sub>2</sub> also decreased the IL-1 $\beta$  stimulated increase in LO mRNA levels to the same extent as 11-deoxy PGE<sub>1</sub>.

#### **EP Receptor Profile in Rat-1 Cells**


To further test the role of PGE<sub>2</sub> or cAMP in regulating COL $\alpha$ 1(I) mRNA levels, another type of cells, Rat-1 cells were tested. These cells actively produce collagen [Granot et al., 1993]. To determine the EP receptor subtype profile in this cell type, total RNA was hybridized with EP1, EP2, EP3 and EP4 receptor probes. Results are shown in Figure 14. Unlike the IMR-90, Rat-1 cells do not express detectable EP1, EP2 or EP3 mRNAs. However, some EP4 receptor mRNA was detected as shown in the figure.

#### **Effect of PGE<sub>2</sub> on COL $\alpha$ 1(I) in Rat-1 Cells**

Rat-1 cells were incubated with PGE<sub>2</sub> (1 $\mu$ M) or forskolin (25 $\mu$ M) and probed for COL $\alpha$ 1(I) mRNA. Results are shown in Figure 15. In contrast to the effects observed in IMR-90 cells, PGE<sub>2</sub> slightly increased COL $\alpha$ 1(I) mRNA levels. However, forskolin (25 $\mu$ M) increased COL $\alpha$ 1(I) mRNA levels by approximately 80%.

Figure 14

Northern Blot Analysis of EP receptors  
in Rat-1 Cells





## Legend for Figure 14

Northern blot analysis was performed on 10 $\mu$ g of total RNA isolated from confluent, untreated Rat-1 cells. Each lane was probed with specific type of cDNA. The positions of the 28s and 18s rRNAs are marked with solid arrow heads. The hybridizing transcript of EP4 receptor is marked with solid-lined arrows ( $\rightarrow$ ). No band was detected when the blot was probed with for EP1, EP2 or EP3 receptor cDNAs. Autoradiography was performed at -80°C. The radiogram is a representative of results obtained from two separate experiments.

Figure 15

Effect of PGE<sub>2</sub> and Forskolin on  
COL $\alpha$ 1(I) mRNA in Rat-1 Cells



## Legend for Figure 15

Rat-1 cells grown to confluence were untreated (control) or treated with PGE<sub>2</sub> (1μM) or forskolin (25μM) for 18 h. Total RNA isolated from 2 separately treated 100x20mm culture plates was combined. 10μg of total RNA was fractionated by electrophoresis and transferred to nylon membrane as described in the methods section. Then, the RNA was hybridized with COLα1(I) cDNA probe. The northern blot analysis is a representative of two separate experiments. 18s rRNA represents the loading control.


**Cyclic AMP Assay for Rat-1 Cells with Forskolin and PGE<sub>2</sub>**

The effects of forskolin and PGE<sub>2</sub> in Rat-1 cells were also examined with respect to cAMP production (Figure 16). PGE<sub>2</sub> (1 $\mu$ M) had no effect on cAMP levels when compared to the control levels. However, forskolin increased the cAMP levels by approximately 10-fold from the control level.

In summary, EP2 receptor expression was predominant in IMR-90 and only EP4 was detected in Rat-1 cells. In IMR-90, 11-deoxy PGE<sub>1</sub> mimicked the effects of PGE<sub>2</sub> on the three genes to include the cytokine induced effects on COL $\alpha$ 1(I) and LO mRNA levels. Further, forskolin, an adenylyl cyclase activator, demonstrated very similar effects on the three genes as PGE<sub>2</sub> and 11-deoxy PGE<sub>1</sub>. However, the effects of PGE<sub>2</sub> and forskolin on cAMP production in IMR-90 were different than in Rat-1 cells. Also, the maximal level of cAMP produced by PGE<sub>2</sub> was different from that of forskolin.

Figure 16

Effect of Forskolin and  $\text{PGE}_2$   
on cAMP Production in Rat-1 Cells



## Legend for Figure 16

Rat-1 cells were labeled with  $^3\text{H}$ -adenine as described in the methods section. The cells were treated with IBMX and with or without (control)  $\text{PGE}_2$  ( $1\mu\text{M}$ ) or forskolin ( $25\mu\text{M}$ ) for 30 minutes.  $^3\text{H}$ -cAMP was eluted and quantitated as described in the methods section. Values are the means  $\pm$  standard error of three experiments.

## **DISCUSSION**

The results illustrate that human embryonic lung fibroblasts, IMR-90, express EP2, EP3 and EP4 receptor mRNAs, but no detectable level of EP1 mRNA. Among these, the EP2 receptor mRNA is predominant while EP4 and EP3 receptor mRNAs are expressed to a lesser degree with EP3 showing the least amount. The receptor types were stimulated with agonists, sulprostone for EP3/EP1 receptors and 11-deoxy PGE<sub>1</sub> for EP2/EP4 receptors, to determine their functions with respect to PGE<sub>2</sub> inhibition/stimulation of COL $\alpha$ 1(I), LO and COX-1 mRNA expressions. Sulprostone did not show any effect on the expression of the three genes of interest. On the other hand, 11-deoxy PGE<sub>1</sub> at 1 $\mu$ M decreased the basal levels of COL $\alpha$ 1(I) and LO mRNA, and increased the COX-1 mRNA basal level. These results were comparable to the effects observed with PGE<sub>2</sub> at 1 $\mu$ M. Interestingly, 11-deoxy PGE<sub>1</sub> at a lower concentration increased the basal level of COL $\alpha$ 1(I) mRNA, only. Additionally, 11-deoxy PGE<sub>1</sub> abrogated the TGF- $\beta$  and IL-1 $\beta$  induced COL $\alpha$ 1(I) and LO mRNA levels, respectively. These results strongly suggest that EP2 and possibly EP4 receptors are key mediators of PGE<sub>2</sub> in the regulation of COL $\alpha$ 1(I), LO and COX-1 gene expressions in IMR-90 cells.

The EP2 [Regan et al., 1994] and EP4 [An et al., 1993; Bastien et al., 1994] receptors have been reported to increase cAMP. Therefore, we examined the effects of a known cAMP stimulator, forskolin, on COL $\alpha$ 1(I), LO and COX-1 mRNA expressions. Forskolin not only limited the basal COL $\alpha$ 1(I) and LO mRNA

levels, but increased the basal COX-1 mRNA levels. The effect was comparable to the effect of PGE<sub>2</sub>. Interestingly, the co-incubation of forskolin and PGE<sub>2</sub> in IMR-90 limited COL $\alpha$ 1(I) mRNA production to a greater degree. However, PGE<sub>2</sub> and forskolin together did not have any additive effects on LO or COX-1 mRNA levels.

It is possible that the synergy between PGE<sub>2</sub> and forskolin on COL $\alpha$ 1(I) mRNA expression is due to the sub-optimal stimulation of cAMP by either agent alone. This hypothesis appears to be true according to results shown in Figure 10. In this figure, 1 $\mu$ M of PGE<sub>2</sub> or 25 $\mu$ M of forskolin produced sub-optimal levels of cAMP while their combined effects produced the maximal level. However, when cAMP levels were measured in response to increasing PGE<sub>2</sub> concentrations, 1 $\mu$ M of PGE<sub>2</sub> produced the maximal levels of cAMP as shown in Figure 11.

The comparison of the two results suggests that forskolin and PGE<sub>2</sub> may be inducing cAMP production and subsequent downstream effectors in different manner. This may be possible by targeting different isoforms of cAMP signaling components such as adenylate cyclase, cAMP PDE or PKA. To date, nine isoforms of adenylate cyclases and approximately 30 isoforms of cAMP PDE have been reported in mammalian cells [Houslay and Milligan, 1997].

Forskolin has been reported to activate all but one isoform of adenylate cyclases. The adenylate cyclase isoform(s) involved in EP2 or EP4 receptors signaling have not been determined yet. It is possible that the two receptors

interact with more limited number of adenylate cyclase isoforms, which could be co-located with the receptors, than forskolin. Further, it is possible that the cAMP signaling components (adenylate cyclase, cAMP PDE, PKA) that are specific for the EP receptors could be maintaining a certain threshold level of cAMP for the receptors' signaling mechanisms. Such a threshold level may be reason for EP receptors to produce less cAMP levels than forskolin.

In 1996, Jurevicius and Fischmeister reported the co-localization of phosphodiesterase with the cAMP signaling cascade of beta-adrenergic receptors in frog ventricular cells. Consequently, the intracellular cAMP level was not uniform throughout the cell whenever the receptor signaling cascade was activated. This suggests that the cAMP signaling components are not only specific but also spatially organized. Further, the additive effect of PGE<sub>2</sub> and forskolin in suppression of COL $\alpha$ 1(I) mRNA level, shown in Figure 7, seems to indicate that inhibition of COL $\alpha$ 1(I) gene expression could be induced by more than one type of cAMP signaling mechanism.

Recent publications indicate that EP4 and EP2 similarly bind at high affinity with PGE<sub>1</sub>, PGE<sub>2</sub>, 11-deoxy PGE<sub>1</sub> and 16, 16-dimethyl-PGE<sub>2</sub> [Kiriya et al., 1997]. Therefore, it is possible that in IMR-90, both EP2 and EP4 receptors are active in the mechanism limiting the production and deposition of collagen. However, there are some important structural and functional differences between these two receptors. For example, in the cloned mouse EP4 receptors, there are about 3.5 fold more serine and threonine residues in the C-terminal than in the

mouse EP2 receptors [Nishigaki et al., 1996]. Also, EP4 receptor contains potential PKA phosphorylation sites whereas none has been found in EP2 receptor [Nishigaki et al., 1996]. Such phosphorylation sites may be critical in receptor desensitization [Liggett et al., 1993] and subsequently, the termination of signal transduction. Nishigaki et al (1996) demonstrated a difference in PGE<sub>2</sub> induced-desensitization between EP4 and EP2 receptors in CHO cells. In this experiment, PGE<sub>2</sub> induced EP4 receptors to undergo rapid short-term desensitization while it had no such effect on EP2 receptors. Importantly, EP2 receptors generated a higher adenylate cyclase activity than EP4 receptors a in time dependent manner. Also, EP2 receptor proved more responsive to 15-keto-PGE<sub>2</sub>, a PGE<sub>2</sub> metabolite, than EP4 in stimulating adenylate cyclase. Interestingly, 15-keto-PGE<sub>2</sub> is metabolized by 15-hydroxy PG dehydrogenase which is an enzyme that is abundant in the lung [Anggard et al., 1971]. Hence, EP2 receptor mRNA being the predominately expressed gene, and more capable in producing cAMP than EP4 receptor, is most likely responsible for the inhibitory effect of PGE<sub>2</sub> on COL $\alpha$ 1(I) and LO mRNA levels in IMR-90.

Rat-1 cells are active producers of the COL $\alpha$ 1(I) peptide [Granot et al., 1993]. Examination of PGE<sub>2</sub> receptor types in these cells shows that they express EP4 mRNA but lack detectable EP1, EP2 and EP3 mRNA. These cells were treated with PGE<sub>2</sub> or forskolin to examine whether either effected COL $\alpha$ 1(I) mRNA and cAMP levels. Surprisingly, forskolin increased the COL $\alpha$ 1(I) mRNA levels. PGE<sub>2</sub> also had a different effect in Rat-1 cells than in IMR-90. PGE<sub>2</sub>

slightly increased COL $\alpha$ 1(I) mRNA level in Rat-1 cells. Furthermore, PGE<sub>2</sub> had no effect on cAMP levels with 30 min incubation, while forskolin increased it by 10-fold. Interestingly in the Rat-1 cells, EP4 receptors are predominant. These results further suggest that EP4 is a poor inducer of cAMP production which could not be detected after 30m incubation with PGE<sub>2</sub>. These results strengthen the hypothesis that the EP2 receptors are the receptors mediating PGE<sub>2</sub> effects on the COL $\alpha$ 1(I), LO and COX1 gene expression.

Further, 11-deoxy PGE<sub>1</sub>, at low concentration (10nM) in IMR-90 (Figure 3) increased COL $\alpha$ 1(I) mRNA level. One explanation for this is that at low concentrations, 11-deoxy PGE<sub>1</sub> could be more potent for EP4 receptor which may be linked to a mechanism involved in increasing the COL $\alpha$ 1(I) mRNA level. In fact, the K<sub>i</sub> of EP4 and EP2 for 11-deoxy PGE<sub>1</sub> is 23nM and 45nM, respectively, while K<sub>i</sub> for PGE<sub>2</sub> is 1.9nM and 12nM, respectively [Kiriyaama et al., 1997]. These lower K<sub>i</sub>'s could possibly cause the EP4 receptors become activated at low concentrations.

Although both EP2 and EP4 receptors stimulate cAMP synthesis, they may effect different isoforms of cAMP signaling components such as PKA isoforms, thus eliciting different cellular responses. The type II regulatory subunit (RII) of PKA is widely known to be associated with an A-kinase anchoring protein, AKAP, and is insoluble [Johnson et al., 1997; Visconti et al., 1997; Feliciello et al., 1997; Pawson and Scott, 1997; Dell'Acqua and Scott, 1997]. According to Visconti et al. (1997), dibutyryl cAMP, a cAMP analog, did not

dissociate the catalytic subunit associated with the RII in cauda epididymal mouse sperm. If EP2 receptor generated cAMP targets RII, this may explain why dibutyryl cAMP did not induce any decrease in COL $\alpha$ 1(I) mRNA level, reported by Fine et al. (1992).

In addition, previous studies using MC3T3-E1, a mouse osteoblast-like cell line, showed that forskolin and other agents which increase cAMP levels increase COL synthesis [Hakeda et al., 1987]. On the other hand, an elevation of cAMP in UMR 106-06 rat osteosarcoma cells was shown to decrease collagen synthesis [Iida-Klein et al., 1992; Pun 1989]. The understanding of these apparently contradictory effects of cAMP clearly warrants further investigation and may ultimately prove important in the role of PGE<sub>2</sub> as a regulator of collagen synthesis and the fibrotic response.

It appears that limitation by PGE<sub>2</sub> of collagen production and its deposition as an extracellular matrix involves the EP2 receptor with cAMP at the core of this signal transducing mechanism. These results may be pertinent to mechanisms related to tissue scarring or connective tissue-linked diseases. In fact, the limiting by PGE<sub>2</sub> of cytokine stimulated collagen deposition and the failure or reversal of this mechanism could prove critical to the fibrotic sequel of inflammation and in certain connective tissue diseases.

## **REFERENCES**

1. Alheim K, Chai Z, Fantuzzi G, Hasanvan H, Malinowsky D, Di Santo E, Ghezzi P, Dinarello CA, Bartfai T (1997): Hyperresponsive febrile reactions to interleukin (IL) 1alpha and IL-1beta, and altered brain cytokine mRNA and serum cytokine levels, in IL-1beta-deficient mice. *Proc Natl Acad Sci U S A* 94:2681-2686.
2. An S, Yang J, Xia M, Goetzl EJ (1993): Cloning and expression of the EP2 subtype of human receptors for prostaglandin E2. *Biochem Biophys Res Commun* 197:261-270.
3. An S, Yang J, So SW, Zeng L, Goetzl EJ (1994): Isoforms of the EP3 subtype of human prostaglandin E2 receptor transduce both intracellular calcium and cAMP signals. *Biochemistry* 33:14496-14502.
4. Anggard E, Larsson C, Samuelsson B (1971): The distribution of 15-hydroxy prostaglandin dehydrogenase and prostaglandin-delta 13-reductase in tissues of the swine. *Acta Physiol Scand* 81:396-404.
5. Appleby SB, Ristimaki A, Neilson K, Narko K, Hla T (1994): Structure of the human cyclo-oxygenase-2 gene. *Biochem J* 302( Pt 3):723-727.
6. Arend WP, Welgus HG, Thompson RC, Eisenberg SP (1990): Biological properties of recombinant human monocyte-derived interleukin 1 receptor antagonist. *J Clin Invest* 85:1694-1697.
7. Arend WP, Malyak M, Smith MF Jr, Whisenand TD, Slack JL, Sims JE, Giri JG, Dower SK (1994): Binding of IL-1 alpha, IL-1 beta, and IL-1 receptor antagonist by soluble IL-1 receptors and levels of soluble IL-1 receptors in synovial fluids. *J Immunol* 153:4766-4774.
8. Armendariz-Borunda J, Katayama K, Seyer JM (1992): Transcriptional mechanism of type I collagen gene expression are differentially regulated by interleukin-1 $\beta$ , tumor necrosis factor  $\alpha$ , and transforming growth factor- $\beta$  in Ito cells. *J Biol Chem* 267:14316-14321.
9. Auron PE, Webb AC, Rosenwasser LJ, Mucci SF, Rich A, Wolff SM, Dinarello CA (1984): Nucleotide sequence of human monocyte interleukin 1 precursor cDNA. *Proc Natl Acad Sci U S A* 81:7907-7911.
10. Bastien L, Sawyer N, Grygorczyk R, Metters KM, Adam M (1994): Cloning, functional expression, and characterization of the human prostaglandin E2 receptor EP2 subtype. *J Biol Chem* 269:11873-11877.

11. Baum BJ, Moss J, Breul SD, Crystal RG (1978): Association in normal human fibroblasts of elevated levels of adenosine 3':5'-monophosphate with a selective decrease in collagen production. *J Biol Chem* 253:3391-3394.
12. Bhatnagar R, Penfornis H, Mauviel A, Loyau G, Saklatvala J, Pujol JP (1986): Interleukin-1 inhibits the synthesis of collagen by fibroblasts. *Biochem Int* 13:709-720.
13. Bienkowski RS and Gotkin MG (1995): Control of collagen deposition in mammalian lung. *Proc Soc Exp Biol Med* 209:118-140.
14. Birk DE and Silver FH (1984): Collagen fibrillogenesis in vitro: comparison of types I, II, and III. *Arch Biochem Biophys* 235:178-185.
15. Black RA, Kronheim SR, Cantrell M, Deeley MC, March CJ, Prickett KS, Wignall J, Conlon PJ, Cosman D, Hopp TP (1988): Generation of biologically active interleukin-1 beta by proteolytic cleavage of the inactive precursor. *J Biol Chem* 263:9437-9442.
16. Bornstein P (1974): The biosynthesis of collagen. *Annu Rev Biochem* 43:567-603.
17. Brand T and Schneider MD (1995): Inactive type II and type I receptors for TGF beta are dominant inhibitors of TGF beta-dependent transcription. *J Biol Chem* 270:8274-8284.
18. Breen E, Shull S, Burne S, Absher M, Kelley J, Phan S, Cutroneo KR (1992): Bleomycin regulation of transforming growth factor-beta mRNA in rat lung fibroblasts. *Am J Respir Cell Mol Biol* 6:146-152.
19. Brilla CG, Zhou G, Rupp H, Maisch B, Weber KT (1995): Role of angiotensin II and prostglandin E sub 2 in regulating cardiac fibroblast collagen turnover. *Am J Cardiol* 76:8D-13D.
20. Broadley KN, Aquino AM, Hicks B, Ditesheim JA, McGee GS, Demetriou AA, Woodward SC, Davidson JM (1988): Growth factors bFGF and TGB beta accelerate the rate of wound repair in normal and in diabetic rats. *Int J Tissue React* 10:345-353.
21. de Caestecker MP, Hemmati P, Larisch-Bloch S, Ajmera R, Roberts AB, Lechleider RJ (1997): Characterization of functional domains within smad4/DPC4. *J Biol Chem* 272:13690-13696.

22. Cannon JG, Friedberg JS, Gelfand JA, Tompkins RG, Burke JF, Dinarello CA (1992): Circulating interleukin-1 beta and tumor necrosis factor-alpha concentrations after burn injury in humans. *Crit Care Med* 20:1414-1419.
23. Coleman RA, Smith WL, Narumiya S (1994): International Union of Pharmacology Classification of Prostanoid Receptors: Properties, Distribution and Structure of the Receptors and Their Subtypes. *Pharmacol Rev* 46:205-229.
24. Colotta F, Re F, Muzio M, Bertini R, Polentarutti N, Sironi M, Giri JG, Dower SK, Sims JE, Mantovani A (1993): Interleukin-1 type II receptor: a decoy target for IL-1 that is regulated by IL-4. *Science* 261:472-475.
25. Crofford LJ (1997): COX-1 and COX-2 tissue expression: Implications and predictions. *J Rheumatol* (suppl 49)24:15-19.
26. Dell'Acqua ML and Scott JD (1997): Protein kinase A anchoring. *J Biol Chem* 272: 12881-12884.
27. Derynck R, Jarrett JA, Chen EY, Eaton DH, Bell JR, Assoian RK, Roberts AB, Sporn MB, Goeddel DV (1985): Human transforming growth factor-beta complementary DNA sequence and expression in normal and transformed cells. *Nature* 316:701-705.
28. DeWitt DL and Meade EA (1993): Serum and glucocorticoid regulation of gene transcription and expression of the prostaglandin H synthase-1 and prostaglandin H synthase-2 isozymes. *Arch Biochem Biophys* 306:94-102
29. Diaz A, Varga J, Jimenez SA (1989): Transforming growth factor-beta stimulation of lung fibroblast prostaglandin E2 production. *J Biol Chem* 264:11554-11557.
30. Diaz A, Munoz E, Johnston R, Korn JH (1993): Regulation of human lung fibroblast alpha 1(I) procollagen gene expression by tumor necrosis factor alpha, interleukin-1 beta, and prostaglandin E2. *J Biol Chem* 268:10364-10371.
31. Dinarello CA (1988): Biology of interleukin-1. *FASEB J* 2:108-115.
32. Dinarello CA (1996): Biologic basis for interleukin-1 in disease. *Blood* 87:2095-2147.

33. Doskeland SO, Maronde E, Gjertsen BT (1993): The genetic subtypes of cAMP-dependent protein kinase-functionally different or redundant? *Biochem Biophys Acta.* 1178:249-258.
34. Dripps DJ, Brandhuber BJ, Thompson RC, Eisenberg SP (1991): Interleukin-1 (IL-1) receptor antagonist binds to the 80-kDa IL-1 receptor but does not initiate IL-1 signal transduction. *J Biol Chem* 266:10331-10336.
35. Elias JA, Freundlich B, Adams S, Rosenbloom J (1990): Regulation of human lung fibroblast collagen production by recombinant interleukin-1, tumor necrosis factor, and interferon-gamma. *Ann NY Acad Sci* 580:233-244.
36. El Maradny E, Kanayama N, Halim A, Maehara K, Sumimoto K, Terao T (1996): Biochemical changes in the cervical tissue of rabbit induced by interleukin-8, interleukin-1beta, dehydroepiandrosterone sulphate and prostaglandin E2: a comparative study. *Hum Reprod* 11:1099-1104.
37. Eyre DR, Paz MA, Gallop PM (1984): Cross-linking in collagen and elastin. *Annu Rev Biochem* 1984;53:717-748.
38. Fedyk ER and Phipps RP (1996): Prostaglandin E2 receptors of the EP2 and EP4 subtypes regulate activation and differentiation of mouse B lymphocytes to IgE-secreting cells. *Proc Natl Acad Sci USA* 93:10978-10983.
39. Feliciello A, Li Y, Avvedimento EV, Gottesman ME, Rubin CS (1997): A-kinase anchor protein 75 increases the rate and magnitude of cAMP signaling to the nucleus. *Curr Biol* 7:1011-1014.
40. Fine A, Poliks CF, Smith BD, Goldstein RH (1990): The accumulation of type I collagen mRNAs in human embryonic lung fibroblasts stimulated by transforming growth factor-beta. *Connect Tissue Res* 24:237-247.
41. Fine A, Matsui R, Zhan X, Poliks CF, Smith BD, Goldstein RH (1992): Discordant regulation of human type I collagen genes by prostaglandin E2. *Biochim Biophys Acta* 1135:67-72 .
42. Funk CD, Furci L, FitzGerald GA, Grygorczyk R, Rochette C, Bayne MA, Abramovitz M, Adam M, Metters KM (1993): Cloning and expression of a cDNA for human prostaglandin E receptor EP1 subtype. *J Biol Chem* 268:26767-26772.
43. Gallivan K, Alman B, Moriarty K, Pajerski M, O'Donnell C, Crombleholme T (1997): Differential collagen I gene expression in fetal fibroblasts. *J Pediatr Surg* 32:1033-1036.

44. Gauldie J, Jordana M, Cox G (1993): Cytokines and pulmonary fibrosis. *Thorax* 48:931-935.
45. Goldstein RH and Polgar P (1982): The effect and interaction of bradykinin and prostaglandins on protein and collagen production by lung fibroblasts. *J Biol Chem* 257:8630-8633.
46. Granot I, Halevy O, Hurwitz S, Pines M (1993): Halofuginone: an inhibitor of collagen type I synthesis. *Biochim Biophys Acta* 1156:107-112.
47. Greenfeder SA, Nunes P, Kwee L, Labow M, Chizzonite RA, Ju G (1995): Molecular cloning and characterization of a second subunit of the interleukin 1 receptor complex. *J Biol Chem* 270:13757-13765.
48. Grotendorst GR, Grotendorst CA, Gilman T (1988): Production of growth factors (PDGF & TGF-beta) at the site of tissue repair. *Prog Clin Biol Res* 266:47-54.
49. Hakeda Y, Nakatani Y, Yoshino T, Kurihara N, Fujita K, Maeda N, Kumegawa M (1987): Effect of forskolin on collagen production in clonal osteoblastic MC 3T3-E1 cells. *J Biochem (Tokyo)* 101:1463-1469.
50. Hannum CH, Wilcox CJ, Arend WP, Joslin FG, Dripps DJ, Heimdal PL, Armes LG, Sommer A, Eisenberg SP, Thompson RC (1990): Interleukin-1 receptor antagonist activity of a human interleukin-1 inhibitor. *Nature* 343:336-340.
51. Heldin CH, Miyazono K, ten Dijke P (1997): TGF-beta signalling from cell membrane to nucleus through SMAD proteins. *Nature* 390:465-471.
52. Hopp TP (1995): Evidence from sequence information that the interleukin-1 receptor is a transmembrane GTPase. *Protein Sci* 4:1851-1859.
53. Horwitz AL, Hance AJ, Crystal RG (1977): Granulocyte collagenase: selective digestion of type I relative to type III collagen. *Proc Natl Acad Sci U S A* 74:897-901.
54. Houslay MD and Milligan G (1997): Tailoring cAMP-signalling responses through isoform multiplicity. *TIBS* 22:217-224.
55. Ichikawa A, Sugimoto Y, Negishi M (1996): Molecular aspects of the structures and functions of the prostaglandin E receptors. *J Lipid Mediat Cell Signal* 14: 83-87.

56. Iida-Klein A, Yee DC, Brandli DW, Mirikitani EJ, Hahn TJ (1992): Effects of calcitonin on 3',5'-cyclic adenosine monophosphate and calcium second messenger generation and osteoblast function in UMR 106-06 osteoblast-like cells. *Endocrinology* 130:381-388.
57. Irie A, Segi E, Sugimoto Y, Ichikawa A, Negishi M (1994): Mouse prostaglandin E receptor EP3 subtype mediates calcium signals via Gi in cDNA-transfected Chinese hamster ovary cells. *Biochem Biophys Res Commun* 204:303-309.
58. Jackson BA, Goldstein RH, Roy R, Cozzani M, Taylor L, Polgar P (1993): Effects of transforming growth factor- $\beta$  and interleukin-1 $\beta$  on expression of cyclooxygenase 1 and 2 and phospholipase A2 mRNA in lung fibroblasts and endothelial cells in culture. *Biochem Biophys Res Commun* 197:1465-1474.
59. Johnson LR, Foster JA, Haig-Ladewig L, VanScoy H, Rubin CS, Moss SB, Gerton GL (1997): Assembly of AKAP82, a protein kinase A anchor protein, into the fibrous sheath of mouse sperm. *Dev Biol* 192:340-350.
60. Jurevicius J and Fischmeister R (1996): cAMP compartmentation is responsible for a local activation of cardiac Ca<sup>2+</sup> channels by beta-adrenergic agonists. *Proc Natl Acad Sci USA* 93:295-299.
61. Kagan HM (1986): Characterization and regulation of lysyl oxidase. In *Biology of Extracellular Matrix*, Vol 1 (Edited by Mecham R. P.), pp. 321-398. Academic Press, New York.
62. Kahari VM, Heino J, Vuorio E (1987): Interleukin-1 increases collagen production and mRNA levels in cultured skin fibroblasts. *Biochim Biophys Acta* 929:142-147.
63. Kang AH, Bornstein P, Piez KA (1967): The amino acid sequence of peptides from the cross-linking region of rat skin collagen. *Biochemistry* 6:788-795.
64. Katoh H, Watabe A, Sugimoto Y, Ichikawa A, Negishi M (1995): Characterization of the signal transduction of prostaglandin E receptor EP1 subtype in cDNA-transfected Chinese hamster ovary cells. *Biochim Biophys Acta* 1244:41-48.
65. Kiriya M, Ushikubi F, Kobayashi T, Hirata M, Sugimoto Y, Narumiya S (1997): Ligand binding specificities of the eight types and subtypes of the mouse prostanoid receptors expressed in Chinese hamster ovary cells. *Br J Pharmacol* 122:217-224.

66. Kobayashi Y, Yamamoto K, Saido T, Kawasaki H, Oppenheim JJ, Matsushima K (1990): Identification of calcium-activated neutral protease as a processing enzyme of human interleukin 1 alpha. *Proc Natl Acad Sci U S A* 87:5548-5552.
67. Kobayashi Y, Oppenheim JJ, Matsushima K (1991): Human pre-interleukin 1 alpha and beta: structural features revealed by limited proteolysis. *Chem Pharm Bull (Tokyo)* 39:1513-1517.
68. Kol S, Ben-Shlomo I, Ando M, Payne DW, Adashi EY (1997): Interleukin-1 beta stimulates ovarian phospholipase A2 (PLA2) expression and activity: up-regulation of both secretory and cytosolic PLA2. *Endocrinology* 138:314-321.
69. Kotani M, Tanaka I, Ogawa Y, Usui T, Mori K, Ichikawa A, Narumiya S, Yoshimi T, Nakao K (1995): Molecular cloning and expression of multiple isoforms of human prostaglandin E receptor EP3 subtype generated by alternative messenger RNA splicing: multiple second messenger systems and tissue-specific distributions. *Mol Pharmacol* 48:869-879.
70. Kraemer SA, Meade EA, DeWitt DL (1992): Prostaglandin endoperoxide synthase gene structure: identification of the transcriptional start site and 5'-flanking regulatory sequences. *Arch Biochem Biophys* 293:391-400.
71. Krebs EG (1989): Role of the cyclic AMP-dependent protein kinase in signal transduction. *JAMA* 262:1815-1818.
72. Krupsky M, Fine A, Kuang PP, Berk JL, Goldstein RH (1996): Regulation of type I collagen production by insulin and transforming growth factor-beta in human lung fibroblasts. *Connect Tissue Res* 34: 53-62.
73. Kuivaniemi H (1985): Partial characterization of lysyl oxidase from several human tissues. *Biochem J* 230:639-641.
74. Kuno K and Matsushima K (1994): The IL-1 receptor signaling pathway. *J Leukoc Biol* 56:542-547.
75. Laiho M, Weis MB, Massague J (1990): Concomitant loss of transforming growth factor (TGF)-beta receptor types I and II in TGF-beta-resistant cell mutants implicates both receptor types in signal transduction. *J Biol Chem* 265:18518-18524.
76. Lands WE and Samuelsson B (1968): Phospholipid precursors of prostaglandins. *Biochim Biophys Acta* 164:426-429.

77. Lapiere CM, Lenaers A, Kohn LD (1971): Procollagen peptidase: an enzyme excising the coordination peptides of procollagen. *Proc Natl Acad Sci U S A* 68:3054-3058.
78. Lefkowitz RJ (1993): G protein-coupled receptor kinases. *Cell* 74:409-412.
79. Leibovich SJ and Ross R (1975): The role of the macrophage in wound repair. A study with hydrocortisone and antimacrophage serum. *Am J Pathol* 78:71-100.
80. Lenaers A, Ansay M, Nusgens BV, Lapiere CM (1971): Collagen made of extended chains, procollagen, in genetically defective dermatosparaxic calves. *Eur J Biochem* 23:533-543.
81. Liggett SB, Freedman NJ, Schwinn DA, Lefkowitz RJ (1993): Structural basis for receptor subtype-specific regulation revealed by a chimeric beta 3/beta 2-adrenergic receptor. *Proc Natl Acad Sci USA* 90:3665-3669.
82. Lomedico PT, Gubler U, Hellmann CP, Dukovich M, Giri JG, Pan YC, Collier K, Semionow R, Chua AO, Mizel SB (1984): Cloning and expression of murine interleukin-1 cDNA in *Escherichia coli*. *Nature* 312:458-462.
83. Maier JA, Statuto M, Ragnotti G (1994): Endogenous interleukin 1 alpha must be transported to the nucleus to exert its activity in human endothelial cells. *Mol Cell Biol* 14:1845-1851.
84. Massague J (1990): The transforming growth factor-beta family. *Annu Rev Cell Biol* 6:597-641.
85. McDonald JA, Broekelmann TJ, Matheke ML, Crouch E, Koo M, Kuhn C III (1986): A monoclonal antibody to the carboxyterminal domain of procollagen type I visualizes collagen-synthesizing fibroblasts. Detection of an altered fibroblast phenotype in lungs of patients with pulmonary fibrosis. *J Clin Invest* 78:1237-1244.
86. Miller EJ and Matukas VJ (1969): Chick cartilage collagen: a new type of alpha 1 chain not present in bone or skin of the species. *Proc Natl Acad Sci U S A* 64:1264-1268.
87. Miyazono K, Hellman U, Wernstedt C, Heldin CH (1988): Latent high molecular weight complex of transforming growth factor beta 1. Purification from human platelets and structural characterization. *J Biol Chem* 263:6407-6415.

88. Mori K, Tanaka I, Kotani M, Miyaoka F, Sando T, Muro S, Sasaki Y, Nakagawa O, Ogawa Y, Usui T, Ozaki S, Ichikawa A, Narumiya S, Nakao K (1996): Gene expression of the human prostaglandin E receptor EP4 subtype: differential regulation in monocytoid and lymphoid lineage cells by phorbol ester. *J Mol Med* 74:333-336.
89. Munger JS, Harpel JG, Gleizes PE, Mazziere R, Nunes I, Rifkin DB (1997): Latent transforming growth factor-beta: structural features and mechanisms of activation. *Kidney Int* 51:1376-1382.
90. Myers BA, Dubick MA, Reynolds RD, Rucker RB (1985): Effect of vitamin B-6 (pyridoxine) deficiency on lung elastin cross-linking in perinatal and weanling rat pups. *Biochem J* 229:153-160.
91. Narumiya, S (1994): Prostanoid receptors. Structure, Function and Distribution. *Ann N Y Acad Sci* 744:126-138.
92. Nesterova M, Yokozaki H, McDuffie E, Cho-Chung YS (1996): Overexpression of RII $\beta$  regulatory subunit of protein kinase A in human colon carcinoma cell induces growth arrest and phenotypic changes that are abolished by site-directed mutation of RII $\beta$ . *Eur J biochem* 235:486-494.
93. Nishigaki N, Negishi M, Ichikawa A (1996): Two Gs-coupled prostaglandin E receptor subtypes, EP2 and EP4, differ in desensitization and sensitivity to the metabolic inactivation of the agonist. *Mol Pharmacol* 50:1031-1037.
94. Padgett RW, Savage C, Das P (1997): Genetic and biochemical analysis of TGF- $\beta$  signal transduction. *Cytokine Growth Factor Rev* 8: 1-9.
95. Pardo A, Selman M, Ramirez R, Ramos C, Montano M, Stricklin G, Raghu G (1992): Production of collagenase and tissue inhibitor of metalloproteinases by fibroblasts derived from normal and fibrotic human lungs. *Chest* 102:1085-1089.
96. Pastan I (1972): Cyclic AMP. *Sci Am* 227:97-105.
97. Pasternak RD, Hubbs SJ, Caccese RG, Marks RL, Conaty JM, DiPasquale G (1986): Interleukin-1 stimulates the secretion of proteoglycan- and collagen-degrading proteases by rabbit articular chondrocytes. *Clin Immunol Immunopathol* 41:351-367.
98. Pawson T and Scott JD (1997): Signaling through scaffold, anchoring and adaptor proteins. *Science* 278:2075-2080.

99. Phan SH, Gharaee-Kermani M, Wolber F, Ryan US (1991): Stimulation of rat endothelial cell transforming growth factor-beta production by bleomycin. *J Clin Invest* 87:148-154.
100. Pierce KL, Gil DW, Woodward DF, Regan JW (1995): Cloning of human prostanoid receptors. *Trends Pharmacol Sci.* 16:253-256.
101. Piez KA, Eigner EA, Lewis MS (1963): The chromatographic separation and amino acid composition of the subunits of several collagens. *Biochemistry* 2:58-66.
102. Pinnell SR and Martin GR (1968): The cross-linking of collagen and elastin: enzymatic conversion of lysine in peptide linkage to alpha-amino adipic-delta-semialdehyde (allysine) by an extract from bone. *Proc Natl Sci USA* 61:708-714.
103. Pun KK (1989): The importance of parathyroid hormone in inhibition of collagen synthesis and mitogenesis of osteoblastic cells. *J Biochem (Tokyo)* 106:1090-1093.
104. Purchio AF, Cooper JA, Brunner AM, Lioubin MN, Gentry LE, Kovacina KS, Roth RA, Marquardt H (1988): Identification of mannose 6-phosphate in two asparagine-linked sugar chains of recombinant transforming growth factor-beta 1 precursor. *J Biol Chem* 263:14211-14215.
105. Raisz LG, Fall PM, Petersen DN, Lichtler A, Kream BE (1993): Prostaglandin E2 inhibits alpha 1(I)procollagen gene transcription and promoter activity in the immortalized rat osteoblastic clonal cell line Py1a. *Mol Endocrinol* 7:17-22.
106. Regan JW, Bailey TJ, Donello JE, Pierce KL, Pepperl DJ, Zhang D, Kedzie KM, Fairbairn CE, Bogardus AM, Woodward DF (1994): Molecular cloning and expression of human EP3 receptors: evidence of three variants with differing carboxyl termini. *Br J Pharmacol* 112:377-385.
107. Reed MJ, Vernon RB, Abrass IB, Sage EH (1994): TGF- $\beta$ 1 induces the expression of type I collagen and SPARC, and enhances contraction of collagen gels, by fibroblasts from young and aged donors. *J Cell Physiol* 158:169-179.
108. Ricard-Blum S and Ville G (1989): Collagen cross-linking. *Int J Biochem* 21:1185-1189.

109. Ritzenthaler JD, Goldstein RH, Fine A, Smith BD (1993): Regulation of the  $\alpha 1(I)$  collagen promoter via a transforming growth factor- $\beta$  activation element. *J Biol Chem* 268:13625-13631.
110. Robins SP, Duncan A (1983): Cross-linking of collagen. Location of pyridinoline in bovine articular cartilage at two sites of the molecule. *Biochem J* 215:175-182.
111. Roy R, Polgar P, Wang YY, Goldstein RH, Taylor L, Kagan HM (1996): Regulation of Lysyl Oxidase and Cyclooxygenase Expression in Human Lung Fibroblasts: Interactions Among TGF- $\beta$ , IL-1 $\beta$ , and Prostaglandin E. *J Cell Biochem* 62:411-417.
112. Rucker RB, Murray J (1978): Cross-linking amino acids in collagen and elastin. *Am J Clin Nutr* 31:1221-1236.
113. Saltzman LE, Moss J, Berg RA, Hom B, Crystal RG (1982): Modulation of collagen production by fibroblasts. Effects of chronic exposure to agonists that increase intracellular cyclic AMP. *Biochem J* 204:25-30.
114. Sandell LJ and Boyd CD (1990): Conserved and divergent sequence and functional elements within collagen genes. In: Sandell LJ, Boyd CD, Eds. *Extracellular Matrix Genes*. San Diego: Academic Press, pp1-56.
115. Samuelsson B, Granstrom E, Green K, Hamberg M, Hammarstrom S (1975): Prostaglandins. *Annu Rev Biochem* 44:669-695.
116. Scheule RK, Perkins RC, Hamilton R, Holian A (1992): Bleomycin stimulation of cytokine secretion by the human alveolar macrophage. *Am J Physiol* 262(4 Pt 1):L386-L391.
117. Schmid A, Thierauch K, Schleuning W, Dinter H (1995): Splice variants of the human EP3 receptor for prostaglandin E2. *Eur J Biochem* 228:23-30.
118. Scutt A, Zeschnigk M, Bertram P (1995): PGE<sub>2</sub> Induces the Transition from Non-Adherent to Adherent Bone Marrow Mesenchymal Precursor Cells Via a cAMP/EP2-Mediated Mechanism. *Prostaglandins* 49:383-395.
119. Seckinger P, Williamson K, Balavoine JF, Mach B, Mazzei G, Shaw A, Dayer JM (1987): A urine inhibitor of interleukin 1 activity affects both interleukin 1 alpha and 1 beta but not tumor necrosis factor alpha. *J Immunol* 139:1541-1545.

120. Seibert K, Zhang Y, Leahy K, Hauser S, Masferrer J, Perkins W, Lee L, Isakson P (1994): Pharmacological and biochemical demonstration of the role of cyclooxygenase 2 in inflammation and pain. *Proc Natl Acad Sci U S A* 91:12013-12017.
121. Seyer JM, Hutcheson ET, Kang AH (1976): Collagen polymorphism in idiopathic chronic pulmonary fibrosis. *J Clin Invest* 57:1498-1507.
122. Shapiro SD, Kobayashi DK, Welgus HG (1992): Identification of TIMP-2 in human alveolar macrophages. Regulation of biosynthesis is opposite to that of metalloproteinases and TIMP-1. *J Biol Chem* 267:13890-13894.
123. Siegel RC, Pinnell SR, Martin GR (1970): Cross-linking of collagen and elastin. Properties of lysyl oxidase. *Biochemistry* 9:4486-4492.
124. Siegel RC (1979): Lysyl oxidase. *Int Rev Connect Tissue Res* 8:73-118.
125. Smith WL and Dewitt DL (1996): Prostaglandin endoperoxide H synthases-1 and -2. *Adv Immunol* 62:167-215.
126. Stevenson FT, Torrano F, Locksley RM, Lovett DH (1992): Interleukin 1: the patterns of translation and intracellular distribution support alternative secretory mechanisms. *J Cell Physiol* 152:223-231.
127. Stevenson FT, Bursten SL, Fanton C, Locksley RM, Lovett DH (1993): The 31-kDa precursor of interleukin 1 alpha is myristoylated on specific lysines within the 16-kDa N-terminal propeptide. *Proc Natl Acad Sci U S A* 90:7245-7249.
128. Takii T, Akahoshi T, Kato K, Hayashi H, Marunouchi T, Onozaki K (1992): Interleukin-1 up-regulates transcription of its own receptor in a human fibroblast cell line TIG-1: role of endogenous PGE2 and cAMP. *Eur J Immunol* 22:1221-1227.
129. Taylor L, Ricupero D, Jean J, Jackson BA, Navarro J, Polgar P (1992): Functional expression of the bradykinin-B2 receptor cDNA in Chinese hamster lung CCL39 fibroblasts. *Biochem Biophys Res Commun* 188:786-793.
130. Taylor SS, Buechler JA, Yonemoto W (1990): cAMP-dependent protein kinase: framework for a diverse family of regulatory enzymes. *Annu Rev Biochem.* 59: 971-1005.

131. Vane JR, Mitchell JA, Appleton I, Tomlinson A, Bishop-Bailey D, Croxtall J, Willoughby DA (1994): Inducible isoforms of cyclooxygenase and nitric-oxide synthase in inflammation. *Proc Natl Acad Sci U S A* 91:2046-2050.
132. Visconti PE, Johnson LR, Oyaski M, Fornes M, Moss SB, Gerton GL, Kopf GS (1997): Regulation, localization, and anchoring of protein kinase A subunits during mouse sperm capacitation. *Visconti Dev Biol* 192:351-363.
133. Walker NP, Talanian RV, Brady KD, Dang LC, Bump NJ, Ferenz CR, Franklin S, Ghayur T, Hackett MC, Hammill LD (1994): Crystal structure of the cysteine protease interleukin-1 beta-converting enzyme: a (p20/p10)<sub>2</sub> homodimer. *Cell* 78:343-352.
134. Whitby DJ and Ferguson MWJ (1991): The extracellular matrix of lip wounds in fetal, neonatal, and adult mice. *Development* 112:651-658.
135. Wilborn J, Crofford LJ, Burdick MD, Kunkel SL, Stierter RM, Peters-Golden M (1995): Cultured lung fibroblasts isolated from patients with idiopathic pulmonary fibrosis have a diminished capacity to synthesize prostglandin E<sub>2</sub> and to express cyclooxygenase-2. *J Clin Invest* 95:1861-1868.
136. Woodward DF, Regan JW, Lake S, Ocklind A (1997): The molecular biology and ocular distribution of prostanoid receptors. *Surv Ophthalmol* 41 Suppl 2:S15-S21.
137. Wrana J and Pawson T (1997): Mad about SMADS. *Nature* 388:28-29.
138. Zhang K, Gharaee-Kermani M, McGarry B, Phan SH (1994): In situ hybridization analysis of rat lung alpha 1(I) and alpha 2(I) collagen gene expression in pulmonary fibrosis induced by endotracheal bleomycin injection. *Lab Invest* 70:192-202.
139. Zhang Y, Lee TC, Guillemin B, Yu MC, Rom WN (1993): Enhanced IL-1 beta and tumor necrosis factor-alpha release and messenger RNA expression in macrophages from idiopathic pulmonary fibrosis or after asbestos exposure. *J Immunol* 150:4188-4196.