

AD _____

GRANT NUMBER DAMD17-96-1-6181

TITLE: Role of Bcl-2 in Breast Cancer Progression

PRINCIPAL INVESTIGATOR: Hyeong-Reh Kim, Ph.D.

CONTRACTING ORGANIZATION: Wayne State University
Detroit, Michigan 48202

REPORT DATE: September 1998

TYPE OF REPORT: Annual

PREPARED FOR: Commanding General
U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

**Reproduced From
Best Available Copy**

19981204 056

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE September 1998	3. REPORT TYPE AND DATES COVERED Annual (1 Aug 97 - 31 Jul 98)	
4. TITLE AND SUBTITLE Role of Bcl-2 in Breast Cancer Progression		5. FUNDING NUMBERS DAMD17-96-1-6181	
6. AUTHOR(S) Hyeong-Reh C. Kim, Ph.D.			
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Wayne State University Detroit, Michigan 48202		8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012		10. SPONSORING / MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES			
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited		12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) The anti-apoptotic gene bcl-2 is frequently overexpressed in many human tumors including invasive breast cancer. Involvement of bcl-2 in cancer development was believed to result from its ability to prevent cell death (thereby increasing cell number). We previously showed that bcl-2 deregulates G ₁ /S checkpoint in MCF10A cells, which involves induction of cyclin D ₁ associated kinase activity. Increasing evidence suggests that overexpression of cyclin D ₁ contributes to the oncogenic transformation of cells <i>in vitro</i> and <i>in vivo</i> . In the present study, we examined the effects of bcl-2 overexpression on cyclin D ₁ expression in human breast epithelial cells. Bcl-2 overexpression in MCF10A cells significantly induced expression of cyclin D ₁ during the cell cycle as determined by an immunoblot analysis. We also show that bcl-2 induction of cyclin D ₁ expression occurs at the transcriptional level as determined by an assay of cyclin D ₁ promoter activity. Cyclin D ₁ reporter activity was induced in a bcl-2 dependent manner in human breast carcinoma cell lines, MCF7 and BT549 as well as in MCF10A. Our previous and present studies suggest that bcl-2 may serve as an oncogene in the development of human breast cancer, which involves induction of cyclin D ₁ expression.			
14. SUBJECT TERMS Breast Cancer		15. NUMBER OF PAGES 10	16. PRICE CODE
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited

FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the U.S. Army.

____ Where copyrighted material is quoted, permission has been obtained to use such material.

____ Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

Jgm Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.

Jgm In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and use of Laboratory Animals of the Institute of Laboratory Resources, national Research Council (NIH Publication No. 86-23, Revised 1985).

____ For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.

Jgm In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

Jgm In the conduct of research utilizing recombinant DNA, the investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules.

Jgm In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories.

Jgm

PI - Signature

8-31-98

Date

TABLE OF CONTENTS

	Page Number
Introduction	5
Body of Report	5-9
Conclusion	9
References	9-10

Introduction

The proto-oncogene *bcl-2* is frequently overexpressed in many human tumors including invasive breast cancers (1, 2). *In vitro* studies clearly demonstrate that the *bcl-2* gene product prevents apoptosis following a variety of stimuli including radiation, hyperthermia, growth factor withdrawal and chemotherapeutic drugs. In transgenic mice, overexpression of *bcl-2* under the immunoglobulin promoter induces follicular hyperplasia, suggesting a role for *bcl-2* as an oncogene. However, high levels of *bcl-2* expression have shown a positive correlation in clinicopathological studies such as tumor grade, and better response to hormone treatment and chemotherapy (3). The long-term goal of our studies is to investigate the *in vivo* functions of *bcl-2* in human breast cancer development. To this end, we have used MCF10A and MCF10AT models.

MCF10A cell line was established in our institute without viral or chemical intervention from mortal diploid human breast epithelial cells (4). This cell line has been utilized by many laboratories to study sequential development of differentiated or malignant states of breast epithelial cells.

Dr. Miller (consultant in this application) and his colleagues have developed an *in vivo* model system to study human breast cancer progression (5-7); Whereas MCF10A cells do not survive *in vivo* in immune deficient mice, *c-Ha-ras* oncogene transfected MCF10A cells (MCF10AneoT) form small nodules in Nude/Beige mice which persist for at least one year and sporadically progress to carcinomas. Cell lines have been established in cultures from lesions representing 4 successive transplant generations (designated MCF10ATG1, MCF10ATG2, MCF10ATG3 and MCF10ATG4). With each generation, cells progress to high risk lesions resembling human proliferative breast disease. Thus, the MCF10AT model provides a setting in which the steps in the conversion of the breast ductal epithelial cell to malignant disease can be studied.

Body of Report

During the 1997-1998 funding period, we have continued to investigate potential oncogenic activities of *bcl-2* in breast cancer development.

Methods

Immunoblot analysis

MCF10A and *bcl-2* overexpressing MCF10A clones (MCF10A *bcl-2-6*, MCF10A *bcl-2-8*, MCF10A *bcl-2-30* and MCF10A *bcl-2-40*) were cultured as previously described (8). Cells were growth-arrested by culturing in serum-free medium for 48 hours at confluence. Cells were then treated with regular MCF10A medium containing 5% horse serum and growth factors to induce the cell cycle. At various times between 0-24 hr after serum-stimulation, whole-cell extracts were prepared using SDS lysate buffer. Protein concentrations were measured using bicinchoninic acid protein assay reagents (Pierce, Rockford, IL). Cell lysates (20 µg/lane) were denatured, subjected to SDS-PAGE analysis, and then electrophoretically transferred to a nitrocellulose membrane. Membranes were incubated with anti-cyclin D₁ (Ab2, Oncogene Research, MA), anti-*bcl-2* or β-actin antibodies. Proteins were visualized using HRP-conjugated goat anti-mouse IgG (1:3000 dilution) and chemiluminescence reagent (Dupont, Boston MA 02118). The membranes were exposed to X-ray film from 1 to 15 minutes.

Cyclin D₁ promoter activity assay

MCF10A, MCF7 and BT549 cells were transfected with -964 CD1 promoter, PCH110

(MDVlacZ), and bcl-2 expression plasmid using FuGENE6 transfection reagent (Boehringer Mannheim). Fifty percent confluent cells in a 60-mm dish were transfected in 3 ml culture medium into which 500 μ l FuGENE6 reaction mixture was added. The FuGENE mixture was prepared as follows; 2 μ g of -964CD1 plasmid, 0.2 μ g of PCH110 and increasing amounts of bcl-2 expression plasmid were mixed in 250 μ l of serum free medium. The DNA solution was mixed with 250 μ l of serum free-medium which contains 50 μ l FuGENE6 reagent. The FuGENE6 mixture was incubated at room temperature for 30 minutes before transfection. Cells were harvested after 48 hours of transfection with 1X reporter lysis buffer. After mixing 20 μ l of cell lysate and 100 μ l luciferase substrate, the luciferase activity was measured using a luminometer. The β -galactosidase activity was measured by a chemiluminescent reporter assay (Tropix) to normalize transfection efficiencies. Protein concentration of the cell lysate was measured using the BCA reagent (Pierce).

Results

We previously showed that bcl-2 deregulates G₁/S check point through modulation of cyclin D₁-associated kinase activity. In the present study, we examined whether bcl-2 regulates cyclin D₁ expression. When the levels of cyclin D₁ expression were examined in growing cells of MCF10A and bcl-2 overexpressing MCF10A cells, it appeared that bcl-2 induced expression of cyclin D₁ (Fig 1). To exclude the possibility that the differences in the levels of cyclin D₁ reflect the differences in the cell cycle distribution between the control and bcl-2 overexpressing cells, we examined the effects of bcl-2 overexpression on cyclin D₁ expression during the cell cycle. The control and bcl-2 overexpressing MCF10A cells were synchronized at G₀ by culturing in serum-free medium for 48 hours and the cell cycle was induced by culturing them in complete MCF10A culture medium. At various times between 0-24 hr after cell cycle induction, whole-cell extracts were prepared and the levels of cyclin D₁ were determined by immunoblot analysis. As shown in Fig. 2, the levels of cyclin D₁ expression were significantly higher throughout the cell cycle in bcl-2 overexpressing cells than in the control cells.




Fig. 1. Bcl-2 induces cyclin D₁ expression in MCF10A cells. The levels of cyclin D₁ (top panel) and bcl-2 (middle panel) proteins were determined by immunoblot analyses. Protein samples were prepared from neo-resistant vector transfected MCF10A (lane 1) and bcl-2 overexpressing MCF10A clones (lanes 2-5). To confirm the amount and quality of proteins loaded in each lane, the identical blot was probed with anti- β -actin antibody (bottom panel).





Fig. 2. Bcl-2 induces cyclin D₁ expression during the cell cycle in MCF10A cells. Confluent control (top panel) and bcl-2 overexpressing MCF10A clone 30 (bottom panel) were cultured in serum free DMEM/F12 medium for 48 hours. At various time points after serum treatment, cells were lysed using SDS sample buffer. Levels of cyclin D₁ protein were determined by immunoblot analyses.

We then examined whether bcl-2 induces transcription of cyclin D₁ in MCF10A cells. The effects of bcl-2 on cyclin D₁ promoter activity were determined by cotransfection experiments using -964CD1LUC plasmid (containing the human cyclin D₁ promoter linked to the luciferase reporter provided by Dr. Pestell, *see ref. 12*) and bcl-2 expression vector. As shown in Fig 3A, cyclin D₁ reporter activity was induced in a bcl-2 dependent manner. It was examined whether bcl-2 induction of cyclin D₁ transcription occurs in other human breast epithelial cells. To this end, we used human breast carcinoma cell lines, MCF7 and BT549. As shown in Fig 3A and B, bcl-2 effectively induced promoter activities of cyclin in these cell lines. These results show that bcl-2 induces expression of cyclin D₁ at the transcriptional level in human breast epithelial cells.

A. **BCL-2 Induction of Cyclin-D1 Transcription**



B. **Bcl2 Induction of CyclinD1 Transcription**




Fig. 3. Bcl-2 induces cyclin D₁ promoter activity in human breast epithelial cells.

A. -964CD1LUC reporter was transfected with increasing amounts of bcl-2 expression vector into MCF10A (hatched bar in panel A), BT549 (dotted bar in panel A) and MCF 7 (panel B). All experiments were performed in triplicates and the folds of induction represent the luciferase activity/LacZ activity/mg protein.

We are currently investigating the regions of the cyclin D₁ promoter responsible for transcriptional activation by bcl-2 overexpression using control and bcl-2 overexpressing cells. Our preliminary study showed that mutation of the AP1 binding site in the cyclin D₁ promoter does not have any effect on bcl-2 activation of cyclin D₁ promoter activity. However, deletion of the SP-1 binding site in -163 significantly reduced bcl-2 activation of cyclin D₁ promoter activity (data not shown). Further studies are in demand to determine the mechanisms by which bcl-2 activates cyclin D₁ promoter activity.

Conclusion

Increasing evidence suggests that overexpression of cyclin D₁ contributes to the oncogenic transformation of cells *in vitro* and *in vivo* (9-11). Involvement of bcl-2 in cancer development was believed to result from its ability to prevent cell death (thereby increasing cell number). However, our previous and present studies suggest that bcl-2 may serve as an oncogene in the development of human breast cancer, which involves induction of cyclin D₁ expression. We now wish to determine the *in vivo* oncogenic role of bcl-2 using bcl-2 overexpressing MCF10ATG3B (we previously showed that bcl-2 overexpression in MCF10ATG3B induces a transformed phenotype as determined by a soft agar assay). Since the Career Development Award does not support any research expense, it is difficult to perform the proposed study which includes animal experiments.

References

1. Silvestrini, R., Veneroni, S., Daidone, M. G., Benini, E. B., Boracchi, P., Mezzetti, M., Di Fronzo, G., Rilke, F. and Veronesi, U. The bcl-2 protein: a prognostic indicator strongly related to p53 protein in lymphoma node-negative breast cancer patients. *J. Natl Cancer Inst.* 86, 499-504, 1994
2. Siziopikou, K. P., Prioleau, J. E., Harris, J. R. and Stuart, J. S. Bcl-2 expression in the spectrum of preinvasive breast lesions. *Cancer* 77 (3), 499-506, 1996
3. Visscher, D. W., Sarkar, F., Tabaczka, P., Crissman, J. Clinicopathologic analysis of bcl-2 immunostaining in breast carcinoma. *Modern Pathology* 9, 642-646, 1996
4. Soule, H., Maloney, T. M., Wolman, S. R., Peterson, W. D., Brenz, R., McGrath, C. M., Russo, J., Pauley, R. J., Jones, R. F. and Brooks, S. C. Isolation and characterization of a spontaneously immortalized human breast epithelial cell line, MCF-10. *Cancer Res.* 50, 6075-6086, 1990
5. Miller, F. R., Soule, H. D., Tait, L., Pauley, R. J., Wolman, S. R., Dawson, P. J., Heppner, G. H. Xenograft model of progressive human proliferative breast disease. *JNCI* 85, 1725-1732,

1993

6. Dawson, P. J., Wolman, S. R., Tait, L., Heppner, G. H., Miller, F. R. MCF10AT: A model for the evolution of cancer from proliferative breast disease. *Am. J. Path* 148, 313-319, 1996
7. Miller, F. R. Models of progression spanning preneoplasia and metastasis: The human MCF10AneoT.TGn series and a panel of mouse mammary tumors. In: *Mammary tumor cell cycle, differentiation and metastasis*, ed. M.E. Lippman and R, B, Dickon, pp243-263. Kluwer Academic Publication, Boston, 1996
8. Upadhyay, S., Li, G., Liu, H., Chen, Y., Sarkar, F and Kim, H.-R. C. Bcl-2 suppresses expression of p21^{WAF1/CIP1} in human breast epithelial cells. *Cancer Res.* 55, 4520-4524, 1995
9. Bodrug, S. E., Warner, B. J., Bath, M. L., Lindeman, G. J., Harris, A. W. and Adams, J. M. Cyclin D₁ transgene impedes lymphocyte maturation and collaborates in lymphomagenesis with the myc gene. *EMBO J.*, 13, 2124-2130, 1994
10. Hinds, P. W., Dowdy, S. F., Eaton, E. N., Arnold, A. and Weinberg, R. A. Function of a human cyclin gene as an oncogene. *Proc. Natl. Acad. Sci. USA*, 91, 709-713, 1994
11. Wang, T. C., Cardiff, R. D., Zukerberg, L., Lees, E., Arnolds, A and Schmidt, E. V. Mammary hyperplasia and carcinoma in MMTV-cyclin D1 transgene mice. *Nature*, 369, 669-671, 1994
12. Watanase, G., Howe, A., Lee, R. J., Albanese, C., Shu, I.-W., Karnezis, A. N., Zon, L., Kyriakis, J., Rundell, K. and Pestell, R. G. Induction of cyclin D₁ by simian virus 40 small tumor antigen. *Proc. Natl. Acad. Sci.* 93, 12861-12866, 1996