

**UNITED STATES AIR FORCE  
ARMSTRONG LABORATORY**

**THE BIOLOGICAL EFFECTS OF ADN  
ON HEPATOCYTES: AN  
EPR STUDY**

**Sara Berty  
Linda Steel-Goodwin  
Kenneth Dean**

**OCCUPATIONAL AND ENVIRONMENTAL  
HEALTH DIRECTORATE TOXICOLOGY DIVISION  
ARMSTRONG LABORATORY  
WRIGHT-PATTERSON AFB OH 45433-7400**

**Alasdair Carmichael  
ARMED FORCES RADIOBIOLOGY  
RESEARCH INSTITUTE  
BETHESDA MD**

**October 1995**

19990316 045

**Occupational and Environmental Health  
Directorate  
Toxicology Division  
2856 G Street  
Wright-Patterson AFB OH 45433-7400**

Approved for public release; distribution is unlimited.

## NOTICES

When US Government drawings, specifications or other data are used for any purpose other than a definitely related Government procurement operation, the Government thereby incurs no responsibility nor any obligation whatsoever, and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise, as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use, or sell any patented invention that may in any way be related thereto.

Please do not request copies of this report from the Air Force Armstrong Laboratory. Additional copies may be purchased from:

National Technical Information Service  
5285 Port Royal Road  
Springfield, Virginia 22161

Federal Government agencies and their contractors registered with the Defense Technical Information Center should direct requests for copies of this report to:

Defense Technical Information Service  
8725 John J. Kingman Rd., Ste 0944  
Ft. Belvoir, Virginia 22060-6218

## DISCLAIMER

This Technical Report is published as received and has not been edited by the Technical Editing Staff of the Air Force Armstrong Laboratory.

## TECHNICAL REVIEW AND APPROVAL

**AL/OE-TR-1995-0173**

The animal use described in this study was conducted in accordance with the principles stated in the "Guide for the Care and Use of Laboratory Animals", National Research Council, 1996, and the Animal Welfare Act of 1966, as amended.

This report has been reviewed by the Office of Public Affairs (PA) and is releasable to the National Technical Information Service (NTIS). At NTIS, it will be available to the general public, including foreign nations.

This technical report has been reviewed and is approved for publication.

**FOR THE DIRECTOR**



**STEPHEN R. CHANNEL**, Maj, USAF, BSC  
Branch Chief, Operational Toxicology Branch  
Air Force Armstrong Laboratory

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE October 1995	3. REPORT TYPE AND DATES COVERED Interim Report - August 1994-September 1995		
4. TITLE AND SUBTITLE The Biological Effects of ADN on Hepatocytes: An EPR Study			5. FUNDING NUMBERS Contract PE 61102F PR 2300 TA 23000T WU 23000T51	
6. AUTHOR(S) S.E. Berty, K.W. Dean, L. Steel-Goodwin, and A.J. Carmichael				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Armstrong Laboratory, Occupational and Environmental Health Directorate Toxicology Division, Human Systems Center Air Force Materiel Command Wright-Patterson AFB, OH 45433-7400			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Armstrong Laboratory, Occupational and Environmental Health Directorate Toxicology Division, Human Systems Center Air Force Materiel Command Wright-Patterson AFB, OH 45433-7400			10. SPONSORING/MONITORING AGENCY REPORT NUMBER  AL/OE-TR-1995-0173	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION AVAILABILITY STATEMENT  Approved for public release; distribution is unlimited.			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) This project investigated the biological effects of ammonium dinitramide (ADN) on hepatocytes. It was hypothesized that ADN decomposes to form free radicals which would be deleterious to the body. The effects of ADN on the liver were studied because regardless of the route of exposure, once inside the body it will enter the bloodstream and ultimately pass through the liver. The leakage of the enzymes aspartate aminotransferase (AST), alanine aminotransferase (ALT), and lactate dehydrogenase (LDH) was measured to ascertain the viability of WB 344 hepatocytes after a 24 h exposure to ADN. Electron paramagnetic resonance (EPR) spectroscopy was used to determine if ADN induced the production of free radicals. As free radicals are highly reactive, a-phenyl-tert-butyl nitron (PBN) and 5,5-dimethyl-1-pyrroline-1-oxide (DMPO) were used to trap the radicals produced in the experiments. Incubation of hepatocytes with 2.8 mM ADN for 24 h was toxic to 50% of the cells. Cells exposed to ADN produced free radicals in the presence of both PBN and DMPO. The generation of free radicals using PBN seemed to be pH dependent. Further studies are necessary to determine the effects of ADN on the possible target organs, the lungs and skin.				
14. SUBJECT TERMS Ammonium Dinitramide                      Free Radicals Electron paramagnetic resonance        Spectroscopy			15. NUMBER OF PAGES 24	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT  UNCLASSIFIED	18. SECURITY CLASSIFICATION OF THIS PAGE  UNCLASSIFIED	19. SECURITY CLASSIFICATION OF ABSTRACT  UNCLASSIFIED	20. LIMITATION OF ABSTRACT  UL	

## PREFACE

This study was supported by the Air Force Office of Scientific Research and the Pharmacodynamics Group, Armstrong Laboratory, Toxicology Division. The authors wish to gratefully acknowledge the assistance they received from SrA Gerri Miller and SrA Stacie Southwell with clinical analysis, Mrs. Cressance Booher for secretarial assistance, along with Wright State University and Dr. P. Serve for the use of EPR.

## TABLE OF CONTENTS

SECTION	PAGE
PREFACE.....	iii
TABLE OF CONTENTS.....	iv
LIST OF FIGURES.....	v
INTRODUCTION.....	1
METHODOLOGY.....	5
Cell Culture.....	5
Cell Viability.....	5
Cell Proliferation.....	5
EPR Spectroscopy.....	5
RESULTS.....	6
DISCUSSION.....	10
REFERENCES.....	11

## LIST OF FIGURES

FIGURE		PAGE
1	Currently used high energy solid fuel .....	2
2	Possible pathways for ADN decomposition .....	3
3	Viability and proliferation assay results .....	6
4	EPR spectra with PBN .....	7
5	EPR spectra with DMPO .....	8
6	The EPR/spin trapping spectra over time .....	9

# THE BIOLOGICAL EFFECTS OF ADN ON HEPATOCYTES: AN EPR STUDY

## Introduction

One of the major objectives of occupational health and environmental toxicology research is to determine and quantify risks that may occur as a result of exposure to experimental chemicals (1). These risks must be ascertained as early as possible so that proper safety measures may be taken throughout the research and development of new chemicals. The main goals of studying the biological effects of exposure to the oxidizer ammonium dinitramide (ADN) is to ensure that the current or past exposure of workers is "safe" (ie does not permit an unacceptable health risk) and to detect potential excessive exposure before the occurrence of detectable adverse health effects. It is essentially a preventative medical activity.

The results of this biological effects program could also potentially be used to make a biological monitoring device. Such a device could be used to interpret exposure on an individual basis, which could then be used to estimate for each examined worker the amount of exposure absorbed during a specific time interval or the amount retained in the organism or bound to critical cells in the body or even to soil particles at waste sites. The information may also be used to appreciate the overall work hygiene conditions by analyzing the distribution of the biological results in a group of workers. It evaluates the internal dose received and hence helps to estimate health risks. The greatest advantage of understanding the biological effects of ADN is the fact that the biological

parameter of exposure is more directly related to the adverse health effects which the U.S. Air Force attempts to prevent than any environmental measurement.

ADN is a new type of energetic material (2). Currently, several chemicals including TNT, HMX, RDX, and AP (ammonium perchlorate) are being used as high energy solid fuels, Fig. 1.




Figure 1. Currently used high energy solid fuels

Recent research is leading to the development of replacements for these compounds. ADN is being investigated as an alternative compound to AP. It is predicted that ADN will be an improvement over AP in weapons systems development or as an oxidizer in the solid fuel of the booster rockets used to put the space shuttle in orbit (2). First, ADN contains no chlorine atoms. Chlorine is a potential pollutant. Second, ADN would not result in the contrail currently produced by AP (2). The absence of this trail will make the detection of rocket launches powered by ADN more difficult. Third, ADN will permit an increased payload capacity which is important for space launches. ADN is an oxide of nitrogen and its chemical formula is  $\text{NH}_4\text{N}(\text{NO}_2)_2$  (3). Based on the chemical formula of ADN it can decompose to nitrogen dioxide,  $\text{NO}_2$  (4).  $\text{NO}_2$  is a free radical.

Free radicals can be defined as a molecule or ion containing an unpaired electron (5). Free radicals are very reactive and can cause injury to biological tissue (6-10).

ADN has been experimentally shown to produce  $\text{NO}_2$  on exposure to gamma-radiation (4). Although the possible ADN induced free radical reactions can be shown chemically (Fig. 2), it is not known whether they can occur within living cells.




Figure 2. Possible pathways for ADN decomposition (Ref. 4)

The best technique to study free radicals is electron paramagnetic resonance spectroscopy (EPR). Radicals in concentrations down to about  $10^{-10}$  M can be detected by EPR (11). In this technique a sample placed in a magnetic field is subjected to microwave radiation. The unpaired

electron acting as a magnet can take up two orientations with respect to the external field corresponding to two energy levels. The energy difference between these levels induced by the microwave radiation produces an absorption peak which is detected by the spectrometer (12). As free radicals react very quickly, one way of detecting them is by spin trapping. Spin trapping consists of reacting short-lived free radicals with a spin trap (usually a nitron or nitroso compound) yielding a longer-lived nitroxide spin adduct which can be detected by EPR (13). There are a number of spin traps which can be used in biological systems (9,13). The most commonly used spin traps are *a*-phenyl-*tert*-butyl nitron (PBN) and 5,5-dimethyl-1-pyrroline-1-oxide (DMPO).

It was hypothesized that the main routes of exposure to ADN would be through the lungs or skin. Study of the biological effects of ADN have to take into consideration absorption by all routes. Regardless of the route of entry of ADN, once inside the body it will enter the bloodstream and will ultimately pass through the liver. The liver is the largest gland in the body (14) and is often the target organ of chemical-induced tissue injury, a fact recognized for over 100 years (15-16). Hazard assessment studies often focus on the liver because it is the organ largely responsible for the detoxification and metabolism of chemicals in the body. While, the biological effects of exposure to ADN in the liver can be studied in many ways, the initial study of the biological effects in cultured hepatocytes is the most logical because it is economical, provides large supplies of samples and requires no animals. The objective of this project was to study the biological effects of ADN on the viability and proliferation of hepatocytes and to measure the free radical decomposition products of ADN by EPR and EPR/spin trapping techniques.

## METHODOLOGY

### Cell Culture

WB 344 hepatocytes were isolated and cultured in DMEM (10% fetal bovine serum, 1% penicillin/streptomycin, pH 7.4). The cells were allowed to become confluent, and diluted to a concentration of  $5 \times 10^5$  cells/mL before use in the cell viability assay.

### Cell Viability

Cell viability was measured on WB 344 hepatocytes to determine the integrity of the cell membrane. Using the Kodak Ektachem 700XR, the leakage of aspartate aminotransferase (AST) and alanine aminotransferase (ALT) was ascertained. The lactate dehydrogenase (LDH) assay was conducted on the Dupont ACA discrete clinical analyzer.


### Cell Proliferation

A CellTiter 96 Non-Radioactive Cell Proliferation Assay was conducted using a Molecular Devices Thermo max microplate reader. This assay was used to determine the absorbance of the sample which is directly proportional to the number of viable cells.

### EPR Spectroscopy


Cell preparations ( $1 \times 10^6$  cells/ml) were packed in quartz aqueous cells. Using a Varian E4 EPR spectrometer, the EPR spectra were recorded under the following conditions: microwave power, 20 mW; microwave frequency, 9.54 GHz; scan range, 200 G; field set, 3430 G; time constant, 0.5 sec; modulation amplitude, 1 G; and modulation frequency, 100 kHz.

## Results



**Figure 3.** Viability (A-C) and proliferation (D) assay results of WB 344 hepatocytes following a 24 hr. exposure to ADN.


After a 24 hr. exposure to ADN, viability assays were taken to determine the leakage of the enzymes AST, ALT, and LDH. Figures 3A and 3B show the effect of ADN on the leakage of AST and ALT. In both cases, an increase in the ADN concentration is reflected in the greater percentage of enzyme leakage. Figure 3C shows the results of the LDH viability assay. Irrespective of ADN concentration, the assay gave the same LDH leakage value ( $37 \pm 10\%$ ). The results of the cell proliferation test are displayed in Figure 3D. As ADN concentration (mM) in the media was increased, there was a decrease in the number of surviving cells.



**Figure 4.** EPR spectra of hepatocytes and PBN (4A), ADN and PBN (4B), and PBN and hepatocytes with ADN (4C).

Figure 4 displays EPR spectra gathered under the following conditions: scan range, 200 G; field set, 3430 G; time constant, 0.5; modulation amplitude, 1 G; receiver gain,  $10 \times 10^4$ , microwave power, 20 mW; and microwave frequency, 9.54 GHz. All tests were conducted at room temperature.


Figures 4A and 4B, show the EPR spectra obtained when the spin trap 0.02 M PBN was added to WB344 hepatocytes and incubated for 30 min at 37°C, and when 1 M ADN is added to PBN without cells, respectively. For both of these samples, the spectra represent random noise. Figure 4C displays the EPR spectra produced after the 30 minute incubation of ADN in PBN with cells ( $1 \times 10^6$  cells/mL). In this spectrum, the presence of spin adducts are not clear although their formation is beginning.



**Figure 5.** EPR spectra of hepatocytes and DMPO (5A), ADN and DMPO (5B), and cells and DMPO with ADN (5C).

With the exception of receiver gain, the conditions in Figure 5 remained the same as those explained in Figure 4. Figure 5A (receiver gain  $1 \times 10^4$ ) shows the EPR spectrum of the spin trap 0.02 M DMPO and WB 344 hepatocytes ( $1 \times 10^6$  cells/mL) after a 30 min. incubation at  $37^\circ\text{C}$ . Unlike the spectrum of the cells and spin trap from the previous figure (4A), this figure shows a nitroxide triplet of hyperfine coupling constant  $a_N=15.0$ . The hyperfine coupling constants were measured directly from the spectra as the separation in peaks measured in mT. Figure 5B (receiver

gain  $8 \times 10^4$ ) is the spectrum drawn when ADN and DMPO were tested immediately after mixing. Figure 5B consists of two DMPO adducts. The first consists of a nitroxide triplet with similar hyperfine coupling constants as those described in Figure 5A. The second DMPO spin adduct consists of a triplet of triplets suggesting the addition of a nitrogen center to the DMPO. The hyperfine coupling constants of these spin adducts are  $a_N=12.0$  for the primary nitrogen,  $a_{N_2}=5.0$  for the secondary nitrogen. Figure 5C is identical to Figure 5B, but less intense, suggesting that the cells compete for the free radicals of ADN in the presence of DMPO.



**Figure 6.** The EPR/spin trapping spectra over time (6A) and (6B) graph of 10  $\mu$ L NaOH with 1 M ADN in .02 M PBN

Figure 6A shows the spectra from a run in which 10 uL of NaOH was added to .02 M PBN and 1 M ADN. Over time, the peaks increased in intensity. The relative intensity, calculated from the peak height divided by the receiver gain, reached a maximum at  $5.64 \times 10^{-4}$ . Figure 6B is the graph of the relative intensity of the EPR signal over 50 minutes, for three different concentrations (5.2 uL, 10 uL, and 20 uL) of NaOH in PBN and ADN (pH=8.6, 9.3, and 9.7 respectively). In all cases except the 5.2 uL concentration, the intensity of the EPR signal grew significantly over time.

### Discussion

The effects of ADN on cell viability and free radical production in the liver have been studied using EPR/spin trapping and various viability assays. Based on the literature search, this is the first study on the effects of ADN on hepatocytes. Figure 3 demonstrated the effect of ADN on cell viability as determined by AST, ALT, LDH, and proliferation assays. Three of these four tests showed that increasing concentrations of ADN caused decreased cell viability, while in the LDH assay no trend was visible. Figures 4 and 5 show the EPR spectra gathered under different spin traps. The use of PBN as the spin trap indicated the presence of free radicals only in the sample containing ADN, PBN, and hepatocytes. Peaks were identified in all the samples in which DMPO was used as the spin trap. This is due to the fact that DMPO detects oxygen-centered radicals which are formed naturally in the body, yielding DMPO-adducts with a distinctive characteristic pattern. Figure 6 is the result of an experiment in which 10 uL of NaOH was added to .02 M PBN and 1 M ADN. Over time, the intensity of the peaks grew larger, indicating the growing presence of free radicals as time

## References

1. HQ HSC Planning, Requirements and Engineering Marketing Office  
1993. In "Human Systems Center Products and Progress". p117. AFMC, Brookes AFB, Tx.
2. Borman, S., "Advanced energetic materials emerge for military and space applications,"  
Chemical and Engineering News, 1994, 18-22.
3. SRI International, Menlo Park, CA. (1992). Material Safety Data Sheet.
4. Steel-Goodwin, L., Pace, D.W., and Carmichael, A.J., "Ammonium Dinitramide: an  
EPR/ENDOR Study," Conference on Temporal Aspects in Risk Assessment for Noncancer  
Endpoints, 1994, 74.
5. Rice-Evans, C.A., Diplock, A.T., and Symons, M.C.R., "Techniques in free radical research,"  
Laboratory Techniques in Biochemistry and Molecular Biology, ed. Burden, R.H. and van  
Knippenbey, P.H., Elsevier, N.Y., 1991.
6. Halliwell, B. and Gutteridge, J.M.C. Free radicals in biology and medicine. 2nd. ed.  
Clarendon Press Oxford 1989 pp 418.
7. Pryor, W.A., "The Role of Free Radical Reactions in Biological Systems," Free Radicals in  
Biology, Academic Press, Inc., N.Y., 1976.
8. Grisham, M.B., Reactive Metabolites of Oxygen and Nitrogen in Biology and Medicine,  
Medical Intelligence, R.G. Landes Company, Austin, T.X., 1992.
9. Aust, S.D., Chignell, C.F., Bray, T.M., Kalyanaraman, B., and Mason, R.P., "Free Radicals in  
Toxicology," Toxicology and Applied Pharmacology, 1993, 168-178.

elapsed. The pH of the solution appears to be important in determining free radicals in ADN with the spin trap PBN. All tests involving cells had a pH  $6.9 \pm 0.29$  and the spin adducts formed by ADN were unclear.

The tests conducted in this study (Figure 3D) indicate that ADN is toxic to 50% of cells at a concentration of 2.8 mM. EPR/spin trapping data indicates that production of free radicals occurs in hepatocytes in the presence of ADN, and this production increases over time. Further studies must be conducted with various routes of exposure (eg. inhalation, dermal absorption, and ingestion) in order to determine exposure limits and establish safety standards regarding the use of ADN in the workplace.

10. Mosler, M.T. and Smith, C.V., Free Radical Mechanisms of Tissue Injury, CRC Press, Inc., Boca Raton, F.L., 1992.
11. Grundon, M.F. and Henbest, H.B., Organic Chemistry, an Introduction, MacDerald Technical and Scientific, London, 1971.
12. Wertz, J.E. and Bolton, J.R. Electron spin resonance. Elementary theory and practical application, Mc Graw-Hill, NY. 1972.
13. Knecht, K.T. and Mason, R.P., "In Vivo Spin Trapping of Xenobiotic Free Radical Metabolites," Archives of Biochemistry and Biophysics, 1993, 185-194.
14. Gaudin, A.J. and Jones, K.C. 1989. In Human Anatomy and Physiology Chapter 18. The digestive system pp 440-477., Harcourt, Brance, Jovanovich Publ., NY.
15. James, R.C., "Hepatotoxicity: Toxic Effects in the Liver," In Industrial Toxicology, Safety and Health Applications in the Workplace, Van Nostrand Reinhold Co., N.Y., 1985, 78-105.
16. Pathology of the Liver, ed. MacSween, R.M.N., Anthony, P.P., and Scheuer, P.J., Churchill Livingstone, N.Y., 1979.