

**UNITED STATES AIR FORCE
ARMSTRONG LABORATORY**

**TRICHLOROETHYLENE: FREE
RADICAL STUDIES IN
B6C3F1 MOUSE LIVER SLICES**

**Linda Steel-Goodwin
Tasha Pravecek
Barry Hancock
William Schmidt
Stacie Trivunovic
Gerri Miller
Christel Bishop
David Bartholomew
Stephen Channel**

**OCCUPATIONAL AND ENVIRONMENTAL HEALTH
DIRECTORATE TOXICOLOGY DIVISION
ARMSTRONG LABORATORY
WRIGHT-PATTERSON AFB, OH 45433-7400**

**Marcia Ketcha
MANTECH GEO-CENTERS JOINT VENTURE
DAYTON, OH 45437**

**Alasdair Carmichael
ARMED FORCES RADIOBIOLOGY RESEARCH INSTITUTE
BETHESDA, MD 20991**

October 1996

**Occupational and Environmental Health
Directorate
Toxicology Division
2856 G Street
Wright-Patterson AFB OH 45433-7400**

Approved for public release; distribution is unlimited.

19990316 050

NOTICES

When US Government drawings, specifications or other data are used for any purpose other than a definitely related Government procurement operation, the Government thereby incurs no responsibility nor any obligation whatsoever, and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise, as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use, or sell any patented invention that may in any way be related thereto.

Please do not request copies of this report from the Air Force Armstrong Laboratory. Additional copies may be purchased from:

National Technical Information Service
5285 Port Royal Road
Springfield, Virginia 22161

Federal Government agencies and their contractors registered with the Defense Technical Information Center should direct requests for copies of this report to:

Defense Technical Information Service
8725 John J. Kingman Rd., Ste 0944
Ft. Belvoir, Virginia 22060-6218

DISCLAIMER

This Technical Report is published as received and has not been edited by the Technical Editing Staff of the Air Force Armstrong Laboratory.

TECHNICAL REVIEW AND APPROVAL

AL/OE-TR-1996-0079

The animal use described in this study was conducted in accordance with the principles stated in the "Guide for the Care and Use of Laboratory Animals", National Research Council, 1996, and the Animal Welfare Act of 1966, as amended.

This report has been reviewed by the Office of Public Affairs (PA) and is releasable to the National Technical Information Service (NTIS). At NTIS, it will be available to the general public, including foreign nations.

This technical report has been reviewed and is approved for publication.

FOR THE DIRECTOR



STEPHEN R. CHANNEL, Maj, USAF, BSC
Branch Chief, Operational Toxicology Branch
Air Force Armstrong Laboratory

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE October 1996	3. REPORT TYPE AND DATES COVERED Interim Report - February 1994-September 1994	
4. TITLE AND SUBTITLE Trichloroethylene: Free Radical Studies in B6C3F1 Mouse Liver Slices			5. FUNDING NUMBERS Contract F41624-96-C-9010 PE 61102F PR 231Z TA 231ZA2 WU 231ZA202	
6. AUTHOR(S) L. Steel-Goodwin, T.L. Pravecsek, B. Hancock, W.J. Schmidt, S. Trivunovic, G. Miller, C. Bishop, D.G. Bartholomew, and S.R. Channel				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Man Tech Geo-Centers Joint Venture P.O. Box 31009 Dayton, OH 45437-0009			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Armstrong Laboratory, Occupational and Environmental Health Directorate Toxicology Division, Human Systems Center Air Force Materiel Command Wright-Patterson AFB, OH 45433-7400			10. SPONSORING/MONITORING AGENCY REPORT NUMBER AL/OE-TR-1996-0079	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION AVAILABILITY STATEMENT Approved for public release; distribution is unlimited.			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) Electron paramagnetic resonance EPR/spin techniques were used to test the hypothesis exposure of liver to trichloroethylene (TCE) causes increased formation of free radicals. Precision cut liver slices from B6C3F1 mice were incubated for 20 min. with TCE at headspace concentrations ranging from 0-10,000 ppm with and without the spin trap N-tert-butyl-a-phenyl nitron (PBN). PBN (10 mM) did not interfere with K+ levels, lactate dehydrogenase, aspartamine transaminase or alanine transaminase. Free radicals were measured using Bruker ESP300E spectrometer. A solution of 3-carbamoyl-2,2,5,5-tetra methyl-1-yloxy free radical was used as a standard to quantitate radicals. All data was normalized by liver wet weight. Total radicals increased linearly with increasing TCE headspace concentration (TR = {4.9E-04[TCE] + 6.84} X 10E14, r=0.99 P<0.001). EPR/spin trapping results were corroborated by conjugated diene measurements and analysis by gas chromatography.				
14. SUBJECT TERMS trichloroethylene B6C3F1 liver slices free radicals EPR/spin trapping			15. NUMBER OF PAGES 30	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED	18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED	19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED	20. LIMITATION OF ABSTRACT UL	

PREFACE

This is one of a series of technical reports generated for the Free Radical Studies of Trichloroethylene funded by the Air Force Office of Scientific Research. The research presented here began in February 1994 and was completed in September 1994. The results were partially presented at the International Society of Free Radicals Conference in Sydney, Australia in November 1994 and at the Society of Toxicology Meeting in Baltimore in March 1995. This research was part of the Environmental Initiative project, work unit # 231ZA202 managed by Maj. SR Channel, USAF, BSC. Lt Col Terry Childress served as Contract Technical Monitor for the U.S. Air Force, Toxicology Division.

The animals used in this study were cared for under the *Guide for the Care and Use of Laboratory Animals*, prepared by the committee on Care and Use of Laboratory Animals in the Institute of Laboratory Resources, National Research Council, Department of Health and Human Services, National Institute of Health Publication # 86-23, 1985, and the Animal Welfare Act of 1966, as amended.

The authors wish to thank the following for their assistance. C. Booher, OETA, and Capt. B. Garrity, USAF and Maj. J. Williams, USAF, BSC for TCE partition coefficients and Dr. William Jackson, AFRRRI for statistical analysis.

TABLE OF CONTENTS

SECTION	PAGE
PREFACE.....	iii
TABLE OF CONTENTS.....	iv
LIST OF TABLES.....	v
LIST OF FIGURES.....	vi
ABBREVIATIONS.....	vii
INTRODUCTION.....	1
MATERIALS AND METHODS.....	3
Free radicals.....	3
Conjugated dienes.....	3
Viability.....	4
Gas chromatograph.....	4
Statistics.....	4
RESULTS.....	5
DISCUSSION.....	13
REFERENCES.....	16

LIST OF TABLES

TABLE		PAGE
I	Viability Determination of liver slices exposed to 0-10,000 ppm TCE over 20 min. ± PBN.....	10
II	Gas chromatograph data of metabolism of TCE by liver Slices ± 10 mM PBN	11

LIST OF FIGURES

FIGURE		PAGE
1	EPR spectra of (A) Lyophilized liver incubate after 20 min. exposure to 2500 ppm TCE (B) Spectrum after dissolving incubate in water (C) Computer simulated spectrum of B	5
2	Comparison of PBN-spin adducts detected and TCE dose.....	6
3	Absorbance spectrum of conjugated dienes of liver exposed to 0 or 5000 ppm TCE	7
4	Conjugated dienes measure from 0-10,000 ppm in liver slices incubated \pm PBN	8
5	Comparison of free radicals trapped with conjugated dienes	9

ABBREVIATIONS

ANOVA	Analysis of variance analysis
3-CAR	2,2,5,5,-tetramethyl-1-pyrrolidinyloxy-3-carboxamide
dB	decibel
EPR	electron paramagnetic resonance
G	Gauss
mM	millimole/liter
ms	millisecond
mW	milliwatt
PBN	N-tert-butyl- α phenyl nitron
TCE	trichloroethylene

INTRODUCTION

Trichloroethylene (TCE) is a widely used industrial solvent and degreasing agent and is an environmental contaminant in the ground water near landfills and in highly industrialized areas (Atlas 1995). TCE is lipophilic and can diffuse quickly through hepatic parenchyma cell membranes where it is extensively metabolized. Ingestion of this halogen compound causes liver toxicity in B6C3F1 mice (NCI 1976, Elcombe 1985, Dekant et al 1986) and the mechanism (or mechanisms) responsible for this liver toxicity are currently being explored (Channel 1994). The toxic effects of TCE are important from the standpoint of human health and environmental cleanup requirements. Our aim was to perform as many tests as possible (free radical production, viability tests, TCE metabolism) in the same sample of mouse liver to reduce numbers of animals and experimental variation from one experiment to another. This is possible by preparing slices of liver of equal size and thickness and placing it in incubation media containing the chemical of interest (Smith et al 1987). Electron paramagnetic resonance (EPR)/spin trapping techniques are the only definitive methods to study free radicals (Carmichael et al 1993, Carmichael and Steel-Goodwin 1994). EPR/spin trapping techniques were used to examine free radical formation by liver metabolism of TCE in aqueous media. Spin trapping is a technique in which short-lived free radicals are accumulated by an addition reaction to a spin trap generating a long-lived radical product (Mason 1984). This spin adduct is detected by EPR. Understanding of TCE-induced radicals in aqueous media has

potential application for TCE risk assessment and for applied technology for removal of TCE from soil and ground water (Atlas 1995).

METHODS AND MATERIALS

Liver slice preparation

Precision cut liver slices from B6C3F1 mice were pre-incubated for 2 h in Waymouth's control media prior to incubation in control media or media supplemented with 0.01M N-tert-butyl- α -phenyl nitron, PBN (Sigma, St. Louis MO.). TCE (Aldrich, Milwaukee, WI) was added to the headspace of the media at concentrations of 0, 2500, 5000, 7500, or 10,000 ppm TCE. After 20 min. the liver slices and incubate were harvested for free radical, conjugated dienes, viability and gas chromatography analysis.

Free radicals

Free radicals were measured using a Bruker ESP300E spectrometer. The magnetic field was set at: 335.5 mT; microwave frequency: 9.4 GHz; microwave power: 20 mW; modulation amplitude: 0.1 mT; time constant: 0.5 s; scan time: 4 min.; scan range 10.0 mT. The samples were quantitated using a standard solution of 3-carbamoyl-2,2,5,5-tetra methyl-1-yloxy (3-CAR) radical dissolved in bicarbonate solution and lyophilized in the same manner as the liver incubate media.

Conjugated dienes

Conjugated diene concentration was calculated from a standard curve of absorbance of known octadecadienoic acid concentrations (Sigma Chemical Co., St. Louis, MO) read at 235 nm.

Viability

Viability was determined by measuring intracellular potassium content and leakage of the liver enzymes lactate dehydrogenase (LDH), aspartamine transaminase (AST) and alanine transaminase (ALT).

Gas Chromatography

Gas chromatography was used to measure TCE added to the headspace, the partition coefficient of TCE between the headspace and the media, TCE in the media, trichloroethanol and trichloroacetic acid by a modification of the method of Garrett and Lambert 1966).

Statistics

All results were normalized by liver wet weight and analyzed by analysis of variance.

RESULTS

Figures 1-4 describe the formation of radicals after liver slices are exposed to TCE and the subsequent formation of conjugated dienes which suggests TCE induced lipid peroxidation. Tables I & II show the data obtained using established enzymatic and photometry techniques in the laboratory for viability (liver enzymes and K^+) and for TCE and TCE-metabolite quantitation.




Figure 1. EPR spectra . (A) Lyophilized liver incubate after 20 min. exposure to 2500ppm TCE. (B) Spectrum after dissolving lyophilized incubate in water (C) Computer simulated spectrum of B.

Figure 1A is typical of a nitroxide EPR spectrum in solid matrix. This shows the PBN spin adduct EPR spectrum of lyophilized liver incubate after 20 min. exposure to 2500 ppm TCE. The PBN spin adduct EPR spectrum after dissolving the lyophilized liver incubate in water is shown in Figure 1B. Samples generated EPR spectra consisting of a triplet of doublets characteristic of PBN spin adducts. The computer generated EPR spectrum for the EPR spectrum in B is shown in Figure 1C. This spectrum was generated using hyperfine coupling constants $a_N = 1.61$ mT and $a_H = 0.325$ mT.




Figure 2 *Correlation analysis of PBN-spin adducts between TCE concentration and free radicals generated.*

The radicals generated in the incubate of liver samples exposed to 0-10,000 ppm TCE were quantitated using 3-CAR as a standard are shown in Figure 2. Correlation analysis of PBN-spin adducts quantitated using 3-CAR determined a significant link between TCE concentration and free radicals generated ($r = 0.99$ $P < 0.001$).




Figure 3 *Absorbance spectrum of conjugated dienes of liver exposed to 0 or 5000 ppm trichloroethylene.*

Conjugated dienes were measured in the liver slices in these experiments. The typical spectra is shown in Figure 3. The absorbance at 235 nm was used to quantitate the conjugated dienes in the liver slices following exposure to 0-10,000 ppm TCE.




Figure 4 *Conjugated diene measured from 0-10,000 ppm in liver slices incubated \pm 10 mM PBN.*

Generation of free radicals in liver is known to cause lipid peroxidation which can be determined by measurement of conjugated dienes or malondialdehyde (MDA). In these experiments attempts to measure MDA were abandoned because the spin trap

reduced the levels of MDA to levels below detection for the assay. However the levels of conjugated diene measured indicated that as TCE concentration increased, there was a concomitant increase in the concentration of TCE in the control media. The diene concentrations remained the same in the PBN-media suggesting sufficient PBN was present to trap any radicals produced by the liver slices.




Figure 5 *Comparison of free radicals trapped with conjugated dienes.*

Figure 5 is the comparison of the data for the free radicals trapped by PBN (Figure 2) with the difference in conjugated dienes \pm spin trap (Figure 4). The resulting line was linear ($b[0] = -1.1644$, $b[1] = 9.5679$, $r^2 = 0.9289$)

This data was dependent on the assumption that the TCE used in the experiments did not cause any adverse changes in the liver during the course of the experiment. The viability of the liver was determined by measurement of liver enzyme leakage and K^+ leakage. Table I is a summary of the measurements of viability.

TCE ppm	Media	Slices	% ALT	% AST	%LDH	K^+
0	-PBN	n = 22	6 ± 2	1 ± 1	9 ± 3	54 ± 16
0	+PBN	n = 12	7 ± 2	1 ± 0.2	10 ± 2	53 ± 7
2500	-PBN	n = 12	6 ± 1	1 ± 0.3	8 ± 1	63 ± 14
2500	+PBN	n = 12	6 ± 3	1 ± 0.5	10 ± 4	59 ± 5
5000	-PBN	n = 12	6 ± 4	2 ± 1	9 ± 6	49 ± 22
5000	+PBN	n = 12	5 ± 2	1 ± 0.3	8 ± 3	60 ± 4
7500	-PBN	n = 12	7 ± 2	2 ± 0.3	10 ± 4	58 ± 18
7500	+PBN	n = 12	6 ± 1	1 ± 0.2	10 ± 1	57 ± 5
10000	-PBN	n = 18	7 ± 2	2 ± 0.6	10 ± 2	59 ± 13
10000	+PBN	n = 12	6 ± 1	2 ± 0.4	10 ± 2	51 ± 2

Table I *Viability determination of liver slices exposed to 0-10,000 ppm TCE over 20 min. \pm PBN.*

Based on enzyme and K^+ leakage tests the liver slices are viable regardless of the TCE concentration and incubation media. Figures 1 & 2 and Table I suggest radicals formed by liver metabolism of TCE in control media induce lipid peroxidation at levels which are not cytotoxic.

TCE ppm	Media	Slices	TCE ppm added	TCE ug/g liver	TCOH ug/g liver	TCA ug/g liver
0	-PBN	n = 8	0	0	0	0
0	+PBN	n = 8	0	0	0	0
2500	-PBN	n = 8	2460	27±3	116±16	49±4
2500	+PBN	n = 8	2339	120±38*	142±14	31±4*
5000	-PBN	n = 8	5057	98±1	130±12	82±4
5000	+PBN	n = 8	5169	274±27*	141±8	32±4*
7500	-PBN	n = 8	7505	177±10	181±13	54±5
7500	+PBN	n = 8	7646	302±39*	177±10	44±2
10000	-PBN	n = 8	9977	171±18	155±15	61±7
10000	+PBN	n = 8	9846	500±3*	163±23	42±2

* $P < 0.05$

Partition Coefficient -PBN_{media:air} 2.45

Partition Coefficient +PBN_{media:air} 2.24±0.24 (mean±SD)

Table II. Metabolism of TCE by liver slices ± 10 mM PBN.

Table II is the data of TCE metabolism in the control and PBN-supplemented media measured by gas chromatography (GC). Although TCE added to the headspace and the partition coefficients determined in both media were the same, the amount of TCE recovered at the end of 20 min. was greater in the PBN-media than in the control. TCE is believed to be metabolized through free radical pathways and despite being at the limits of detection of the metabolites there was significantly more ($P < 0.05$) TCA measured in the control media than the PBN media at headspace concentrations of 2500

and 5000 ppm TCE. In these experiments there was no difference in the TCOH recovered.

DISCUSSION

The objectives of this study were: (1) determine if free radicals and their subsequent biological effect (lipid peroxidation) could be quantitated in B6C3F1 mice after TCE exposure and (2) corroborate data with established laboratory techniques.

The results in Figures 1 and 2 show that free radicals are generated in aqueous media when liver slices are exposed to TCE and that there is a correlation between these radicals and oxidative stress. It is important to stress that radicals were measured in aqueous media. Previous studies of radical effect of TCE have been carried out in non-polar solvents (Gronthier et al 1989 and Stevens 1994). The majority of biological reactions occur in the aqueous phase of the cell. (von Sonntag 1991). It is therefore important to study radical induced reactions in this phase. In this study 3-CAR was used to quantitate the radicals trapped by PBN. However the radicals trapped by PBN are at most only 20 % of the total radicals in the liver incubate (Steel-Goodwin and Carmichael 1995). In this project it is concluded that liver metabolism of TCE generates PBN spin adducts in aqueous media with hyperfine coupling constants of $a_N = 1.61$ mT and $a_H = 0.325$ mT.

Lipid peroxidation in biological systems involves complicated reactions in which conjugated dienes are only part of the cascade (Rice-Evans et al 1991). Attempts to establish a relationship between TCE addition and malondialdehyde, the main defined lipid peroxidation reaction product measurable in biological systems were not pursued but initial experiments showed no detectable radicals in the PBN-media (results not shown). Radical-induced lipid peroxidation (oxidation of polyunsaturated fatty acids) has

mainly been studied in organic solvents but the difference of radicals quantitated with and without TCE in the aqueous media correlates with the oxidative stress measured from the dienes with and without the spin trap.

The results in Table I and II suggest TCE-generated radicals are selective in their reactions. For example, the TCE-induced radicals were not reactive enough to alter the viability of liver cell membranes at the TCE head space concentrations 2500-10,000 ppm. However the TCE concentration was sufficient to increase in conjugated dienes.

Halogenated hydrocarbons: carbon tetrachloride, trichloroethylene and bromobenzene do induce toxic effects on cells (Steup et al 1991, Ashby et al 1994) but lipid peroxidation (Comporti, 1987., Recknagel et al 1987., Fraga et al 1989) may only be a secondary event, because of cell damage induced by the majority of toxins (Halliwell et al 1992., Gutteridge 1995).

The results shown in Table II suggest free radicals are involved in the liver metabolism of TCE by B6C3F1 mice. The disappearance of TCE at concentrations of 0-10,000 ppm was inhibited when the liver slices were incubated in the spin trap PBN. The results further suggest that the slices exposed to spin trap produced less TCA. Free radical pathways of TCA and DCA production are currently being studied by Stevens, 1994 and there study will not be duplicated. In the present study, the spin trap did not alter the production of TCOH. B6C3F1 mice are a strain which is sensitive to adverse TCE effects, however, free radical studies in human slices could provide important information on potential adverse effects free radicals may cause following TCE exposure. Humans are normally exposed to TCE in low doses in drinking water. Drinking water

resources are finite and TCE-contaminated water, tastes bad. As TCE-induced radicals are generated in aqueous media, radical forming agents could be used to rehabilitate former industrial sites and remove TCE from ground water. A number of techniques are currently being used to reduce TCE in drinking water (Atlas 1995) and these methods may be successful because of their effects on free radicals. Information on free radical effects of TCE could prove useful for studies of potential risks to humans. The risk of potential adverse health effects to humans following exposure to TCE contaminated water supplies is currently based on PD models (Fisher et al 1991 and Fisher and Allen 1993). These models are based on measurements of parent compound and metabolites which have been determined by GC techniques so the GC data, Table II, can in theory be compared to these PD models. In addition, the new parameters: free radicals and conjugated dienes can be added to the model.

Finally, as this study determined that the liver slices were viable, Table I, the results, Figures 1-5, strongly suggest TCE breaks down in liver slices producing free radicals which induce lipid peroxidation. Should this occur *in vivo* in the mouse liver, the resulting lipid peroxidation cascade could lead to impairment of cell junctions or cell death or malignant transformations.

REFERENCES

Ashby, J., A. Brady, C.R. Elcombe., B.M. Elliott, J. Ishmael, J. Odum, J.D. Tugwood, S. Kettle and I.F.H. Purchase 1994 Mechanistically-based human hazard assessment of peroxisome proliferator-induced hepatocarcinogenesis. *Human & Experimental Toxicology* 13 Supplement 2 S1-S117.

Atlas, R .M. 1995 Bioremediation, *Chemical & Engineering News* 73 (14): 32-42.

Carmichael, A.J., L. Steel-Goodwin, B. Gray., and C.M. Arroyo 1993 Reactions of active oxygen and nitrogen species studied by EPR and spin trapping. *Free Rad Res Comms* 19 (1) S1-S16.

Carmichael, A.J. and L. Steel-Goodwin 1994 Decomposition of peroxynitrite studied by EPR/spin trapping. *J. Free Radicals in Biology & Medicine* 2 (3) Q5.

Channel, S.R. 1994 Trichloroethylene: Free radical production, oxidant damage and cell proliferation in B6C3F1 mouse liver. OL/AL Work Unit # 2312A202-Basic Environmental Initiative.

Comporti, M 1987 Glutathione-depleting agents and lipid peroxidation. *Chem Phys Lipids* 45: 143-169.

Dekant, W., A. Schutz, M. Metzler and D. Henschler 1986 Absorption, elimination and metabolism of trichloroethylene: a quantitative comparison between rats and mice. *Xenobiotic* 16: 143-152.

Elcombe, C.R. 1985 Species differences in carcinogenicity and peroxisome proliferation due to trichloroethylene. A biochemical human hazard assessment. Receptors and other targets for toxic substances. *Arch. Toxicol Suppl* 8: 6-17.

Fisher, J.W. and B.C. Allen 1993 Evaluating the risk of liver cancer in humans exposed to trichloroethylene using physiological models. *Risk Analysis* 13 (1) 87-95

Fisher, J.W., M.L. Gargas, B.C. Allen and M.E. Anderson 1991 Physiologically based pharmacokinetic modeling with trichloethylene and its metabolite, trichloroacetic acid in the rat and mouse. *Toxicol & Appl Pharmacol* 109 183-195

Fraga, C.G., B.E. Leibovitz and A.L. Tappel 1988 Lipid peroxidation measured as thiobarbituric acid reactive substances in tissue slices: characterization and comparison with homogenates and microsomes. *Free Rad. Biol & Med* 4: 155-161

Garrett, E.R. and H.J. Lambert 1966 Gas chromatographic analysis of trichloroethanol, chloral hydrate, trichloroacetic acid and trichloroethanol glucuronide. *J. Pharmac. Sci* 55: 812-817.

Gonthier, B.P. and L.G. Barret 1989 *In vitro* spin trapping of free radicals produced during trichloroethylene and diethyl ether metabolism *Tox Lett* 47: 225-234

Gutteridge, J.M.C. 1995 Lipid peroxidation and antioxidants as biomarkers of tissue damage. *Clin Chem* 41: B1819-B1828.

Halliwell, B., J.M. Gutteridge and C.E. Cross 1992 Free radicals, antioxidants and human disease: Where are we now? *J. Lab Clin Med* 119: 598-620

Mason, R.P. 1984 Spin trapping free radical metabolites of toxic chemicals. In *Spin Labeling in Pharmacology* Chapter 2 pp 87-129

National Cancer Institute. 1976 Carcinogenesis bioassay of trichloroethylene CAS No 79-01-6 DHEW Publication No (NIH) 76-802

Ni, Y-C., T.Y. Wong, F.F. Kadlubar, and P.P Fu 1994 Hepatic metabolism of chloral hydrate to free radical(s) and induction of lipid peroxidation *Biochem & Biophys Res Comm* 204 (2): 937-943

Ogino, K., T. Hobara, H. Kobayashi, H. Ishiyama, M. Gotoh, A. Imamura and N. Egami 1991 Lipid peroxidation induced by trichloroethylene in rat liver. *Bull Environ Contam Toxicol* 46: 417-421.

Recknagel, RO, E.A. Glende Jr., J.A. Dolak, R.L. Waller 1987 Mechanisms of carbon tetrachloride toxicity. *Pharmacol Ther* 43: 139-154.

Rice -Evans, C.A., A. T. Diplock and MCR Symons 1991 Techniques in free radical research In *Laboratory Techniques in Biochemistry and Molecular Biology* (RH Burdon & PH van Knippenberg Ed) Elsevier, NY

von Sonntag, C and H-P Schuchmann 1991 The elucidation of peroxy radical reactions in aqueous solution with the help of radiation-chemical methods. *Angew Chem Int Ed Engl* 30: 1229-1283

Steel-Goodwin, L and A. J. Carmichael 1995 Free radicals of TCE: EPR spin trapping study AL/OE-TR1995-0138 .

Stevens, D.K. 1994 Origin, role and fate of dichloroacetic acid in the metabolism of trichloroethylene Status report OL/AL Contract # F33615-91-0539 Mar 94.

Steup, D.R., D. Wiersman, D.A. McMillan and I.G. Sipes 1991 Pretreatment with drinking water solutions containing trichloroethylene or chloroform enhances the hepatotoxicity of carbon tetrachloride in Fischer 344 rats. *Fund. Appl. Toxicol.* 16: 798-809.