




UNITED STATES AIR FORCE ARMSTRONG LABORATORY

USES OF EPR/SPIN LABELING AS A BIOMARKER

L. Steel-Goodwin

K. W. Dean

OCCUPATIONAL AND ENVIRONMENTAL
HEALTH DIRECTORATE TOXICOLOGY DIVISION
ARMSTRONG LABORATORY
WRIGHT-PATTERSON AFB, OH 45433-7400

C. Flemming

MANTECH ENVIRONMENTAL TECHNOLOGY, INC.
DAYTON, OH 45437

F. C. Music

NATIONAL NAVAL MEDICAL CENTER
BETHESDA, MD 20991

A. J. Carmichael

ARMED FORCES RADIOBIOLOGY RESEARCH INSTITUTE
BETHESDA, MD 20991

May 1996

**Occupational and Environmental Health
Directorate**

Toxicology Division

2856 G Street

Wright-Patterson AFB OH 45433-7400

Approved for public release; distribution is unlimited.

19990316 057

NOTICES

When US Government drawings, specifications or other data are used for any purpose other than a definitely related Government procurement operation, the Government thereby incurs no responsibility nor any obligation whatsoever, and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise, as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use, or sell any patented invention that may in any way be related thereto.

Please do not request copies of this report from the Armstrong Laboratory. Additional copies may be purchased from:

National Technical Information Service
5285 Port Royal Road
Springfield, Virginia 22161

Federal Government agencies and their contractors registered with the Defense Technical Information Center should direct requests for copies of this report to:

Defense Technical Information Service
8725 John J. Kingman Rd., Ste 0944
Ft. Belvoir, Virginia 22060-6218

DISCLAIMER

This Technical Report is published as received and has not been edited by the Technical Editing Staff of the Air Force Armstrong Laboratory.

TECHNICAL REVIEW AND APPROVAL

AL/OE-TR-1996-0078

The voluntary, informed consent of the subjects used in this research was obtained as required by Air Force Instruction 40-402.

This report has been reviewed by the Office of Public Affairs (PA) and is releasable to the National Technical Information Service (NTIS). At NTIS, it will be available to the general public, including foreign nations.

This technical report has been reviewed and is approved for publication.

FOR THE DIRECTOR



STEPHEN R. CHANNEL, Maj, USAF, BSC
Branch Chief, Operational Toxicology Branch
Air Force Armstrong Laboratory

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE May 1996		3. REPORT TYPE AND DATES COVERED Interim Report - October 1993-April 1996
4. TITLE AND SUBTITLE Uses of EPR/Spin Labeling as a Biomarker			5. FUNDING NUMBERS Contract PE 61102F PR 2300 TA 2300OT WU 2300OT51	
6. AUTHOR(S) L. Steel-Goodwin, K.W. Dean, C. Flemming, F.C. Music, and A.J. Carmichael				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Armstrong Laboratory, Occupational and Environmental Health Directorate Toxicology Division, Human Systems Center Air Force Materiel Command Wright-Patterson AFB, OH 45433-7400			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Armstrong Laboratory, Occupational and Environmental Health Directorate Toxicology Division, Human Systems Center Air Force Materiel Command Wright-Patterson AFB, OH 45433-7400			10. SPONSORING/MONITORING AGENCY REPORT NUMBER AL/OE-TR-1996-0078	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION AVAILABILITY STATEMENT Approved for public release; distribution is unlimited.			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) Recently protein receptor changes has attracted interest for use as biomarkers in environmental toxicology. Insulin and transferrin which control cell growth were chosen in this study. There were two aims: first, to measure insulin and transferrin receptor binding using spin-labeled insulin and second; basic research on changes in receptor binding of human erythrocytes after exposure to radiation and chemicals likely to be encountered by military personnel in operational or combat scenarios. Insulin and transferrin were spin-labeled and incubated with red blood cells exposed to y-radiation or chemicals (ammonium dinitramide or mustard gas), the cells were centrifuged and receptor binding measured from the EPR spectra using a Bruker ESP300E and EMS104 spectrometer. Transferrin and insulin receptor binding was decreased with ammonium dinitramide. Transferrin receptors are only present in immature erythrocytes. Insulin receptor binding was decreased by y-irradiation and increased by mustard gas. Irradiated mustard gas did not increase insulin receptor binding in red blood cells. As most cells contain insulin receptors, insulin receptor binding is likely to be useful for risk assessment efforts and the spin label technique could be developed as a clinical tool not just for insulin but for other compounds with specific protein receptors on cell membranes.				
14. SUBJECT TERMS EPR/spin labeling transferrin receptors insulin receptor			15. NUMBER OF PAGES 30	
Human ammonium dinitramide and mustard gas biomarker			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED		18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED		19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED
			20. LIMITATION OF ABSTRACT UL	

PREFACE

This is one of a series of technical reports generated from the experimental laboratory programs conducted in the Electron Paramagnetic Resonance (EPR) laboratory of the Armed Forces Radiobiology Research Institute with collaboration with the EPR laboratory at Occupational and Environmental Toxicology Division, Armstrong Laboratory. The collaborative research described in this report began in October 1993 and was completed in April 1996. This work has been presented at the Society of Armed Forces Medical Laboratory Sciences Conference, Washington DC in March 1996 and at the Conference on Advances in Toxicology and Applications to Risk Assessment. This research has been sponsored by the Armed Forces Research Institute work unit 04630 since 1991 and partially by the Air Force Office of Scientific Research, Independent Laboratory Investigator Research, 2300OT51 since 1995. Inquires on details in spin labeled protein methodology and mustard gas data should be directed to AFRRI. Lt Col Terry A. Childress served as Contract Technical Monitor for the U.S. Air Force, Armstrong Laboratory, Toxicology Division.

TABLE OF CONTENTS

SECTION	PAGE
PREFACE.....	iii
TABLE OF CONTENTS.....	iv
LIST OF FIGURES	v
ABBREVIATIONS	vi
1 INTRODUCTION	1
2 MATERIALS AND METHODS.....	4
Chemicals.....	4
EPR/Spin trapping	4
EPR/Spin labeling.....	4
3 RESULTS & DISCUSSION	6
4 REFERENCES	15
5 APPENDIX A.....	17

LIST OF FIGURES

FIGURE	PAGE
1	EPR spectra of (A) Irradiated ammonium dinitramide (B) Irradiated hydrogen mustard (C) Irradiated half mustard dissolved in MNP6
2	(A) Cartoon of the transferrin receptor in a typical membrane. (B) Table of regression analysis of five kinds of doxyl stearic acid and spin labeled transferrin in nucleated red blood cells with and without exposure to ammonium dinitramide7
3	EPR signal intensity versus spin labeled insulin per cell in □ control red blood cells, • ammonium dinitramide treated red blood cells and Δ γ-irradiated red blood cells9
4	EPR signal intensity versus spin labeled insulin per cell in □ control red blood cells, • half-mustard treated red blood cells and Δ γ-irradiated red blood cells11
5	EPR signal intensity versus spin labeled insulin per cell in □ control red blood cells, • half-mustard treated red blood cells, ° γ-irradiated red blood cells treated with half mustard, Δ γ-irradiated red blood cells and red blood cells exposed to Δ γ-irradiated half-mustard12

ABBREVIATIONS

ADN	Ammonium dinitramide
EPR	electron paramagnetic resonance
G	Gray
H-MG	Half mustard gas
rbc	red blood cells
nm	nanometers
m	milli
min	minute
M	mole
MNP	2-methyl-2-nitroso-propane
P	probability
r	correlation coefficient

SECTION 1

INTRODUCTION

Cell membranes can be categorized into different types according to their function and method of permeability. For example, liver cells have type I membranes. The typical type I membrane is about 5 nm thick and consists mainly of lipoid material mixed with some protein (Albert, 1985). Initial studies for this project were performed using liver cells (Carmichael et al., 1995 and Steel-Goodwin & Dean., 1995). Liver cells were excellent for our initial studies because as Type I membranes they represent the commonest type of membrane found in living organisms. However, liver cell receptor isolation is impractical for screening occupational and environmental risks in field studies. Based on our initial experiments, we assumed for this study that spin labels would be potential biomarkers of cell membrane damage.

The goal of this study was to express possible transitions of a protein receptor assay for occupational and environmental health studies. For transition studies we used human red blood cells which are classified as Type II membranes. Type II membranes are adapted to hasten the absorption of metabolites which could not be expected to penetrate a Type I membrane because of ionization or an excessively hydrophilic structure.

Red blood cells are the most abundant cells in the body and red blood cells contain many different protein receptors depending on their age and functional needs. For example, transferrin receptors which bind to the iron binding protein transferrin are

present in immature blood cells. However, when red cells mature and lose their nucleus, they lose the transferrin receptor. Insulin receptors are also present in red blood cells. Insulin receptors are present in almost every cell of the body. Unlike transferrin receptors, they are present in mature red blood cells. Many hazardous chemicals exert their deleterious effects through receptors or by dissolving into the lipid portions of the cell membrane and acting by altering the biophysical properties of those membranes. This study focused on the effects of ammonium dinitramide (ADN), and it transitioned to studying the effects of half mustard (H-MG) on the red blood cell membrane.

The Air Force is currently determining the suitability of using ADN, a high energy oxidizer, to improve solid rocket fuel performance. The Army is exploring neutralization and neutralization plus biodegradation as an alternative to incineration of schedule I chemicals such as mustard gas.

This study had three aims:

1. To determine if ADN exerts a more predictive effect on the lipid or the protein portion of the red cell membrane when the effects of ADN on labeled probes which bind to the lipid portion of membranes and the protein receptors of transferrin in red cells are compared.
2. To compare the effects of ammonium dinitramide on the type of labeled probe which provides the best prediction in (1) to other deleterious effects such as cell irradiation and H-MG.
3. To see if the labeled probe used in (2) can demonstrate that radiation of H-MG neutralizes H-MG effects on red blood cell membranes.

The probes that were used were non-radioactive to decrease the use of hazardous materials in the laboratory. The probes were spin labels which are stable free radicals. Free radicals are paramagnetic and can be detected by electron paramagnetic resonance spectroscopy (EPR). Spin labels have been used for over twenty years to determine the physical structure of biological compounds. Spin labels are available commercially.

SECTION 2

METHODS

Chemicals

Insulin, transferrin and the spin label succinimidyl 2,2,5,5 tetra methyl-3-pyrroline-1-oxyl-3 were purchased from Calbiochem, Sigma and Molecular Probes respectively. Insulin and transferrin were spin labeled by a modification of the technique of Twining et al (1981). The concentration of the spin labeled proteins were determined by spectrophotometry. Other spin labels: 5-, 7-, 10-, 12-, 16, and 20-doxyl stearic acid were purchased from Sigma and Aldrich. Ammonium dinitramide (ADN) was obtained from SRI, CA., and half mustard (H-MG) was obtained from the USA.

EPR and EPR/Spin trapping

ADN and H-MG was irradiated in a ^{60}Co γ -ray source. The dose rate was 1 Gy per minute. The total dose was 25 Gy for ADN and 50 Gy for H-MG. The samples were immediately measured in a Bruker EMS 300E spectrometer. H-MG was spin trapped using the trap 2-methyl-2-nitroso-propane (MNP). MNP was purchased from Aldrich and was prepared as described previously (Carmichael & Reisz 1984).

EPR/Spin labeling

The experimental details for preparation of red blood cells and EPR/spin labeling have been described previously (Steel-Goodwin 1991 and Carmichael et al 1995). Briefly,

packed human red blood cells, prepared at the National Naval Medical Center Blood Bank, Bethesda MD or the clinical pathology laboratory OET, were exposed to ADN, 15 Gy γ -radiation from a ^{60}Co -gamma ray source, H-MG, or irradiated H-MG and then receptor binding was determined using a spin labeled probe. The amount of spin-label bound to the erythrocytes was measured using a Bruker EMS104 or a Bruker ESP300E spectrometer using techniques described previously (Steel-Goodwin & Hutchens 1995). Red cell numbers were measured using a clinical blood cell counter. The relative EPR signal intensity was measured and statistically monitored by analysis of variance.

Statistics: All data was analysed by regression analysis. Scatter plots were performed on spin labeled insulin.

SECTION 1

RESULTS & DISCUSSION

Figure 1 shows the spectra of solid ADN and H-MG following exposure to 25 Gy γ -irradiation. High frequency electromagnetic radiation, from UV via X-rays to γ -rays and also high energy particles such as neutrons, protons or α -particles (helium nuclei) if they have sufficient energy, have the capability to eject electrons from chemicals such as ADN and H-MG. In solid-state studies, EPR spectroscopy is the best detection method for studying radical intermediates following radiolysis (Rice-Evans et al 1991). Figure 1 shows a signal was detected by each chemical studied.





Figure 1 EPR spectra of (A) Irradiated ammonium dinitramide (B) Irradiated hydrogen mustard (C) Irradiated half mustard dissolved in MNP.

The radicals generated by irradiation of ADN, Figure 1A have been described elsewhere (Steel-Goodwin et al 1995). This is the first demonstration of the radical

generated by H-MG, Figure 1B. On dissolution in the spin trap, MNP, the spectrum shown in Figure 1C was obtained. ADN



A



B

Label	r^2	F-Value	P-value
5-doxyl-	0.64	85.9	$4.05E^{-22}$
7-doxyl-	0.61	50.2	$1.54E^{-22}$
10-doxyl-	0.45	25.7	$2.91E^{-22}$
12-doxyl-	0.65	87.9	$2.03E^{-22}$
16-doxyl-	0.55	38.3	$2.61E^{-16}$
transferrin	0.92	288	$1.75E^{-37}$

Figure 2 (A) Cartoon of the transferrin receptor in a typical membrane. (B) Table of regression analysis of 5-, 7-, 10-, 12-, and 16-doxyl stearic acid and spin labeled transferrin in red blood cells with and without exposure to ammonium dinitramide

Figure 2A shows a cartoon of the structure of the transferrin receptor and Figure 2B is the table of the regression analysis results of the changes in EPR signal intensity of

each spin label tested with and without ADN. The regression data shows that the commercially available spin labels are not as specific as the transferrin spin label synthesized in-house. The doxyl spin labels enter the lipid portion of the membrane while the spin labeled transferrin is specific for the transferrin receptor. This experiment suggested that the spin labeled proteins synthesized provided more specific data than the non-developmental items already available commercially. Spin labeled stearic acid has been used to study membrane fluidity. Our assumption for the stearic acid studies was that ADN would affect membrane fluidity of red blood cells as well as their protein receptors because it is an oxidizer. ADN is a hydrophilic chemical and the regression analysis of the spin labeled stearic acid study, Figure 2B suggests ADN has a more selective effect on the proteins in cell membranes, such as transferrin receptors, than lipids. In these studies, transferrin receptor changes only measure the effects of ADN on immature red blood cells. The data results are given in Appendix A.

Both mature and immature red blood cells have insulin receptors. Figure 3 shows the curve of spin-labeled insulin binding in non-exposed red blood cells. The insulin binding curve was depressed following exposure to 1 mM ADN or γ -radiation. The dose of radiation, 15 Gy, is the dose approved by the Food and Drug Administration for preparation of red blood cells prior to transfusion into neonates or immunocompromised patients. At present there is no biomarker to ascertain if irradiated blood has any adverse effects on these patients. Irradiation of blood is necessary for patient survival to prevent transfusion associated graft versus host disease (Leitman, 1989).




Figure 3 *EPR signal intensity versus spin labeled insulin per cell in $\square$ control red blood cells, $\bullet$ ammonium dinitramide treated red blood cells and Δ γ -irradiated red blood cells.*

The major function of red blood cells is to transport oxygen, a paramagnetic molecule. Oxygen transport is carried out in red cells by hemoglobin. Hemoglobin is carried bound in the red blood cells because it is toxic. Hemolysis occurs when the red blood cell membrane is damaged. Hemolysis of red cells can occur as a result of exposure of cells to radiation, physical damage, chemical damage or aging of the cell membrane. This spin label assay could be tailored to a clinical setting to determine if red blood cells are functioning correctly and are suitable for human transfusion.

ADN will cause hemolysis to red blood cells at concentrations of greater than 10 mM (results not shown). In these experiments, ADN had the same effect as irradiation on insulin receptor binding. This assay could be used as a biomonitor to determine risk of

ADN exposure in the work environment. The assay could even be tailored to mitigate risk of adverse effects on wild life exposed to ADN environmentally.

The mission boundaries of occupational and environmental toxicology are medical, safe work practices and environmental risk assessment related to health and safety of the environment and the worker. Health risks to workers can be much greater in a disposal or a clean-up environment than in a production site or laboratory. In production and the laboratory, the same work is performed in the same location and this work is repeated day after day, week after week and sometimes year after year. The health risks to the worker in production sites and laboratories are more likely to be well defined because of this repetition. However, at a disposal facility or in a clean-up site, workers could continually face new health hazards as the chemicals sent for disposal and the clean-up sites vary. This poses many challenges to occupational health risk assessment and also challenges to the type of assay chosen to determine a biological effect. A spin label assay could be used to determine changes in membranes. As the assay involves receptor effects even artificial membranes with purified receptors worn as a dosimeter or set at certain locations could be used. This could remove the requirement to draw blood and the receptor would be purified so binding should theoretically be more specific. As the technique of spin labeling requires the use of an external magnetic field and a means to detect the change in the magnetic field reading, this technique, in theory, could lend itself to remote monitoring or sensing.




Figure 4 *EPR signal intensity versus spin labeled insulin per cell in $\square$ control red blood cells, $\bullet$ half-mustard treated red blood cells and Δ γ -irradiated red blood cells*

Figure 4 shows the erythrocyte binding response to spin labeled insulin after exposure to H-MG and γ - irradiation. In Figure 4, the relative EPR signal intensity was recorded with the concentration of spin-labeled insulin/cell. All data was plotted on the same curve. Binding of red blood cells to unexposed cells is curvilinear showing maximal binding to the spin labeled insulin. Equilibration studies between red blood cells and H-MG were not found in the literature but in vivo injections of ^{35}S -labeled MG suggest rapid diffusion and distribution of the radioactivity to all tissues were it becomes firmly fixed or bound to protein complexes (Bournsnel et al 1946). In the present experiments

binding to insulin receptors in irradiated and non-irradiated red blood cells was increased with H-MG exposure but this excess binding was depressed when non-irradiated red blood cells were incubated with γ -irradiated H-MG. This experiment shows that H-MG does alter insulin binding to red blood cells. Also, there was no synergistic effect if red blood cells were exposed to γ -irradiation and then H-MG.




Figure 5 *EPR signal intensity versus spin labeled insulin per cell in $\square$ control red blood cells, $\bullet$ half-mustard treated red blood cells, $\circ$ γ -irradiated red blood cells treated with half mustard, Δ γ -irradiated red blood cells and red blood cells exposed to Δ γ -irradiated half-mustard.*

However, if the half mustard is irradiated and then exposed to red blood cells the biological effect of the H-MG on the red blood cells was drastically reduced, Figure 5. Therefore, the spin-labeled insulin did provide support to the theory that radiation will

mitigate the effects of H-MG. This assay could therefore potentially have uses as a biomonitor if radiation is used to destroy MG.

Defense of armed forces against the effects of chemical weapons is, nowadays, an essential component of defense in an international context. The requirements are numerous, complex, expensive but a necessity. Many Iranian victims of mustard gas attacks were treated in hospitals all over Europe in 1984-1986. The biological effects of mustard gas have been recently reviewed by Dacre and Goldman 1995 and Marrs et al., 1996. Foreign governments, including the U.S., recognizing the danger of unlimited and uncontrolled exportation of chemicals, have initiated means to control the export of chemicals which can be used for the production of chemical weapons. The mustard gases were chosen in this study because they are by far the easiest of chemical warfare agents to manufacture (Franse et al 1986) and there is an interest in finding ways of monitoring their destruction.

In conclusion, this poster shows the data obtained using spin-labeled probes to study red blood cell membrane damage. Spin-labeled probes are a more environmentally safe method to study receptor binding than radiolabelled assays, they are inexpensive to make, and the results can be read within 60 seconds even using the portable EMS104 spectrometer. Thus, there is potential to use the technique of spin labeling to study biological effects for occupational and environmental toxicology. The spin-label technique also has the potential to be adapted to other biomolecules suggesting it can be tailored to the needs of customers in the fields of environmental chemistry and medicine. The spin label assay could be used to investigate ways to reduce, eliminate, recycle and dispose of

hazardous chemicals, as an alternative tool to radiolabeled insulin (Gambir et al 1977., Ogunwole et al 1986., MacDonald et al 1991 and Grunberger 1993) in clinical medicine and for quality control of irradiated blood.

The technique of EPR/spin labeling:

- is a cost effective
- an alternative to radiolabels
- allows multilabeling
- is a method that provides extreme sensitivity of measurement
- reagents are easy to use
- system is environmentally safe
- is a proven technology.

REFERENCES

- Albert, A.** (1985) *Selective Toxicity. The Physico-chemical basis of therapy.* 7th edition Chapman & Hall London
- Boursnell, J.C. , J.A. Cohen, M. Dixon, G.E. Francis, G.D. Greville, D.M. Needham and O Wormall** 1946 Studies on mustard gas ($\beta\beta'$ -Dichloroethyl sulfide) and some related compounds. 5. The fate of injected mustard-gas (containing radioactive sulfur) in the animal body. *Biochem.J.* 40: 756-764.
- Carmichael, A.J. & P Reisz** (1984) *Photoinduced reactions of anthraquinone antitumor agents with peptides and nucleic acid bases: An electron spin resonance and spin trapping study.* *Arch Biochem and Biophys* 237 (2) 433-444
- Carmichael, A. J., Armengol, J., Young, A., Dean, K.W., Pace, M.D., and Steel-Goodwin, L.** (1995) *Biological effects of the nitric oxide releasing agent ADN: An EPR/spin labeling study* International Free Radical Meeting, Denver CO.
- Dacre, J.C. and M. Goldman** 1995 *Toxicology and Pharamacology of the chemical warfare agent sulfur mustard-A review* Technical Report 9202 US Army Biomedical Research and Development Laboratory Fredrick, MD AD A294927
- Franse M., H. Westerbroek, W. Hielkema, S. Jahromi, D. Bonn and F. Stunnenberg** (1986) *Chemical Weapons: Chaos and Misunderstanding* p 71-76. Second World Congress, New Compounds in Biological and Chemical Warfare: Toxicological Evaluation Industrial Chemical Disasters, Civil Protection and Treatment , The International Association of Forensic Toxicologists, 23rd European International Meeting
- Gambir, K.K. & Carter, C** (1977) *Insulin radioreceptor assay for human erythrocytes* *Clin Chem* 23 (9) 1590-1595
- Grunberger G.** (1993) *Insulin and IGF-I receptor function among Type II diabetic responders and nonresponders to glybriide* *J. Lab. Clin. Med* 121,551-61
- Leitman, S.F.** (1989) *Use of blood cell irradiation in the prevention of graft versus host disease.* *Transfusion Sci.* 10 212-232
- MacDonald, R.S., L. Steel-Goodwin and R.J. Smith** (1991) *Influence of dietary fiber on insulin receptors in rat intestinal mucosa.* *Ann Nutr Metab* 35: 328-338

Marrs, TC., R.L. Maynard, F.R. Sidell (1996) *Mustard Gas* In Chemical Warfare Agents: Toxicology and Treatment Chapter 6. J. Wiley & Sons Ltd, England

Ogunwole, J.O., E.M. Knight, J.C. Adkins, R.H. Ponter (1986) *Effects of dietary fiber and protein on rat erythrocyte insulin receptor interaction*. Nutr Res 6, 673-688.

Rice -Evans, C.A., A. T. Diplock and MCR Symons (1991) *Techniques in free radical research* In Laboratory Techniques in Biochemistry and Molecular Biology (RH Burdon & PH van Knippenberg Ed) Elsevier, NY

Steel-Goodwin, L (1991) *Mouse erythrocyte insulin receptor analysis* EP03 Principal Investigator Radionucleotide Authorization # A91-25-002 Nuclear Regulatory Commission License 19-08330-02.

Steel-Goodwin, L and Hutchens B 1995 *Efficiency of PBN to trap 3-CAR in B6C3F1 mouse liver slices: An EPR study* AL/OE-TR-1995-0139 Technical Report

Steel-Goodwin, L. and Dean, K.W. *Apoptosis in Liver Cells on Exposure to Ammonium Dinitramide* Ann Clin and Lab Sci 25 (3) 1995

Steel-Goodwin, L., M.D. Pace and A.J. Carmichael (1995) *Ammonium Dinitramide: An EPR/ENDOR Study* AL-OE-TR-1995-0160

Twining, S.S., R.C. Sealy and D.M. Glick (1981) *Preparation and activation of a spin labeled pepsinogen*. Biochem 20 1267-1272

APPENDIX A

Chemical Codes for Appendix A

1 = 5-doxyl-; 2 = 7-doxyl-; 3 = 10-doxyl-; 4 = 12-doxyl-; 5 = 16-doxyl-;

6 = transferrin

chemical	LABEL	READING
1	20	1110.1
1	20	1111
1	20	1108.6
1	18	658.8
1	18	680.8
1	18	685.2
1	16	616.3
1	16	621.7
1	16	575.9
1	14	221.4
1	14	232
1	14	234.5
1	12	233
1	12	243.1
1	12	243.9
1	10	143.1
1	10	153
1	10	152.4
1	8	213.3
1	8	217.8
1	8	218.8
1	6	116.5
1	6	120
1	6	120.1
1	4	66.47
1	4	68.54
1	4	69.18
1	2	29.6
1	2	29.26
1	2	29.08
1	0	8.6
1	0	9.01
1	0	8.24
2	20	414.1
2	20	418.2
2	20	420.5
2	18	260.2
2	18	282.3
2	18	287
2	16	195.5
2	16	202.8

2	16	204.8
2	14	140.8
2	14	138.1
2	14	138.8
2	12	72.97
2	12	77.88
2	12	78.72
2	10	48.21
2	10	49.65
2	10	49.14
2	8	50.19
2	8	50.76
2	8	50.54
2	6	39.85
2	6	40.87
2	6	40.06
2	4	35.06
2	4	35.34
2	4	34.95
2	2	29.38
2	2	28.22
2	2	28.39
2	0	8.44
2	0	8.78
2	0	8.79
3	20	1177.7
3	20	1208.6
3	20	1222
3	18	839.8
3	18	933.2
3	18	975.8
3	16	737
3	16	788.7
3	16	808.5
3	14	570.1
3	14	611.7
3	14	631.9
3	12	329
3	12	362.6
3	12	350
3	10	305.6
3	10	313.3
3	10	322.3
3	8	352.3
3	8	369.5
3	8	382.7
3	6	333.7
3	6	336.5
3	6	335.9
3	4	358.6
3	4	363
3	4	351.6
3	2	227.7
3	2	244.2

3	2	236.6
3	0	109.8
3	0	109.7
3	0	103
4	20	806.7
4	20	825.6
4	20	830.7
4	18	922.6
4	18	927.3
4	18	927.2
4	16	390.5
4	16	426.5
4	16	432.6
4	14	406.2
4	14	446.7
4	14	452.4
4	12	308.6
4	12	335.1
4	12	336.8
4	10	268.3
4	10	284.6
4	10	287.7
4	8	227.9
4	8	238.6
4	8	239.6
4	6	190
4	6	197.1
4	6	197
4	4	171
4	4	174.2
4	4	174
4	2	11.75
4	2	12.88
4	2	11.67
4	0	8.12
4	0	8.58
4	0	8.05
5	20	244.2
5	20	245.7
5	20	246
5	18	175.2
5	18	170.9
5	18	174.6
5	16	101.1
5	16	105.6
5	16	107.8
5	14	133.5
5	14	142
5	14	142.3
5	12	103.4
5	12	96.91
5	12	106.2
5	10	80.91
5	10	92.17

5	10	96.08
5	8	71.29
5	8	78.47
5	8	80.64
5	6	102.8
5	6	106.2
5	6	106.5
5	4	65.64
5	4	69.76
5	4	69.84
5	2	8.33
5	2	8.33
5	2	8.49
5	0	9.56
5	0	8.32
5	0	8.4
6	20	1006.5
6	20	1004.3
6	20	1002.3
6	18	867.9
6	18	880.4
6	18	879
6	16	725.2
6	16	732.4
6	16	730.8
6	14	716.5
6	14	721.7
6	14	718.9
6	12	598.7
6	12	601.6
6	12	605.8
6	10	371.6
6	10	373.7
6	10	370.2
6	8	373
6	8	350
6	8	357.2
6	6	209.6
6	6	210.5
6	6	217.1
6	4	89.29
6	4	88.21
6	4	85.16
6	2	47.85
6	2	44.95
6	2	46.4
6	0	34.36
6	0	31.64
6	0	30.86