

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this

1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE APRIL 5 1999	3. REPORT TYPE AND DATES COVERED FINAL REPORT JULY 1993 - SEPT. 30, 1998	
4. TITLE AND SUBTITLE DIGESTIVE KINETICS DETERMINES BIOAVAILABILITY OF POLLUTANTS		5. FUNDING NUMBERS N00014-93-1-1210	
6. AUTHOR(S) LAWRENCE M. MAYER			
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) UNIVERSITY OF MAINE ORONO ME 04469		8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) OFFICE OF NAVAL RESEARCH 800 N. QUINCY ST. ARLINGTON VA 22217-5000		10. SPONSORING / MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES			
12a. DISTRIBUTION / AVAILABILITY STATEMENT DISTRIBUTION UNLIMITED		12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) The bioavailability of sedimentary contaminants to animals in harbor sediments was addressed by studying the mechanisms by which animals solubilize contaminants during feeding and digestion. Digestive physiology work on many different animal species revealed patterns of enzymes, surfactants and dissolved organic matter that correlate with feeding mode, phyletic position, and diet. Incubation of digestive fluids to dissolve contaminants from polluted sediments was developed to provide numerical estimates of bioavailability, and showed that much higher fractions of total contaminant loading are available than predicted by currently established, aqueous equilibrium approaches. The kinetics of reactions are slow enough that variations in feeding rates will influence overall bioavailability. Experimental manipulations showed mechanisms of bioavailability. Dissolved amino acids, in the form of enzyme proteins and hydrolyzed food, are responsible for solubilization of metals such as copper. At high levels, copper can inactivate digestive enzymes. Metals in sedimentary sulfide minerals were largely impervious to digestive fluid attack. Surfactants are responsible for most solubilization of polycyclic aromatic hydrocarbons (PAH), though other agents also appear to play a role. Bioavailability of both metals and PAH can be limited by saturating the digestive agents responsible for their dissolution.			
14. SUBJECT TERMS BIOAVAILABILITY SEDIMENTS		CONTAMINANTS METALS	PAH
		15. NUMBER OF PAGES 4	
		16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED	18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED	19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED	20. LIMITATION OF ABSTRACT UL

Grant #: N00014-93-1-1210

PRINCIPAL INVESTIGATOR: Dr. Lawrence M. Mayer

INSTITUTION: University of Maine

PROJECT TITLE: Digestive Kinetics Determines Bioavailability of Pollutants

REPORTING PERIOD: 1 July 1993 - 30 September 1998

AWARD PERIOD: 1 July 1993 - 30 September 1998

OBJECTIVES: (1) Determine digestive physiology of deposit feeders; (2) Determine the fraction of total pollutants in sediments released during incubation of polluted sediments with digestive fluid of deposit feeders; (3) Examine partitioning of pollutants in sediments, mechanism of digestive fluid solubilization, and design an *in vitro* method to measure their bioavailability.

APPROACH: (1) Use enzyme and surfactant assays to study digestive capability of a variety of benthic invertebrate animals for lipid component of sediments; identify surfactant compounds in animal guts and digestive products of lipid digestion; (2) Extract gut juices from benthic invertebrates and incubate them with polluted sediments, followed by measurement of pollutants released during the incubation; (3) After determining patterns of digestive agents in various animals, design cocktail of commercially available enzymes and surfactants that can be used to mimic the pollutant release kinetics found in part (2), and to explore phase associations of sedimentary pollutants.

ACCOMPLISHMENTS: We surveyed the digestive enzyme and surfactant activities of 18 benthic invertebrate animals. We focused on deposit feeders from the U.S. east and west coasts, but included carnivores, herbivores, and filter feeders for context. Enzyme and surfactant activities vary across both functional and phyletic categories. Polychaetes have higher enzyme activities than echinoderms, and detritivorous polychaetes have higher protease:lipase ratios than herbivorous and carnivorous species. Intense surfactant activities are found in deposit feeders. Less intense surfactancy is found in carnivores, herbivores and filter feeders. The surfactants from *Arenicola marina* were found at mM concentrations, consisting of a C9 fatty acid connected via amide bond to an amino acid. The fatty acids show branching and saturation. Feeding experiments with an omnivorous polychaete showed that varying diet induces changes in protease:lipase ratio similar to cross-phyletic patterns. Sediment in the diet had no effect on the surface tension of digestive fluid. Sediment did induce higher surfactant concentrations, as indicated by micelle formation, which can affect contaminant solubilization. Gut fluids have high dissolved organic matter concentrations, with some polychaetes showing dissolved amino acids alone of 5-10%. Amino acids make up roughly half of the dissolved organic matter. Lipids are much less concentrated. Circumstantial and experimental data showed that animals can concentrate dissolved materials (e.g., amino acids, PAH, and radiolabelled polymers) in guts by retaining fluids relative to solids, which can affect subsequent absorption.

19990426 000

Dissolved trace metal concentrations in digestive fluids are very high, with several species having ppm levels. Levels of some metals are proportional to amino acid concentrations, with correlations evident across species, gut section and molecular weight fraction. Metal enrichments in gut fluids, relative to sediments, showed the Irving-Williams order, consistent with soft ligands (e.g., amino acids) as the solubilizing agents.

Metals in contaminated sediments were released during incubation with digestive fluid to a much greater extent than in incubation with seawater. Kinetics of release to digestive fluid were complex among metals and sediments, including even biphasic patterns with opposite signs (early adsorption followed by release). For highly contaminated sediments, the release rate for a metal like Cu is slow enough that incomplete solubilization will occur during gut passage of sediment through the animal. These kinetic patterns will affect the animals' ability to bioaccumulate metals.

The amount of Cu solubilized by different molecular weight fractions of digestive fluid is related to their peptide content. Solubilization is due to complexation by proteins rather than their enzymatic activity. We could mimic Cu release with solutions of off-the-shelf proteins at similar concentrations as are found in digestive fluid. Histidine residues are critical for Cu binding by digestive fluids. Both high molecular weight proteins, representing enzymes secreted by the animal, and lower molecular weight peptides, representing nutritional material solubilized from sediment, were important in releasing Cu from contaminated sediments. Low molecular weight fractions were found to be more effective solubilizing agents, per amino acid residue, likely due to greater exposure to solution. Ion-specific electrodes were used to determine conditional binding constants between Cu and gut ligands, and ranged from 10^{-4} to 10^{-14} , consistent with known histidine affinities for Cu.

Comparison of digestive fluid extraction with the acid-volatile sulfide (AVS) method showed agreement. Digestive solubilization of borderline metals (Cu, Cd, Ni, Zn, Pb) occurred only when these metals were present in concentrations in excess of the AVS, consistent with the relative binding affinities of histidine vs. sulfide.

Solubilization of Cu from sediment can result in inactivation of digestive fluid enzymes. Inactivation occurs at similar ratios of Cu to gut amino acids, across enzyme type and animal species.

Polycyclic aromatic hydrocarbons (PAH) were also found to be solubilized by digestive fluids much more than by seawater. PAH solubilization was found to depend strongly on the solid-solution ratio and sedimentary organic carbon content. At in vivo solid-solution ratios, PAH solubilized by digestive fluid was inversely dependent on sedimentary OC concentration. Increasing the solid-solution ratio decreased the proportion of total PAH extractable by digestive fluid. These two sets of results imply a strong, fugacity-driven partitioning between digestive fluid and sediment. Using dilution experiments that tested for PAH solubility above and below the critical micelle concentration (CMC) of the surfactants in gut fluids, we established an important role of surfactants in PAH solubilization. Solubilities of heavy and light PAH are similar in gut fluid micelles, presumably due to similar limitation by micellar volume. Proteins may also solubilize PAH and other hydrophobic materials (e.g., methyl Hg). PAH solubilization from contaminated sediments is much less than from pure PAH solids, implicating competition for micellar space from other sedimentary lipids. PAH solubilization from highly contaminated sediment is also limited by saturation of the micelles. Hence

bioavailability from such sediments will be more a function of surfactant secretion by the animal than of inherent solubility from the sediment matrix. Repeatedly incubating sediment with fresh batches of gut fluid released similar amounts of PAH, consistent with saturation behavior. The kinetics of PAH solubilization are rapid, reaching completion within a gut residence time. Absorption of PAH by gut walls is dominantly by passive diffusion.

CONCLUSIONS: Digestive fluid contains agents - e.g., amino acids and surfactants - that significantly increase exposure of benthic animals to contaminants during digestion. Variations in bio/chemical properties among animal species and among sediments can explain patterns of bioavailability among animal-sediment combinations. The interactive chemistry of these two "reactants" governs bioavailability.

SIGNIFICANCE: Lack of understanding of bioavailability is the major hindrance in the application of science to contaminant management issues. This study provides a scientific basis for the failure of current regulatory models to explain animal exposure to sedimentary contaminants in harbors. It provides a basis for more accurate understanding and routine determination of contaminant bioavailability and hence risk assessment.

PUBLICATIONS:

- Mayer, L., Z. Chen, R. Findlay, J. Fang, S. Sampson, L. Self, P. Jumars, C. Quet el, & O. Donard, 1996. Bioavailability of sedimentary contaminants subject to deposit-feeder digestion, *Envir. Sci. Tech.*, 30:2641-2645.
- Mayer, L.M., L.L. Schick, R.F.L. Self, P.A. Jumars, R. H. Findlay, Z. Chen, S. Sampson, 1997. Digestive environments of benthic macroinvertebrate guts: Enzymes, surfactants, and dissolved organic matter, *J. Mar. Res.*, 55:1-30.
- Chen, Z. and L. Mayer, 1998. Mechanisms of Cu solubilization during deposit-feeding *Envir. Sci. Tech.*, 32:770-775.
- Chen, Z. and L. Mayer, 1998, Digestive proteases of the lugworm, *Arenicola marina*, inhibited by Cu from contaminated sediments, *Envir. Toxic. Chem.*, 17:433-438.
- Weston, D. and L. Mayer, 1998. *In vitro* digestive fluid extraction as a measure of the bioavailability of sediment-associated polycyclic aromatic hydrocarbons: Sources of variation and implications for partitioning models, *Envir. Toxic. Chem.*, 17:820-829.
- Weston, D. and L. Mayer, 1998. Comparison of *in vitro* digestive fluid extraction and traditional *in vivo* approaches as measures of polycyclic aromatic hydrocarbon bioavailability from sediments, *Envir. Toxic. Chem.*, 17:830-840.
- Chen, Z. and L. Mayer, 1999. Assessment of sedimentary Cu availability: A comparison of biomimetic and AVS approaches, *Envir. Sci. Tech.*, 33:650-652.
- Chen, Z., and L. Mayer 1999. Sedimentary metal bioavailability determined by the digestive constraints of marine deposit feeders: gut retention time and dissolved amino acids, *Mar. Ecol. - Prog. Ser.*, 176:139-151.
- Bock, M. & L. Mayer, in press. Digestive plasticity of the marine benthic omnivore *Nereis virens*, *J. Exp. Mar. Biol. Ecol.*
- Lawrence, A.L., K.M. McAloon, R.P. Mason, and L.M. Mayer, in press. Intestinal solubilization of particle-associated organic and inorganic mercury as a measure of bioavailability to benthic invertebrates, *Envir. Sci. Tech.*

ABSTRACTS

- Plante, C., Mayer, L., and Watling, L. (1994) The role of surfactants in deposit-feeder digestion. AGU-ASLO Ocean Sci. Mtg.
- Mayer L., Z. Chen, L. Self, S. Sampson, R. Findlay, J. Fang, C. Plante, P. Jumars & L. Watling (1995) Digestive physiology and contaminant bioavailability. Benthic Ecol. Mtg.
- Chen Z. & L. Mayer (1996) Assessing heavy metal bioavailability in sediments. Asia-Pacific Conf. Science & Mgmt. Coastal Environ.
- Mayer L. (1995) The role of the biota in controlling bioavailability of contaminants. DoD Tri-Service Workshop on: Bioavailability of Organic Contaminants in Soils and Sediments.
- Mayer, L., Z. Chen, R. Findlay, J. Fang, S. Sampson, L. Self, P. Jumars, C. Quétel, and O. Donard (1996) Bioavailability of harbor sedimentary contaminants subject to deposit feeder digestion. AGU-ASLO Ocean Sci. Mtg..
- Weston D.P. and L.M. Mayer (1996) Digestive solubilization of sediment-associated pollutants: In vitro extraction versus in vivo bioavailability. AGU-ASLO Ocean Sci. Mtg..
- Chen Z. and L. Mayer (1996) Probing the mechanisms in sedimentary copper bioavailability. Soc. Environ. Toxicol. Chem. Ann. Mtg.
- Weston D. and L. Mayer (1996) Digestive solubilization of sediment-associated pollutants: *in vitro* extraction vs. *in vivo* bioavailability. Soc. Env. Toxic. Chem. Ann. Mtg..
- Chen Z. and L. Mayer (1997) Solubilization of metals from sediments by deposit feeders: The role of the physiological ecology of protein. Amer. Soc. Limnol. Oceanog. Oc. Sci. Mtg.
- Chen Z. and L. Mayer (1997) Inhibition of digestive proteases by copper. Benthic Ecology Mtg.
- Mayer L.M. (1997) Digestive processes and bioavailability. Navy Copper Workshop, San Diego.
- Weston D. and L. Mayer (1997) In vitro digestive fluid extraction as a measure of contaminant bioavailability from sediments. Soc. Environ. Toxicol. Chem. Ann. Mtg.
- Bock M. and L. Mayer (1998) Digestive plasticity of the marine benthic omnivore *Nereis virens*, Benthic Ecology Mtg.
- Voparil I. and L. Mayer (1998) Mechanism of PAH solubilization during digestive attack. Benthic Ecology Mtg.
- Chen Z. and L. Mayer (1998) Bioavailability of sedimentary metals to deposit-feeders as a function of kinetics of solubilization within animal guts. Soc. Environ. Toxicol. and Chem.-Europe Ann. Mtg.
- Weston D., L. Mayer, L. Gravitz, and M. Bock (1998) A cross-phyletic comparison of contaminant bioavailability. Soc. Environ. Toxicol. Chem. Ann. Mtg.
- Chen Z., L. Mayer, M. Bock, D. Weston, R. Self, P. Jumars (1998) Cu inhibition of digestive enzymes of deposit feeders: Implications for changing benthic community structure. Soc. Environ. Toxicol. Chem. Ann. Mtg.
- Voparil I. and L. Mayer (1998) Surfactant control of PAH solubility in *Arenicola marina* gut fluids. Soc. Environ. Toxicol. and Chem. Ann. Mtg.
- Bock M., L. Mayer and D. Weston (1999) Dietary effects on the digestive chemistry of benthic polychaetes: Enzymes and surfactants. Amer. Soc. Limnol. Oceanog. Oc. Sci. Mtg.
- Chen Z, L. Mayer, S Sampson, C Quétel, O. Donard, R. Self and P. Jumars (1999) Enhanced solubilization of trace metals in deposit feeder Guts. Amer. Soc. Limnol. Oceanog. Oc. Sci. Mtg.