

**STRATEGY
RESEARCH
PROJECT**

The views expressed in this paper are those of the author and do not necessarily reflect the views of the Department of Defense or any of its agencies. This document may not be released for open publication until it has been cleared by the appropriate military service or government agency.

SIMULTANEOUS STRENGTH AND ENDURANCE TRAINING

BY

**LIEUTENANT COLONEL (P) ROY C. WAGGONER
United States Army**

**DISTRIBUTION STATEMENT A:
Approved for public release.
Distribution is unlimited.**

USAWC CLASS OF 1999

19990607 048



U.S. ARMY WAR COLLEGE, CARLISLE BARRACKS, PA 17013-5050

DTIC QUALITY INSPECTED 1

USAWC STRATEGY RESEARCH PROJECT

Simultaneous Strength and Endurance Training

by

LTC (P) Roy C. Waggoner

COL Jack Ellertson
Project Advisor

The views expressed in this paper are those of the author and do not necessarily reflect the views of the Department of Defense or any of its agencies. This document may not be released for open publication until it has been cleared by the appropriate military service or government agency.

U.S. Army War College
CARLISLE BARRACKS, PENNSYLVANIA 17013

DISTRIBUTION STATEMENT A:
Approved for public release.
Distribution is unlimited.

ABSTRACT

AUTHOR: LTC (P) Roy C. Waggoner

TITLE: Simultaneous Strength and Endurance Training

FORMAT: Strategy Research Project

DATE: 01 April 1998 PAGES: 135 CLASSIFICATION: Unclassified

The purpose of this study is to identify the exact workouts required to simultaneously achieve optimum improvement in 1 Repetition Maximum (RM) strength, muscle endurance and cardiovascular endurance. The first section identifies the physiological adaptations to each type of training, and using these, the second and third sections identify the optimum application of the acute variables to a periodized training program. The fourth section reviews studies done to date on simultaneous strength and endurance training and identifies how to apply that research to an optimum solution. The fifth section or conclusion presents a combined training program periodizing both strength and cardiovascular workouts using 4-week mesocycles inside a 12-16 week macrocycle. The sixth section identifies the application of this solution on a limited amount of training time.

TABLE OF CONTENTS

ABSTRACT	III
LIST OF TABLES	IX
SECTION I: EXERCISE PHYSIOLOGY	1
1. Fuel Sources	1
a. ATP Fuels all Muscle Contractions	1
b. The Body Stores Fuel for ATP Production in Several Ways	2
c. Energy Systems	4
d. Phosphocreatine	4
e. Aerobic Glycolysis	4
f. Aerobic Oxidation	6
g. Anaerobic Metabolism	7
h. Lactic Acid	8
2. Types of Muscle Fiber	9
a. Type 1 Fibers	9
b. Type 2 Fibers	10
c. Fiber Transformation	11
3. Central Nervous System	11
a. Motor Units	11
b. Motor Unit Recruitment	12
SECTION II: AEROBIC (ENDURANCE) TRAINING	14
1. Energy Costs Change Based on the Type of Event	14
2. Measuring activation of Energy Systems	15
a. VO ₂ max	16
b. Lactate Threshold	16
3. Acute Variables	18
a. Frequency	18
b. Duration	19
d. Aerobic Conditioning	23
e. Anaerobic Conditioning	25
f. Aerobic Capacity	28
g. Anaerobic Capacity	29
h. Strength Runs	30

4. Over-training	32
a. Frequency	32
b. Volume Vs Intensity	33
c. 3-Week Barrier	33
d. Summary.....	34
5. Cross Training	34
6. Periodization	35
7. Summary	36
SECTION III: STRENGTH TRAINING.....	38
1. Acute Variables.....	38
a. Choice of Exercise	38
b. Order of Exercise.	41
c. Number of Sets.	42
d. Rest between Exercises, Sets	45
e. Resistance Used (Number of Repetitions)	48
f. Frequency	50
2. Central Nervous System	53
3. Periodization	55
a. Weight Training Systems	55
b. Periodization Works.	56
c. Periodization Cycles.....	58
4. Over-Training	59
a. What is Over-Training	59
b. Application of Acute Variables in Over-Training.....	60
5. Summary	62
SECTION IV: COMBINED STRENGTH AND ENDURANCE TRAINING	65
1. Initial Studies	65
2. Power, not Strength May be Affected	66
3. A More Realistic Protocol	67
4. The Cellular Level.....	69
SECTION V: CONCLUSION	72

1. Summary	72
2. Recommendation	73
SECTION VI: DETRAINING: TRAINING ON LIMITED TIME	76
1. Cardiovascular Training.....	76
a. Intensity.....	76
b. Duration	78
c. Frequency	78
2. Strength Training.....	79
a. Acute Variables.....	79
b. Frequency	79
3. Recommendation	80
ENDNOTES.....	83
BIBLIOGRAPHY	103
1. Periodicals: Strength Training Studies.....	103
2. Periodicals: Cardiovascular Training Studies	110
3. Periodicals: Combined Strength and Cardiovascular Training	117
4. Periodicals: Over-training Studies	121
5. Books Lectures, Interviews and Studies	122

LIST OF TABLES

Table II-1: Frequency of Cardiovascular Training.....	19
Table II-2: Weekly Mileage	20
Table II-3: Specific Workouts for Specific Energy Systems	22
Table II-4: Duration of Aerobic Conditioning Workouts.....	24
Table II-5: Frequency of Aerobic Conditioning Workouts	25
Table II-6: Duration of Anaerobic Conditioning Workouts.....	27
Table II-7: Frequency of Anaerobic Conditioning Workouts.....	28
Table II-8: Duration of Aerobic Capacity Workouts.....	29
Table II-9: Frequency of Aerobic Capacity Workouts	29
Table II-10: Frequency of Anaerobic Capacity Workouts	30
Table II-11: Frequency of Strength Runs.....	31
Table III: 12: Neural Activation for Resistance Exercises	40
Table III-13: Number of Sets.....	44
Table III-14: Rest Intervals between Sets	47
Table III-15: Number of Repetitions.....	49
Table III-16: Frequency	52
Table III-17: Frequency: Hypertrophy Vs Strength/Power	53
Table III-18: Periodization: Stowers et al	56
Table III-19: Periodization: Willoughby	58
Table V-20: Periodization	73
Table V-21: Schedule	73
Table V-22: Acute Variables: Strength	74

Table V-23: Frequency: Strength.....	74
Table V-24: Acute Variables: Cardiovascular	75
Table V-25: Periodization	80
Table V-26: Schedule	80
Table V-27: Acute Variables: Strength	81
Table V-28: Frequency: Strength.....	81
Table V-29: Acute Variables: Cardiovascular	82

SECTION I: EXERCISE PHYSIOLOGY

"If an athlete could run aerobically without sacrificing muscle fibers for blood vessels and if slow twitch fibers could be persuaded to contract as fast and with as much power output as fast twitch-oxidative fibers, the 10K race would be won in about 17 minutes."¹

1. FUEL SOURCES

a. ATP Fuels all Muscle Contractions

The production of energy in muscles involves the break down of fuel (carbohydrates and fats) in the presence of oxygen by the mitochondria in the muscles to produce adenosine tri-phosphate (ATP). For the energy in foods to be used, it must first be converted into ATP. This conversion, the oxidation or glycolysis of fuels such as glucose and fat is known as cellular respiration and occurs in every living thing from bacteria to humans.²

ATP is the universal energy intermediate in cellular biology. When the ATP content of a cell falls below about 50-60% the cell will probably die, thus the production of ATP must keep pace with its usage.

Each time a sprinter takes a stride something like 1×10^{18} molecules of ATP are converted into ADP.³ This means that a marathon runner must have something like 75 kilograms of ATP to get through a race. Obviously, this would be hard since most elite marathon runners barely weigh 75 kilograms.

Seventy-five kilograms of ATP are obviously not stored in the body. In fact, the human body has as little as 100g at any given time.⁴ As soon as an ATP molecule is used for energy, the body makes more. Luckily, the body has several different energy-producing reactions, all of which generate the same intermediate, ATP. These energy

pathways provide critical insight into the types and amounts of training that an athlete can and should conduct.

ATP undergoes a chemical reaction with water to split into one molecule of adenosine di-phosphate (ADP) and one of phosphate. The catalysts that enable this chemical reaction are the oxidative enzymes in the muscle cell. The energy produced is enough to bend a cross bridge linking the thick and thin filaments within the myofibril. If enough cross bridges bend, the filaments slide over each other and if enough filaments slide, the muscle contracts.

b. The Body Stores Fuel for ATP Production in Several Ways

In their book, "Keep on Running; The Science of Training and Performance", Eric Newsholme et al. describe the perfect fuel storage tank for the human body. ⁵ Ideally, it would have the following characteristics:

- * A high chemical energy content, so that energy can be stored without being too heavy.
- * Chemical Stability so it does not break down before needed.
- * Capacity for rapid mobilization
- * Ability to generate ATP without production of toxic end-products
- * High molecular weight or insolubility to avoid osmotic complications (i.e. the movement of water in or out of a cell in response to a change in the concentration of dissolved materials in the cell).

Unfortunately, no single source of energy for ATP scores highly in all of the criteria listed above. The body's solution therefore, is store more than one kind of fuel and switch from one to another as circumstance change.

There are three important fuel storage sources in the body: glycogen (carbohydrate), triglyceride (fat) and phosphocreatine (phosphagen).⁶ (To a lesser degree, protein or more specifically amino acids can also be used as a fuel source since the liver can manufacture glucose using amino acids).⁷ Each has a role to play in the athlete's energy usage.

Fuel is stored in the muscles and liver as glycogen (carbohydrate) and fat cells or in the blood stream as triglycerides. The body can store a very large amount of fat, but the supply of glycogen is limited. In addition, a substance called phosphocreatine is stored in the muscles for immediate use before aerobic metabolism has had a chance to begin. Phosphocreatine is already present in muscles and requires only one reaction to form ATP, but is depleted very quickly.

As early as 1934, scientists knew that carbohydrates are the body's preferred fuels for distance running.⁸ The body uses a mixture of carbohydrates and stored fat as fuel sources to produce ATP, but the oxidation of fat requires oxygen so when you increase the intensity of exercise, significantly more demands are made on muscle glycogen and blood glucose than on stored body fat. During exercise, the body will switch to the fuel source that it can best use given the supply of oxygen. That is why when exercise intensity is low, the body burns fat and when you turn up the intensity, the body switches to carbohydrates.

As you begin exercising, muscle glycogen is the primary source of energy. In his book, "Inside Running", David Costill demonstrates the change in muscle glycogen content of a runners' gastrocnemius muscle during three hours of treadmill running, at the runner's best marathon pace. Although the test was run at a constant pace, the rate

of muscle glycogen use in the calf muscle was greatest in the first 90 minutes.

Thereafter, the amount of glycogen in the muscle approached zero. As the muscle glycogen approached zero, the runner became severely fatigued.

c. Energy Systems

The body can use glycogen, fatty acids, phosphocreatine or even protein in certain circumstances for ATP production. The significance of this is that there are five major options for fueling the production of ATP in the muscle:

1. Phosphocreatine(CP): stored in the muscle
2. Aerobic Glycolysis: using glycogen stored in the muscles or glycogen created from blood glucose
3. Aerobic Oxidation: using glycogen created from fatty acids
4. Anaerobic Glycolysis: using glycogen without the presence of oxygen.
5. Lactic Acid

d. Phosphocreatine

Glycogen stores in the muscles take about 25 seconds to become available as ATP. Something else has to supply fuel before glycogen can kick in. phosphocreatine is used for this maximum power output, when glycolysis cannot occur rapidly enough to supply sufficient ATP. At the beginning of any exertion, regardless of intensity the first source of fuel is phosphocreatine, until aerobic metabolism can begin production.

To generate ATP from phosphocreatine and ADP requires only one reaction and since phosphocreatine is stored in the muscles and ready for instant use it is by far the fastest source of ATP.⁹ The CP system can supply energy for 6-8 seconds; thus, the supply of phosphocreatine in the muscles is quickly exhausted. The body quickly recovers its stores of phosphocreatine after usage. After 22 seconds, the body has replaced about half of that used and after 44 seconds, about 75% have been replaced.

e. Aerobic Glycolysis

Aerobic metabolism as noted above can use fatty acid or glycogen for fuel. Since the stores of glycogen are limited and the stores of fats are essentially unlimited, the two systems work simultaneously, but the proportional share of fuels varies based on the intensity of the exercise and the level of conditioning of the runner.

Glycogen is a carbohydrate made up of thousands of glucose units polymerized into a gigantic tree-like molecule. The significance of the structure is that large amounts of glucose can be rapidly released from the branches of the tree. The muscles and liver use this glucose to build glycogen.¹⁰ Glycogen is stored in the muscles. To produce ATP, glycogen is broken down into glucose by the enzyme phosphorylase and enters the ATP cycle within about 25 seconds of exertion.

Glycogen, however is not an ideal storage fuel. The actual amount of glycogen in the body is limited. A normal human body can hold up to 500g of glycogen in the muscles. (The remaining 100g or so of glycogen in the body is stored in the liver and used to supply tissues that cannot use other fuels like the brain, thus cannot be used safely).¹¹ The attached water is nearly 1500g, thus adding 3 lb. of body weight. The biological significance of this is that if a human stored all of his fuel reserve as glycogen, vice fat, he would weigh twice as much!¹²

In addition, the energy content of glycogen is 16 kJ/g or less than half that of fat. A more significant comparison is that aerobic glycolysis can produce 36 molecules of ATP for each molecule of glucose used.¹³ This is 12 times less than that of fat. Secondly, pure glycogen is not found in the cells. Glycogen granules in muscle cells contain water and proteins that must be broken down with chemical reactions.

During simulated marathon running, elite marathoners consume about 5g per minute. With the 500grams stored in the muscles, a runner can carry enough to last about 100 minutes. Since the world record for the marathon is longer than 100 minutes, another fuel source must assist.

f. Aerobic Oxidation

Aerobic oxidation, the use of triglycerides as a fuel can provide 441 molecules of ATP for every molecule of glucose.¹⁴ Triglyceride molecules consist of a glycerol unit and three fatty acid units. These fatty acids provide the chemical energy for muscles. Since most fat is stored in, adipose cells vice in the muscle it must be transported through the bloodstream to be used by working muscles.

Triglyceride molecules react with water through the enzyme lipase to produce glycerol and three fatty acid molecules. Although the fatty acid molecules can pass into the blood, they are insoluble in water or blood plasma. To pass through the blood these fatty acids attach to the protein albumin and are oxidized in the muscles to begin the ATP cycle.¹⁵

The triglycerides stored in the body have energy content of 35kJ/g, more than twice that of glycogen.¹⁶ Triglycerides are stored dry and thus are by far the most efficient fuel in the body. By way of comparison, aerobic oxidation, the use of fatty acid as a fuel can produce 441 molecules of ATP for every molecule of glucose, 220 time that of anaerobic glycolysis and 12 times that of aerobic glycolysis.¹⁷

A 70kg male will normally store 8kg of fat and an average woman of 60kg, nearly twice as much. This means that the average man could run non-stop for 3 days and nights at an elite marathoner's pace without the need for food. Obviously, this is not

possible. The problem is that fat cannot be used by the muscle fast enough to sustain the pace. The reason is that fat cells are not soluble in water so to be carried in the blood they must be combined with protein thus requiring additional chemical reactions to become ATP¹⁸

g. Anaerobic Metabolism

There is enough ATP in a human leg muscle to support about 1 second of sprinting. Without oxygen almost none of the energy released in the breakdown of carbohydrates or fat could be used to create ATP. ATP can, however, be made without oxygen, or anaerobically.¹⁹

Anaerobically creating ATP is not easy. First, only glycogen can be used in the process. Fat cannot be broken down unless oxygen is present. Obviously, there is a lot less available glycogen in the human body than fat.

The second problem is that anaerobically, the body can produce only a small fraction of the ATP that it can aerobically. A glucose unit of glycogen can provide only 2-3 molecules of ATP anaerobically, compared to 36-37 molecules from aerobic glycolysis. In their book on the science of training, Newsholme, et al. Sited a study which compared the percentage of stored glycogen used through aerobic processes vice that of the anaerobic process for various track events. In a 5K race, although the anaerobic pathway contributed only 13% of the ATP produced, it used 46% of the glycogen depleted in the race.²⁰

The third problem with anaerobic metabolism is the accumulation of waste products. There are several by-products of anaerobic metabolism, but the most significant one is lactic acid. The accumulation of lactic acid in a working muscle

decreases its pH to a point where ultimately all of its enzymes would cease functioning and the cells would die. Fatigue is the safety valve that prevents lactic acid from building to this point. We simply slow down as our bodies catch up with lactic acid removal.²¹

h. Lactic Acid

As a runner approaches the top end of aerobic capacity, anaerobic metabolism becomes increasingly important. In the initial minutes of aerobic exercise, oxygen consumption rises rapidly. If the energy demands can be met aerobically, oxygen consumption climbs to a plateau and stays at this plateau, often called steady state.

Steady state reflects equilibrium between energy demands and the rate of aerobic ATP production.²² Little lactic acid accumulates at steady state. When oxygen consumption reaches a point where it does not rise, despite ever-increasing workload this is known as your VO_2 max. Since the workload is certainly increasing, the additional energy demands must be met somehow. Above your VO_2^{max} , energy demands must be met through anaerobic metabolism.

The reason is called the lactate threshold. As you increase the pace, anaerobic metabolism takes on an increasingly higher percentage of energy production until, at some point the accumulation of lactate in the muscles causes you to stop or slow. The accumulation of 4 mmol / liter of lactic acid in the blood has been termed the onset of blood lactic acid (OBLA). OBLA is one of the key factors that limit running pace.

The body uses part of lactic acid for fuel. It separates lactic acid into hydrogen ions and lactate. It is actually the hydrogen ions that increase acidity in the muscles and cause the "lactic acid burn". Lactate is used for fuel. In fact muscles, the liver, the

kidneys, the heart and even the brain can metabolize lactate. Sixty percent of accumulated lactate is metabolized for fuel. Most of the remaining forty percent is converted to glucose and protein. Only a small portion is excreted in the urine.²³

The half-life of the lactaid portion of the by-products of anaerobic metabolism is about 25 minutes. Ninety five percent is removed in 1 hour and 15 minutes.

2. TYPES OF MUSCLE FIBER

Inside a human muscle (excluding the heart) there are two types of muscle fibers. The thicker the muscle, the more myofibrils it contains and the greater is its capability to generate force. The thinner the muscle, the more oxidative enzymes, mitochondria and capillary density it contains.

a. Type 1 Fibers.

Type 1 fibers are the thin type. They are often labeled red or slow twitch fibers when looked at with a microscope. Type I fibers look red due to the higher capillary density and oxidative capability. They are relatively slow to contract with relatively little force but they can continue to contract for a relatively long time. Given these characteristics, they are obviously designed for efficient use of oxygen to produce ATP. Elite distance runners have been found to have a predominance of type 1 fibers.²⁴ Type 1 fibers contain the highest density of mitochondria (those cells within muscles that produce energy aerobically from glycogen). They have a much greater capillary density, higher levels of oxidative enzymes and large intra-muscular triglyceride stores.²⁵ All of these factors mean a greater oxygen flow to the muscle and better waste removal capability. Type 1 fibers have been shown to experience significant improvement in aerobic function with aerobic training.²⁶

A type 1C fiber has recently been identified. There are very few of these fibers and they are less oxidative than normal type 1 fibers. There is some evidence that with resistance training, type 1 fibers may become 1C fibers, taking on some of the characteristics of type 2 fibers.²⁷

b. Type 2 Fibers.

Type 2 fibers are often called fast-twitch or white fibers. They are relatively fast to contract with relatively high force but they only continue to contract for a relatively short time. Given these characteristics, they are obviously designed for generation of force. These fibers have high levels of the enzymes that break down ATP and release energy for the shortening of muscle fibers to produce force. Type 2 fibers have a low ability for aerobic work, as evidenced by their low capillary density, low intra-muscular stores of triglyceride, low mitochondria density and low levels of oxidative enzymes. Since these fibers rely predominantly on anaerobic metabolism and have a poor ability to supply ATP aerobically, they are very susceptible to fatigue.²⁸

There is recent evidence that type 2B fibers may be just a pool of unused fibers with low oxidative capability that begin a transformation to type 2A fiber type on recruitment. This is supported by the fact that they are the last to be recruited and observed dramatic reductions of type 2B fibers with heavy resistance training.²⁹ There are also reports of a type 2C fiber. There are relatively few of these in humans (less than 5%). They are more oxidative than type 2A or 2B fibers. With aerobic training, these may take on the characteristics of type 1 fibers. Type 2AB fibers, also recently identified, represent a combination of type 2A and 2B fibers and are transitional fibers.

Thus, a fiber might be transformed from 2B to 2AB to 2A because of heavy resistance exercise.³⁰

Fast twitch or type 2 motor units have more muscle fibers for each motor neuron; thus, they can simultaneously fire more muscle fibers than slow twitch motor units.

c. Fiber Transformation

Although a number of studies indicate that there may be fiber transformation between the type 1 and type 2 fibers, more recent evidence indicates that the changes occur only within the subtypes (e.g. 2B-2AB-2C). The significance of this is that it appears that fiber transformations may occur along a continuum of fiber sub-types based on the intensity, duration and type of exercise.³¹ This concept is critical to the design of simultaneous aerobics and strength training programs. All muscle fibers respond to endurance training by increasing their capacity for energy production. This means that type 2A and even type 2B fibers can take on greater oxidative capability (or become 2C fibers). All muscle fibers also respond to resistance training (lifting weights or running hills for instance) by growing larger and increasing their capacity to produce force.

3. CENTRAL NERVOUS SYSTEM

a. Motor Units.

A single nerve fiber controls as few as 5 or as many as 1000 muscle fibers. Each nerve fiber and its attendant muscle fibers are known as a motor unit. The force production of the muscle increases as more motor units fire simultaneously. Resistance training has been proven to increase the synchronous firing of motor units thus increasing the strength of the muscle.

Initial increases in strength are due to increases in the synchronous firing of motor units, or the efficiency of the neuro-muscular system. The neuro-muscular system is “learning a pattern to get stronger. Simultaneously, the antagonist muscle is learning to decrease its neural activity with the training of that particular movement.”³²

b. Motor Unit Recruitment.

When a muscle begins to exercise, not all of the motor units are recruited at once. During slow, low intensity, aerobic exercise type 1 fibers are generating most of the force. As the intensity increases, type 2A fibers are added to the workload. At sprinting speeds, where maximal force production is needed the type 2B fibers are recruited. In running, once the type 2A fibers kick in, you are operating at or slightly below your lactate threshold.³³

During long distance events like a marathon, as the type 1 fibers become fatigued, the type 2A and eventually the type 2B fibers are recruited, although the pace remains constant.³⁴ (This may explain the “wall” in a marathon). Since most of the type 1 and type 2A fibers must already be working for the body to recruit type 2B fibers, these fibers are difficult to train.

During resistance training, the type 2A fibers begin to kick in at somewhere around 70% of one repetition maximum in weight lifting.³⁵ Heavier resistance (e.g., 3-5 RM) requires the recruitment of higher threshold motor units composed of primarily type 2 fibers however, just as in aerobic exercise, the smaller type 1 fibers are recruited first. From a training standpoint, this size order of recruitment is important. In order to recruit type 2 fibers (or more specifically motor units composed mostly of type 2 fibers), the exercise must, be intense.

Recruitment order within a muscle is fixed for a specific movement³⁶ and changes if the body position is changed.³⁷ Recruitment order in the quadriceps for instance would be different for the performance of the leg extension and the squat. In their book, "Designing Resistance Training", Fleck and Kraemer hypothesis that variation in recruitment pattern may be one of the factors responsible for the specificity of strength gains. This is evidence that supports anecdotal beliefs long held by strength coaches that to completely develop a particular muscle it must be exercised using several different exercises.³⁸

Resistance training builds up the myofibrils in all muscle types, but appears to affect type 2 fibers more than type 1. As the size of the muscle fiber increases, simultaneously the mitochondrial volume and the oxidative enzymes in the muscle are decreasing.³⁹ At the cellular level, aerobic and strength training cause opposing actions. This balancing act must be overcome if an athlete is to simultaneously train for endurance and strength.

Training cannot change the relative balance of fast twitch and slow twitch fibers (discounting the unproved theory of hyperplasia). This is determined by genetics. Sprinters and power lifters have a higher percentage of type 2 fibers and marathoners have a higher proportion of type 1 fibers. You can, however cause one fiber type to take on the characteristics of the other, based on the training that you conduct.⁴⁰

SECTION II: AEROBIC (ENDURANCE) TRAINING

The goal of this study is to identify the best combination of training to get simultaneous improvement in 1RM strength and middle distance running (3-5 kilometers) or more specifically the 2-mile run. Because the physiological alterations are specific to the type of training, this range is the toughest of the distance continuum. The energy costs of middle distance running include some aerobic and some anaerobic metabolism. Because of this balance, the limiting factors on this training are probably the conversion of fibers and the limits of over-training.

The purpose of this chapter is to first examine the specific energy costs and how specific workouts affect each of these energy systems. The results of this review will indicate how to apply the acute and chronic variables in the design of a cardiovascular training program.

1. ENERGY COSTS CHANGE BASED ON THE TYPE OF EVENT.

The proportion of ATP derived from aerobic metabolism has been extensively catalogued for different distances. The aerobic contribution ranges from less than 5% for a 100-meter sprint to almost 100% for a marathon. For the 5K race, it is about 87% and the 1500-meter it is about 65%.⁴¹ From those figures, we can deduce that during a 2 mile run, the body gets about 76% of its ATP from aerobic metabolism. Since these distances are run at or just below the anaerobic threshold glycogen is the most significant fuel in races from 3k to 5K. However the 25% contribution from anaerobic metabolism is still critical to performance and becomes more critical as a runner approaches the lactate threshold.

In middle distance running, you have a combination of high intensity and relatively long duration. Neither the aerobic or anaerobic systems can independently provide the energy required.⁴² Thus, runs in this range are dependent on the simultaneous contribution of both aerobic and anaerobic metabolism. In fact, the systems are not mutually exclusive, as both can contribute simultaneously. The percentage of contribution changes not only for the distance run, but also by the physiological ability of the runners involved.⁴³

2. MEASURING ACTIVATION OF ENERGY SYSTEMS.

How do you predict when an energy system kicks in? Each type of training (or training zone) has a specific purpose for optimizing one or more of the physiological factors that impact running performance (fiber type and the energy pathways). To stress a certain energy pathways or a certain type of muscle fibers you must train at a specific intensity for a specific amount of time with a specific amount of recovery. This has been shown to help you train optimally and gradually improve race performance.⁴⁴

If you were to build to a steady pace and maintain that pace, the energy pathways would begin to contribute in the following sequence: Stored ATP and phosphocreatine are the primary energy sources for the first 25 seconds of a run until the muscle can begin aerobic glycolysis. At about 30 seconds until about 3 minutes, aerobic glycolysis is the primary energy system, using stored muscle glycogen until the body can deliver glycogen and stored triglycerides through the blood to the muscle. After about 3 minutes, the blood is delivering glycogen from the blood. At about 10

minutes, aerobic oxidation begins to contribute; starting to burn stored fat as a fuel that begins to be the primary fuel source after about 15-20 minutes.

There are several physiological variables an athlete can monitor to identify when the body has shifted to the various energy systems. The two most common are VO₂max and Lactate Threshold.

a. VO₂max

The physiological variable that best describes the capability of the cardiorespiratory system to transport oxygen to the muscles for generation of ATP is VO₂max. Because 76% of the 2-mile run is fueled by aerobic metabolism, VO₂max is important to performance. Values for VO₂max average from 40 to 50 milliliters / kilogram / minute (ml/kg/min) to 94 ml/kg/min for elite cross-country skiers. Average values in elite middle distance runners range from 68 to about 77 ml/kg/min. Average values in elite long distance runners average from 75-85 ml/kg/min.⁴⁵

Middle distance runners have generally lower VO₂max than long distance runners but compete at a higher percentage of that VO₂max and incur a greater energy cost for distance run. The correlation between middle distance performance and VO₂max is generally lower than the correlation between long distance running and VO₂max. Brandon and Boileau found that the correlation of VO₂max and performance was a more important predictor of 10K races than shorter distances.⁴⁶

b. Lactate Threshold

The running pace that most people can sustain is not nearly at their VO₂^{max}. As you increase the pace, anaerobic metabolism takes on an increasingly higher percentage of energy production until, at some point the accumulation of lactate in the

muscles causes you to stop or slow. The accumulation of 4 mmol / liter of lactic acid in the blood has been termed the onset of blood lactic acid (OBLA). OBLA is one of the key factors that limit running pace. The body can no longer buffer the acid in the blood and muscles as fast as it is being produced.

The pace at which excess lactate begins to accumulate has been defined as the anaerobic threshold (AT).⁴⁷ The heart rate at which this generally is about 85% of maximum heart rate. The anaerobic threshold is, however different for every runner. To compound the problem, there is not a direct or linear correlation between percentage of maximum heart rate and percentage of VO₂max. Science dictates that we train at a given percentage of VO₂max, but it would be hard to measure VO₂max outside of a laboratory.

There is, however, a direct and linear correlation between the percentage of reserve heart rate and the percentage of reserve VO₂max. The reserve heart rate and reserve VO₂max are the differences between the maximum value for an individual and the value for that individual at complete rest. This is the reason that the Karnoven method of determining training heart rate zones is significantly more effective.⁴⁸ You simply subtract your resting heart rate from your maximum heart rate, calculate the percentage of the difference at which you wish to train (e. g. 60% for longer runs) and add the resting heart rate back to that percentage.

The anaerobic threshold (AT) is considered a good estimation of the pace that a runner can maintain during a longer race. Maffulli, et al. evaluated 112 endurance athletes for run speed over 800m, 1500m, 3000m 5000m and 10,000m. They concluded that there was a high correlation between the run performance and the AT

for races of 5000m and above. There was no significant correlation between AT and race performance at 800m or less.⁴⁹

3. ACUTE VARIABLES

Acute variables define all possible conditions within a single training session. In cardiovascular training, the acute variables are frequency, intensity, duration, and rest periods between intervals and periods of exercise.

By varying the acute variables, each workout targets a specific energy system. To get the optimum results from a workout, each workout must have a specific purpose targeting a specific aspect of cardiovascular performance. “Junk miles” or running the same pace and same distance each workout would be grossly ineffective for anything except general fitness and health.

Planned manipulation of these acute variables causes significant changes in physiological adaptations to the workout. Although the percentage is specific to the event, all middle distance events derive a small portion of their energy needs from phosphocreatine, a moderate portion from anaerobic glycolysis and the largest portion from aerobic glycolysis. The lactate removal system is also significant to middle distance races. Planned variation of the acute variables based on the specific energy costs of a given event allows coaches to design cardiovascular training programs.

a. Frequency

How often should you train? In 1998, the American College of Sports Medicine (ACSM) recommended a frequency of 3-5 days per week in order to maintain cardiorespiratory health and fitness in adults.⁵⁰ These guidelines are not intended for athletic performance, but the maintenance of health. Interestingly, their 1998

recommendations on frequency stated that the ACSM “now views exercise/physical activity for health and fitness in the context of an exercise dose continuum... programs involving higher intensities and/or greater frequency/duration provide additional benefits”.⁵¹

The current guidelines are similar to those that running have been recommending for several decades. When one reviews various program recommendations for 5-K race performance the frequency of training is amazingly consistent (See Table II-1). The significance of all of these recommendations, including ACSM, is that cardiovascular improvement for competition is best attained by training 5-6 days each week.

Source	Robinson 52*	Burke 53*	Newsholme 54*	Lawrenc e 55*	Anderson 56*	McMillan 57*
Days/Week	5-6	6	6	6	5-6	5

Table II-1: Frequency of Cardiovascular Training

*Note: Recommends alternating hard and easy days.

b. Duration

Another aspect of frequency is the age-old question of how many miles a week should you run. There appears to be a point of diminishing returns where further increases in weekly mileage fail to increase performance. One study showed that training 40-80 kilometers (24-48 miles) a week elicited significant improvements in VO2max, but greater distances produced only slightly greater increases, but significantly greater incidences of over-use and injury.⁵⁸

Dudley et al Found that longer exercise bouts produce greater increases in mitochondrial density, however there is a point of diminishing returns (see figure 1). After 1 hour the rate of increase drops off , and after 2 hours it drops off significantly. By 3

hours the rate of increase is insignificant. It is important to note that after 1 week of detraining, almost 50% of the increase in mitochondrial content is lost. After 5 weeks all of the adaptations had been lost.⁵⁹

Elite distance athletes have a proven requirement for the marginal additional increases in performance provided by additional mileage. However, the muscle fiber adaptations created by these levels of mileage creates a specific effect and virtually guarantees a significant loss of upper body strength and eventual overuse injuries.

Below is a comparison of recommendations from coaches and running authorities on how much weekly mileage is optimal for novice competitors training for a 5-K race.

Coach / Category	Glover⁶⁰	Lawrence⁶¹	Anderson⁶²	Collins⁶³	Higdon^{64**}	Burfoot⁶⁵
Weekly Mileage	25-40	35-46	25-35	30	32	30

Table II-2: Weekly Mileage

The significance of these recommendations is that for 5-K performance below the elite level duration of 25-40 miles a week appears to be optimum.

c. Intensity

In a series of studies, Hickson et al had subjects execute identical training programs and then reduced either the duration or the intensity by one thirds or two thirds. They found that it is possible to maintain almost all of the performance (less long-term endurance) increases with up to a two-thirds decrease in duration (from 40minutes to 26 minutes.⁶⁶ Their conclusion was that intensity is an essential requirement for maintaining increased aerobic power.⁶⁷

Mitochondria are the cells that aerobically produce ATP in muscles. It is well known that long duration aerobic exercise induces significant increases in mitochondrial

content of muscle fibers. Figure 2 displays improvement in muscle fiber mitochondrial content during running programs of shorter duration. The peak adaptation appears to occur, in shorter runs as the intensity increases with very short intervals at 100% of VO₂max producing the greatest improvement.⁶⁸

Different running intensities have been shown to produce different hormonal responses. In 1989, Kraemer et al. studied the hormonal responses to training of 3 different running programs over 10 weeks.⁶⁹

One group (sprint interval- SI) performed sprint intervals on 3 days per week. They did 2 sets of intervals separated by 5 minutes of rest. Each set consisted of 4 (20-second) maximal sprints with one-minute rest intervals. Heart rates were monitored and after the second sprint, subjects were determined to be training at 96% of their directly determined maximal heart rate.

The second group (endurance-E) trained 3 days a week and ran as far as they could in 30 minutes. Heart rate was again monitored and these subjects were determined to be training at a mean exercise intensity of 80.0% of their treadmill VO₂max. This is at or slightly above the lactate threshold.

The combined group trained six days a week and performed the workouts of the endurance group and the sprint interval groups on alternating days. Heart rate determinations during exercise showed that they trained at 93.4% of their directly determined maximal heart rate during sprints and 79.7% of treadmill VO₂max during the endurance runs.

VO₂max improved 6.6% for the sprint group, 18.3% for the endurance group and 14.1% for the combined group. Interestingly, the combined group showed significant

pre to post training reduction in the magnitude of increase of beta-endorphins, lactic acid and cortisol at 5 minutes after workouts.⁷⁰ The authors noted that the combined group probably exhibited some over training towards the end of the study, but the significance is that the combined group showed a significant improvement in the ability to flush lactic acid as compared to the other groups.

In 1996, in a pair of studies, Tabata, et al explored the effects of moderate-intensity endurance and high intensity intermittent (interval) training on both anaerobic capacity and VO_{2max} . While the moderate intensity (70% VO_{2max}) exercise improved VO_{2max} , it had no effect on anaerobic capacity. The interval training, however, improved both VO_{2max} and anaerobic capacity, probably through imposing intensive stimuli on both systems.⁷¹

Each category of intensity corresponds to one of the energy systems and the corresponding aspect of cardiovascular performance. Those five aspects of cardiovascular performance although named differently by different sources are essentially endurance, stamina, speed, sprinting and strength.^{72,73,74}

Primary Physiological System	Type Workout	Aspect Improved
Aerobic Oxidation & Aerobic Glycolysis	Aerobic Conditioning	Endurance
Lactate Removal	Anaerobic Conditioning	Stamina
Anaerobic Glycolysis	Aerobic Capacity	Speed
phosphocreatine	Anaerobic Capacity	Sprinting
Lactate Removal and Muscle Fiber Composition	Strength	Strength

Table II-3: Specific Workouts for Specific Energy Systems

d. Aerobic Conditioning

Aerobic conditioning improves endurance or the body's ability to use oxygen.

VO₂max is the common measure of the body's ability to use oxygen and is increased by longer runs at relatively slow pace (60-75% of maximum heart rate). These workouts increase the muscles' ability to consume oxygen, increase the capillary density in the working muscles, the amount of mitochondria and the oxidative enzymes in the muscle fibers. They also teach the muscles to use glycogen and fatty acids more efficiently improving aerobic glycolysis and oxidation. In other words, they improve your VO₂max.

How much can training affect VO₂max. Most studies in the past have indicated that a minimum increase in VO₂max of 10-15% can be attained using the ACSM guidelines.⁷⁵ Improvement is, however directly related to frequency, duration and intensity.

R. C. Hickson et al showed an improvement of VO₂max of 44% (average) over ten weeks induced by training 6 days a week. For 3 days a week, the eight subjects performed 6 5-minute intervals on a bicycle ergometer that elicited VO₂max separated by 2-minute intervals of exercise requiring 50-60% of VO₂max. On the alternate days, they ran as far as they could in 40 minutes. Four of the Eight subjects achieved VO₂max approaching or exceeding 60 ml/kg/min.

K. A. Mikesell and G. A. Dudley repeated this study in 1984 for 5 weeks using seven well conditioned male runners to determine its effect on 10-K run times. The mean run time dropped by over 81 seconds.^{76 77} The significance of this study is that it showed VO₂max can increase more rapidly and to a greater extent than was previously thought.

Duration of exercise at low intensity has been shown to increase the body's ability to oxidize fat for fuel. Phillips, et al. did a study in 1996 to determine the effects of training duration on substrate turnover and oxidation during exercise. Active, but untrained males trained for 31 days on a bicycle ergometer for 2 hours a day at 60% of pre-training VO₂max. Exercise increased the rate of glucose production and utilization. Interestingly, at day 5, the authors noted a training-induced 10% increase in total fat oxidation due primarily to an increase in intramuscular triglyceride oxidation and a decreased glycogen oxidation. By day 31, the total fat oxidation increased a further 58%. The pattern of fat oxidation at day 31 showed a decreased reliance on free fatty acids in the blood and an increased reliance of intramuscular triglyceride.⁷⁸ This study supports previous evidence that long-term aerobic conditioning induces a progressive increase in fat utilization and reduction in glucose oxidation.

How far must an athlete run to improve VO₂max? Most sources indicate that if an athlete is training for longer races the longest run should be about two-thirds as long (time, not distance) as the intended race, but at a slower pace.⁷⁹ For shorter races (10K to 5K), most authors recommend no more than 10-12 miles for a longest run.⁸⁰ Table II-4 shows a review of these recommendations.

Coach	Glover 81	Lawrence 82	Anderson 83	Collins 84	Higdon 85**	Burfoot 86
Weekly Mileage	25-40	35-46	25-35	30	32	30
Long Run	8-10	7-10	8-10	8	10	10

Table II-4: Duration of Aerobic Conditioning Workouts

Despite the great improvements in VO₂max to be had by longer runs, they pose an imminent danger of over training. To avoid over training, the intensity or pace for

these runs is commonly cited as 55-65% of maximum heart rate or up to 3-minutes slower than marathon pace. In addition, unanimously, coaches and athletes agree that athletes should execute long runs no more frequently than once a week.

Source	Robinson 87*	Burke 88*	Newsholme 89**	Lawrence 90*	Anderson 91*	McMillan 92*
Aerobic Conditioning	2-7 d/w (1 long)	4 d/w (1 long)	3 d/w (1 long)	4 d/w (1 long)	4 d/w (1 long)	3 d/w (1 long)

Table II-5: Frequency of Aerobic Conditioning Workouts

*Note: Recommends alternating hard and easy days.

e. Anaerobic Conditioning

Anaerobic conditioning builds affects stamina. The purpose of anaerobic conditioning is to raise the pace that a runner can sustain at anaerobic threshold. Remember; steady state reflects equilibrium between energy demands and the rate of aerobic ATP production. Little lactic acid accumulates at steady state. Anaerobic threshold is slightly above steady state. Lactic acid begins to accumulate, but not so fast that the athlete cannot hold the intensity. Typically the lactate threshold occurs between 40-85% of VO₂max. That pace (anaerobic threshold) occurs at around 85% of maximum heart rate.

The lactate threshold may be the most significant factor in middle distance performance. Robinson and Carrino, in their book Max O₂, state: "In fact, the lactate threshold is so significant in determining performance, it's possible for someone with a lower VO₂max to have a greater aerobic performance capacity."⁹³

In a longitudinal study, Tanaka et al. found significant correlation at every assessment point over nearly 5 months between 10,000m run performance and percentage of VO₂max at the AT. They hypothesized that a 20-second improvement in

the 10,000-m race time correlates to a 1-ml/kg/min increase in VO_2 at the AT. The relationship did not change as run performance increased.⁹⁴ The significance of this is that raising the percentage of VO_2max a runner can hold at the anaerobic threshold has the most direct impact on middle distance performance.

In other words, anaerobic conditioning primarily serves to increase the percentage of VO_2max an athlete can sustain at or near the lactate threshold. Additionally, they improve the oxidative capability of type 1 fibers not recruited during aerobic conditioning runs and begin to increase the oxidative capacity of type 2A fibers. Anaerobic Conditioning workouts improve aerobic glycolysis, lactic acid buffering and removal and, to a lesser degree, anaerobic glycolysis. At these increased speeds aerobic oxidation is no longer significant (there is not enough oxygen present to use fat as a fuel source). Lactate threshold can be increased independently of VO_2max and can improve 10-20% in response to training.

A significant question to the application of lactate threshold runs is whether to run at or above the LT in training. A study by Joanne Henritze et al from the Human Performance Laboratory of the University of Colorado compared the effects of training at and above lactate threshold. Subjects were randomly assigned to 3 groups for a 12-week training program. One group trained above the lactate threshold, a second group trained at the lactate threshold and the third was a control group. Subjects were assessed for VO_2max , VO_2^{LT} , and $\text{VO}_2\text{max}/\text{VO}_2^{\text{LT}}$ before and after the study. Training loads were equated so that each subject expended approximately 1465 kJ per training session. This effectively removed training volume as a variable. The study concluded that training above LT produces significant increases in VO_2^{LT} (48%) as compared to

training at LT (18%). Interestingly, no significant change was found in VO₂max in any group. (5.6% for >LT and .9% for =LT).⁹⁵ The significance of this is that it supports other findings that indicate that VO₂max at LT can be increased independent of improvements in VO₂max. The clear indication is that to improve race pace, you must train above LT.

Anaerobic conditioning workouts typically take the form of cruise intervals, tempo runs or fartlek runs, all of which increase your body's lactate threshold running speed (anaerobic threshold).

Tempo runs are fast continuous runs at 80-85% of maximum heart rate, aiming to run at just below the anaerobic threshold. These runs are typically between 20 and 40 minutes for middle distance runners. Cruise intervals are 800-m to 2-mile repeats at 85% of maximum heart rate with 1-2 minutes in between each interval. Fartlek runs are periods of faster running introduced into slower runs. When used to affect anaerobic conditioning, the pickups are run at about 85% of maximum heart. These are normally used to prepare runners for more demanding aerobic capacity intervals.

Coaches are again consistent on their recommendations for the duration of anaerobic conditioning workouts:

Source	Robinson 96*	Burke 97*	Lawrence⁹⁸ *	Anderson 99*	McMillan 100*
Tempo Runs	20-40 min	20-40 min	20-40 min	20-40 min	20-40 min
Cruise Intervals	1200m- 1 mile	1200m- 1 miles	1200m- 1 mile	1200m- 1 mile	1200m- 1 mile

Table II-6: Duration of Anaerobic Conditioning Workouts

Anaerobic conditioning workouts are intense. They require anaerobic metabolism and produce the resulting by-products that the body must flush. As an athlete trains in the anaerobic range too often, overtraining will set in. As a result, most coaches recommend lactate threshold runs only once a week. See table II-7:

Source	Burke 101*	Newsholme 102**	Lawrence 103*	Anderson 104*	McMillan 105*
Anaerobic Conditioning	1 d/w	1 d/w	1 d/w	1 d/w	1 d/w

Table II-7: Frequency of Anaerobic Conditioning Workouts

*Note: Recommends alternating hard and easy days.

f. Aerobic Capacity

Aerobic capacity builds speed. They accomplish this by increasing the body's ability to consume and utilize oxygen at speeds close to the anaerobic threshold and increase the ability to buffer lactic acid and flush it from the system. These type workouts increase not only the lactate buffering of the muscles but of the kidneys, liver and blood. They teach a runner to relax and hold concentration while holding a challenging pace. Holding the pace steady involves not only removing lactic acid from the system, but also learning to tolerate higher concentrations in the blood. These workouts improve the high end of aerobic glycolysis, anaerobic glycolysis and lactic acid removal.

Aerobic capacity workouts are often called intervals. They include intervals lasting from two to six minutes at 90-95% of the maximum heart rate. Recovery jogs between intervals are equal to or longer than the interval. Intervals are one of the three key workouts that form the base of literally every running program currently in publication for

3000-m to 10,000-m. Below are comparisons of coaches' and running authorities' recommendations on percentage of aerobic capacity (greater than 90% of MHR) miles can be run each week.

Source	Glover 106	Lawrence 107	Anderson 108	Collins 109	Higdon 110	Burfoot 111
% of weekly miles	10-20% *	9%	10%	10%	9.3%	7.5%

Table II-8: Duration of Aerobic Capacity Workouts

* Experienced runners preparing for a 10K or less can do 25%.

These workouts are intense and executing them too frequently will cause over training. There is some anecdotal evidence that even once a week may be too often, but coaches remain consistent in their guidelines:

Source	Burke 112*	Newsholme 113**	Lawrence 114*	Anderson 115*	McMillan 116*
Aerobic Capacity Intervals	1 d/w	1 d/w	2 d/w	1 d/w	1 d/w

Table II-9: Frequency of Aerobic Capacity Workouts

*Note: Recommends alternating hard and easy days.

g. Anaerobic Capacity

Anaerobic capacity workouts are truly anaerobic work and build sprinting ability. They reach a level of intensity that recruits type 2A and finally type 2B fibers. These workouts will further increase your VO₂max and psychologically makes your race pace feel easier. Most importantly anaerobic capacity workouts train the body's ability to produce energy anaerobically by specifically targeting the phosphocreatine system. This energy system fuels true sprints and only lasts up to about 15 seconds.

Most coaches recommend sprints of 50-100m that start at 800m pace and build to top speed with at least two-minute recoveries to replenish the CP system. Because these are very demanding workouts, most authors and coaches recommend relatively few of them, and most only in a peaking phase:

Variable/ Source	Robinson 117*	Burke 118*	Newsholme 119*	Lawrence 120*	Anderson 121*	McMillan 122*
Anaerobic Capacity Intervals	1 d/w to 2 d/ m	0 d/w except peaking	0 d/w except peaking	2 d/m peaking	0 d/w except peaking	1 d/w (peaking) vice AC Intervals

Table II-10: Frequency of Anaerobic Capacity Workouts

h. Strength Runs

These are hill repeats. Anecdotal information would indicate that hill running helps increase the strength or force generation of muscle fibers. In a study at the Department of Kinesiology at University of Waterloo investigators studied the effects of an intense hill running protocol on middle aged (34-37) endurance-trained men. For six weeks, the training program was conducted four times a week. The program consisted of a 3-km warm up run at low intensity followed by 3 maximal runs on a long paved hill of 3.3% grade for 60 seconds each with two minutes of recovery jogging between repeats. Immediately afterwards, the subjects ran 5 sprints of 6 seconds on a grassy 44% grade with 24 seconds recovery and 2 more maximal runs of 90 seconds on the 3.3% grade with 3 minutes of recovery. After running, subjects performed leg presses at 15-RM.

The results were surprising. Significant improvements were noted in training distance for each interval and resistance used for the leg press. Anaerobic capacity, as judged by a treadmill run to exhaustion at 20% grade improved by an average of 16.7%. Terminal blood lactate level increased by 14%. VO₂max improved by an average of

only 4.1% (ml/kg/min). At the cellular level, there was a 3.3% increase in the proportion of fast twitch muscle fibers. This is not statistically significant, but till a relatively large change over 6-weeks! This study, however was conducted prior to the identification of fiber sub-types, so this was not evaluated, but probably occurred to a significant degree based on more recent research.

What is interesting is that although resting Creatine Phosphate concentration increased by 4.5% and resting ATP concentrations increased by a large amount (14.8%). Given more recent research on hyperplasia, we know that type 1C fibers are a type of low twitch fiber that takes on some of the characteristics of fast twitch fibers. These fibers have a larger amount of ATP concentration

Strength runs appear to cause physiological adaptations by increasing the stored concentrations of ATP indicating fiber sub-type transformation. They also help the muscles absorb the pounding of training and allow a runner to maintain the pace up hills. These are similar to the effects of weight training on muscle fiber type.

Table II-11 shows frequency recommendations of various 5-km running programs. As both types of training are anaerobic in nature, over-training is a primary concern. Most indicate that strength runs can be cut back if weight training is used simultaneously as they target the same physiological adaptations.

Source	Robinson 123*	Burke 124*	Newsholme 125*	Anderson 126*	McMillan 127*
Strength (Hills)	0-1 d/w	1 d/w	1 d/w (cross country run)	0-1 d/w	1 d/w (base only) vice tempo run

Table II-11: Frequency of Strength Runs

*Note: Recommends alternating hard and easy days.

4. OVER-TRAINING

a. Frequency

How much cardiovascular training causes over training. W. J. Kraemer et al. did a study training 6 days each week. The participants did 3 days of intervals and 3 days of fast continuous running (anaerobic threshold) on a treadmill. Based on the guidelines above, performing intervals three times a week and anaerobic threshold runs three times a week would seem to invite over training. The hormonal responses of this group in fact indicated that the training intensity and frequency might have elicited over-training.¹²⁸

Another study by Lehmann et al studied recreational athletes on a 6 day/week program for 6 weeks. They performed lactate threshold and interval training on cycle ergometers training for 40-60-minutes each day ceasing at muscular exhaustion. The seventh day was a regeneration day of 30-40 minutes of low intensity training. The duration was increased in weeks 3 and 4 eliminating the regeneration day. Weeks 5 and 6 were used for regeneration (2h/week). Power input increased through three weeks, but decreased at week 6 despite two regeneration weeks. Obviously training seven days a week caused over training that not even two weeks of reduced training could repair.

A series of studies by M. L. Pollock et al showed that the amount of improvement in VO₂max increases with the frequency of training. The slope of the improvement graphed over time became minimal after 5 days a week, but the incidence of injury increases significantly beyond 5 days per week.¹²⁹ The significance of this study is that there appears to be a point of diminishing returns.

b. Volume Vs Intensity

In a pair of companion, studies Lehmann et al studied the effects of over-training with an increase in training volume and an increase in training intensity. In the volume study, volume was increased 100% over 4 weeks. Almost all of the training consisted of long slow distance runs. All athletes failed to equal their personal records in a three month follow up. In the companion study, the total volume was increased by only 37%, but the proportion of intense training (tempo runs and intervals) was increased by 150%, following the concept of alternating hard and easy days. All athletes recorded personal bests during the three-month follow up.¹³⁰ The significance here is that increased volume has a more drastic and lasting affect of over training than an increase in intensity.

c. 3-Week Barrier

A series of tests on special forces soldiers and experienced athletes showed a decrease in performance after 2-3 h/d of training for 10-14 days, but super-compensation was exhibited after 1-2 weeks of recovery (tapering).¹³¹ The conclusion was that athletes are at a risk of over-training after three weeks of intensive or prolonged endurance training (2-3 h/d). They further concluded that alternating hard and easy days and one complete day of rest each week limits the potential for over-reaching or over-training.¹³² Interestingly, this study supported the anecdotal evidence that coaches have believed for several decades. This is important to the design of training programs.

d. Summary

The consensus is quite clear on four key points to avoid over training.. To avoid over training, at least one day a week must be complete rest. Hard days and easy days of cardiovascular training should be alternated and training crossing into the anaerobic spectrum (greater than 85-90% of reserve heart rate) should be limited to no more than once a week. After 3 weeks of hard training, additional recovery should be planned into the program.

5. CROSS TRAINING

Cross training is universally considered an excellent way to control the stress of over-training by shifting the stress from joints and soft tissue overuse in running. As cited above, several different studies have shown significant improvement in both VO₂max and run times when mixing running with cycle ergometer training despite training 6 days a week. These mixed training programs produced significant results in run times although the participants only ran 3 days a week and cross-trained on a cycle ergometer for the 3 additional days.

In a 1993 study, Edward Eyestone et al, compared water running, cycle ergometer and running for maintenance of VO₂max and 2-mile run times. Thirty-two trained subjects were tested for VO₂max and 2-mile run time. They were then separated into ability groups and then randomly assigned to three different groups so that each group had a spread of ability groups. All three groups exercised at similar frequency, intensity and duration over a 6-week period. The study concluded that there was no significant difference in changes in VO₂max or 2 mile run time between the three groups, thus

both water running and cycle ergometer training can maintain or even improve VO₂max in injured runners. ¹³³

In another study subjects who exercised on a stair climbing machine 30-45 minutes, four days per week for nine weeks at 70-90% of MHR improved their 1.5-mile run time trial by an average of 1 minute. Even at an 8 minute pace, this equals an 8.5% increase in nine weeks! ¹³⁴

There are some differences in heart rate at VO₂max for various forms of training. For instance, research has shown you must increase your heart rate range by 5-10 beats per minute on a cycle ergometer and decrease your heart rate range by 8-11 beats per minute in water running to equal the same percentage of VO₂max you would achieve when running on a track. ¹³⁵

6. PERIODIZATION

Training in cycles provides guidelines for the times in the training program when regeneration should be complete and limits over-training. Exposing an athlete to increasing levels of stress has been called overload training. In order for the athlete to recover and adapt (super-compensation), periods of recovery must be built in and severe increases in amount of stress must be avoided. This state of lack of adaptation has been called overreaching. ¹³⁶

A training program must provide for overload training within tolerance and regeneration periods that optimize performance improvements, but avoid over-training. The cycling of these training and regeneration periods has been called periodization. ¹³⁷ Fry, et al. cites studies indicate that several sessions in sequence can create a valley of fatigue necessary to induce training adaptation. It is unclear at present what the exact

number of sessions to create the optimum valley and what is the appropriate regeneration time.¹³⁸

Periodization is nothing more than a training plan that changes your workout protocol at regular intervals and builds in regeneration cycles. Both aerobic and strength periodization plans generally use microcycles (7-10 days long) within mesocycles (generally 4-6 weeks), within macrocycles (6 months to a year). The running and power-lifting communities have used cycles geared to the competitive season. Generally, runners use a base phase, a strength phase, a stamina phase, a speed phase and a peaking phase.

7. SUMMARY

Summarizing the points from above and their application to program design:

a. Frequency: The consensus of coaches and experts in the field of running suggest that you must run 5-6 days a week. . Obviously, the lower end of this range is more appropriate if the subjects are also involved in high intensity resistance training.

b. Duration: Again, the experts are amazingly consistent in their recommendations. In preparation for a 5-K run, the base phase should build to an 8/10-mile weekly long run and 25-35 miles a week. As athletes transition to a stamina phase the mileage should drop as the intensity increases.

c. Intensity: There are three key workouts each week to sustain the energy systems most prevalent in the 5-K run; 1 tempo run or cruise interval session, one aerobic capacity interval session and one long run alternating each of these with easy days of recovery runs or rest. During the base phase, coaches often substitute fartlek

runs in lieu of intervals and during the peaking phase, add some anaerobic capacity as you decrease the amount of aerobic capacity intervals.

d. Over-training: The consensus is quite clear on four key points to avoid over training. To avoid over training, at least one day a week must be complete rest. Since we are combining, high intensity resistance training 2 days is a logical progression. Hard days and easy days of cardiovascular training should be alternated and training crossing into the anaerobic spectrum (greater than 85-90% of reserve heart rate) should be limited to no more than once a week. After 3 weeks of hard training, additional recovery should be planned into the program.

e. Cross-training: Previous studies that have combined bicycle ergometer intervals with running have had lesser problems with over-training, probably due to muscle recruitment order. Research consistently shows that cycle ergometer training can be substituted for running with no degradation in improvement of VO₂max or run times. It therefore seems prudent to substitute this for at least one of the days of easy recovery runs to prevent soft tissue injuries particularly for middle aged runners.

f. Periodization: The most significant finding of all over-training studies is that you cannot increase intensity and distance simultaneously. Periodization is frequently cited as a solution to this problem. By increasing the mileage at no more than 10% a week during the base phase and gradually reducing the mileage as we increase the intensity in the stamina and peaking phases, you limit the opportunity for over-training. During a simultaneous high intensity endurance and strength program, the strength phase could logically be eliminated, as strength training is ongoing.

SECTION III: STRENGTH TRAINING

1. ACUTE VARIABLES

In 1983, using statistical analysis, Dr. W. J. Kraemer developed a set of variables that contribute the most significant change in results for a specific workout. These acute program variables describe all possible combinations within a single training session. They are exercise selection, order of exercise, number of sets, rest periods between sets, exercises & workouts and resistance used.¹³⁹

Exercise variables play a significant role in the serum responses of human growth hormone. Kraemer et al found it readily apparent that even keeping total work constant; changing one program variable significantly alters the serum growth hormone response pattern.¹⁴⁰

a. Choice of Exercise

The effectiveness of a program is strongly affected by the choice of exercise. Kraemer and Fleck, in their seminal work, "Designing Resistance Training Programs" categorize resistance exercises into two categories: structural, (Multi-joint), and single joint. Structural movements are those that require the coordinated action of several muscle groups, such as the squat, dead lift or power clean. Structural exercises are also called Multi-joint exercises or those exercises where movement occurs at multiple joints. Single joint exercises are those that isolate a specific muscle group. Examples are the sit-up, biceps curls, or leg extensions.¹⁴¹

Electromyography is a scientific method of measuring the level of excitement (electrical signal) in a working muscle group. Muscle contraction is initiated by electrical charges (ions) that move across muscle fibers. Scientists can measure this flow of ions

on the skin using a surface electromyogram (SEMG).^{142,143} An SEMG is representative of the entire electrical activity in all of the motor units in a given muscle and the frequency of their firing rates during a given movement.¹⁴⁴

The purpose of these studies was to find, using EMG recordings which exercises caused the greatest amount of stimulation within a muscle group. Exercises that produce the greatest amount of electrical activity during muscular contraction will produce the greatest amount of muscular efficiency. In their book " Serious Strength Training" Bompa and Cornacchia list exercises in order of IEMG activity for different muscle groups.

Coaches have know for years that whole-body exercises, or Multi-joint exercises are the best for building strength for sports such as football, wrestling or field events. The reason is that they stress the neuro-muscular links between muscles and require the generation of power. Because they involve the largest muscles of the body, they recruit the most muscle fibers. The studies cited above bore this out. The exercise that produced the greatest IEMG signal for the quadriceps (rectus femoris) muscle was the squat (90 degrees with a shoulder width stance). For the latissimus dorsi (back) muscle, it was bent-over rows and for the pectoralis major (chest), it was the decline and flat bench press. Dumbbells produced slightly greater recruitment than barbell.¹⁴⁵

For the various muscle groups, Table II-12 lists the most effective exercises and the percentage of IEMG stimulation:¹⁴⁶

There is some concern that Multi-joint exercises like the squat are dangerous. In a study at the Department of Health and Human Performance at Auburn University, Chandler, et al specifically studied the effect of the squat exercise on knee stability. In

this study, the authors measured one hundred male and female college students on nine tests of knee stability using a knee ligament arthrometer. Over an eight-week training period, full (below parallel) or half squats. In addition, the same measurements were collected on male powerlifters and male weightlifters. They found that not only did the 8 weeks of squat training (both full and half) not affect anterior and posterior knee stability, but that powerlifters and weightlifters with years of squatting had tighter knee joints (quadriceps active drawer measured at 90 degrees of knee flexion) than untrained individuals. The authors concluded that the squat had no affect on knee stability.¹⁴⁷

Muscle	Best (% IEMG)	Second (%IEMG)
Pectoralis Major (Chest)	Decline Dumbbell and Barbell Press (93 & 89%)	Flat Bench Dumbbell and Barbell Press (87 & 85 %)
Pectoralis Minor	Incline Dumbbell Press (91%)	Incline Barbell Press (85%)
Medial Deltoids	Incline Dumbbell Side Laterals (66%)	Standing Dumbbell Side Laterals (63%)
Anterior Deltoids	Seated Front Dumbbell Press (79%)	Seated Front Dumbbell Raises (73%)
Biceps Brachii (Long Head)	Biceps Preacher Curl (Olympic Bar) (90%)	Incline Seated Dumbbell Curls (88%) and Standing Barbell Curl (86%)
Triceps Brachii (Outer Head)	Triceps Extension (Olympic Bar) (92%)	Triceps Pushdown (angled bar) (90%)
Latissimus Dorsi (Back)	Bent over Barbell Rows (93%)	1 Arm Dumbbell Rows (91%)
Quadriceps	Squats (90degree angle, shoulder-width stance) (88%)	Seated Leg Extensions (86%)
Biceps Femoris (Hamstring)	Standing Leg Curls (82%)	Lying Leg Curls (71%)

Table III: 12: Neural Activation for Resistance Exercises

The significance of these studies is that the choice of exercise does make a difference in the overall improvement in strength and muscle hypertrophy.

b. Order of Exercise.

For years, strength coaches have advised that the order of exercise be from largest muscle group to smallest group. The reasoning behind this concept was to avoid limiting the resistance that could be used in a large muscle group exercise (e.g. bench press for the chest) by first fatiguing the triceps muscles which assist in the bench press. To test this concept, Kraemer and Fleck compared the workout logs of 50 football players, some of whom placed squats at the beginning of the workout, and some of whom placed squats at the end of the workout. The results were that players who executed squats at the beginning of the workout were using significantly heavier resistance.¹⁴⁸

Other authors refer to structural Multi-joint exercises as functional strength exercises. Unanimously, they call for these to precede single-joint exercises. The reason is that in doing functional strength exercises, you are also strengthening ligaments and thickening bones. Only substantial resistance places sufficient stress on the body to strengthen these ligaments and bones.¹⁴⁹ If you have placed these functional strength exercises at the end of the workout, as cited above, you will not lift enough weight to increase the stress on the muscular-skeletal system or recruit the type 2 fibers.

Health for Life, in their book Secrets of Advanced Bodybuilding has developed a principle called the interdependency of muscle groups. Their research has shown that one specific sequence of a given sequence of exercises is more effective than the same exercises performed in a random order. Their theory is that the body works as an integrated whole. It is impossible to activate one muscle without involving others.

During an exercise, the work is done by a “prime mover”. These muscles are helped by “synergists”. Other muscles keep you balanced by providing a foundation for stable movement. These muscles are called “stabilizers”.¹⁵⁰

In the example we used earlier, the bench press, the pectoralis muscles are prime movers, the anterior deltoids and the triceps are the synergists and the latissimus, rhomboids, posterior deltoids and abdominal are the stabilizers.

From this principle, Health for Life defines a set of applications which can increase the effectiveness of a sequence of exercises by maximizing the sequence. Those applications are:¹⁵¹

1. Never work a muscle as a prime mover before that muscle is called on to work as a synergist.
2. Do compound sets using the same prime mover, but different synergists to further fatigue the prime mover.
3. Pre exhaust the prime mover first before doing an exercise where the synergist will tire first. The idea is for the exhaustion of the prime mover to be the limiting factor, not the exhaustion of the synergist.
4. Sequence exercises so that a muscle that acts as a synergist or stabilizer in one exercise acts as the prime mover in the next exercise.

c. Number of Sets.

There has been an ongoing debate for several decades over how many sets you need to achieve maximum hypertrophy and strength gains. It is apparent that there is some threshold of fatigue, tension¹⁵² or strain that must be met to create the optimum stimulus for growth.¹⁵³ The stimulus must be sufficient to produce increased protein synthesis and thus hypertrophy in the muscle fibers.

Numerous studies have shown that in untrained individuals, almost any reasonable resistance-training program will initially induce growth. In fact, the American

College of Sports Medicine (ACSM) cites one set per exercise as its recommendation for quantity of exercise to promote health.¹⁵⁴ (The 1998 ACSM guidelines further state the multiple sets may promote greater gains if time is available).

In order to gain size, you must recruit more motor units. A single set can only recruit 80-85% of the motor units in a muscle. You must work your way through all of the motor units to get hypertrophy. After the initial period, gains using a one set format ultimately causes a plateau in performance gains. Studies bear this out. In those studies of more than 6 weeks, all have shown that multiple sets produce better gains than single sets, even in untrained individuals.^{155,156,157,158,159}

Multiple sets have also been shown to promote greater endurance adaptations. In 1992, McGee, et al studied the effect of three different weight training programs on leg and hip endurance as measured by cycle endurance at 265 watts and parallel squat endurance. The authors found that one set to failure did not increase high intensity exercise endurance as effectively as the use of multiple sets. This supports research in cardiovascular exercise that equates higher loads (volume x intensity) with greater gains in endurance.¹⁶⁰

That is not to say that one-set programs are not useful. They are frequently used by athletes who use multiple set programs in the off-season and a one-set program in-season to slow the loss of strength and muscular size while competing. The other clear instance is individuals who are time constrained. Business professionals who have competing priorities and have modified their goals from optimum strength or muscle size to general health or maintenance.

When you poll strength coaches on the optimum number of sets to elicit maximum strength and hypertrophy you find that they are consistent in recommending 2/3-6 sets for optimum strength and hypertrophy.

Author	# of Sets
Fleck & Kraemer 161	3-6
Stone et al. 162	3-5
Applewhite 163	2-6 (8 for leg press/squat)
Hatfield 164	2-5
Georgia State 165	4-6
Berger 166	3

Table III-13: Number of Sets

The total length of a workout, endurance or strength should not exceed 45-60 minutes total time. The hormonal response to exercise is for the body to begin secreting growth hormone shortly after the beginning of a workout. The body also begins secreting catabolic substances like cortisol, but the start is delayed relative to the production of growth hormone and testosterone. The difference in levels of growth hormone and cortisol is the net amount of protein synthesis that the body can optimally produce. Since the catabolic hormone curve is behind the anabolic curve, the difference begins to get smaller as you approach an hour. After an hour, the total amount of catabolic production begins to exceed anabolic production and growth potential is limited.

d. Rest between Exercises, Sets

Controlling the rest between sets or rest interval (RI) is probably the least used variable in workout design. Recent research has shown that the rest periods between sets and exercises control how much Phosphocreatine has been restored in the muscles and the level of lactate in the muscles. A series of studies by Kraemer, et al.^{167,168,169,170,171} demonstrated the significant affect of rest periods on blood lactate, hormonal concentrations and metabolic responses to resistance training in both men and women.

The significance of the findings is that the amount of work and duration of force demands on the muscle determine the blood lactate levels, not the amount of resistance used. In this case 10 repetitions at 10 RM with one-minute rest periods caused significantly higher lactate concentrations in the blood than 5 repetitions at 5 RM with 3-minute rest intervals.¹⁷²

The growth hormone response to a lifting session is directly related to the amount of lactic acid produced in the workout. Dr. Steve Fleck argues that the lactic acid response to a workout is directly correlated to the amount of work done in the total work session including rest periods.¹⁷³ In a study by Kraemer, et al. the researchers studied the growth hormone response to changing the load or rest period. In every combination, changing the rest period had a significantly greater affect on growth hormone response, both during exercise and up to 2 hours afterwards.¹⁷⁴

High blood lactate levels must be slowly introduced into the workout program so the body has time to build up the buffering mechanisms. In some sports like wrestling,

where the athlete must perform under conditions of high lactate concentrations, this form of training should ultimately reach a very high level.

Interestingly, the same mechanisms that buffer lactate during these resistance programs are the ones used when running at anaerobic threshold pace. This concept has obvious application to increasing middle distance running performance.

An inadequate rest interval obviously puts a lot of stress on the lactic acid system for energy because less phosphocreatine has been restored to the muscles. Reliance on lactic acid as a fuel means a greater accumulation of it in the blood and muscles. This, in turn reduces the ability of the body to produce force.

In their book, "Serious Strength Training", Bompa and Cornacchia define several physiological cues for rest intervals:¹⁷⁵

1. A 30-second RI restores approximately 50% of the depleted ATP/CP.
2. A RI of 3-5 minutes or longer allows almost entire restoration of ATP/CP
3. After working to exhaustion, a 4-minute RI is not sufficient to eliminate lactic acid or replenish glycogen stores.

In his address to the international summit of personal trainers, Mitchell S. Simon, MS, RTS, CPT, noted that the half-life of PCr is 20-28 seconds. In other words, after about 30 seconds, the body has restored 50%. After the next 30 seconds 75% has been restored. The body has almost completely restored its PCr stores in 8 minutes. That is the reason you watch a power lifter in the gym do a very heavy set and sit on the end of the bench for 8-10 minutes before the next set.

In addition, the nervous system must recover. When the brain sends a to a muscle signal to contract it transmits a nerve impulse. The stronger the nerve impulse, the stronger the contraction. When the muscle is tired, chemical disturbances are

signaled back to the central nervous system and the brain sends weaker nerve impulses to protect the body. Following a set, a 2-3 minute RI usually returns the nerve impulses to full strength.¹⁷⁶

Bompa and Cornacchia offer the following guidelines on rest intervals:¹⁷⁷

Load (% 1RM)	Rest Interval (minutes)	Applicability
>105% (eccentric)	4-5/7	Maximum Strength
80-100%	3-5/7	Maximum Strength
60-80%	< 2	Muscle Hypertrophy
50-80%	4-5	Power
30-50%	1-2	Muscle Endurance (definition)

Table III-14: Rest Intervals between Sets

These guidelines are almost identical to those offered by other strength coaches and scientific studies. For muscle hypertrophy, there is a great deal of evidence that lower rest intervals produce greater hypertrophy. In a series of studies by Kraemer, et al. a medium length rest period (1-3 minutes) resulted in a blood profile (increased growth hormone and testosterone) that leads to muscle growth.^{178,179,180,181} A medium length rest period allows enough rest to replenish phosphocreatine and allows the use of a heavy enough weight to increase strength.

Heavier resistance requires longer rest periods because the body has had to recruit the type 2 fibers and further exhausted the muscle. Long rest period (longer than 5 minutes) allow for almost complete recovery of phosphocreatine and allow near-

maximal resistance to be used.¹⁸² Thus when the goal is to increase strength and power heavier resistance and longer rest periods are appropriate.

Health for Life describes the difference as tension level and fatigue level. The greater the weight lifted, the greater the tension level and the greater the strength gains. The shorter the rest between sets, the greater the cumulative fatigue level. The greater the fatigue level, the greater the potential for hypertrophy. They hypothesize a fatigue/tension threshold which you must cross to cause growth. The fatigue/tension threshold can be crossed by a combination of greater resistance and/or shorter rest intervals.¹⁸³

e. Resistance Used (Number of Repetitions)

The resistance used is probably the most important variable used in the design of a training program. The scientific community has coined the term “repetition maximum” or the exact resistance that allows for the execution of a specific number of repetitions, but not one more repetition. The maximum weight that a lifter can lift one time in a given exercise would be a 1RM. The maximum weight that he could lift 10 times would be 10RM.

In what is probably the most cited study on number of repetitions, Richard Berger studied 199 college students to determine the optimum number of repetitions for strength gain. Nine groups were tested for bench press strength before and after 12 weeks of strength training. The author used resistance which allowed 2RM, 4RM, 6RM, 10RM and 12 RM for one set to failure. By using one set in all groups, the variable of training volume was controlled. Resistance was increased when any subject could do one more than the specified number of repetitions. The author found that training with

less than 2 repetitions and more than 10 repetitions will not improve strength as rapidly as training with 4,6,or 8 repetitions. They concluded that the optimum for strength gains was 3-9 number of repetitions.¹⁸⁴ Research supports a scale of repetition maximum ranges that correlate to training effects that they produce. In general, six or fewer repetitions target strength gains. The strength gains above 25 repetitions are insignificantly small or non-existent. Any strength gains produced with these higher levels of repetitions are probably due to neuro-muscular pattern efficiency.

A resistance of at least 60% of 1-RM is considered the minimum for optimum development of strength. Many strength coaches and popular strength books define resistance lower than that as a "warm-up" set and do not count it in program design.

Fleck and Kraemer, in their seminal work "Designing Resistance Training Programs" quantify the RM continuum. These recommendations are very consistent with all other findings on the subject.

Objective of Training / Number of Reps	Fleck & Kraemer 185	Hatfield 186	Georgia State U. 187 188	Bompa & Cornacchia 189
Strength	< 6	3-8	1-4	1-7
Hypertrophy	6-12	4-12	6-12	6-12
Local Muscular Endurance	12-20	25-40		30-150

Table III-15: Number of Repetitions

These findings are supported by a number of studies in which low repetition training (3-5 RM) produces significantly greater strength, intermediate repetition training (6-12 RM) produces the greatest hypertrophy and high repetition training (22-26 RM)

produces significantly greater local muscular endurance.^{190,191,192} Six to eight repetitions appears to be the optimum compromise between hypertrophy and strength.

At some point in a workout, a lifter must go to temporary concentric failure. You have not reached overload and will not cause the muscle to grow until you do so. Beyond temporary concentric failure, however, a lifter may actually defeat the body's ability to hypertrophy. Once you go beyond failure by using forced repetitions or negative repetitions, the strength of the muscle fiber is already gone. The additional effort is made up by the nervous system. During recovery, the body will preferentially recover the nervous system over the muscles. This makes sense, as the nervous system is the "conductor" of recovery. To allow hypertrophy, the body must send all of its recovery into protein synthesis not recovery of the central nervous system. It may be more appropriate to recruit additional motor units in the next set as opposed to extending sets with heavy forced repetitions.

f. Frequency

To a great degree, the frequency of workouts depends on the experience and recoverability of the lifter. Early studies and the current guidelines from the American College of Sports Medicine all indicated that 3 days of lifting, with a day of rest between each workout allowed adequate recovery for novice lifters and produced strength gains.

What the studies are really saying is that a muscle can be trained every other day. Muscles gain strength and size during the rest period between exercise periods. This does not mean that another muscle cannot be worked while one is resting.

Exercising every muscle in the body on the same day can lead to marathon workouts. Focusing a workout on one or two specific muscle groups and alternating the

muscles to be worked each day can and has led to development of programs involving 5-7 days a week without over training a given muscle group.

In one study, a significantly greater improvement in strength was found on a 5d/w schedule vice 2 and 3 days per week.¹⁹³ Other studies also indicated that 4(consecutive) or 5 times a week will produce greater strength gains,^{194,195} but in many cases, the workout designs in these studies were flawed.

In many cases, the participants executed the same exercises every day of the week vice splitting the muscle groups up over several days. No one designing resistance-training programs in the sports world would ever suggest this type of scheme. In addition, the number of exercises performed was very small. This certainly accomplishes the scientific requirement to isolate variables, but ignores the realities of training the entire body for athletic performance.

In a much larger study using a proven resistance-training program, Hoffman and Kraemer et al, studied the effects of self selection of a training frequency on sixty-one members of a division 1AA football team in the off-season.

All participants used a periodized approach including structural exercises and assistance exercises. The set/rep schemes for the structural exercises progressed from 4 x 8 in the first 4 weeks, 5 x 6 in the second four weeks and (10,8,6,4,2,10,10,10) in the last two weeks. Assistance exercises were held constant at 3 x 10.

In addition to the weight training, the players participated in a two-day a week conditioning program consisting of flexibility, agility, endurance, speed training and plyometrics. The options are listed in the table below:¹⁹⁶

Frequency	Frequency / Muscle Group	Sets / Week	Days	Muscle Groups
3d/week	3/week	150- 162	M,W,F	Entire Body
4d/week	3/week	118- 126	M,TH T,F	Chest, Shoulders, Triceps, Neck Legs, Back, Biceps, Forearm
5d/week	3/week	148- 156	M,W,F T,TH	Chest, Triceps, Legs, Neck, Back, Shoulders, Biceps, Forearms
6d/week	2/week	154- 162	M,TH T,F W,SA	Chest, Triceps Legs, Shoulders, Neck Back, Biceps, Forearms

Table III-16: Frequency

The results of the study showed a distinct lack of improvement in any strength measure for the 3d/week group. All groups significantly decreased 2-mile run times and body fat. The 6 d/week group had the highest average improvements in bench press (5.8 kg) and squat (12.5 kg) but the differences between players made the bench press measure for the 6/d week group statistically insignificant. Part of the reason may have been that there were an extraordinarily high number of exercises for chest and shoulders (7 and 8 respectively). This could have caused over-training. The 5d/week format was concluded to elicit the best performance in strength.

A review of the suggested frequencies from popular and scientific strength experts indicates remarkable consistency:

Author / Objective	Hypertrophy	Strength
Fleck and Kraemer ¹⁹⁷	6d/week Body Split:3	4d/week Body Split:2
Georgia State University ^{198,199}	5-6 d/week Body Split:3	2-4 d/week Body Split: 2
Tesch in "Strength & Power in Sport" ²⁰⁰	6d/week (4 On/ 1 off w/ 4 way split) or (6 on/ 1 off w/ 3 way split)	Soviet Model: 5d/week Body Split 1-2

Table III-17: Frequency: Hypertrophy Vs Strength/Power

2. CENTRAL NERVOUS SYSTEM

In 1988, Hakkinen, et al. did a 2 year study of elite male weight lifters. Over two years of well designed training, they gained a 37.9-kg (2.8%) in maximum weight-lifting total. There was a slight increase in muscle fiber size (.7 cm) but the magnitude of size increase would not account for this relatively large (for trained athletes) jump in strength. What made the difference? The averaged IEMG activity of the examined muscles went up by a large, but not statistically significant 4.2%.²⁰¹

Dr. Steve Fleck interpreted this data to mean; that once you approach or hit your genetic ceiling on physiological strength, the only remaining variable that you can improve is neural activation or IEMG activity.²⁰² As you approach your genetic potential and the bulk of the gains in strength are from increasing neural pattern efficiency or motor unit recruitment, the gains become smaller and smaller.

This interpretation is supported by numerous studies that found large initial increases (10-20%) in maximal strength in previously untrained subjects. As the upper

limit of strength is approached, however, increases in muscle size and consequently strength are very difficult to achieve.²⁰² When working with United States Olympic Weight Lifters, Dr. Fleck noted that a gain of 2 Lb. of lean body weight in one year was a significant gain.²⁰³

At the higher levels of neuro-muscular efficiency, the antagonist muscles actually decrease in IEMG activity when a muscle is firing, thus increasing the neural signals to the motor neurons in the firing muscle and lessening the wasted force vectors from antagonist muscles, often found in untrained subjects.

In a study by Moritani and DeVries a 30%, increase in performance (strength) was studied to determine the neural factors and hypertrophic factors of the gain. The study concluded that 9% of the gain was caused by hypertrophy of the muscle fibers, and 12% of the gain was caused by increased IEMG activity in the working muscle.²⁰⁴ (Increased IEMG activity in a working muscle means that more motor units have been recruited; that motor units are firing at higher rates or some combination of the two adaptations is occurring). They found the remaining 9% to be the relaxing or lessening of IEMG in the antagonist muscles.

In a recent study, researchers found a 23% gain in bench press in 48 hours from one training session to another. Since it takes 7-10 days for protein synthesis to add muscle, this is clearly not hypertrophy, but neural gains.²⁰⁵

3. PERIODIZATION

a. Weight Training Systems

Prior to the recent studies on periodization, in both the weight training and scientific communities, there has been continuous debate over the relative efficiency of one or another systems of weight training. In a landmark study Leighton et al studied over 300 college students, using 10 different weight- training systems over eight weeks. This study is significant in that the number of participants is unusually large for an anaerobic exercise format. For safety, most studies involving anaerobic exercise are kept small. The ten different systems are briefly described below.²⁰⁶

1. Power: Increase loads and decrease the number of repetitions on each set.
2. Superset: Using 2 exercises for the same muscle group or antagonistic groups with little or no rest between exercises.
3. Cheating: Bringing in other muscle groups to assist in the exercise in order to handle heavier poundage i. e. rocking with the lower back to help with a barbell curl.
4. Bulk: Multiple sets using the same repetitions and load for each set.
5. Tri-set: 3 exercises for the same body segment- can include different muscles- with little or no rest between exercises.
6. Double-progressive: Increasing the resistance by first reaching a specified number of repetitions and then increasing the load and decreasing the repetitions to the original number.
7. Heavy-Light: Using maximal poundage for one workout period each week and 2/3 maximal poundage during the other periods that week.
8. DeLorme: Three sets with 10 repetitions each. The first with $\frac{1}{2}$ maximal load, the second with $\frac{2}{3}$ maximal load and the last with maximal load (10RM).
9. Oxford: The same poundage and number of sets as DeLorme, but in reverse order.

10. Isometric: A single maximal contraction held for 6 seconds.

The cheating and tri-set routines elicited the greatest improvement in arm flexor and arm extensor strength, although the heavy-light and Oxford systems were nearly equal. In leg strength, the bulk system was clearly superior. Interestingly, the isometric system caused actual but insignificant losses in strength.²⁰⁷

b. Periodization Works.

The sports science community began comparing periodized resistance protocols to conventional protocols in the early 1980's. In 1983, Stowers et al. compared the short-term (7 weeks) affects of three different strength-power training methods on trained weight lifters. Each group trained 3 days a week. They compared 1 set of 10-12 repetitions, 3 sets of 10 repetitions and a periodized approach. After warm up sets, the 1 set group executed 1 set of each exercise using 10-12 repetitions to exhaustion. After warm up sets, the 3 set group did 3 sets of 10-12 repetitions to exhaustion, lowering the weight after each set as appropriate to stay within the repetition scale. The periodized group did 5 sets of 10 for 2 weeks, 3 sets of 5 for 3 weeks and 3 sets of 3 for the last two weeks. The results are listed in table:

Group/measure	Bench Press	Squat	Vertical Jump
1 set	6.6 Kg	15.2 kg	.2 cm
3 sets	8.3kg	20.3 kg	.7 cm
Periodized	7.4 kg	27.o kg	5.7 cm

Table III-18: Periodization: Stowers et al

Of interest is that the 3 set group showed relatively equal improvements to the periodized group in all three areas, but reached a plateau in the final two weeks. The periodized group was still improving when the test ended. In addition, of significance is

that the improvement in vertical jump of the periodized group was over 800% of the 3 set group and 2800% of the one set group. This indicates that motor unit recruitment or neural performance was significantly affected by the periodized protocol.

In a more recent study by Darryn S. Willoughby from the Department of Health and Kinesiology at Texas A&M University studied 92 participants, again a large population for a study on anaerobic exercise. This study is also significant because it used participants with previous weight training. It has been clearly shown that untrained subjects will gain significant strength and hypertrophy on almost any program. All participants for this study had to have at least 3 years of previous weight training experience and have abstained for 6 months prior to the study.²⁰⁸ To participate lifters had to be able to bench press 120% of body weight and squat 150% of their body weight. This implies a great deal of previously developed strength. The study is also significant in that it lasted 16 weeks, much longer than previous studies, and it partially equated the volume of training among the different groups.

All groups trained 3 times each week on Monday, Wednesday and Friday and were tested on free weight bench press with an Olympic bar and parallel back squat.

Group 1 trained with 5 sets of 10-RM using 78.9% of 1-RM. Group 2 trained with 6 sets of 8-RM using a resistance of 83.3% of 1-RM. The resistance for both groups was increased if appropriate after each 4-week test. Group 3 used a periodized approach with 4 weeks at 5 sets of 10-RM with 78.9% of 1-RM; 4 weeks at 4 sets of 8-RM with 83.3% of 1-RM; 4 weeks at 3 sets of 6-RM using 87.6% of 1-RM; and 4 weeks of 3 sets of 4-RM using 92.4% of 1-RM.

This protocol was an obvious progression of decrease in volume and increase in intensity. The volume of training was nearly equal until after week 8 when group 3 decreased below the other 2 groups. Group 4 was a control group that met all of the prior experience prerequisites, but abstained from weight training during the 16 weeks.²⁰⁹

A 4 x 5 MANOVA with repeated measures on test showed pre-test bench and squat values were statistically equal when the study began.²¹⁰ At the end of the weeks 8,12 and 16, group 3 showed significantly greater strength gains over the other three groups. Groups 1 and 2 showed lesser gains in strength that were statistically equal at weeks 8,12 and 16.²¹¹ The results are listed in table x:

Group/Measure	Bench Press	Squat
3 sets of 10	8%	13%
3 sets of 6-8	17%	26%
Periodized	28%	48%

Table III-19: Periodization: Willoughby

The implications are clear that some variation in training improves strength gains. In comparative studies and reviews the results consistently found that in short term training, periodization produces significantly greater strength gains than traditional methods.^{212,213,214,215,216}

c. Periodization Cycles.

The terms used to define training phase length in a periodized program have become relatively standard in the strength and scientific communities.²¹⁷

(1) Macrocycle: Although it is normally used to describe one entire year, it can describe a period of as little as 6 months or as long as 4 years in the case of Olympic

athletes. It is generally thought of as beginning and ending after the last competition of a season.

(2) Mesocycle: Originally used to define the major training cycles of an entire year of training. Recent evidence clearly shows that to elicit continuing gains, changes must be made every 4-6 weeks. This term has thus been altered from a period of 2-3 months to one of 4-6 weeks.

(3) Microcycle: Most coaches and scientists use microcycle to refer to a period of seven days to correspond to the week. There is some ongoing evidence that a cycle of 10 or 14 days may be ideal to prevent over training, but in general a microcycle today still defines a 7-day period.

Lifters and lifting coaches from strength and power sports use five phases or mesocycles: Hypertrophy phase; Strength phase; Power phase; Peaking phase and Active rest phase.

4. OVER-TRAINING

a. What is Over-Training

Progressive overload is the foundation of successful adaptation to training. Overload is the stress that causes fatigue (temporary decrease in exercise ability), and adaptation is the improved performance (following recovery from that fatigue).²¹⁸

If you were to draw a graph of responses to training in relation to genetic potential the over-reaching phenomenon would stand out clearly. A planned suppression in performance caused by an increase in training is followed by a rebound closer to

optimum potential. If the suppression continues too long, the athlete is in a state of over-training. True acute over-training requires months of recovery.

Callister et al. increased the volume of resistance training, running and judo practice in the U. S. National Judo Team over six weeks. The expected increase of resting heart rate or resting blood pressure did not occur. What they did observe was significantly decreased times in 300-m sprints and decreased isokinetic force production.²¹⁹

The first controlled model to create and study over-training in a lab was done by Fry et al. They specifically identified suppressed testosterone and elevated cortisol endocrine profiles as markers of anaerobic over-training.

b. Application of Acute Variables in Over-Training

Manipulation of the acute variables, described by Fleck and Kraemer are the means of progression in strength training. Mistakes in the application of these variables lead to over-training. The use of periodization is an attempt to plan the progression for optimum adaptive response while preventing over-training.

(1) Choice of Exercise

You can classify weight training equipment as fixed form or free form. Free form involves equipment like free weights that allow movement through multiple planes of motion and require balance. Fixed form fixes the position and range of motion, normally isolating the muscle at a specific joint. A lack of variation in the pattern of motor unit recruitment means lack of development in the target muscle. In an isolation exercise, you deprive the muscle of the requirement for balance thus not training the synergist muscles. In addition, not providing for balance in the growth of antagonist muscle

groups around a joint (like the quadriceps and hamstring muscles) can create dangerous muscle imbalances. The choice of exercise is critical to the balanced development of muscles in order to prevent over-training through injury.

(2) Order of Exercise.

A plateau in the adaptive training response is possible if you exercise muscles in a fatigued state. The use of pre-exhaustion techniques, using isolation exercises prior to Multi-joint exercises clearly causes this fatigue. A maximum load must be used if you want to achieve maximum gains in strength. This implies placing structural exercises at the beginning of a workout.

(3) Number of Sets

Work is generally defined as sets x repetitions x load. If total work is increased too quickly, the muscles do not recover, causing over-training. The clear implication is a planned progression of workload over time to achieve optimum adaptation.

(4) Amount of Rest between Sets and Exercises

This variable has dramatic effects on the acid-base status of muscle and energy substrate utilization and depletion patterns.²²⁰ Rest periods of less than 1-minute have been shown to create blood lactate concentrations of greater than 10 mmol/L in untrained individuals and as high as 20 mmol/L in elite lifters. Since direct links to muscle fatigue have recently been attributed to hydrogen ions as well as the lactate itself, about 8-weeks is required to develop the body's lactate buffering systems. The implication for training is building lactate levels too high, too early in training will cause over-training. A planned progression in the interactive variables (e. g. number of sets;

number of repetitions; resistance used and rest between sets) is critical to allow for progressive adaptation.

(5) Rest between Workouts

Even one day between workouts has been shown to significantly reduce the markers of over-training. In two studies by Fry, et al. recreationally trained athletes performed 8 sets of machine squats at 95% of 1-RM, six days a week. No reduction of 1-RM was observed however several other signs of over-training were observed, specifically sprint times and agility to the non-dominant side. The conclusion was that the body protects high force production of 1-RM and even one day of recovery a week allowed tolerance of this protocol.²²¹

In the follow up study, the day of rest was eliminated resulting in significant decrease in 1-RM strength in 73% of the participants. Interestingly, some participants were still making progress and were not yet in an over-trained state. Clearly, the adaptation to stress is an individual response, but some clear guidelines have emerged for program design.²²²

5. SUMMARY

Summarizing the points from above and their application to program design:

a. Choice of Exercise: Electromyography clearly indicates that some exercises are better than others for recruiting motor units within a muscle. Structural or Multi-joint exercises are generally more effective than isolation exercises. In the development of a resistance training program too many exercises can lead to over-training, thus in attempting to recruit as many motor units as possible, the selection of exercises must include those that are more effective (see table III-1).

b. Order of Exercise: The preponderance of evidence indicates that structural or Multi-joint exercises should precede isolation exercises in a program. This allows the use of maximal resistance thus greater gains in strength. In general, you should sequence exercises so that a muscle does not act as a synergist before it is called on to become a prime mover so the point of failure is the prime mover and not a synergist or assistance muscle.

c. Number of Sets: Multiple sets are better than a single set and the optimum number of sets for hypertrophy and strength improvement is somewhere between three and six.

d. Rest between Sets. A muscles needs between 3 and 5 minutes to almost entirely replenish its phosphocreatine stores, however after 30 seconds, almost half of the CP is restored. The nervous system needs 2-3 minutes to return nerve impulses to full strength. The fatigue-tension principle indicates that a combination of load and rest period creates a fatigue/tension threshold that must be crossed in multiple sets to create the conditions for optimum growth. Medium-length rest intervals (1-3 minutes) creates a blood profile (increased growth hormone and testosterone) that causes hypertrophy. Heavier resistance, required for optimum increase in strength require longer rest periods (3-5 minutes) to fully replenish the CP stores in a muscle and revive the nervous system.

e. Resistance: A load of 60% of 1-RM is the minimum required for developing strength. Less than this primarily enhances muscle endurance by increasing lactic acid buffering capacity in the muscle. The strength gains beyond 20-25 repetitions are insignificant. Six or less repetitions appear to be optimum for increases in strength. Six

to twelve repetitions affect muscle hypertrophy and repetitions greater than 12 tend to enhance muscular endurance more than strength.

f. Frequency: A muscle needs 24-48 hours to recover from a resistance-training workout. The minimum number of days each week to train for optimum strength gains appears to be about three. Several major studies and the preponderance of popular strength and body-building coaches indicate that 5-6 days a week, splitting the body into groups of muscles is the most effective approach to hypertrophy and strength.

g. Periodization: It works. A planned variation in the acute variables over time produces significantly greater muscle gains than any combination of variables. In strength sports, the cycles should be 4-6 weeks long, progressing from hypertrophy to strength to power to peaking to active rest. If power lifting is not one of the program goals, the strength phase can be extended slightly.

h. Over-Training: The body must have at least one day of recovery each week from resistance training. The planned periodization of acute variables provides a clear advantage in avoiding over-training in strength athletes.

SECTION IV: COMBINED STRENGTH AND ENDURANCE TRAINING

Training is specific. More particularly, the cellular changes in the muscle are specific to the type of exercise. Are strength and endurance exercise compatible? Exercise compatibility studies have focused on endurance and strength compatibility in the last 2 decades.

1. INITIAL STUDIES

The question exercise compatibility first surfaced in the exercise science community in 1980, when Hickson compared three groups over 10 weeks of training. Group 1 did both high intensity resistance and high intensity aerobic training. Group 2 did only high intensity resistance training. Group three did only high intensity aerobic training. His conclusion was that although strength training did not compromise improvement in aerobic capacity, endurance training compromised simultaneous strength training.

The design of the study has some potential flaws as a model for combined training. The strength component, used by both the strength group and the combined group focused solely on leg strength. Three days a week they performed 5 sets of parallel squats at 5-RM, and 3 sets of knee extensions and knee flexion at 5-RM. On alternate days they 3 sets of leg presses at 5-RM and 3 sets of calve raises at 20-RM. Rest intervals were 3 minutes for all sets.

In other words, they worked legs five days a week. What is interesting is for the first 7 weeks, the rate of improvement for the strength and combined groups was identical, with both groups improving by an average of 26-kg in the parallel squat to that point. The strength only group continued to improve to an end-point of 42-kg by the

tenth week. The combined group's improvement leveled out during week 8 and actually declined during weeks 9 and 19 to an overall average of 22-kg above start.²²³

The endurance-training program combined 6 days of intense training. Three days a week the [participants performed bicycle ergometer intervals (six 5-min intervals at VO₂max separated by 2-min rest intervals). On the alternate 3 days, they ran continuously as fast as possible building from 30 minutes a day during the 1st week to 40 minutes a day by week 3. There was no difference in the improvement of VO₂max in the strength and combined groups, with both improving 17% over 10 weeks.²²⁴

Since the time of this study, significant evidence has shown that alternating hard and easy days will help to prevent over-training. In this study, they trained hard every day. The gains were beginning to level off by the tenth week for both the endurance group and the combined group. The significance of this study is that endurance training was not compromised by simultaneous strength training, despite the extremely high volume of intensity in both protocols.

2. POWER, NOT STRENGTH MAY BE AFFECTED

In 1985, Dudley and Djamil studied a significantly lower volume of combined training for seven weeks. Strength training consisted of 2 sets of knee extensions 3 days a week for 30 seconds (26-28 contractions) separated by 5 minutes. Aerobic training involved 5 sets of 5 minutes achieving VO₂max in the 4th or 5th minute, separated by 5-minute rest intervals. The combined group alternated endurance training days with strength training days. As in the Hickson study, the improvement of aerobic capacity was not affected by the combined training. Combined training did not

affect the high force, slow velocity 1-RM strength. The high velocity component of strength (angle specific peak torque at fast velocity) was compromised. This was the first study to suggest that combined strength and endurance training might affect power, vice strength.²²⁵

3. A MORE REALISTIC PROTOCOL

In 1987, Hunter, Demment and Miller examined what coaches would consider a much more realistic training program of combined exercise. Participants executed 8 workouts a week for 12 weeks. The strength portion consisted of 6 exercises: bench press, seated press, lat. pull down, two-hand curl, squat and leg curl. They executed 3 sets of 7-10 repetitions, adding weight when they were able to do 3 sets of 10 repetitions. The endurance portion consisted of continuous runs at 75% of heart rate reserve increasing from 20 minutes the first week to 40 minutes in the eight-week and beyond. The subjects trained 6 days each week, with two days including both strength and endurance workouts.²²⁶

In addition to the 3 standard groups, they included a group of trained endurance athletes who executed combined training. As in previous studies the improvement of VO2max for the untrained endurance and combined groups were identical (33%). The trained endurance athletes had, as would be expected a lesser increase in VO2max.

Although the combined group improved similarly to the strength group in the parallel squat, the strength group improved vertical jump by almost twice as much. Interestingly, the endurance athletes executing combined training had the greatest overall improvement in bench press, squat and equaled the strength group in vertical

jump. Hunter et al. concluded that power, but not strength may be compromised in untrained individuals training simultaneously for strength and endurance. If the endurance component is maintained at previously established levels (32-K/week in this case) power was not compromised.²²⁷

In a more recent 10-week study, McCarthy et al. used a much more typical training routine with sedentary adult males. The strength group performed 3 sets to failure of 8 exercises (bench press, squat, barbell curl, knee extension, leg curl, wide grip pull-downs, overhead press and heel raises). The range of repetitions was 5-7 with the goal being a 6-RM on each set. The strength protocol was executed 3 days a week. The endurance protocol involved 3 days a week of continuous cycle ergometry, building from 30 to 50 minutes at 70% of reserve heart rate. The intensity was recalculated every two weeks. The combined group performed the same protocols on the same day. The order of training was rotated and the training sessions were separated by 10-20 minutes.²²⁸

The results indicated that the magnitude of increase in both strength and VO₂max were the same for the combined protocol as the strength and aerobic protocols performed separately.²²⁹ The significance of this study was that the subjects in the combined group had 4 days a week of recovery, vice one day in previous studies. Obviously, for even untrained subjects, combining training effects is possible.

These results were supported in a study by Volpe et al. Previously sedentary college females were placed on a 9 week periodized weight training program. The combined group ran prior to their weight training. They started at 20 minutes of run-walk and progressed to 25 minutes of running at 75% of reserve heart rate. The

combined group performed significantly better than the strength only group in the 12-minute run after training, thus there was a significant improvement in aerobic capacity. There were no statistical differences in the strength improvements between the strength and combined groups.²³⁰

4. THE CELLULAR LEVEL

There have been several studies attempting to explain the cellular affects of simultaneous strength and endurance training. The muscle fiber is faced with a dilemma. As explained in the physiology section, the muscle must try to increase mitochondria, oxidative enzymes, triglyceride stores and capillary density to improve aerobic function or to increase the enzymes that assist in force production. In other words, do muscle fibers move towards type 1 characteristics or type 2 characteristics?

Kraemer et al. studied these cellular changes over a 3-month period in 1995. The training protocol involved 4 days a week. The strength component was 2 days of 5 sets of 5-RM and 2 days of 3 sets of 10-RM. The running involved 2 days of intervals at 95-100% of VO₂max and 2 days of continuous running for 40 minutes at 80-85% of VO₂max. The workouts for the combined groups were planned such that the high lactic acid workouts (10-Rm and Intervals) were not executed on the same day. This study included a 4th exercise group that did only upper body exercises and running.²³¹

The results supported previous studies in that VO₂max increased significantly for both combined groups and the endurance only group, although the endurance only group improved slightly more than either combined group. The improvement in the combined groups was almost identical.

Muscle fiber transformation was unique to the groups. The combined group had a significant increase in both size and percentage of type 2a fibers. The upper body combined group saw a significant increase in percentage of 2c and 2a muscle fibers along with a significant decrease in percentage of type 2b fibers. No increase in fiber size was noted. The strength only group realized significant increases in percentage of type 2a fibers and significant decreases in percentage of type 2b fibers. In addition, the strength only group had significant increase in fiber area in type 1, 2c and 2a fibers. The endurance only group had a significant increase in percentage of type 2c and 2a fibers and a significant decrease in the percentage of type 2b fibers. The endurance only group also saw a significant decrease in the size of type 1 and 2c fibers.

These findings support Dudley and Djamil in that power was compromised in the combined groups. The 1-RM strength of the combined group appears to be similar to that of the strength only group. The upper body only combined group had similar improvements to the strength only group in upper body strength, but not lower body strength.

A study by Craig et al. in 1991 also supports these findings. Craig also found that a program of running and lifting could produce the same gains in VO₂max and upper body strength. Craig found, however that combined training did not elicit similar lower body strength. In Craig's study, the lifting program immediately followed the running. They hypothesize that this was the cause of the lack of lower body strength improvement.

The protocol is, however, at odds with the findings on frequency of both resistance and aerobic protocols developed in the last two sections. In the endurance protocol of

this study, there were four hard runs in a week and the strength protocol worked the same muscle groups 4 days a week. The unanswered question is the effect of combining popular protocols. For instance one involving a 5-K running program with only one day of intervals and one day of lactate threshold runs each week and a strength program which splits the body up and works each muscle group only 1 ½ – 2 times a week.

SECTION V: CONCLUSION

1. SUMMARY

To summarize the results from the previous 4 sections, a combined training program would include each of the points in the individual summaries of each section. The tables below represent those summaries in the form of a recommended training program. This is a program designed specifically for increasing cardiovascular performance in middle distance running and 1-RM strength. A program tailored for a specific sport would obviously be different. The program outlined above uses 11 workouts a week with 5 days a week being two workouts a day and one day completely off for regeneration. The next chapter will deal with time constrained programs.

The strength program will split the muscle groups into 4 days, executing 6 workouts each week. In this way, each muscle group is worked 3 times in two weeks with Sunday completely off.

2. RECOMMENDATION

Table V-20: Periodization

	Mesocycle 1: (4-8 weeks)	Mesocycle 2: (4 weeks)	Mesocycle 3: (4 weeks)
Aerobic	Base Phase	Stamina Phase	Peaking Phase
Strength	Hypertrophy Phase	Strength Phase	Peaking Phase

Table V-21: Schedule

Week / Day	Workout	Week 1	Week 2	Week 3	Week 4
Monday	1	Run: Tempo	Run: Tempo	Run: Tempo	Run: Tempo
	2	Chest	Legs	Chest	Legs
Tuesday	1	Bike	Bike	Bike	Bike
	2	Shoulders, Triceps	Back, Biceps	Shoulders, Triceps	Back, Biceps
Wednesday	1	Intervals	Intervals	Intervals	Intervals
	2	Legs	Chest	Legs	Chest
Thursday	1	OFF	OFF	OFF	OFF
	2	Back, Biceps	Shoulders, Triceps	Back, Biceps	Shoulders, Triceps
Friday	1	Run: Easy	Run: Easy	Run: Easy	Run: Easy
	2	Chest	Back, Biceps	Chest	Back, Biceps
Saturday	1	Run: Long	Run: Long	Run: Long	OFF
	2	Shoulders, Triceps	Legs	Shoulders, Triceps	OFF
Sunday		OFF			

Table V-22: Acute Variables: Strength

Phase	Exercises/ muscle group	Sets (MJ) per exercise	Reps (MJ)	Rest (MJ)	Sets (SJ) per exercise	Reps (SJ)	Rest (SJ)	Total Sets
Base	4 large 3 small	3	10	1	3	10-12	1	21
Strength	3 large 2 small	4	6	3	3	8	2	18
Peaking	3 large 1 small	4	6-1	4	3	6	2	15

1. MJ = Multi joint
2. SJ = single joint
3. Large muscle groups e.g. chest, back, legs
4. Small muscle groups e.g. biceps, triceps
5. Rest between sets in minutes

Table V-23: Frequency: Strength

Day	Week 1	Week 2	Week 3	Week 4
Monday	Chest	Legs	Chest	Legs
Tuesday	Shoulders, Triceps	Back, Biceps	Shoulders, Triceps	Back, Biceps
Wednesday	Legs	Chest	Legs	Chest
Thursday	Back, Biceps	Shoulders, Triceps	Back, Biceps	Shoulders, Triceps
Friday	Chest	Back, Biceps	Chest	Back, Biceps
Saturday	Shoulders, Triceps	Legs	Shoulders, Triceps	OFF
Sunday	OFF	OFF	OFF	OFF

Table V-24: Acute Variables: Cardiovascular

Type workout /Phase	Base Phase 4-8 weeks	Stamina Phase 4 weeks	Speed/Peaking Phase 4 weeks
Monday	Tempo Run (5) increasing from 2@80% to 4 miles @ 85% RRHR	Tempo Run (5) 4 miles @ 85% RRHR	Tempo Run (5) 4 miles @ 85% RRHR
Tuesday	Bike: (Aerobic Capacity Intervals) increasing from 3-6 sets of 3:00 min at 85% RRHR	Bike: Aerobic Conditioning 45 min @ 55-65% RRHR	Bike: Aerobic Conditioning 45 min @ 55-65% RRHR
Wednesday	Fartlek Run increasing from 2-4: ½ mile pick ups @ 80-85%. The base running is at @ 55-65% MHR	Aerobic Capacity Intervals: 220, 440 warm ups and 880 intervals building to 6-8x880 @ 90% RRHR	Aerobic Capacity Intervals: same as stamina decreasing to 2x880 and adding anaerobic intervals building to 4 (110m) @ 95%
Thursday	OFF	OFF	OFF
Friday	Run: Aerobic Conditioning increasing from 3-6 miles @ 55-65 % RRHR	Run: Aerobic Conditioning 3 miles @ 55-65 % RRHR	Run: Aerobic Conditioning 3 miles @ 55-65 % RRHR
Saturday	Run: Aerobic Conditioning (Long) increasing from 8-10 miles @ 55-65% RRHR. Every 4 th week is OFF	Run: Aerobic Conditioning (Long) alternating 6/8 miles @ 55-65% RRHR. Every 4 th week is OFF	Run: Aerobic Conditioning (Long) alternating 6/8 miles @ 55-65% RRHR with the 4 th week off
Sunday	OFF	OFF	OFF
Total Intensity Mileage	None above 85% RRHR	Building from 8-9% the first week to 10% the last week none above 90% RRHR	Decreasing to 8%. 880's remain at 90% and 110's increase to 95% RRHR
Total Running Mileage	Increasing to 30 over 4 weeks or 40 over 8 weeks (10%/week)	22-28 miles/week depending on length of base phase	Run 18-24 miles/week.

1. RRHR = Reserve/Resting Heart Rate calculated with Karnoven Method
2. Assumes a base of 20 miles / week before starting base phase

SECTION VI: DETRAINING: TRAINING ON LIMITED TIME

The program outlined above incorporates 11 workouts a week, which take about 1.5 hours a day on most days. Most people are probably not willing or able to commit this amount of time to a workout program. This chapter will use the principles outlined above to construct the optimum program for one hour a day, 5-6 days a week.

1. CARDIOVASCULAR TRAINING

a. Intensity

Detraining studies unanimously indicate that intensity not duration is key to sustaining performance during periods of reduced training.

Hickson, et al studied the effects of both reduced training duration and reduced training intensity in previously trained subjects. In both studies, subjects participated in an intense cycling and running program for 10 weeks. The training was performed 40 minutes a day, 6 days a week, and consisted of intervals on a cycle ergometer and continuous running. Following the training period, both studies studied the results of 15 weeks of detraining where intensity and frequency remained constant, but duration was reduced by one-third or two-third.²³²

In the reduced duration program two groups continued to train for 26 and 13 minutes a day. This corresponded to one-third and two-third reductions. Calculated left ventricular mass increases 15-20% after training and remained elevated during detraining. VO₂max increased 10-20% during training and did not change during the detraining. Short-term endurance was measured by running and cycling at maximal work rates that resulted in exhaustion in 4-8 minutes. Short-term endurance remained

the same in both reduced duration groups after 15 weeks of reduced training. Long term endurance, as measured by cycling to exhaustion at 80% of VO₂max increased by more than two-fold during training. Interestingly, after 15 weeks of detraining, the one-third reductions group showed no loss in long term endurance. The two-thirds reduction group, however showed a significant loss in long term endurance.²³³

In a companion study of reduced training intensity, Hickson et al , similar groups executed an identical training program followed by 15 weeks of reduced intensity. Duration and Frequency were held constant, but work load was reduced by one-thirds and two thirds respectively. On the cycle, the pedal resistance was reduced and in running workouts, the speed was reduced.²³⁴

The results indicated that intensity is much more significant in maintaining cardiovascular performance. As in the reduced duration study, left ventricular mass increased by about 15%, but with reduced intensity, it returned to control levels in both groups. VO₂max increased from 11-20% after 10 weeks of training, In the one-thirds reduction group, VO₂max was significantly lower after 10 and 15 weeks of reduced training, but still slightly higher than pre-training level. In the two-thirds reduction group, significant decline was evident after only 5 weeks and remained low through 15 weeks. In running, the 15-week VO₂max was actually lower than pre-training level.²³⁵

The significance of these two studies is that in situations of reduced training, it is more important to retain intensity than duration. The exception being long term endurance to exhaustion. The two most important forms of intensity are lactate threshold runs and intervals. Advice from coaches supports the contention that one of

each of these in a 7-day microcycle is adequate to maintain or slightly improve cardiovascular performance.

b. Duration

For maintenance of long term endurance, duration appears to be important. Dudley et al found that longer exercise bouts produce greater increases in mitochondrial density, however there is a point of diminishing returns (see figure 1). It is important to note that after 1 week of detraining, almost 50% of the increase in mitochondrial content were lost. After 5 weeks all of the adaptations had been lost (see figure 2).²³⁶ The significance of this finding is that to maintain the full range of cardiovascular capability an athlete must devote at least one day in each microcycle, to a long run. Advice from running coaches supports this evidence.

The significance of this finding is that an athlete must maintain at least one longer run each week to sustain the mitochondrial content of the muscle fibers.

c. Frequency

The American College of Sports Medicine recommends training cardiovascular systems 3-5 days each week. If you only have a limited amount of time to devote, then each workout must have a specific purpose.

The evidence clearly supports 3 specific workouts over 7 days; one long run to maintain long term endurance, one lactate threshold run to maintain endurance at the anaerobic threshold, and one day of intervals to maintain anaerobic glycolysis and recruitment of type 2 fibers.

2. STRENGTH TRAINING

a. Acute Variables

The application of the acute variables will not change with the reduction in frequency. A proven formula is a periodized program using at least 3 sets, varying the number of repetitions and resistance by mesocycle. Those recommendations are covered in the section on Strength Training.

b. Frequency

To a great degree, the frequency of workouts depends on the experience and recoverability of the lifter. Early studies and the current guidelines from the American College of Sports Medicine all indicated that 3 days of lifting, with a day of rest between each workout allowed adequate recovery for novice lifters and produced strength gains.

What the studies are really saying is that a muscle can be trained every other day. Once resistance workouts reach a high level of intensity, then more than 48 hours is appropriate between workouts. Muscles gain strength and size during the rest period between exercise periods. This does not mean that another muscle cannot be worked while one is resting.

McCarthy et al cited significant results from a 6 workout per week format (3 cardiovascular and 3 strength).²³⁷ Hunter et al proved that 8 workouts (4 cardiovascular and 4 strength) a week over 6 days was also effective.²³⁸ These studies provide a framework for training options. Both 3 and 4 days a week are proven to be effective when combined with cardiovascular training. The optimum frequency per

muscle group appears to be 1.5 –2 times per week when combined with high intensity cardiovascular exercise.

3. RECOMMENDATION

Table V-25: Periodization

	Mesocycle 1: (4-8 weeks)	Mesocycle 2: (4 weeks)	Mesocycle 3: (4 weeks)
Aerobic	Base Phase	Stamina Phase	Peaking Phase
Strength	Hypertrophy Phase	Strength Phase	Peaking Phase

Table V-26: Schedule

Week / Day	Week 1	Week 2	Week 3	Week 4
Monday	Chest, Shoulders, Triceps	Legs, Back, Biceps	Chest, Shoulders, Triceps	Legs, Back, Biceps
Tuesday	Run: Tempo	Run: Tempo	Run: Tempo	Run: Tempo
Wednesday	Legs, Back, Biceps	Chest, Shoulders, Triceps	Legs, Back, Biceps	Chest, Shoulders, Triceps
Thursday	Intervals	Intervals	Intervals	Intervals
Friday	Chest, Shoulders, Triceps	Legs, Back, Biceps	Chest, Shoulders, Triceps	Legs, Back, Biceps
Saturday	Run: Long	Run: Long	Run: Long	OFF
Sunday	OFF	OFF	OFF	OFF

Table V-27: Acute Variables: Strength

Phase	Exercises/ muscle group	Sets (MJ) per exercise	Reps (MJ)	Rest (MJ)	Sets (SJ) per exercise	Reps (SJ)	Rest (SJ)	Total Sets
Base	3 large 3 small	3	10	1	3	10-12	1	18
Strength	3 large 2 small	3	6	3	3	8	2	15
Peaking	3 large 1 small	3	6-1	4	3	6	2	12

1. MJ = Multi joint
2. SJ = single joint
3. Large muscle groups e.g. chest, back, legs
4. Small muscle groups e.g. biceps, triceps
5. Rest between sets in minutes

Table V-28: Frequency: Strength

Day	Week 1	Week 2	Week 3	Week 4
Monday	Chest, Shoulders, Triceps	Legs, Back, Biceps	Chest, Shoulders, Triceps	Legs, Back, Biceps
Tuesday	OFF	OFF	OFF	OFF
Wednesday	Legs, Back, Biceps	Chest, Shoulders, Triceps	Legs, Back, Biceps	Chest, Shoulders, Triceps
Thursday	OFF	OFF	OFF	OFF
Friday	Chest, Shoulders, Triceps	Legs, Back, Biceps	Chest, Shoulders, Triceps	Legs, Back, Biceps
Saturday	OFF	OFF	OFF	OFF
Sunday	OFF	OFF	OFF	OFF

Table V-29: Acute Variables: Cardiovascular

Type workout /Phase	Base Phase 4-8 weeks	Stamina Phase 4 weeks	Speed/Peaking Phase 4 weeks
Monday	OFF	OFF	OFF
Tuesday	Tempo Run (5) increasing from 2@80% to 4 miles @ 85% RRHR	Tempo Run (5) 4 miles @ 85% RRHR	Tempo Run (5) 4 miles @ 85% RRHR
Wednesday	OFF	OFF	OFF
Thursday	Fartlek Run increasing from 2-4: ½ mile pick ups @ 80-85%. The base running is at @ 55-65% MHR	Aerobic Capacity Intervals: 220, 440 warm ups and 880 intervals building to 6-8x880 @ 90% RRHR	Aerobic Capacity Intervals: same as stamina decreasing to 2x880 and adding anaerobic intervals building to 4 (110m) @ 95%
Friday	OFF	OFF	OFF
Saturday	Run: Aerobic Conditioning (Long) increasing from 8-10 miles @ 55-65% RRHR. Every 4 th week is OFF	Run: Aerobic Conditioning (Long) alternating 6/8 miles @ 55-65% RRHR. Every 4 th week is OFF	Run: Aerobic Conditioning (Long) alternating 6/8 miles @ 55-65% RRHR with the 4 th week off
Sunday	OFF	OFF	OFF
Total Intensity Mileage	None above 85% RRHR	Building from 8-9% the first week to 10% the last week none above 90% RRHR	Decreasing to 8%. 880's remain at 90% and 110's increase to 95% RRHR
Total Running Mileage	Increasing to 24 over 4 weeks or 30 over 8 weeks (10%/week)	18-24 miles/week depending on length of base phase	Run 16-21 miles/week.

1. RRHR = Reserve/Resting Heart Rate calculated with Karnoven Method
2. Assumes a base of 20 miles / week before starting base phase

Word Count: 21,497

ENDNOTES

¹ Newsholme, E.; Leech, T.; and Duester, G. Keep on Running: The Science of Training and Performance. West Sussex, England: John Wiley & Sons LTD, 1994: 106.

² Ibid, 64.

³ Newsholme, E.; Leech, T.; and Duester, G. Keep on Running: The Science of Training and Performance. West Sussex, England: John Wiley & Sons LTD, 1994: 63-64.

⁴ Ibid, 64.

⁵ Edwards, H.T; Margaria, R. and Dill, D.B. 1984. Metabolic rate, blood sugar and the utilization of carbohydrate. The American Journal of Physiology, 108 (1984): 70-71.

⁶ Newsholme, E.; Leech, T.; and Duester, G. Keep on Running: The Science of Training and Performance. West Sussex, England: John Wiley & Sons LTD, 1994: 72.

⁷ Costill, David L. 1986. Inside Running: Basics of Sports Physiology. Indianapolis, IN: Benchmark Press, 1986: 43.

⁸ Edwards, H.T; Margaria, R. and Dill, D.B. 1984. Metabolic rate, blood sugar and the utilization of carbohydrate. The American Journal of Physiology, 108 (1984): 209.

⁹ Ibid, 90-92.

¹⁰ Newsholme, E.; Leech, T.; and Duester, G. Keep on Running: The Science of Training and Performance. West Sussex, England: John Wiley & Sons LTD, 1994: 72.

¹¹ Ibid, 75.

¹² Ibid, 73.

¹³ McMillan, G. (editor) 1997. "How running physiology impacts on optimal training. Peak Running Performance". (Internet: <http://www.peakrun.com/articles>):2-3.

¹⁴ McMillan, G. (editor) 1997. "How running physiology impacts on optimal training. Peak Running Performance". (Internet: <http://www.peakrun.com/articles>): 2.

¹⁵ Newsholme, E.; Leech, T.; and Duester, G. Keep on Running: The Science of Training and Performance. West Sussex, England: John Wiley & Sons LTD, 1994: 83.

¹⁶ Ibid, 74.

¹⁷ McMillan, G. (editor) 1997. "How running physiology impacts on optimal training. Peak Running Performance". (Internet: <http://www.peakrun.com/articles>): 2.

¹⁸ Newsholme, E.; Leech, T.; and Duester, G. Keep on Running: The Science of Training and Performance. West Sussex, England: John Wiley & Sons LTD, 1994: 77-78.

¹⁹ Ibid, 88.

²⁰ Newsholme, E.; Leech, T.; and Duester, G. Keep on Running: The Science of Training and Performance. West Sussex, England: John Wiley & Sons LTD, 1994: 89.

²¹ Ibid, 88.

²² Robinson, J. and Carrino, F. Max O2: The Complete Guide to Synergistic Aerobic Training. Los Angeles, CA: Health for Life, 1993: 27.

²³ Fleck, S.J.; and Kraemer, W.J. Designing Resistance Training Programs. Champaign, IL: Human Kinetics Publishers, Inc., 1997:57.

²⁴ Costill, D.L.; Fink, W.J. and Pollock, M. L. 1976. "Muscle fiber composition and enzyme activities of elite distance runners." Medicine and Science in Sports, 8 no. 2(1976): 96.

²⁵ Fleck, S.J.; and Kraemer, W.J. Designing Resistance Training Programs. Champaign, IL: Human Kinetics Publishers, Inc., 1997:59.

²⁶ Gollnick, P. D.; Armstrong, R.B.; Saubert IV, C. W. Piehl, K. and Saltin, B. "Enzyme activity and fiber composition in skeletal muscle of trained and untrained men". Journal of Applied Physiology, 33 (1972): 312.

²⁷ Fleck, S.J.; and Kraemer, W.J. Designing Resistance Training Programs. Champaign, IL: Human Kinetics Publishers, Inc, 1997: 61.

²⁸ Ibid, 59.

²⁹ Staron, R. S.; Karapondo, D. L. ; Kraemer, W. J. ; Fry, A. C. ; Gordon, S. E. ; Falkel, J. E. ; Hagerman, F. C. ; and Hikida, R. S. "Skeletal muscle adaptations during the early phase of heavy resistance training in men and women." Journal of Applied Physiology, 76 (1994):1255.

³⁰ Adams, G. R.; Hather, B. M.; Baldwin, K. M.; and Dudley, G. A. "Skeletal muscle myosin heavy chain composition and resistance training." Journal of Applied Physiology, 74 (1993):915.

³¹ Fleck, S.J.; and Kraemer, W.J. Designing Resistance Training Programs. Champaign, IL: Human Kinetics Publishers, Inc. 1997: 61.

³² Kirksey, Brett. "Strength Training Breakthrough, The First International Conference on Weight Lifting and Strength Training." Muscular Development, April (1999): 96-99.

³³ Robinson, J.; and Carrino, F. Max O2: The Complete Guide to Synergistic Aerobic Training. Los Angeles, CA: Health for Life, 1993: 39.

³⁴ Costill, David L. Inside Running: Basics of Sports Physiology. Indianapolis, IN: Benchmark Press, 1986: 6.

³⁵ Ibid, 39.

³⁶ Desmedt, J. E. and Godaux, E. "Ballistic contractions in man: Characteristic recruitment pattern of single motor units of the tibialis muscle". Journal of Physiology. 264 (1977): 673-694.

³⁷ Grimby, L. and Hannerz, J. "Firing rate and recruitment order of toe extensor motor units in different modes of voluntary contraction". Journal of Physiology. (London) 264 (1977): 879.

³⁸ Fleck, S.J.; and Kraemer, W.J. Designing Resistance Training Programs. Champaign, IL: Human Kinetics Publishers, Inc. 1997: 62.

³⁹ Ibid, 59-60.

⁴⁰ Robinson, J.; and Carrino, F. Max O2: The Complete Guide to Synergistic Aerobic Training. Los Angeles, CA: Health for Life, 1993: 40.

⁴¹ Newsholme, E.; Leech, T.; and Duester, G. Keep on Running: The Science of Training and Performance. West Sussex, England: John Wiley & Sons LTD, 1994: 97.

⁴² Lacour, J. R.; Magunacelaya, A. P; and Barthelemy, J. C. et al. "The Energetics of Middle Distance Running". European Journal of Applied Physiology. 60 (1990): 38-43.

⁴³ Maffulli, N.; Capasso, G.; and Lancia, A. "Anaerobic threshold and performance in middle and long distance running". Journal of Sports Medicine & Physical Fitness. 31 (1991):332-338.

⁴⁴ McMillan, G. (editor) 1997. "How running physiology impacts on optimal training". Peak Running Performance. (Internet: <http://www.peakrun.com/articles>). p. 1.

⁴⁵ Brandon, L.J. "Physiological factors associated with middle distance running performance". Sports Medicine 19 no. 4(1995): 270.

⁴⁶ Brandon, L.J. and Boileau, R.A. "The contribution of selected variables to middle and long distance run performance". Journal of Sports Medicine and Physical Fitness. 27 (1987): 157-164.

⁴⁷ Costill, David L. Inside Running: Basics of Sports Physiology. Indianapolis, IN: Benchmark Press, 1986: 6.

⁴⁸ Robinson, J. and Carrino, F. Max O2: The Complete Guide to Synergistic Aerobic Training. Los Angeles, CA: Health for Life (1993): 68.

⁴⁹ Maffulli, N.; Capasso, G.; and Lancia, A. "Anaerobic threshold and performance in middle and long distance running". Journal of Sports Medicine & Physical Fitness. 31 (1991):332-338.

⁵⁰ American College of Sports Medicine. "The recommended quantity and quality of exercise for development and maintaining cardiorespiratory muscular fitness in healthy adults." Medicine and Science in Sports and Exercise, 30 no. 6 (1998): 975.

⁵¹ Ibid, 967.

⁵² Robinson, J. and Carrino, F. Max O2: The Complete Guide to Synergistic Aerobic Training. Los Angeles, CA: Health for Life (1993): 96-99.

⁵³ Burke, E. R. (editor). Precision Heart-rate Training. Champaign, IL: Human Kinetics Publishers, Inc., 1998: 88-90.

⁵⁴ Newsholme, E.; Leech, T.; and Duester, G. Keep on Running: The Science of Training and Performance. West Sussex, England: John Wiley & Sons LTD, 1994: 332.

⁵⁵ Lawrence, A.; and Scheid, M. Running and Racing after 35. Boston, MA: Little, Brown and Co, 1990: 165-167.

- ⁵⁶ Anderson, Owen. "A perfect week". Runner's World. 32 no. 11(1994) 32
- ⁵⁷ McMillan, G. (editor) "Spring 5K and 10K training schedules: part II". Peak Running Performance. 7 no. 2(1998): 7-10.
- ⁵⁸ Costill, David L. Inside Running: Basics of Sports Physiology. Indianapolis, IN: Benchmark Press, 1986: 95.
- ⁵⁹ Dudley, G. A.; Abraham, W. M. and Terjung, R. L. "Influence of exercise intensity and duration on biochemical adaptations in skeletal muscle". Journal of Applied Physiology, 53 (1982): 844-850.
- ⁶⁰ Glover, Bob and Schuder, Pete. Competitive Runner's Handbook. New York, NY: Penguin Books, 1983: 59.
- ⁶¹ Lawrence, A.; and Scheid, M. Running and Racing after 35. Boston, MA: Little, Brown and Co., 1990: 117
- ⁶² Anderson, O. "The Perfect Pace". Runner's World: May, 1992: 50.
- ⁶³ Collins, J. in Burfoot, A. The Perfect 30-Mile Week. Runner's World. May, 1987: 36
- ⁶⁴ Higdon, Hal. Run Fast: How to train for a 5K or 10K Race. Emmaus, PA: Rodale Press, 1992: 28-29
- ⁶⁵ Burfoot, A. "The Perfect 30-Mile Week". Runner's World. May, 1987: 40.
- ⁶⁶ Hickson, R.C.; et al. "Reduced training duration effects on aerobic power, endurance and cardiac growth". Journal of Applied Physiology. 53 (1982): 225.
- ⁶⁷ Hickson, R.C., et al. "Reduced training intensities and loss of aerobic power, endurance, and cardiac growth". Journal of Applied Physiology. 58 no. 2(1985): 492.
- ⁶⁸ Dudley, G. A.; Abraham, W. M. and Terjung, R. L. "Influence of exercise intensity and duration on biochemical adaptations in skeletal muscle." Journal of Applied Physiology, 53 (1982): 844-850.
- ⁶⁹ Kraemer, W. J. et al. "Training Responses of Plasma Beta-endorphin, Adrenocorticotropin and Cortisol." Medicine & Science in Sports & Exercise, 21 no. 2 (1989): 147.
- ⁷⁰ Ibid, 148.

⁷¹ Tabata, I., et al. "Effects of moderate-intensity endurance and high-intensity intermittent training on anaerobic capacity and VO₂max". Medicine & Science in Sports & Exercise. 28 no. 10(1996), 1327.

⁷² Robinson, J. and Carrino, F. 1993. Max O₂: The Complete Guide to Synergistic Aerobic Training. Los Angeles, CA: Health for Life, 1993: 59-61.

⁷³ McMillan, G. (editor) "Middle distance running: a proven guide". Peak Running Performance. 7 no. 6 (1998): 2.

⁷⁴ Aschmoneit, D. "The 4th important law of training: variety". Peak Running Performance. 6 no. 6 (1997): 3.

⁷⁵ American College of Sports Medicine. "The recommended quantity and quality of exercise for development and maintaining cardiorespiratory muscular fitness in healthy adults." Medicine and Science in Sports and Exercise, 30 no. 6 (1998): 977.

⁷⁶ Mikesell, K. A., and G. A. Dudley. "Influence of intense endurance training on aerobic power of competitive distance runners". Medicine & Science in Sports & Exercise. 16 no. 4 (1984): 375.

⁷⁷ Hickson, R.C.; et al. "Linear increase in aerobic power induced by strenuous program of endurance exercise". Journal of Applied Physiology. 42 (1977): 372.

⁷⁸ Phillips, S. M.; Green, H. J.; Tamopolsky, M. A.; Heigenhauser, G. F.; Hill, R. E.; and Grant, S. M. "Effects of Training Duration on Substrate Turnover and Oxidation during Exercise." Journal of Applied Physiology, 81 no. 5 (1996): 2182.

⁷⁹ Glover, Bob and Schuder, Pete. Competitive Runner's Handbook. New York, NY: Penguin Books, 1983: 75.

⁸⁰ Sparks, Ken and Kuehls, Dave. The Runners Book of Training Secrets. Emmaus, PA: Rodale Press, 1996: 12.

⁸¹ Glover, Bob and Schuder, Pete. Competitive Runner's Handbook. New York, NY: Penguin Books, 1983: 59.

⁸² Lawrence, A.; and Scheid, M. Running and Racing after 35. Boston, MA: Little, Brown and Co 1990: 117.

⁸³ Anderson, O. "The Perfect Pace". Runner's World. May, 1992: 50.

- ⁸⁴ Collins, J. in Burfoot, A. "The Perfect 30-Mile Week". Runner's World. May, 1987: 36
- ⁸⁵ Higdon, Hal. Run Fast: How to train for a 5K or 10K Race. Emmaus, PA: Rodale Press., 1992: 28-29
- ⁸⁶ Burfoot, A. "The Perfect 30-Mile Week". Runner's World. May, 1987: 40.
- ⁸⁷ Robinson, J. and Carrino, F. 1993. Max O2: The Complete Guide to Synergistic Aerobic Training. Los Angeles, CA: Health for Life, 1993: 96-99.
- ⁸⁸ Burke, E. R. (editor). Precision Heart-rate Training. Champaign, IL: Human Kinetics Publishers, Inc., 1998: 88-90.
- ⁸⁹ Newsholme, E.; Leech, T.; and Duester, G. Keep on Running: The Science of Training and Performance. West Sussex, England: John Wiley & Sons LTD, 1994: 332.
- ⁹⁰ Lawrence, A.; and Scheid, M. Running and Racing after 35. Boston, MA: Little, Brown and Co 1990: 165-167.
- ⁹¹ Anderson, Owen. A perfect week. Runner's World. 32 no. 11(1991) 32
- ⁹² McMillan, G. (editor) 1998. Spring 5K and 10K training schedules: part II. Peak Running Performance 7(2): 7-10.
- ⁹³ Ibid, 32.
- ⁹⁴ Tanaka, K.; Matsuura, Y.; Natsuzaka, A. et al. 1984. A longitudinal assessment of anaerobic threshold and distance running performance. Medicine and Science in Sports and Exercise. 16: 278-282.
- ⁹⁵ Henritze, J. ; Weltman, A. ; Schurrer, R. L. ; and Barlow, K. "Effects of Training at and Above the Lactate Threshold on the Lactate Threshold and Maximal Oxygen Uptake." European Journal of Applied Physiology, 54 (1985): 84-86
- ⁹⁶ Robinson, J.; and Carrino, F. 1993. Max O2: The Complete Guide to Synergistic Aerobic Training. Los Angeles, CA: Health for Life. p. 96-99.
- ⁹⁷ Burke, E. R. (editor). 1998. Precision Heart-rate Training. Champaign, IL: Human Kinetics Publishers, Inc. p. 88-90.
- ⁹⁸ Lawrence, A.; and Scheid, M. 1990. Running and Racing after 35. Boston, MA: Little, Brown and Co.: 165-167.

- ⁹⁹ Anderson, Owen. A perfect week. *Runner's World*. 32 no. 11(1991): 32
- ¹⁰⁰ McMillan, G. (editor) 1998. Spring 5K and 10K training schedules: part II. *Peak Running Performance* 7(2): 7-10.
- ¹⁰¹ Burke, E. R. (editor). 1998. *Precision Heart-rate Training*. Champaign, IL: Human Kinetics Publishers, Inc. p. 88-90.
- ¹⁰² Newsholme, E.; Leech, T.; and Duester, G. 1994. *Keep on Running: The Science of Training and Performance*. West Sussex, England: John Wiley & Sons LTD. p. 332.
- ¹⁰³ Lawrence, A.; and Scheid, M. 1990. *Running and Racing after 35*. Boston, MA: Little, Brown and Co.: 165-167.
- ¹⁰⁴ Anderson, Owen. "A perfect week". *Runner's World*. 32 no. 11(1991): 32
- ¹⁰⁵ McMillan, G. (editor) "Spring 5K and 10K training schedules: part II". *Peak Running Performance*. 7 no. 2 (1998): 7-10.
- ¹⁰⁶ Glover, Bob and Schuder, Pete. *Competitive Runner's Handbook*. New York, NY: Penguin Books, 1983: 59.
- ¹⁰⁷ Lawrence, A. and Scheid, M. *Running and Racing after 35*. Boston, MA: Little, Brown and Co., 1990: 117
- ¹⁰⁸ Anderson, O. "The Perfect Pace". *Runner's World*: May, 1992: 50.
- ¹⁰⁹ Collins, J. in Burfoot, A. "The Perfect 30-Mile Week". *Runner's World*. May, 1987: 36
- ¹¹⁰ Higdon, Hal. *Run Fast: How to train for a 5K or 10K Race*. Emmaus, PA: Rodale Press, 1992: 28-29
- ¹¹¹ Burfoot, A. "The Perfect 30-Mile Week". *Runner's World*. May, 1987: 40.
- ¹¹² Burke, E. R. (editor). *Precision Heart-rate Training*. Champaign, IL: Human Kinetics Publishers, Inc., 1998: 88-90.
- ¹¹³ Newsholme, E.; Leech, T.; and Duester, G. *Keep on Running: The Science of Training and Performance*. West Sussex, England: John Wiley & Sons LTD., 1994: 332.

¹¹⁴ Lawrence, A.; and Scheid, M. Running and Racing after 35. Boston, MA: Little, Brown and Co., 1990: 165-167.

¹¹⁵ Anderson, Owen. "A perfect week". Runner's World. 32 no. 11(1991): 32

¹¹⁶ McMillan, G. (editor). "Spring 5K and 10K training schedules: part II". Peak Running Performance. 7 no. 2 (1998): 7-10.

¹¹⁷ Robinson, J.; and Carrino, F. Max O2: The Complete Guide to Synergistic Aerobic Training. Los Angeles, CA: Health for Life., 1993: 96-99.

¹¹⁸ Burke, E. R. (editor). Precision Heart-rate Training. Champaign, IL: Human Kinetics Publishers, Inc., 1998: 88-90.

¹¹⁹ Newsholme, E.; Leech, T. and Duester, G. Keep on Running: The Science of Training and Performance. West Sussex, England: John Wiley & Sons LTD., 1994: 332.

¹²⁰ Lawrence, A.; and Scheid, M. Running and Racing after 35. Boston, MA: Little, Brown and Co., 1990: 165-167.

¹²¹ Anderson, Owen. "A perfect week". Runner's World. 32 no. 11(1991): 32

¹²² McMillan, G. (editor) 1998. "Spring 5K and 10K training schedules: part II". Peak Running Performance. 7 no. 2 (1998): 7-10.

¹²³ Robinson, J.; and Carrino, F. Max O2: The Complete Guide to Synergistic Aerobic Training. Los Angeles, CA: Health for Life., 1993: 96-99.

¹²⁴ Burke, E. R. (editor). Precision Heart-rate Training. Champaign, IL: Human Kinetics Publishers, Inc., 1998: 88-90.

¹²⁵ Newsholme, E.; Leech, T. and Duester, G. Keep on Running: The Science of Training and Performance. West Sussex, England: John Wiley & Sons LTD., 1994: 332.

¹²⁶ Anderson, Owen. "A perfect week". Runner's World. 32 no. 11 (1991): 32

¹²⁷ McMillan, G. (editor) "Spring 5K and 10K training schedules: part II". Peak Running Performance 7 no. 2 (1998): 7-10.

¹²⁸ Kraemer, W.J., et al. "Training responses of plasma beta-endorphin, adrenocorticotropin, and cortisol". Medicine & Science in Sports & Exercise. 21no. 2 (1989), 152.

¹²⁹ Pollock, M. L., et al. 1998. "Position Stand: The recommended quantity and quality of exercise for developing and maintaining cardiorespiratory and muscular fitness and flexibility in healthy adults". Medicine & Science in Sports & Exercise. 30 no. 6 (1998), p. 979.

¹³⁰ Lehmann, M.; Gastmann, U.; Petersen, K.G.; Bachl, N.; Seidel, A.; Khalaf, A. N.; Fischer, S.; and Keul, J. "Training-over-training; performance, and hormone levels, after a defined increase in training volume versus intensity in experienced middle and long distance runners". British Journal of Sports Medicine 26 (1992): 242.

¹³¹ Kreider, R. B.; Fry, A. C.; and O'Toole, M. L. Over-training in Sport. Champaign, IL: Human Kinetics Publishers, Inc. , 1998: 24.

¹³² Ibid, 41.

¹³³ Eyestone, E. D.; Fellingham, G.; George, J. and Fisher, G. "Effect of Water Running and Cycling on Maximum Oxygen Consumption and 2-Mile Run Performance." The American Journal of Sports Medicine, 21 no. 1 (1993): 41-44.

¹³⁴ Loy, S. F., et al. "Effects of Stair-climbing Versus Run Training on Treadmill and Track Running Performance." Medicine and Science in Sports and Exercise, 25 no. 11 (1993): 1275-1278.

¹³⁵ Peak Running Performance. "The most effective cross-training for runners." <<http://www.peakrun.com/articles/x-train.html>> (1997). Internet: Accessed 1-14-99.

¹³⁶ Fry, R.W., et al. "Over-training in athletes: An update". Sports Medicine. 12 no. 1 (1991): 32-65.

¹³⁷ Fry, R. W.; A. R. Morton, D. Keast. "Periodization and the prevention of overtraining". Canadian Journal of Sports Science. 17 (1992): 242.

¹³⁸ Ibid, 245.

¹³⁹ Fleck, S.J.; and Kraemer, W.J. 1997. Designing Resistance Training Programs. Champaign, IL: Human Kinetics Publishers, Inc., 1997: 91.

¹⁴⁰ Kraemer, W. J.; Marchitelli, S. E.; Gordon, S. E.; Harman, E.; Dziados, J. E.; Mello, R.; Frykman, P.; McCurry, D.; and Fleck, S. J. "Hormonal and Growth Factor

Responses to Heavy Resistance Exercise Protocols." Journal of Applied Physiology, 69 no. 4 (1990): 1449.

¹⁴¹ Fleck, S.J.; and Kraemer, W.J. 1997. Designing Resistance Training Programs. Champaign, IL: Human Kinetics Publishers, Inc., 1997: 92.

¹⁴² Moritani, T.; Nagata, A. and Muro, M. "Intramuscular and surface electromyogram changes during muscle fatigue". Journal of Applied Physiology, 60 (1986): 1184-1185.

¹⁴³ Moritani, T.; Nagata, A. and Muro, M. "Electromyographic manifestations of muscle fatigue". Medicine and Science in Sports and Exercise. 14 (1982): 201-202.

¹⁴⁴ Moritani, T. and deVries, H.A. "Reexamination of the relationship between the surface integrated electromyogram (IMEG) and force of isometric contraction." American Journal of Physiological Medicine. 57 (1987):276-277.

¹⁴⁵ Bompa, T.O.; and Cornacchia, L.J. Serious Strength Training. Champaign, IL: Human Kinetics Publishers, Inc., 1998: 123-124.

¹⁴⁶ Ibid, 124.

¹⁴⁷ Chandler, T. J.; Wilson, G. D. and Stone, M. H. "The effect of the squat exercise on knee stability." Medicine and Science in Sports and Exercise, 21 no. 3(1989): 299-303.

¹⁴⁸ Fleck, S.J.; and Kraemer, W.J. 1997. Designing Resistance Training Programs. Champaign, IL: Human Kinetics Publishers, Inc. p. 92.

¹⁴⁹ Health For Life. Secrets of Advanced Body-building. Los Angeles, CA: Health for Life, 1985: 26.

¹⁵⁰ Ibid, 21.

¹⁵¹ Ibid, 23-25.

¹⁵² Ibid, 16-18.

¹⁵³ Lieber, R. L. and Frieden, J. Muscle damage is not a function of muscle force, but active strain. Journal of Applied Physiology, 74 (1993): 526.

¹⁵⁴ American College of Sports Medicine. "The recommended quantity and quality of exercise for development and maintaining cardiorespiratory muscular fitness in healthy adults." Medicine and Science in Sports and Exercise, 22 (1990): 273-274.

¹⁵⁵ Kraemer, W.J., et al. Varied multiple set resistance training program produces greater gains than single set program. Medicine & Science in Sports & Exercise, 30 no. 5(1998)(5): S1096.

¹⁵⁶ Marx, J., et al.. "The effect of periodization and volume of resistance training in women". Medicine & Science in Sports & Exercise, 30 no. 5(1998): S164.

¹⁵⁷ Stowers, T.; McMillian, J.; Scala, D.; Davis, V.; Wilson, D.; and Stone, M. 1983. "The short term effects of three different strength-power training methods". National Strength and Conditioning Association Journal. 5 (1983): 27.

¹⁵⁸ Wagman, Dan, and Jose Antonio. "Growth settings: weighing the evidence between single vs. multiple sets". Muscle & Fitness. February, 1999: 72-75.

¹⁵⁹ Stone, Michael H., Steven Scott Plisk, Margaret E. Stone, Brian K. Schilling, Harold S O'Bryant, and Kyle C. Pierce. "Athletic performance development: volume load – 1 set vs. multiple sets, training velocity and training variation". Strength and Conditioning. December, 1998: 26.

¹⁶⁰ McGee, D.; Jessee, T. C.; Stone, M. H. and Blessing, D. "Leg and Hip Endurance Adaptations to Three Weight-training Programs." Journal of Applied Sports Science Research, 6 no. 2 (1992): 92-95.

¹⁶¹ Fleck, S.J.; and Kraemer, W.J. 1997. Designing Resistance Training Programs. Champaign, IL: Human Kinetics Publishers, Inc. 92.

¹⁶² Stone, Michael H., Steven Scott Plisk, Margaret E. Stone, Brian K. Schilling, Harold S O'Bryant, and Kyle C. Pierce. "Athletic performance development: volume load – 1 set vs. multiple sets, training velocity and training variation." Strength and Conditioning, December: 28. 1998.

¹⁶³ Applewhite, Roger. Personal Trainer. Golden CO: Muscle Media, 1997. 49.

¹⁶⁴ Hatfield, Frederick C. Powerlifting: A Scientific Approach. Chicago, IL: Contemporary Books, 1981. 134-5.

¹⁶⁵ Georgia State University. "Hypertrophy." Atlanta: Department of Kinesiology and Health, 1998. http://www.gsu.edu/~hpriad/kh_755/hyper.html. Internet: Accessed 9 November 1998.

¹⁶⁶ Berger, R. A. "Effect of varied weight training programs on strength." Research Quarterly, 33 (1962a): 181.

¹⁶⁷ Kraemer, W.J.; Noble, B.J.; Clark, M.J. and Culver, B.W. "Physiologic responses to heavy-resistance exercise with very short rest periods." International Journal of Sports Medicine, 8 (1987): 251.

¹⁶⁸ Kraemer, W. J.; Dziados, J. E.; Marchitelli, L.J.; Gordon, S. E.; Harman, E.A.; Mello, R.; Fleck, S. J.; Frykman, P. N.; and Triplett, N. T. "Effects of different heavy-resistance exercise protocols on plasma B-endorphin concentrations." Journal of Applied Physiology, 74 no. 1 (1993): 458.

¹⁶⁹ Kraemer, W.J.; Marchitelli, S. E.; Gordon, S. E.; Harman, E.; Dziados, J.E.; Mello, R.; Frykman, P.; McCurry, D.; and Fleck, S. J. "Hormonal and Growth Factor Responses to Heavy Resistance Exercise Protocols." Journal of Applied Physiology, 69 no. 4 (1990): 1450.

¹⁷⁰ Kraemer, W.J.; Fleck, S.J.; Dziados, J.E.; Harman, E.A.; Marchitelli, L. J.; Gordon, S. E.; Mello, R.; Frykman, P. N.; Koziris, L. P. and Triplett, N. T. "Changes in hormonal concentrations following different heavy resistance exercise protocols in women." Journal of Applied Physiology, 75 (1990): 594.

¹⁷¹ Kraemer, W.J.; Gordon, S. E.; Fleck, S.J.; Marchitelli, L.J.; Mello, R.; Dziados, J. E.; Friedl, K.; Harman, E.; Maresh, C.; and Fry, A. C. "Endogenous anabolic hormonal and growth factor responses to heavy resistance exercise in males and females." International Journal of Sports Medicine, 12 (1991): 233-4.

¹⁷² Fleck, S.J. and Kraemer, W.J. Designing Resistance Training Programs. Champaign, IL: Human Kinetics Publishers, Inc., 1997. p. 95-96.

¹⁷³ Fleck, Steve J. "Periodization Breakthrough Part I," lecture, Baltimore Maryland, at the International Summit of Personal Trainers, 5 March 1999, cited with permission of Dr. Fleck.

¹⁷⁴ Kraemer, W.J.; Marchitelli, S. E.; Gordon, S. E.; Harman, E.; Dziados, J.E.; Mello, R.; Frykman, P.; McCurry, D.; and Fleck, S. J. "Hormonal and Growth Factor Responses to Heavy Resistance Exercise Protocols." Journal of Applied Physiology, 69 no. 4 (1990): 1445-1448.

¹⁷⁵ Bompa, T. O.; and Comacchia, L. J. Serious Strength Training. Champaign, IL: Human Kinetics Publishers, Inc., 1998. 30-31.

¹⁷⁶ Ibid, 32.

- ¹⁷⁷ Ibid, 33.
- ¹⁷⁸ Kraemer, W.J.; Marchitelli, S. E.; Gordon, S. E.; Harman, E.; Dziados, J.E.; Mello, R.; Frykman, P.; McCurry, D.; and Fleck, S. J. "Hormonal and Growth Factor Responses to Heavy Resistance Exercise Protocols." Journal of Applied Physiology, 69 no. 4 (1990): 1442-1450.
- ¹⁷⁹ Kraemer, W.J; Fleck, S.J.; Dziados, J.E.; Harman. E.A.; Marchitelli, L.J.; Gordon, S. E.; Mello, R.; Frykman, P. N.; Koziris, L. P. and Triplett, N. T. "Changes in hormonal concentrations following different heavy resistance exercise protocols in women." Journal of Applied Physiology, 75 (1990): 594-604.
- ¹⁸⁰ Kraemer, W. J.; Gordon, S. E.; Fleck, S. J.; Marchitelli, L. J.; Mello, R.; Dziados, J. E.; Friedl, K.; Harman, E.; Maresh, C.; and Fry, A. C. "Endogenous anabolic hormonal and growth factor responses to heavy resistance exercise in males and females. International Journal of Sports Medicine, 12 (1991): 228-235.
- ¹⁸¹ Kraemer, W.J.; Noble, B.J.; Clark, M.J. and Culver, B.W. "Physiologic responses to heavy-resistance exercise with very short rest periods." International Journal of Sports Medicine, 8 (1987): 247-252.
- ¹⁸² Bompa, T. O.; and Cornacchia, L. J. Serious Strength Training. Champaign, IL: Human Kinetics Publishers, Inc., 1998. 30-31.
- ¹⁸³ Health For Life. 1985. Secrets of Advanced Body-building. Los Angeles, CA: Health for Life. 16-19.
- ¹⁸⁴ Berger, R. A.; "Optimum Repetitions for the Development of Strength." Research Quarterly, 33 no. 2 (1962):334-338.
- ¹⁸⁵ Fleck, S.J. and Kraemer, W.J. Designing Resistance Training Programs. Champaign, IL: Human Kinetics Publishers, Inc., 1997. 101.
- ¹⁸⁶ Hatfield, Frederick C. Powerlifting: A Scientific Approach. Chicago, IL: Contemporary Books, 1981. 25.
- ¹⁸⁷ Georgia State University. "Hypertrophy." Atlanta: Department of Kinesiology and Health, 1998. http://www.gsu.edu/~hprjad/kh_755/hyper.html. Internet: Accessed 9 November 1998.

- ¹⁸⁸ Georgia State University. "Maximum Strength." Atlanta: Department of Kinesiology and Health, 1998. http://www.gsu.edu/~hpriaj/kh_755/max_str.html Internet: Accessed 9 November, 1998.
- ¹⁸⁹ Bompa, T. O.; and Cornacchia, L. J. Serious Strength Training. Champaign, IL: Human Kinetics Publishers, Inc., 1998. 29.
- ¹⁹⁰ Wendeln, T., et al. "Effects of three different resistance training protocols on muscular strength and endurance." Medicine & Science in Sports & Exercise, 30 no. 5 (1998): S941.
- ¹⁹¹ Berger, R. A. "Effect of varied weight training programs on strength." Research Quarterly, 33 (1962a):168.
- ¹⁹² Stone, M. H. ; Butterfield, G.; Kleiner, S. and Lemon, Methods of Weight Gain in Athletes" Gatoraide Sports Science Institute Reference Desk. Round Table 21 (5), no. 1 (1994). <http://www.gssiweb.com/references.htm>. Internet. Accessed 17 September 1998.
- ¹⁹³ Gilliam, G. M. "Effects of frequency of weight training on muscle strength enhancement." Journal of Sports Medicine, 21(1981):434.
- ¹⁹⁴ Hoffman, J.R.; Kraemer, W.J.; Fry, A. C.; Deschenes, M. and Kemp, M. "The effects of self selection for frequency of training in a winter conditioning program for football." Journal of Applied Sport Science Research, 4 (1990):76.
- ¹⁹⁵ Gilliam, G. M. "Effects of frequency of weight training on muscle strength enhancement." Journal of Sports Medicine, 21(1981):432.
- ¹⁹⁶ Hoffman, J.R.; Kraemer, W.J.; Fry, A. C.; Deschenes, M. and Kemp, M. "The effects of self selection for frequency of training in a winter conditioning program for football." Journal of Applied Sport Science Research, 4 (1990): 77.
- ¹⁹⁷ Fleck, S.J. and Kraemer, W.J. Designing Resistance Training Programs. Champaign, IL: Human Kinetics Publishers, Inc., 1997. 241-243.
- ¹⁹⁸ Georgia State University. "Hypertrophy." Atlanta: Department of Kinesiology and Health, 1998. http://www.gsu.edu/~hpriaj/kh_755/hyper.html. Internet: Accessed 9 November 1998.
- ¹⁹⁹ Georgia State University. "Maximum Strength." Atlanta: Department of Kinesiology and Health, 1998. http://www.gsu.edu/~hpriaj/kh_755/max_str.html Internet: Accessed 9 November, 1998.

²⁰⁰ Tesch, Per A., "Training for Bodybuilding" in Strength and Power in Sport: Volume III of the Encyclopedia of Sports Medicine and International Olympic Committee Medical Commission Publication. ed. P. V. Komi. Blackwell Scientific Publications, Oxford, 1992, 370-380.

²⁰¹ Hakkinen, K.; Parkarinen, A.; Alen, M.; Kauhanen, H. and Komi, P. V. "Neuromuscular and Hormonal Adaptations in Athletes to Strength Training in Two Years. Journal of Applied Physiology, 65 (1988b): 2408.

²⁰² Fleck, Steve J. "Periodization Breakthrough Part I," lecture, Baltimore Maryland, at the International Summit of Personal Trainers, 5 March 1999, cited with permission of Dr. Fleck.

²⁰³ Hakkinen, K. "Factors Influencing Trainability of Muscular Strength During Short Term and Prolonged Exercise." National Strength and Conditioning Association Journal, 7 (1985): 36-37.

²⁰⁴ Fleck, Steve J. "Periodization Breakthrough Part I," lecture, Baltimore Maryland, at the International Summit of Personal Trainers, 5 March 1999, cited with permission of Dr. Fleck.

²⁰⁵ Moritani, C. D. and DeVries, H. A. "Neural Factors vs. Hypertrophy in Time Course of Muscle Strength Gain." American Journal of Physical Medicine and Rehabilitation, 58 (1979): 128-130.

²⁰⁶ Fleck, Steve J. "Periodization Breakthrough Part I," lecture, Baltimore Maryland, at the International Summit of Personal Trainers, 5 March 1999, cited with permission of Dr. Fleck.

²⁰⁷ Leighton, J. R.; Holmes, D.; Benson, J.; Wooten, B.; and Schmerer, R. "A study of the effectiveness of ten different methods of progressive resistance exercise on the development of strength, flexibility, girth and body weight." Journal of Physical and Mental Rehabilitation, 21 (1967): 78-79.

²⁰⁸ Ibid, 81.

²⁰⁹ Willoughby, D. S. "The effects of mesocycle-length weight training programs involving periodization and partially equated volumes on upper and lower body strength." Journal of Strength and Conditioning Research, 7(1993): 3-4.

²¹⁰ Ibid, 4.

²¹¹ Ibid, 2.

²¹² Ibid, 6.

²¹³ Marx, J. O.; Kraemer, W.J.; Nindl, B. C.; Gotshalk, L. A.; Duncan, N. D. ; Volck, J. S.; Hakkinen, K. ; Newton, R. U. "The effect of periodization and volume of resistance training in women. " Medicine & Science in Sports & Exercise, 30 no.5 (1998): S935.

²¹⁴ Stowers, T.; McMillian, J.; Scala, D.; Davis, V.; Wilson, D. and Stone, M. "The short term effects of three different strength-power training methods.: National Strength and Conditioning Association Journal, 5 (1983): 24-27.

²¹⁵ O'Bryant, H. S.; Byrd, R.; and Stone, M. H. "Cycle ergometry performance and maximum leg and hip strength adaptations to two different methods of weight training." Journal of Applied Sport Science Research, 2 no. 2 (1988): 27-30.

²¹⁶ Willoughby, D. S. "The effects of mesocycle-length weight training programs involving periodization and partially equated volumes on upper and lower body strength." Journal of Strength and Conditioning Research, 7(1993): 2-8.

²¹⁷ Hoffman, J.R.; Kraemer, W.J.; Fry, A. C.; Deschenes, M. and Kemp, M. "The effects of self selection for frequency of training in a winter conditioning program for football." Journal of Applied Sport Science Research, 4 (1990): 76-82.

²¹⁸ Fleck, S. J. and Kraemer, W. J. Periodization Break-Through. Ronkonkoma, NY: Advanced Research Press, Inc., 1996. 52.

²¹⁹ Kreider, R.B.; Fry, A. C.; and O'Toole, M. L. Over-training in Sport. Champaign, IL: Human Kinetics Publishers, Inc., 1998. 10.

²²⁰ Callister, R., et al. "Physiological and Performance Responses to Overtraining in Elite Judo Athletes." Medicine & Science in Sports & Exercise, 22 no. 6 (1990): 823-824.

²²¹ Kreider, R.B.; Fry, A. C.; and O'Toole, M. L. Over-training in Sport. Champaign, IL: Human Kinetics Publishers, Inc., 1998. 77.

²²² Fry, A. C.; Kraemer, W. J.; Lynch, J. M.; Triplett, H. T. and Koziris, L. P. "Does short-term near-maximal intensity machine resistance training induce over-training?" Journal of Strength and Conditioning, 8 (1994): 190-191.

²²³ Fry, R.W., et al. "Over-Training in Athletes: An Update." Sports Medicine, 12 no. 1(1994): 62-65.

²²⁴ Hickson, Robert C., "Interference of strength development by simultaneously training for strength and endurance." European Journal of Applied Physiology. 45 (1980): 255.

²²⁵ Ibid., 255-258.

²²⁶ Dudley, G. A., and R. Djmil. "Incompatibility of endurance and strength training modes of exercise." Journal of Applied Physiology. 59 no. 5. (1985): 1446-1451.

²²⁷ Hunter, G.; Demment, R. and Miller, D. "Development of strength and maximum oxygen uptake during simultaneous training for strength and endurance." Journal of Sports Medicine and Physical Fitness, 27no. 3. (1987): 270.

²²⁸ Ibid., 273-4.

²²⁹ McCarthy, J. P., et al. "Compatibility of adaptive responses with combining strength and endurance training." Medicine & Science in Sports & Exercise, 27 no. 3. (1995): 429-430.

²³⁰ Ibid., 433-34.

²³¹ Volpe, S. L.; Walberg-Rankin, J.; Rodman, K. W.; and Sebolt, D. R. "The effect of endurance running on training adaptations in women participating in a weight lifting program." Journal of Strength and Conditioning Research, 7 no.2. (1993): 101-107.

²³² Kraemer, W. J., et al. "Compatibility of high-intensity strength and endurance training on hormonal and skeletal muscle adaptations" Journal of Applied Physiology, 78 no. 3. (1995): 977-78.

²³³ Hickson, R.C. et al. "Reduced Training Duration Effects on Aerobic Power, Endurance and Cardiac Growth." Journal of Applied Physiology, 53 (1982): 225-226.

²³⁴ Ibid, 227-228.

²³⁵ Hickson, R.C., et al. "Reduced Training Intensities and Loss of Aerobic Power, Endurance, and Cardiac Growth." Journal of Applied Physiology, 58 no. 2 (1985): 492-493.

²³⁶ Ibid, 494-495.

²³⁷ Dudley, G. A.; Abraham, W. M. and Terjung, R. L. "Influence of exercise intensity and duration on biochemical adaptations in skeletal muscle." Journal of Applied Physiology, 53 (1982): 844-850.

²³⁸ McCarthy, J. P., et al. "Compatibility of adaptive responses with combining strength and endurance training." Medicine & Science in Sports & Exercise, 27 no. 3. (1995): 429-430.

²³⁹ Hunter, G.; Demment, R. and Miller, D. "Development of strength and maximum oxygen uptake during simultaneous training for strength and endurance." Journal of Sports Medicine and Physical Fitness, 27no. 3. (1987): 270.

BIBLIOGRAPHY

1. PERIODICALS: STRENGTH TRAINING STUDIES

- Adams, G. R.; Hather, B. M.; Baldwin, K. M.; and Dudley, G. A. "Skeletal muscle myosin heavy chain composition and resistance training." Journal of Applied Physiology, 74 (1993): 911-915.
- Baker, M.; Wilson, G.; and Carlyon, R. Periodization: "The effect on strength of manipulating volume and intensity." Journal of Strength and Conditioning Research 8 (1994): 235-242.
- Bauer, T.; Thayer, R.E. and Baras, G. "Comparison of training modalities for power development in the lower extremity." Journal of Applied Sport Science Research, 4 (1990): 115-121.
- Berger, R.A. "Comparative effects of three weight training programs." Research Quarterly, 34 (1962b): 396-398.
- Berger, R.A.; and Hardage, B. "Effect of maximum loads for each of ten repetitions on strength improvement." Research Quarterly, 38 (1967): 715-718.
- Berger, R. A. "Effect of varied weight training programs on strength." Research Quarterly, 33 (1962a): 168-181.
- Berger, R.A. "Optimum repetitions for the development of strength." Research Quarterly, 34 (1962b): 334-338.
- Boyer, B.T. "A comparison of the effects of three strength training programs on women." Journal of Applied Sport Science Research, 4 (1990): 88-94.
- Borst, S., et al. "Six months of high-or-low volume resistance training increases circulating insulin-like growth factor." Medicine & Science in Sports & Exercise, 30 no. 5 (1998): S1556.
- Brown, E.; Lillegard, W.; Henderson, R.; Wilson, D.; Lewis, E.; Hough, D. and Stringer, K. "Efficacy and safety of strength training with free weights in pre-pubescent to early post-pubescent (Abstract)." Medicine and Science in Sports and Exercise, 24 (1992): S82.
- Butterfield, G., S. Kleiner, P. Lemon, and M. Stone. "Roundtable: Methods of Weight Gain in Athletes." Gatorade Sports Science Institute Reference Desk. RT # 21 (5): No. 1 (1994). <http://www.gssiweb.com/references.htm>. Internet. Accessed 17 September 1998.

- Campos, G.R.; Luecke, T.; Toma, K.; Hagerman, F.C.; Hikida, R. S.; Ragg, V.; Kraemer, W.J. and Staron, R. S. "The effects of three different resistance training regimens on muscle fiber type composition." Medicine & Science in Sports & Exercise, 30 no. 5, (1998): S1176.
- Chandler, T.J., et al. "The effect of the squat exercise on knee stability." Medicine & Science in Sports & Exercise, 21 no. 3 (1989): 299-303.
- Crill, M.; Campos, G.; Staron, R. S.; Hagerman, F.C.; Luecke, T.; Bailey, T.; Kraemer, W.J.; and Hikida, R. S. "Different resistance training protocols and skeletal muscle fiber hypertrophy." Medicine & Science in Sports & Exercise, 30 no.5 (1998): S1180.
- Cureton, K. J.; Collins, M. A.; Hill, D.W. and McElhannon, F. M. "Muscle Hypertrophy in men and women." Medicine and Science in Sports and Exercise, 20 (1988): 338-344.
- DeHoyos, D., et al. "Effects of 6 months of high or low-volume resistance training on muscular strength and endurance." Medicine & Science in Sports & Exercise, 30 no. 5 (1998): S165.
- Desmedt, J.E. and Godaux, E. "Ballistic contractions in man: Characteristic recruitment pattern of single motor units of the tibialis muscle." Journal of Physiology, 264 (1977): 673-694.
- Feigenbaum, M. S. and Pollock, M. L. "Prescription of resistance training for health and disease." Medicine and Science in Sports and Exercise, 31 no. 1 (1999): 38-45.
- Fleck, S. J. and Kraemer, W. J. "Resistance Training: Basic Principles (Part 1 of 4)." The Physician and Sports Medicine, 16 no 3 (1988). 160-171.
- Fleck, S. J. and Kraemer, W. J. "Resistance Training: Physiological Responses and Adaptations (Part 2 of 4)." The Physician and Sports Medicine, 16 no 4 (1988). 108-124.
- Fleck, S. J. and Kraemer, W. J. "Resistance Training: Physiological Responses and Adaptations (Part 3 of 4)." The Physician and Sports Medicine, 16 no 5 (1988). 63-74.
- Fleck, S. J. and Kraemer, W. J. "Resistance Training: Exercise Prescription (Part 4 of 4)." The Physician and Sports Medicine, 16 no 3 (1988). 69-81.
- Ford, H. T.; Puckett, J. R.; Drummond, J .P.; Sawyer, K.; Gantt ,K. and Fussell, C. "Effects of Three Combinations of Plyometric and Weight Training Programs on Selected Fitness Test Items." Perceptual and Motor Skills 56 (1983): 919-922.

- Garfinkel, S.; and Cafarelli, E. "Relative changes in maximal force, EMG, and muscle cross sectional area after isometric training." Medicine and Science in Sport and Exercise, 24 (1992): 1220-1227.
- Garhammer, J.; and McLaughlin, T. "Power output as a function of load variation in Olympic and power lifting." Journal of Biomechanics, 3(1980):1980.
- Georgia State University. "Hypertrophy." Atlanta: Department of Kinesiology and Health, 1998. http://www.gsu.edu/~hprjad/kh_755/hyper.html. Internet: Accessed 9 November, 1998.
- Georgia State University. "Maximum Strength." Atlanta: Department of Kinesiology and Health, 1998. http://www.gsu.edu/~hprjad/kh_755/max_str.html Internet: Accessed 9 November, 1998.
- Gilliam, G. M. "Effects of frequency of weight training on muscle strength enhancement." Journal of Sports Medicine, 21(1981): 432-436.
- Gordon, S. E.; Kraemer, W. J.; Vos, N. H.; Lynch, J. M. and Knuttgen, H. G. "Effect of acid-base balance on the growth hormone response to acute high-intensity cycle exercise." Journal of Applied Physiology, 76 (1994): 821-829.
- Grimby, L. and Hannerz, J. "Firing rate and recruitment order of toe extensor motor units in different modes of voluntary contraction." Journal of Physiology (London), 264 (1977): 867-879.
- Hakkinen, K. "Factors Influencing Trainability of Muscular Strength During Short Term and Prolonged Exercise." National Strength and Conditioning Association Journal, 7 (1985): 32-37.
- Hakkinen, K. and A. Pakarinen. "Acute hormonal responses to two different fatiguing heavy-resistance protocols in male athletes." Journal of Applied Physiology, 74 no. 2 (1993): 882-887.
- Hakkinen, K. "Neuromuscular responses in male and female athletes to two successive strength training sessions in one day." Journal of Sports Medicine and Physical Fitness, 32 (1992): 234-242.
- Hakkinen, K., and Pakarinen, A. "Acute hormonal responses to heavy resistance exercise in men and women at different ages." International Journal of Sports Medicine, 16 (1995): 507-513.
- Hakkinen, K.; Pakarinen, A.; Allen, M.; Kauhanen, H. and Komi, P. V. "Relationships between training volume, physical performance capacity, and serum hormone

concentration during prolonged training in elite weight lifters." International Journal of Sports Medicine, 8 (1987): 61-65.

Hakkinen, K.; Parkarinen, A.; Alen, M.; Kauhanen, H. and Komi, P.V. "Neuromuscular and Hormonal Adaptations in Athletes to Strength Training in Two Years." Journal of Applied Physiology, 65 (1988b): 2406-2412.

Hoeger, W.W.K.; Hopkins, D. R.; Barette, S. L. and Hale, D.F. "Relationship between repetitions and selected percentages of one repetition maximum." Journal of Applied Sport Science Research, 4 (1990): 47-54.

Hortobagyi, T., et al. "Acute strength gains are greater with eccentric overload compared to standard resistance training in young and old women." Medicine & Science in Sports & Exercise, 30 no. 5 (1998): S1998.

Hostler, D., et al. "Effectiveness of 0.5 lb increments in progressive resistance exercise." Medicine & Science in Sports & Exercise. 30 no. 5 (1998): S933.

Housh, D.J., et al. "Hypertrophic response to unilateral concentric isokinetic resistance training." Journal of Applied Physiology. 73 no. 1 (1992): 65-70.

Hunter, G. R. "Changes in body composition, body build and performance associated with different weight training frequencies in males and females." National Strength and Conditioning Association Journal, 7 (1985): 26-28.

Jones, D. A. and Rutherford, O. M. "Human muscle strength training: The effects of three different regimes and the nature of the resultant changes." Journal of Physiology 391 (1987): 1-11.

Kirksey, Brett. "Strength Training Breakthrough, The First International Conference on Weight Lifting and Strength Training." Muscular Development, April (1999): 96-99.

Kraemer, W.J., et al. "Varied multiple set resistance training program produces greater gains than single set program." Medicine & Science in Sports & Exercise, 30 no. 5 (1998): S1096.

Kraemer, W. J.; Dziados, J. E.; Marchitelli, L.J.; Gordon, S. E.; Harman, E.A.; Mello, R.; Fleck, S. J.; Frykman, P. N.; and Triplett, N. T. "Effects of different heavy-resistance exercise protocols on plasma B-endorphin concentrations." Journal of Applied Physiology, 74 no. 1 (1993): 450-459.

Kraemer, W.J.; Marchitelli, S. E.; Gordon, S. E.; Harman, E.; Dziados, J.E.; Mello, R.; Frykman, P.; McCurry, D.; and Fleck, S. J. "Hormonal and Growth Factor Responses to Heavy Resistance Exercise Protocols." Journal of Applied Physiology, 69 no. 4 (1990): 1442-1450.

- Kraemer, W.J.; Fleck, S.J.; Dziados, J.E.; Harman, E.A.; Marchitelli, L.J.; Gordon, S. E.; Mello, R.; Frykman, P.N.; Koziris, L. P. and Triplett, N. T. "Changes in hormonal concentrations following different heavy resistance exercise protocols in women." Journal of Applied Physiology, 75 (1990): 594-604.
- Kraemer, W.J.; Gordon, S. E.; Fleck, S.J.; Marchitelli, L.J.; Mello, R.; Dziados, J. E.; Friedl, K.; Harman, E.; Maresh, C.; and Fry, A. C. "Endogenous anabolic hormonal and growth factor responses to heavy resistance exercise in males and females." International Journal of Sports Medicine, 12 (1991): 228-235.
- Kraemer, W.J.; Noble, B.J.; Clark, M.J. and Culver, B.W. "Physiologic responses to heavy-resistance exercise with very short rest periods." International Journal of Sports Medicine, 8 (1987): 247-252.
- Koch, A.J., et al. "Effects of multiple split exercise sessions on post exercise O₂ uptake and resting metabolic rate." Medicine & Science in Sports & Exercise, 30 no. 5 (1998): S1339.
- Kulling, F. A., et al. "The effect of two different rest periods between sets in a resistance training program." Medicine & Science in Sports & Exercise, 30 no. 5 (1998): S939.
- La Chance, P.F. "Effects of exercise cadence on performance design implications for resistance training programs." Medicine & Science in Sports & Exercise. 22 no. 2 (1990): S110.
- Leighton, J. R.; Holmes, D.; Benson, J.; Wooten, B.; and Schmerer, R. "A study of the effectiveness of ten different methods of progressive resistance exercise on the development of strength, flexibility, girth and body weight." Journal of Physical and Mental Rehabilitation, 21 (1967): 78-81.
- Lieber, R.L. and Frieden, J. "Muscle damage is not a function of muscle force, but active strain." Journal of Applied Physiology, 74 (1993): 520-526.
- Li, Y., et al. "Neuromuscular adaptations to lumbar extension strength gain." Medicine & Science in Sports & Exercise, 30 no. 5(1998): S1177.
- Marx, J., et al. "The effect of periodization and volume of resistance training in women." Medicine & Science in Sports & Exercise, 30 no. 5 (1998): S164.
- Marx, J. O.; Kraemer, W.J.; Nindl, B. C.; Gotshalk, L. A.; Duncan, N. D.; Volck, J. S.; Hakkinen, K.; Newton, R. U. "The effect of periodization and volume of resistance training in women." Medicine & Science in Sports & Exercise, 30 no.5 (1998): S935.
- McGee, D.; Jessee, T. C.; Stone, M. H. and Blessing, D. "Leg and Hip Endurance Adaptations to three Weight Training Programs." Journal of Applied Sports Science Research, 6 (1992): 92-95.

- Moritani, T.; Muro, M.; Kijima, A.; Gaffney, F.A. and Parsons, D. "Electro-mechanical changes during electrically induced and maximal voluntary contractions: Surface and intramuscular EMG responses during sustained maximal voluntary contractions." American Journal of Physiological Medicine, 58 (1985): 115-130.
- Moritani, C. D. and deVries, H. A. "Neural Factors vs. Hypertrophy in Time Course of Muscle Strength Gain." American Journal of Physical Medicine and Rehabilitation, 58 (1979): 115-130.
- Moritani, T. and deVries, H. A. "Rexamination of the relationship between the surface integrated electromyogram (IMEG) and force of isometric contraction." American Journal of Physiological Medicine, 57 (1987): 263-277.
- Moritani, T.; Nagata, A. and Muro, M. "Intramuscular and surface electromyogram changes during muscle fatigue." Journal of Applied Physiology, 60 (1986): 1179-1185.
- Moritani, T.; Nagata, A. and Muro, M. "Electromyographic manifestations of muscle fatigue." Medicine and Science in Sports and Exercise, 14 (1982): 198-202.
- Nindl, B., et al. "Growth hormone, testosterone, strength and hypertrophy changes in women after 6 months of periodized resistance training." Medicine & Science in Sports & Exercise, 30 no. 5 (1998): S49.
- Norenberg, K. M., et al. "Effects of resistance exercise on contractile properties of single muscle fiber in rats." Medicine & Science in Sports & Exercise, 30 no. 5 (1998): S1181.
- Ohmori, H., et al. "Very low-frequency strength training causes augmentation of muscle strength in case of untrained subjects." Medicine & Science in Sports & Exercise, 30 no. 5 (1998): S1175.
- O'Shea, P. "Effects of selected weight-training programs on the development of strength and muscle hypertrophy." Research Quarterly, 37 (1966): 95-102.
- Patton, J.F.; Kraemer, W.J.; Knuttgen, H.G. and Harman, E.A. "Factors in maximal power production and in exercise endurance relative to maximal power." European Journal of Applied Physiology, 60 (1990): 222-227.
- Ryan, Alice S., et al. "Resistance training increases fat-free mass and maintains RMR despite weight loss in post-menopausal women." Journal of Applied Physiology, 79 no. 3 (1995): 818-823.
- Staron, R. S.; Karapondo, D. L.; Kraemer, W.J.; Fry, A.C.; Gordon, S. E.; Falkel, J.E.; Hagerman, F.C.; and Hikida, R. S. "Skeletal muscle adaptations during the early

phase of heavy resistance training in men and women." Journal of Applied Physiology 76 (1994): 1247-55.

Starkey, D. B., et al. "Effect of resistance training volume on strength and muscle thickness." Medicine & Science in Sports & Exercise. 28 no. 10 (1996): 1311-1320.

Stelzer, J.E., et al. "Effects of resistance training on fat-free mass and resting metabolic rate in young men." Medicine & Science in Sports & Exercise. 30 no. 5 (1998): S1039.

Stone, M. H.; Plisk S. S; Stone, M. E.; Schilling, B.K.; O'Bryant, H. S. and Pierce, K.C. "Athletic performance development: volume load – 1 set vs. multiple sets, training velocity and training variation." Strength and Conditioning, December (1998): 22-31.

Stone, M. H.; Butterfield, G.; Kleiner, S. and Lemon, Methods of Weight Gain in Athletes" Gatoraide Sports Science Institute Reference Desk. Round Table 21 (5), no. 1 (1994). <http://www.gssiweb.com/references.htm>. Internet. Accessed 17 September 1998.

Stowers, T.; McMillian, J.; Scala, D.; Davis, V.; Wilson, D. and Stone, M. "The short term effects of three different strength-power training methods.: National Strength and Conditioning Association Journal, 5 (1983): 24-27.

Stussi, K., et al. "Effect of non-exhaustive vs. exhaustive strength training on maximum strength." Medicine & Science in Sports & Exercise, 30 no. 5 (1998): S214.

Takase, K., et al. "Effects of eccentric and concentric muscle strength training in long distance runners." Medicine & Science in Sports & Exercise, 30 no. 5 (1998): S936.

Tarnopolsky, M. A.; MacDougall, J. D. and Atkinson, S.A. "Influence of protein intake and training status on nitrogen balance and lean body mass." Journal of Applied Physiology, 64 (1988): 187-193.

Tokeshi, S.A.; Kraemer, W.J. ; Nindl, B. C. ; Gotshalk, L. A. ; Marx, J. O. ; Harman, F. S. ; Lamont, H. S. ; Sebastianelli, W.J. Putukian, ; M. ; Fleck, S. ; Newton, R.U. and Hakkinen, K.. "Power and strength in women: Adaptations following six months of resistance training." Medicine & Science in Sports & Exercise, 30 no. 5 (1998): S937.

Touey, P. R.; Sforzo, G. A.; and McManis, B.G. "Effects of manipulating rest periods on isokinetic muscle performance." Medicine and Science in Sports and Exercise, 26 no. 5 (1994): S170.

- Tucci, J., et al. "Effect of reduced training frequency and detraining on lumbar extension strength." Medicine & Science in Sports & Exercise, 22 no. 2 (1990): S105.
- Wagman, Dan, and Jose Antonio. "Growth settings: weighing the evidence between single vs. multiple sets." Muscle & Fitness, February (1999): 72-75.
- Waldman, R., and Stull, G. "Effects of various periods of inactivity on retention of newly acquired levels of muscular endurance." Research Quarterly, 40 (1969): 393-401.
- Weiss, L., et al. "Comparative effects of deep vs. shallow periodized squat training by novice lifters." Medicine & Science in Sports & Exercise, 30 no. 5(1998): S942.
- Wendeln, T., et al. "Effects of three different resistance training protocols on muscular strength and endurance." Medicine & Science in Sports & Exercise, 30 no. 5 (1998): S941.
- Wentworth, M.C., et al. "Prediction of 1-RM strength from a 5-10 RM submaximal strength test in college age females." Medicine & Science in Sports & Exercise, 30 no. 5 (1998): S1221.
- Willoughby, D. S. "The effects of mesocycle-length weight training programs involving periodization and partially equated volumes on upper and lower body strength." Journal of Strength and Conditioning Research, 7(1993): 2-8.
- Wilson, G.J.; Newton, R.U.; Murphy, A.J.; and Humphries, B.J. "The optimal training load for the development of dynamic athletic performance." Medicine and Science in Sports and Exercise, 25 (1993):1279-1286.
- Winett, Richard A., "The most effective approach to lower body training." Master Trainer May (1995). < <http://ageless-athletes.com/aerobics.htm>>; Internet. Accessed 20 September, 1998.

2. PERIODICALS: CARDIOVASCULAR TRAINING STUDIES

- Anderson, Owen. "A Perfect Week". Runner's World, 32 no. 11 (1993): 32-34.
- Aschmoneit, D. "The 4th Important Law of Training: Variety". Peak Running Performance, 6 no. 6 (1997): 1-5.
- Aschmoneit, D. "The 3rd Important Law of Training: Gradual Adaptation." Peak Running Performance, 6 no. 5 (1997): 1-4.
- Avery, G. "How to Perfectly Peak for a Great 5k or 10K." Peak Running Performance, 6 no. 4 (1997): 1-8.

- Avery, G. "Developing Your Stamina for Faster 5k Or 10K Racing." Peak Running Performance, 6 no, 3 (1997): 1-7.
- Avery, G. "Training by phases: building your conditioning base for the 5K and 10K." Peak Running Performance, 6 no. 1 (1997): 1-7.
- Billat, V.L.; Flechet, B.; Petit, B.; Muriaux, G. and Koralsztein, J. P. "Interval Training At VO2max: Effects On Aerobic Performance And Over-Training Markers." Medicine and Science in Sports and Exercise, 31 no. 1 (1999): 56-163.
- Brandon, L.J. "Physiological Factors Associated with Middle Distance Running Performance." Sports Medicine, 19 no. 4 (1995): 268-277.
- Brandon, L.J. and Boileau, R.A. "The contribution of selected variables to middle and long distance run performance." Journal of Sports Medicine and Physical Fitness, 27 (1987): 157-164.
- Burke, E.; Daniels, J.; Coyle, E. and Maughan, R. "Are Heart Rate Monitors Worth The Bother?" Gatoraide Sports Science Institute Reference Desk. Round Table 16 (5), no. 2 (1994). <http://www.gssiweb.com/references.htm>. Internet. Accessed 17 September 1998.
- Burke, E.J. and Franks, B. D. "Changes In VO2 Max Resulting From Bicycle Training at Different Intensities Holding Total Mechanical Work Constant." Research Quarterly, 46 (1975): 31.
- Burke, J.; Thayer, R. and Belcamino, M. "Comparison of Effects of Two Interval Training Programmes on Lactate and Ventilatory Thresholds." British Journal of Sports Medicine, 28 (1994): 18-21.
- Burfoot, A. "The Perfect 30-Mile Week." Runner's World, May (1987): 33-40.
- Callister, R.; Shealy, M.J.; Fleck, S.J. and Dudley, G. A. "Performance Adaptations to Sprint, Endurance and Both Modes of Training". Journal of Applied Physiology, 2 (1988): 46-51.
- Conley, D. L.; Krahenbuhl, G S. and Burkett, L.N. "Training for aerobic capacity and running economy." Physician and Sports Medicine, 9 no. 4 (1981): 107-114.
- Costill, D. L.; Fink, W.J. and Pollock, M. L. "Muscle Fiber Composition and Enzyme Activities of Elite Distance Runners." Medicine and Science in Sports, 8 no. 2 (1976): 96-103.
- Costill, D.J., et al. "Effects of Repeated Days of Intensified Training on Muscle Glycogen and Swimming Performance." Medicine & Science in Sports & Exercise, 20 no. 3 (1988): 249-254.

- Costill, R.; Thomas, R.; Robergs, R.; Pascoe, D.; Lambert, C.; Barr, S.; and Fink, W. J. "Adaptations to Swimming Training: Influence of Training Volume." Medicine and Science in Sports and Exercise, 23 no. 3 (1991): 371-377.
- Coyle, Edward F. "Fat metabolism during exercise: new concepts." Gatorade Sports Science Institute Reference Desk. SSE # 59 (8) no. 6, (1995).
<http://www.gssiweb.com/references.htm>. Internet. Accessed 17 September 1998.
- Coyle, E.F.; Martin, W.H.; Sinacore, D. R.; Joyner, M.J.; Hagberg, J.M., and Holloszy, J. O. "Time Course of Loss of Adaptations After Stopping Prolonged Intense Endurance Training." Journal of Applied Physiology, 57 no. 6 (1984): 1857-1864.
- Davis, James A. "Anaerobic Threshold: Review of The Concept And Directions For Future Research." Medicine & Science in Sports & Exercise, 17 no. 1 (1985): 6-18.
- Davis, J. A. et al. "Anaerobic Threshold and Maximal Aerobic Power for Three Modes of Exercise." Journal of Applied Physiology, 41 (1976):544.
- Despres, J. P., et al. "Effects of Aerobic Training on Fat Distribution in Male Subjects." Medicine & Science in Sports & Exercise, 17 no. 1 (1985): 113-118.
- Dudley, G. A.; Abraham, W. M. and Terjung, R. L. "Influence of exercise intensity and duration on biochemical adaptations in skeletal muscle." Journal of Applied Physiology, 53 (1982): 844-850.
- Eddy, D. O. et al. "The Effects of Continuous and Interval Running in Women and Men." European Journal of Applied Physiology, 37 (1977): 83-92.
- Edwards, H.T; Margaria, R. and Dill, D. B. "Metabolic Rate, Blood Sugar and the Utilization of Carbohydrate." The American Journal of Physiology, 108 (1934): 203-209.
- Eyestone, E. D.; Fellingham, G.; George, J. and Fisher, G. "Effect of Water Running and Cycling on Maximum Oxygen Consumption and 2-Mile Run Performance." The American Journal of Sports Medicine, 21 no. 1 (1993): 41-44.
- Foster, C.; Hector, L.I.; Welsh, R.; Schrager, M.; Green, M. A.; Snyder, A. C. "Effects of specific vs. cross training on running performance." European Journal of Applied Physiology, 70 (1995): 367-372.
- Fox, E. L.; et al. "Frequency and Duration of Interval Training Programs And Changes In Aerobic Power." Journal of Applied Physiology, 38 (1975): 481.

- Franch, J.; Madsen, K.; Djurhuus, M. S.; Perersen, P.K. "Improved Running Economy Following Intensified Training Correlates With Reduced Ventilatory Demands." Medicine and Science in Sports and Exercise, 30 no. 8 (1998): 1250-1256
- Fry, R. W.; Morton, A. R. and Keast, D. "Periodization and The Prevention of Overtraining." Canadian Journal of Sports Science, 17 (1992): 241-248.
- Gaesser, G. A. and Rich, G. A. "Effects of High and Low Intensity Exercises on Aerobic Capacity and Blood Lipids." Medicine and Science in Sports and Exercise, 16 (1984): 269-274.
- Goforth, H. W.; Jacobs, I. and Prusaczyk, W. K. "Simultaneous Enhancement of Aerobic and Anaerobic Capacity." Medicine and Science in Sports and Exercise, 26 no. 5(1994): S171.
- Gollnick, P. D.; Armstrong, R.B.; Saubert IV, C.W.; Piehl, K. and Saltin, B. "Enzyme Activity and Fiber Composition In Skeletal Muscle Of Trained And Untrained Men." Journal of Applied Physiology, 33 (1972): 312.
- Gollnick, P. D. and Hermansen, L. "Bio-mechanical adaptations to exercise: anaerobic metabolism." Exercise and Sports Science Reviews, ed. J. Wilmore, Academic Press 1 no.1 (1973):
- Henritze, J.; Weltman, A.; Schurrer, R. L. and Barlow, K. "Effects of training at and above the lactate threshold and maximal oxygen uptake." European Journal of Applied Physiology, 54 (1985):84-88.
- Hickson, R.C., et al. "Reduced Training Intensities and Loss of Aerobic Power, Endurance, and Cardiac Growth." Journal of Applied Physiology, 58 no. 2 (1985): 492-499.
- Hickson, R.C. et al. "Linear Increase in Aerobic Power Induced by Strenuous Program of Endurance Exercise." Journal of Applied Physiology, 42 (1977): 372-376.
- Hickson, R.C. et al. "Reduced Training Duration Effects on Aerobic Power, Endurance and Cardiac Growth." Journal of Applied Physiology, 53 (1982): 225-229.
- Hill, D.W. "Energy cost of middle distance running races." Medicine and Science in Sports and Exercise, 27no. 5 (1995): S45.
- Houston, M. E. and Thompson, J. A. "The Response of Endurance Adapted Adults to Intense Anaerobic Training." European Journal of Applied Physiology, 36 (1977): 207-213.

- Izumi, T.; Irisawa, K.; Kouzaki, M.; Nishimura, K.; Ogita, F. and Miyachi, M. "Metabolic Profile of High Intensity Intermittent Exercises." Medicine and Science in Sports and Exercise, 29 no. 3 (1997): 390-395.
- Kraemer, W. J. et al. "Training Responses of Plasma Beta-Endorphin, Adrenocorticotropin and Cortisol." Medicine & Science in Sports & Exercise, 21 no. 2 (1989): 146-153.
- LaFontaine, T.P. et al. "The Maximal Steady State versus Selected Running Events." Medicine and Science in Sports and Exercise, 13 (1981): 190.
- Lacour, J.R.; Magunacelaya, A.P. and Barthelemy, J.C. et al. "The Energetics of Middle Distance Running." European Journal of Applied Physiology, 60 (1990): 38-43.
- Lindsay, F.H., et al. "Improved Athletic Performance in Highly Trained Cyclists after Interval Training." Medicine and Science in Sports and Exercise, 28 no. 11 (1996): 1427-1434.
- Loy, S. F., et al. "Effects of Stair-climbing Versus Run Training on Treadmill and Track Running Performance." Medicine and Science in Sports and Exercise, 25 no. 11 (1993): 1275-1278.
- Maffulli, N.; Capasso, G. and Lancia, A. "Anaerobic Threshold and Performance in Middle and Long Distance Running." Journal of Sports Medicine & Physical Fitness, 31 (1991): 332-338.
- McMillan, G. (editor) "Results of a Recent Study of Masters Running." Peak Running Performance. (1997) <<http://www.peakrun.com/articles>> Internet: Accessed 25 September 1998.
- McMillan, G. (editor). "How Running Physiology Impacts on Optimal Training." Peak Running Performance. (1997) <<http://www.peakrun.com/articles>> Internet: Accessed 25 September 1998.
- McMillan, G. (editor). "Anaerobic Conditioning: Raising Your Lactate Threshold." Peak Running Performance. (1997) <<http://www.peakrun.com/articles>> Internet: Accessed 25 September 1998.
- McMillan, G. (editor). "Aerobic Capacity Training: Tough Intervals to Increase Your Aerobic Power." Peak Running Performance. (1997) <<http://www.peakrun.com/articles>> Internet: Accessed 25 September 1998.
- McMillan, G. (editor). "Anaerobic Capacity Training: Teaching Your Body to Buffer Lactic Acid." Peak Running Performance. (1997) <<http://www.peakrun.com/articles>> Internet: Accessed 25 September 1998.

- McMillan, G. (editor). "How to Raise Your Lactate Threshold." Peak Running Performance. (1997) <<http://www.peakrun.com/articles>> Internet: Accessed 25 September 1998.
- McMillan, G. (editor). "Improving Your Race Performances With Age." Peak Running Performance. (1997) <<http://www.peakrun.com/articles>> Internet: Accessed 25 September 1998.
- McMillan, G. (editor). "Changing your fitness Level." Peak Running Performance, 7 no. 5 (1998): 7-12.
- McMillan, G. ed. "Middle Distance Running: A Proven Guide." Peak Running Performance, 7 no. 6 (1998): 1-7.
- McMillan, G., ed. "Spring 5K and 10K Training Schedules: part II." Peak Running Performance, 7 no. 2 (1998): 7-10. San Diego: Road Runner Sports.
- Medbo, J.I. and Burgers, S. "Effect of Training on the Anaerobic Capacity." Medicine and Science in Sports and Exercise, 22 no. 4(1990): 501-507.
- Medbo, J. I. and Tabata, I. "Relative importance of aerobic and anaerobic energy release during short-lasting exhausting bicycle exercise." Journal of Applied Physiology, 67 no. 5 (1989): 1881-1886.
- Mikesell, K. A., and Dudley, G. A. "Influence of intense endurance training on aerobic power of competitive distance runners." Medicine & Science in Sports & Exercise, 16 no. 4 (1984), 371-375.
- Nagelkirk, P. R., et al. "The Effects Of Detraining, Retraining on Cardiorespiratory Responses After Six Weeks of High-Intensity Cycling." Medicine & Science in Sports & Exercise, 30 no. 5 (1998), S1374.
- Ocel, J., et al. "Adaptation of the slow component of VO₂ following 6 weeks of high or low intensity exercise training." Medicine & Science in Sports & Exercise, 30 no. 5 (1998), S943.
- Paradisis, G. P., et al. "The Effects of Combined Uphill-Downhill Sprint Training of Biomechanical Variables." Medicine & Science in Sports & Exercise. 30no. 5 (1998), S1467.
- Peak Running Performance. "The Most Effective Cross-Training for Runners." <<http://www.peakrun.com/articles/x-train.html>> (1997). Internet: Accessed 1-14-99.
- Peak Running Performance. 1997. "How running physiology impacts optimal training." <http://www.peakrun.com/articles/x-train.html> (1997) Internet: Accessed 1-14-99.

- Phillips, S. M.; Green, H. J.; Tarnopolsky, M. A.; Heigenhauser, G. F.; Hill, R. E.; and Grant, S. M. "Effects of Training Duration on Substrate Turnover and Oxidation during Exercise." Journal of Applied Physiology, 81 no. 5 (1996): 2182-2191.
- Prud'homme, D., et al. "Sensitivity of Maximal Aerobic Power to Training Is Genotype-Dependent." Medicine & Science in Sports & Exercise, 16 no. 5 (1984): 489-493.
- Schmidt, W. D., et al. "The Effects of Continuous Versus Short Bout Exercise on VO₂ (Max), Weight Loss and Adherence." Medicine & Science in Sports & Exercise, 30 no. 5 (1998), S1758.
- Seals, D. R., et al. "Endurance Training in Older Men and Women I: Cardiovascular Responses to Exercise." Journal of Applied Physiology, 57 no. 4(1984): 1024-1029.
- Seals, D. R., et al. "Endurance Training in Older Men and Women II: Blood Lactate Response to Sub-Maximal Exercise." Journal of Applied Physiology, 57 no. 4(1984): 1030-1033.
- Selley, E.A., et al. "Running intensity as determined by heart rate is the same in fast and slow runners in both 10- and 21-km races". Journal of Sports Sciences, 13 (1995): 405-410.
- Shepard, R.J. "Aerobic versus anaerobic training for success in various athletic events". Canadian Journal of Applied Sport Sciences, 3 (1978): 9-15.
- Simoës, H. G., et al. "High and low acidosis training: Effects upon aerobic and anaerobic performance". Medicine & Science in Sports & Exercise, 30 no. 5(1998): S932.
- Strath, S.J., et al. "Effect of hard versus light-moderate intensity exercise training on health and fitness benefits". Medicine & Science in Sports & Exercise, 30 no. 5(1998), S1341.
- Swartz, A.M., et al. "Effects of energy expenditure above and below 1200 kilo-calories /week on health and fitness benefits". Medicine & Science in Sports & Exercise, 30no. 5(1998), S1343.
- Tabata, I., et al. "Effects of moderate-intensity endurance and high-intensity intermittent training on anaerobic capacity and VO₂max". Medicine & Science in Sports & Exercise, 28 no. 10(1996), 1327-1330.
- Tanaka, K.; Matsuura, Y.; Natsuzaka, A. et al. "A longitudinal assessment of anaerobic threshold and distance running performance". Medicine and Science in Sports and Exercise, 16 (1984): 278-282.

- Tanaka, K.; et al. "Critical determinants of endurance performance in middle-aged and elderly endurance runners with heterogeneous training habits". European Journal of Applied Physiology, 59 (1990): 443-449.
- Terjung, Ronald L. "Muscle Adaptation to Aerobic Training". Gatorade Sports Science Institute Reference Desk., SSE # 54 no. 8 (1995): Number 1.
- Womack, C. J., et al. "Slow component of O₂ uptake during heavy exercise: Adaptation to endurance training". Journal of Applied Physiology, 79 no. 3(1995): 838-845.
- Winett, Richard A. "Optimizing cardiovascular training". Master Trainer, August, 1995. Internet: <<http://ageless-athletes.com/aerobics.htm>> Accessed November 1998
- Winett, Richard A. "The essence of cardiovascular training: maximizing oxygen consumption (part 1)". Master Trainer, March, 1998. Internet: <<http://ageless-athletes.com/aerobics.htm>> Accessed November 1998
- Winett, Richard A. "The essence of cardiovascular training: maximizing oxygen consumption (part 2)". Master Trainer, March, 1998. Internet: <<http://ageless-athletes.com/aerobics.htm>>

3. PERIODICALS: COMBINED STRENGTH AND CARDIOVASCULAR TRAINING

- American College of Sports Medicine. "The recommended quantity and quality of exercise for development and maintaining cardiorespiratory muscular fitness in healthy adults." Medicine and Science in Sports and Exercise, 22 (1990): 265-274.
- Bell, G.J., et al. "Physiological adaptations to concurrent endurance training and low velocity resistance training". International Journal of Sports Medicine, 12 no. 4(1991): 384-390.
- Bell, G. J.; Petersen, S. R.; Wessel, J.; Bagnall, K. and Quinney, H. A. "Adaptations to endurance and low velocity resistance training performed in a sequence". Canadian Journal of Sport Science 16 (1991a): 186-192.
- Blessing, D. L., et. al. "The influence of co-activation on the adaptive response to concurrent strength and endurance training in women". Medicine & Science in Sports & Exercise, 30 no. 5(1998): S1097.
- Blessing, D.; Stone, M.; Byrd, R.; Wilson, D.; Rozenek, R.; Pushparani, D.; and Liper, H. "Blood lipid and hormonal changes from jogging and weight training of middle-aged men". Journal of Applied Sport Science Research, 1(1987): 25-29.

- Chromiak, J. A.; and D. R. Mulbaney. "A review: The effects of combined strength and endurance training on strength development". Journal of Applied Sports Science Research, 4 (1990)(2): 55-60.
- Craig, B. W.; Lucas, J.; Pohlman, R. and Schilling, H. "The effect of running, weightlifting and a combination of both on growth hormone release." Journal of Applied Sports Science Research, 5 no. 4 (1991): 198-203.
- Dudley, G. A., and S. J. Fleck. "Strength and Endurance Training. Are they mutually exclusive"? Sports Medicine, 4 (1987): 79-85.
- Dudley, G. A., and R. Djmil. "Incompatibility of endurance and strength training modes of exercise." Journal of Applied Physiology. 59 no. 5. (1985): 1446-1451.
- Fry, A. C.; Kraemer, W. J.; Stone, M .H.; Fleck , S. J.; Kearney, J T.; Triplett, N. T. and Knuttgen, H. G. "Performance decrements with high intensity resistance exercise overtraining." Medicine and Science in Sport and Exercise, 26 (1994): 1165-1173.
- Fry, A. C. and Kraemer, W. J. "Physical performance characteristics of American collegiate football players." Journal of Applied Sport Science Research, 5 (1991): 126-138.
- Frykman, P.N., E. A. Harman, B. C. Nindl, W. J. Kraemer, L .A. Gotshalk, J. O. Marx, E. R. Lammi, and C. J. Palmer. "Changes in female thigh composition resulting from six months of heavy strength/endurance training." Medicine & Science in Sports & Exercise, 30 no. 5 (1998): S1040,
- Gettman, L.R., et al. "A comparison of combined running and weight training with circuit weight training". Medicine & Science in Sports & Exercise, 14 no. 3(1982), 229-234.
- Gravelle, B.L., and D. Blessing. "Physiological adaptation in women concurrently training for strength and endurance". Medicine & Science in Sports & Exercise, 30 no. 5(1998), S1098.
- Hennessy, L.C., and A.W.S. Watson. "The interference effects of training for strength and endurance simultaneously". Journal of Strength and Conditioning Research, 8 no. 1(1994): 12-19.
- Hepple, R.T., et al. "Comparison of resistance vs. aerobic training in older men". Medicine & Science in Sports & Exercise, 30 no. 5(1998), S1100.
- Hertling, Mark P. "Physical training for the modern battlefield". School of Advanced Military Studies, U. S. Army Command and General Staff College, Fort Leavenworth, Kansas, 1 April, 1988.

- Hickson, Robert C., "Interference of strength development by simultaneously training for strength and endurance." European Journal of Applied Physiology. 45 (1980): 255-263.
- Hoff, J., et al. "Effect of endurance performance from introducing maximal strength training in upper body for cross-country skiers". Medicine & Science in Sports & Exercise, 30 no. 5(1998), S1468.
- Hoffman, J.R.; Kraemer, W.J.; Fry, A. C.; Deschenes, M. and Kemp, M. "The effects of self selection for frequency of training in a winter conditioning program for football." Journal of Applied Sport Science Research, 4 (1990): 76-82.
- Hortobagyi, T., et al. "Effects of simultaneous training for strength and endurance on upper and lower body strength and running performance". Journal of Sports Medicine and Physical Fitness, 31 (1991): 20-30.
- Hunter, G., et al. "Development of strength and maximum oxygen uptake during simultaneous training for strength and endurance." Journal of Sports Medicine and Physical Fitness, 27no. 3. (1987): 269-275.
- Hurley, B.F., et al. "Effects of high-intensity strength training on cardiovascular function". Medicine & Science in Sports & Exercise, 16 no. 5(1984), 483-498.
- Johnston, R. E.; Quinn, T.J.; Kertzer, R.; and Vroman, N.B. 1995. Strength training in female distance runners: impact on running economy. . Medicine and Science in Sports and Exercise 27(5): S47.
- Kraemer, W. J., et al. "Compatibility of high-intensity strength and endurance training on hormonal and skeletal muscle adaptations" Journal of Applied Physiology, 78 no. 3. (1995): 976-989.
- Kraemer, W.J., et al. "Compatibility of conventional strength and endurance programs". Medicine & Science in Sports & Exercise. 22(2), S111, 1990.
- Kraemer, W. J.; Vogel, J. A.; Patton, J. F.; Dziados, J. E.; and Reynolds, K. L. "The effects of various physical training programs on short duration high intensity load bearing performance and the Army physical fitness test". USARIEM Technical Report, 30/87 August.
- Larsson, L. "Physical Training Effects on Muscle Morphology in Sedentary Males at Different Ages." Medicine & Science in Sports & Exercise, 14 no. 3(1982), 203-206.
- Lee, A.; Craig, B W.; Lucas, J.; Pohlman, R. and Stelling, H. "The Effect of Endurance Training, Weight Training, and a Combination of Endurance And Weight Training Upon the Blood Lipid Profile of Young Male Subjects." Journal of Applied Sport Science Research, 4 (1990): 68-75.

- Marcinik, E.J., et al. "Effects of Strength Training on Lactate Threshold and Endurance Performance." Medicine & Science in Sports & Exercise. 23 no. 6 (1991), 739-743.
- McCarthy, J. P., et al. "Compatibility of Adaptive Responses with Combining Strength and Endurance Training." Medicine & Science in Sports & Exercise, 27 no. 3. (1995): 429-436.
- Nelson, A. G, et al. "Consequences of Combining Strength and Endurance Training Regimens." Physical Therapy, 70 no. 5(1990): 287-294.
- O'Bryant, H. S.; Byrd, R.; and Stone, M. H. "Cycle Ergometry Performance and Maximum Leg and Hip Strength Adaptations to Two Different Methods of Weight Training." Journal of Applied Sport Science Research, 2 no. 2 (1988): 27-30.
- Peterson, J. A. "Total Conditioning: A Case Study." Athletic Journal 56 (1975): 40-55.
- Pedersen, P. K., et al. "Short-interval run training reduces the slow VO₂ component during running." Medicine & Science in Sports & Exercise, 30 no. 5 (1998), S1079.
- Pollock, M. L., et al. "Position Stand: The Recommended Quantity and Quality of Exercise for Developing and Maintaining Cardio-Respiratory and Muscular Fitness and Flexibility in Healthy Adults." Medicine & Science in Sports & Exercise, 30 no. 6 (1998), 975-991, 1998.
- Rousseau, A. K., et al. "The Effects of Combined Endurance and Strength Training in Women Age 60-75." Medicine & Science in Sports & Exercise, 30 no. 5(1998): S1099.
- Sale, D. G., et al. "Comparison of Two Regimens of Concurrent Strength and Endurance Training." Medicine & Science in Sports & Exercise, 22 no. 3 (1990), 348-356.
- Sale, D. G.; MacDougall, J.; Jacobs, I.; and Garner, S. "Interaction Between Concurrent Strength and Endurance Training." Journal of Applied Physiology, 68 (1990): 260-270.
- Volpe, S. L.; Walberg-Rankin, J.; Rodman, K. W.; and Sebolt, D. R. "The effect of endurance running on training adaptations in women participating in a weight lifting program." Journal of Strength and Conditioning Research, 7 no. 2. (1993): 101-107.

4. PERIODICALS: OVER-TRAINING STUDIES

- Callister, R., et al. "Physiological and Performance Responses to Overtraining in Elite Judo Athletes." Medicine & Science in Sports & Exercise, 22 no. 6 (1990): 816-824.
- Costill, D. L.; Flynn, M. G.; Kirwan, J. P.; Houmard, J. A.; Mitchell, J.B.; Thomas, R.; and Park, S.H. "Effects Of Repeated Days of Intensified Training on Muscle Glycogen and Swimming Performance." Medicine & Science in Sports and Exercise, 20 no. 3(1988): 249-254.
- Fry, R.W., et al. "Over-Training in Athletes: An Update." Sports Medicine, 12 no. 1(1996): 32-65.
- Fry, A. C.; Kraemer, W. J.; Lynch, J. M.; Triplett, H. T. and Koziris, L. P. "Does short-term near-maximal intensity machine resistance training induce over-training?" Journal of Strength and Conditioning, 8 (1994): 188-191.
- Fry, A. C.; Kraemer, W. J.; van Borselen, F.; Lynch, J. M.; Marsit, J. L.; Pierre Roy, E.; Triplett, H. T. and Knuttgen, H. G. "Performance Decrements with High-Intensity Resistance Exercise Over-Training." Medicine and Science in Sports and Exercise, 26 (1994): 1165-1173.
- Kirwan, J. P.; Costill, D. L.; Flynn, M. G.; Mitchell, J.B.; Fink, W.J.; Neuffer, P. D.; and Houmard, J. A. "Physiological Responses to Successive Days of Intense Training in Competitive Swimmers." Medicine & Science in Sports and Exercise, 20 no. 3(1988): 255-259.
- Kuipers, H. and. Keizer, H. A. Over-Training in Elite Athletes: Review and Directions for the Future. Sports Medicine, 6 (1988): 79-92,.
- Lehmann, M.J.; Lormes, W.; Opitz-Gress, A.; Steinacker, J. M.; Netzer, N.; Foster, C. and Gastman, U. "Training and Overtraining: An Overview and Experimental Results in Endurance Sports." Journal of Sports Medicine and Physical Fitness, 37 (1997): 7-17.
- Lehmann, M.; Gastmann, U.; Petersen, K. G.; Bachl, N.; Seidel, A.; Khalaf, A. N.; Fischer, S. and Keul, J. "Training-Over-Training; Performance, and Hormone Levels, After a Defined Increase in Training Volume Versus Intensity in Experienced Middle and Long Distance Runners." British Journal of Sports Medicine, 26 (1992): 233-242.
- Lehmann, M.; Baumgartl, P; Wiesenack, C.; Seidel, A.; Baumann, H.; Fischer, S.; Spori, U.; Gendrisch, G.; Kaminski, R.; Keul, J. "Training-over-training: influence of a defined increase in training volume vs. training intensity on performance, catecholamines and some metabolic parameters in experienced middle and long distance runners." European Journal of Applied Physiology, 64 (1992): 169-177.

Lehmann, M.; Gastmann, U.; Steinacker, J. M. and Keul, J. "Unacustomed high mileage compared to high intensity training-related performance and neuromuscular responses in distance runners." European Journal of Applied Physiology, 70(1995): 457-461.

5. BOOKS LECTURES, INTERVIEWS AND STUDIES

American College of Sports Medicine. Guidelines for Exercise Testing and Prescription. Philadelphia: Lea and Febiger, 1991.

Applewhite, Roger. Personal Trainer. Golden CO: Muscle Media, 1997.

Aaberg, Everett, The Science and Art of Resistance Training: Upper Body," lecture, Baltimore, Maryland at International Summit of Personal Trainers, 5 March 1999.

Aaberg, Everett, The Science and Art of Resistance Training: Lower Body," lecture, Baltimore, Maryland at International Summit of Personal Trainers, 6 March 1999.

Baechle, T.R., eds. Essentials of Strength Training and Conditioning. Champaign, IL: Human Kinetics Publishers, Inc., 1994.

Bompa, T. O.; and Cornacchia, L. J. Serious Strength Training. Champaign, IL: Human Kinetics Publishers, Inc., 1998.

Burke, E. R., eds. Precision Heart-rate Training. Champaign, IL: Human Kinetics Publishers, Inc., 1998.

Burke, Edmund R. Precision Cycling. Finland: Graphic Designing CIS, 1994.

Columbu, Franco with Lydia Fragomeni. The Businessman's Minutes-a-Day Guide to Shaping Up. Chicago, IL: Contemporary Books, 1983.

Columbu, Franco with George Fels. Winning Body-building. Chicago, IL: Contemporary Books, 1977.

Columbu, Franco. Franco Columbu's Book of Bodybuilding. Chicago, IL: Contemporary Books, 1987.

Costill, David L. Inside Running: Basics of Sports Physiology. Indianapolis, IN: Benchmark Press, 1986.

Edwards, Sally. The Heart Rate Monitor Book. Sacramento, CA: Fleet Street Press, 1994.

- Fardy, P. S. "Training for aerobic power." in Toward an Understanding of Human Performance, ed. E. J. Burke, 31-39. Ithaca, NY: Movement Press, 1997.
- Fleck, Steve J. "Periodization Breakthrough Part I," lecture, Baltimore Maryland, at the International Summit of Personal Trainers, 5 March 1999, cited with permission of Dr. Fleck.
- Fleck, Steve J. "Periodization Breakthrough Part II," lecture, Baltimore Maryland, at the International Summit of Personal Trainers, 6 March 1999, cited with permission of Dr. Fleck.
- Fleck, S.J. and Kraemer, W.J. Designing Resistance Training Programs. Champaign, IL: Human Kinetics Publishers, Inc., 1997.
- Fleck, S.J.; and Kraemer, W.J. Periodization Break-Through. Ronkonkoma, NY: Advanced Research Press, Inc., 1996.
- Froelicher, V. F. Exercise Testing And Training. Chicago: Yearbook Medical Publishers, 1983.
- Glover, Bob and Schuder, Pete. Competitive Runner's Handbook. New York, NY: Penguin Books, 1983.
- Hansford, Peter A. "The effects of a periodised 24-week heavy and explosive type resistance training program on the maximal strength, muscular power output and functional capacity of aged subjects." Submitted as a partial requirement for the Bachelor of Human Movement Science with Honours at Southern Cross University. Southern Cross University, Australia, 1998.
- Hatfield, Frederick C. Powerlifting: A Scientific Approach. Chicago, IL: Contemporary Books, 1981.
- Hatfield, Frederick C. "Powerlifting and speed-strength training". Adapted from Fitness: The complete guide (3rd edition). International Sports Sciences Association, 1996.
- Health For Life. Secrets of Advanced Bodybuilding. Los Angeles, CA: Health for Life, 1985.
- Higdon, Hal. Run Fast: How to train for a 5K or 10K Race. Emmaus, PA: Rodale Press, 1992.
- Higdon, Hal. Marathon, The Ultimate Training and Racing Guide. Emmaus, PA: Rodale Press, 1993.
- Janssen, Peter, G.J.M. Training Lactate Pulse Rate. Oy Liitto, Finland: Polar Electro Oy, 1994.

- Komi, P. V. ed. Strength and Power in Sport. Oxford, Blackwell Scientific Publications, 1992.
- Kreider, R.B.; Fry, A. C.; and O'Toole, M. L. Over-training in Sport. Champaign, IL: Human Kinetics Publishers, Inc., 1998.
- Lamb, D.R. Physiology of exercise: Response and adaptations. New York: Macmillan, 1978.
- Lawrence, A.; and Scheid, M. Running and Racing after 35. Boston, MA: Little, Brown and Co., 1990.
- MacDougall, J.D; Wenger, H. A. and Green, H.J. eds. Physiological Testing of the Elite Athlete. Canada: Mutual Pres Limited, 1982.
- Moran, G.T.; and McGlynn, G.H. Cross Training for Sports. Champaign, IL: Human Kinetics Publishers, Inc., 1994.
- Moritani, T. "Time course of adaptations during strength and power training." In Strength and Power in Sport, P. V. Komi ed., 226-278. Oxford;Blackwell, 1992.
- Neff, Peter. Natural Bodybuilding. New York, NY: Avon Books, 1985.
- Newsholme, E.; Leech, T. and Duester, G. Keep on Running: The Science of Training and Performance. West Sussex, England: John Wiley & Sons LTD., 1994.
- Newton, R. U. Doctoral Dissertation: "Expression and Development of Maximal Muscle Power." School of Exercise and Sport Management, Southern Cross University, Lismore Australia, 997.
- Noakes, T. Lore of Running. Cape Town, South Africa: Oxford University Press, 1986.
- Riley, Daniel P. Strength Training by the Experts. West Point, NY: Leisure Press, 1977.
- Robinson, J.; and Carrino, F. Max O2: The Complete Guide to Synergistic Aerobic Training. Los Angeles, CA: Health for Life, 1993.
- Rodgers, Bill and Welsch Priscilla. Masters Running and Racing. Emmaus, PA: Rodale Press, 1991.
- Rodgers, Bill and Douglas, Scott. Lifetime Running Plan. New York, NY: Harper Collins, 1996.
- Schwarzenegger, Arnold with Bill Dobbins. Encyclopedia of Modern Body-Building. New York, NY: Simon & Schuster Inc., 1985.

- Simon, Mitchell S. "The Science of Loading for Hypertrophy." Lecture, Baltimore, Maryland at the International Summit of Personal Trainers, 7 March, 1999.
- Sleamaker, Rob. Serious Training for Serious Athletes. Champaign, IL: Leisure Press, 1989.
- Sparks, Ken and Kuehls, Dave. The Runners Book of Training Secrets. Emmaus, PA: Rodale Press, 1996.
- Stone, M. and O'Bryant, H. Weight Training: A Scientific Approach. Minneapolis: Burgess International, 1987.
- Weider, Joe with Bill Reynolds. Ultimate Body-building. Chicago, IL: Contemporary Books, 1989.
- Westcott, W. Strength Fitness. Dubuque, Iowa: Wm C. Brown Publishers, 1989.
- Zatsiorsky, V.M. Science and Practice of Strength Training. Champaign, IL: Human Kinetics Publishers, Inc.