



**ARMSTRONG
LABORATORY**

**THE RELATIONSHIP BETWEEN STRATUM
CORNEUM DIFFUSION COEFFICIENTS AND
TEMPERATURE FOR HALOGENATED
HYDROCARBONS**

M.C. Caracci

**GEO-CENTERS, INC.
7 WELLS AVENUE
NEWTON CENTRE, MA 02159**

Z. Liron

**ISRAEL INSTITUTE FOR BIOLOGICAL RESEARCH
NESS ZIONA, ISRAEL 70450**

**J.D. McCafferty
G.W. Jepson**

19990830 032

**OCCUPATIONAL AND ENVIRONMENTAL HEALTH DIRECTORATE
TOXICOLOGY DIVISION, ARMSTRONG LABORATORY
WRIGHT-PATTERSON AFB, OH 45433-7400**

April 1995

INTERIM REPORT FOR THE PERIOD AUGUST 1994 THROUGH APRIL 1995

Approved for public release; distribution is unlimited

**AIR FORCE MATERIEL COMMAND
WRIGHT-PATTERSON AIR FORCE BASE, OHIO 45433-6573**

NOTICES

When US Government drawings, specifications or other data are used for any purpose other than a definitely related Government procurement operation, the Government thereby incurs no responsibility nor any obligation whatsoever, and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise, as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use, or sell any patented invention that may in any way be related thereto.

Please do not request copies of this report from the Armstrong Laboratory. Additional copies may be purchased from:

National Technical Information Service
5285 Port Royal Road
Springfield, Virginia 22161

Federal Government agencies and their contractors registered with the Defense Technical Information Center should direct requests for copies of this report to:

Defense Technical Information Center
Cameron Station
Alexandria, Virginia 22314

DISCLAIMER

This Technical Report is published as received and has not been edited by the Technical Editing Staff of the Armstrong Laboratory.

TECHNICAL REVIEW AND APPROVAL


AL/OE-TR-1995-0058

The experiments reported herein were conducted according to the "Guide for the Care and Use of Laboratory Animals," Institute of Laboratory Animal Resources, National Research Council.

This report has been reviewed by the Office of Public Affairs (PA) and is releasable to the National Technical Information Service (NTIS). At NTIS, it will be available to the general public, including foreign nations.

This technical report has been reviewed and is approved for publication.

FOR THE COMMANDER


TERRY A. CHILDRRESS, Lt Col, USAF, BSC
Director, Toxicology Division
Armstrong Laboratory

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503

1. AGENCY USE ONLY (Leave Blank)	2. REPORT DATE April 1995	3. REPORT TYPE AND DATES COVERED Interim - August 1994 - April 1995	
4. TITLE AND SUBTITLE The Relationship Between Stratum Corneum Diffusion Coefficients and Temperature for Halogenated Hydrocarbons		5. FUNDING NUMBERS Contract DAMD17-93-C-3006 PE 61102F PR 2312 TA 2312A2 WU 2312A201	
6. AUTHOR(S) M.C. Caracci, Z. Liron, J.D. McCafferty, and G.W. Jepson		8. PERFORMING ORGANIZATION REPORT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Geo-Centers, Inc. Israel Institute for Biological Research 7 Wells Avenue Ness Ziona, Israel 70450 Newton Centre, MA 02159		10. SPONSORING/MONITORING AGENCY REPORT NUMBER AL/OE-TR-1995-0058	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Armstrong Laboratory, Occupational and Environmental Health Directorate Toxicology Division, Human Systems Center Air Force Materiel Command Wright-Patterson AFB OH 45433-7400		11. SUPPLEMENTARY NOTES	
12a. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution is unlimited.		12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) The relationship between diffusion coefficients and exposure temperature was evaluated in isolated stratum corneum (SC). Due to the difficulty in measuring a diffusion coefficient directly, the movement of chemicals into and across the dermal barrier is often described as a function (the permeability constant) of the diffusion coefficient. The permeability constant also depends on the skin: chemical partition coefficient and the pathlength of diffusion. In the idealized expression of diffusion given in the Stokes-Einstein equation, diffusion is described in terms of kinetic energy and the viscous resistance of diffusion. If this relationship holds for the diffusion of halogenated chemicals in isolated stratum corneum then the observed diffusion coefficients will be dependent on the temperature of the exposure system. Rat or human stratum corneum was removed from whole thickness skin trypsin treatment and stored in a dessicator jar. Approximately 5 Mg of SC was placed into the weighing pan of a thermogravimetric analyzer and exposed to vapor or halogenated hydrocarbons. The increase in tissue weight over time for temperatures ranging from 27-40°C were evaluated. Diffusion coefficients were estimated by fitting the pre-equilibrium data with a non-steady state diffusion equation. The observed diffusion coefficients were dependent on the temperature of the exposure, indicating the importance of careful temperature control in dermal absorption studies using halogenated hydrocarbons.			
14. SUBJECT TERMS Diffusion coefficients Halogenated hydrocarbons Skin			15. NUMBER OF PAGES 12
			16. PRICE CODE
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED	18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED	19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED	20. LIMITATION OF ABSTRACT UL

THIS PAGE INTENTIONALLY LEFT BLANK

PREFACE

This technical report describes the relationship between diffusion coefficients and exposure temperature evaluated in isolated stratum corneum. Due to the difficulty in measuring a diffusion coefficient directly, the movement of chemicals into and across the dermal barrier is often described as a function (the permeability coefficient) of the diffusion coefficient. The permeability constant also depends on the skin:chemical partition coefficient and the pathlength of diffusion. In the idealized expression of diffusion given in the Stokes-Einstein equation, diffusion is described in terms of kinetic energy and the viscous resistance of diffusion. If this relationship holds for the diffusion of halogenated chemicals in isolated stratum corneum then the observed diffusion coefficients will be dependent on the temperature of the exposure system. The information contained in this report was presented in a poster format at the Society of Toxicology Annual Meeting held in Baltimore, MD in March 1995.

The animals used in this study were handled in accordance with the principles in the *Guide for the Care and Use of Laboratory Animals*, prepared by the Committee on Care and Use of Laboratory Animals of the Institute of Laboratory Animals Resources, National Research Council, DHHS, National Institute of Health Publication #86-23, 1985, and the Animal Welfare Act of 1966, as amended.

TABLE OF CONTENTS

PREFACE.....	1
TABLE OF CONTENTS.....	2
LIST OF FIGURES	3
LIST OF TABLES.....	3
INTRODUCTION	4
METHODS AND MATERIALS.....	4
Materials	4
Method.....	4
Data Analysis.....	5
RESULTS	5
DISCUSSION	8
CONCLUSION.....	8
REFERENCES	10

LIST OF FIGURES

Figure 1	
Example of Diffusion Coefficient Determination of Human Stratum Corneum.....	6
Figure 2	
Examples of Diffusion Coefficient Determination of Rat Stratum Corneum	6
Figure 3	
Plots of Human Diffusion Coefficient versus Temperature	7
Figure 4	
Plots of Rat Diffusion Coefficients versus Temperature.....	8

LIST OF TABLES

Table 1	
Diffusion Coefficients for Human and Rat Stratum Corneum.....	6

INTRODUCTION

The Stokes-Einstein Equation describes chemical diffusion as a function of temperature,

$$D = \frac{k_B T}{6\pi nr} \quad \text{Equation 1}$$

where D is the Diffusion Coefficient, k_B is Boltzmann's constant, T is temperature, n is the solvent viscosity, and r is the molecular radius [1,2]. If the diffusional behavior of a chemical in stratum corneum is described by the Stokes-Einstein relationship, the diffusion should be proportional to temperature with a proportional constant of $k_B/6\pi nr$. The influence of temperature on measured diffusion coefficients should be accounted for in descriptions of chemical movement through the dermal barrier. This is particularly evident when viewed in the context of the dermal permeability coefficient, k_p (cm/hr), which is commonly used to describe the chemical movement in skin:

$$k_p = \frac{D \times P_{\text{skin/chemical}}}{x} \quad \text{Equation 2}$$

where $P_{\text{skin/chemical}}$ is the skin:chemical partition coefficient, and x is the pathlength of diffusion. As can be seen in Equation 2, accurate determination of the permeability coefficient depends on the use of a diffusion coefficient determined under appropriate temperature conditions.

Thermal Gravimetric Analysis (TGA) was used to determine the diffusion coefficient for a set of halogenated hydrocarbons in rat and human stratum corneum. Diffusion coefficient versus temperature data was used to determine the quantitative relationship between temperature and diffusion.

METHODS AND MATERIALS

Materials

The chemicals of interest in this study were Dibromomethane (99% purity), Tetrachloroethylene (99.99% purity), and Chloropentafluorobenzene (99% purity). All three chemicals were purchased from Aldrich.

Method

Human cadaver skin samples examined in this study were taken from the thigh. Rat skin samples were also collected and studied from the backs of three different rats. A dermatome was utilized to remove the top 0.025 mm layer from the rat or human whole skin sample. Skin sections were carefully removed from the dermatome catch and placed on a cutting board and sliced into manageable sections approximately 1" by 1". These sections were placed onto a 0.5% solution of trypsin type II in pH 7.4 phosphate buffer-

treated filter paper, making sure the ends were not folded under, or the skin had no air bubbles underneath it. Petri dishes containing samples were placed into a Carbon Dioxide-free incubator for two hours at 37°C. Meanwhile, 100 ml of 0.005% soybean trypsin inhibitor type II-S was prepared and poured into glass dishes and an equal number of dishes was filled with distilled water. Following incubation, the top layer was carefully removed using forceps and spatula, and the stratum corneum was allowed to float on top of the trypsin inhibitor for 10 minutes. The stratum corneum was then removed and placed in the water bath for 30 minutes. The stratum corneum was removed using a wire screen and allowed to dry until all water droplets were gone. Samples were stored in a desiccator until needed.

Approximately 5 mG of rat or human stratum corneum (SC) was placed on the titanium weighing pan of the Thermal Gravimetric Analyzer (TGA) and exposed to a dry air flow of 2 L/min. Once equilibrium was reached, the dry air was simultaneously stopped and replaced with a 10,000 ppm concentration of a halogenated hydrocarbon of interest. The mass of the sample was recorded by the TGA until a chemical equilibrium was obtained and the mass appeared constant. Samples were exposed at temperatures of 27, 30, 32, 35, 37, and 40°C.

Data Analysis

Figures 1 and 2 are samples of the data obtained from the TGA, i.e. the sample mass versus time in minutes. Diffusion coefficients were estimated by fitting the pre-equilibrium data with a non-steady state diffusion equation (Equation 3) [3,4,5]. The terms in equation 3 are described below.

$$\frac{MASS}{MINF} = 1 - \exp(-\beta t) - \sum_{n=0}^{\infty} \left(\frac{16}{(2n+1)^2 \pi^2} \right) \left(\frac{2l^2 \beta}{D(2n+1)^2 \pi^2 - 4l^2 \beta} \right) \left(\exp(-\beta t) - \exp\left(-\frac{D(2n+1)^2 \pi^2 t}{4l^2}\right) \right)$$

Equation 3

Description of terms in Equation 3

<i>MASS</i>	Mass of stratum corneum and chemical at a particular time (mg)
<i>MINF</i>	Mass of stratum corneum and chemical at infinite time (mg)
β	Lag constant. Equal to 0.238
<i>t</i>	Time (seconds)
<i>D</i>	Diffusion Coefficient (cm ² /sec)
<i>l</i>	Half thickness of the stratum corneum. Equal to 0.00034 cm

RESULTS

The diffusion coefficient was individually determined for each data set as described in the method section. Graphical results are shown in figures 1 and 2, with the tabular results of all the studies shown in table 1.




Figure 1: Example of diffusion coefficient determination of human stratum corneum at 32 °C with Tetrachloroethylene (PERC).




Figure 2: Example of diffusion coefficient determination of rat stratum corneum at 32 °C with Tetrachloroethylene (PERC).

Temperature (C)	CPF Human	PERC Human	DBM Human
27	2.50e-09	2.00e-09	3.00e-09
30	3.25e-09	2.50e-09	4.00e-09
32	4.00e-09	3.00e-09	7.00e-09
35	9.00e-09	3.00e-09	8.00e-09
37	5.50e-09	5.00e-09	9.00e-09
40	7.00e-09	6.00e-09	1.50e-08

Temperature (C)	CPF Rat (n=3)	PERC Rat (n=3)	DBM Rat (n=3)
27	4.100e-10 ± 1.493e-10	4.033e-10 ± 2.517e-11	6.333e-10 ± 5.774e-11
30	5.133e-10 ± 1.882e-10	5.233e-10 ± 2.082e-11	9.000e-10 ± 0.000e+00
32	6.333e-10 ± 1.358e-10	7.000e-10 ± 1.000e-10	1.800e-09 ± 3.000e-10
35	5.467e-10 ± 2.730e-10	9.000e-10 ± 2.646e-10	2.011e-09 ± 1.934e-11
37	8.833e-10 ± 3.753e-10	8.167e-10 ± 7.638e-11	2.133e-09 ± 2.309e-10
40	1.150e-09 ± 5.895e-10	1.233e-09 ± 3.786e-10	4.567e-09 ± 2.136e-09

Table 1: Diffusion Coefficients (cm²/sec) for human and rat stratum corneum for Chloropentafluorobenzene (CPF), Tetrachloroethylene (PERC), and Dibromomethane (DBM). Rat Diffusion Coefficients are an average of three different stratum corneum samples.

The rat diffusion coefficients shown are the average of the diffusion coefficients from the three different rats. The human diffusion coefficients ranged from 2.0e-9 to 1.5e-8 cm²/sec, while the average rat diffusion coefficients were slightly lower, ranging from 4.0e-10 to 3.4e-9 cm²/sec. A summary of these data is shown in figures 3 and 4 where the

diffusion coefficient is plotted versus temperature for each chemical and skin type. For each data set, the equation of the linear regression is shown.




Figure 3: Plots of Human Diffusion Coefficients versus Temperature for Chloropentafluorobenzene (CPFB), Tetrachloroethylene (PERC), and Dibromomethane (DBM).




Figure 4: Plots of Rat Diffusion Coefficients versus Temperature for Chloropentafluorobenzene (CPFBRat), Tetrachloroethylene (PERCRat), and Dibromomethane (DBMRat).

DISCUSSION

The Stokes-Einstein equation describes chemical diffusion as directly proportional to temperature (Equation 1) with a proportionality constant of $k_B/6\pi\eta r$. This study clearly shows this relationship in figures 3 and 4. Since the diffusion coefficient is used to calculate the dermal permeability coefficient (Equation 2), this influence of temperature on measured diffusion coefficients should be accounted for in descriptions of chemical movement through the dermal barrier.

CONCLUSION

- Diffusion is proportional to temperature for the dermal absorption of Dibromomethane (DBM), Tetrachloroethylene (PERC), and Chloropentafluorobenzene (CPFEB).
- For accurate determination of the dermal permeability coefficient, k_p , care should be taken to account for the influence of temperature on measured dermal diffusion coefficients.

REFERENCES

1. Chang R. 1977. Physical Chemistry With Applications to Biological Systems. MacMillan Publishing Co., New York. pp 66 - 67.
2. Cussler, E.L. 1984. Diffusion: Mass transfer in fluid systems. Cambridge University Press, Cambridge. pp 118.
3. Crank J. 1975. The Mathematics of Diffusion: Second Edition. Oxford University Press, New York. pp 48.
4. Liron Z, Clewell HJ and McDougal JN 1994. Kinetics of Water Vapor Sorption in Porcine Stratum Corneum. Journal Pharmaceutical Sciences 83 (5):692-698.
5. Quinn, D. 1994. Professor of Mathematics, Air Force Institute of Technology, WPAFB, OH. Personal Communication.