



NAVAL FACILITIES ENGINEERING SERVICE CENTER
Port Hueneme, California 93043-4370

**TECHNICAL REPORT
TR-2109-SHR**

**POTENTIAL FOR USING BIOGEOCHEMICAL
MARKERS TO ASSESS AND MONITOR THE
IMPACT OF MILITARY TRAINING EXERCISES
ON DESERT ECOSYSTEMS: PROOF-OF-
CONCEPT**

by


Frederick E. Goetz, Ph.D., UCSB
Leslie Karr, NFESC

September 1999

19991026 158

Approved for public release; distribution is unlimited.

BTIC QUALITY INSPECTED

 Printed on recycled paper

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to: Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE September 1999		3. REPORT TYPE AND DATES COVERED Final; Jan. 1998 – Sept. 1999	
4. TITLE AND SUBTITLE POTENTIAL FOR USING BIOGEOCHEMICAL MARKERS TO ASSESS AND MONITOR THE IMPACT OF MILITARY TRAINING EXERCISES ON DESERT ECOSYSTEMS: PROOF-OF-CONCEPT			5. FUNDING NUMBERS	
6. AUTHOR(S) Frederick E. Goetz, Ph.D., UCSB and Leslie Karr, NFESC				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Commanding Officer Naval Facilities Engineering Service Center Port Hueneeme. CA 93043-4371			8. PERFORMING ORGANIZATION REPORT NUMBER TR-2109-ENV	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) SERDP 901 North Stuart Street Arlington, VA 22203			10. SPONSORING/MONITORING AGENCY	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION/AVAILABILITY STATEMENT Distribution statement			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) The rate of nitrogen fixation and nitrogen flux in soil samples collected from a pristine and heavily disturbed site at Marine Corps Air Ground Combat Center (MCAGCC) Twenty-Nine Palms were 6 ± 2.05 and 3 ± 1.04 mM ethylene/g-h and 30.1 ± 6.47 and 21.9 ± 6.95 $\mu\text{g}/\text{m}^2 - \text{d}$ respectively ($P\leq 0.05$). In addition, significantly more bacteria are present in the disturbed soil samples and principle component analysis demonstrates distinct differences in the bacteria at each site. These data can be interpreted in terms of the destruction of vegetation and release of nutrients that accompany military training exercises. Decay of damaged vegetation releases nutrients that support bacterial growth and suppresses nitrogen-fixing bacteria. The bacteria at the disturbed site also appear to be experiencing an increase in environmental stress that may be associated with the damaged vegetation, soil compaction, reduced percolation of water, and/or a decrease in oxygen in the soil gas. A solid state ammonia sensor constructed by American Research Corporation of Virginia and tested in desert soil detected ammonia concentrations as low as 20 ppb which is the concentration reported for desert soils in this and other studies.				
14. SUBJECT TERMS Biogeochemical, military training, land management, environmental stress, nitrogen, ammonia sensor			15. NUMBER OF PAGES 23	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT U	18. SECURITY CLASSIFICATION OF THIS PAGE U	19. SECURITY CLASSIFICATION OF ABSTRACT U	20. LIMITATION OF ABSTRACT U	

EXECUTIVE SUMMARY

The Naval Facilities Engineering Service Center (NFESC) was tasked to investigate the feasibility of using biogeochemical characteristics of a desert ecosystem as one measure of the impact of military training exercises. Once developed, these measurements have the potential to be used in conjunction with other tools being developed under the Strategic Environmental Research and Development Program (SERDP) for monitoring and assessing the impact of military training exercises on a variety of ecosystems, such as land use management systems.

This Proof-of-Concept consisted of three experiments. They included studies on bacterial community structure in a desert training area and control site; the rate of nitrogen fixation from a desert training area and control site; and demonstrate that a solid state ammonia sensor can measure the flux of ammonia when placed in desert soil.

The results show the rate of nitrogen fixation and nitrogen flux in soil samples collected from a pristine and heavily disturbed site at Marine Corps Air Ground Combat Center (MCAGCC) Twenty-Nine Palms were 6 ± 2.05 and 3 ± 1.04 mM ethylene/g-h and 30.1 ± 6.47 and 21.9 ± 6.95 $\mu\text{g}/\text{m}^2 - \text{d}$ respectively ($P \leq 0.05$). In addition, significantly more bacteria are present in the disturbed soil samples and principle component analysis demonstrates distinct differences in the bacteria at each site. These data can be interpreted in terms of the destruction of vegetation and release of nutrients that accompany military training exercises. Decay of damaged vegetation releases nutrients that support bacterial growth and suppresses nitrogen-fixing bacteria. The bacteria at the disturbed site also appear to be experiencing an increase in environmental stress that may be associated with the damaged vegetation, soil compaction, reduced percolation of water, and/or a decrease in oxygen in the soil gas. A solid state ammonia sensor constructed by the American Research Corporation of Virginia, and tested in desert soil detected ammonia concentrations as low as 20 ppb which is the concentration reported for desert soils in this and other studies.

These results validate the premise that biogeochemical markers can be used to monitor the ecological impact of military training. The data presented confirm that there are statistically significant ($P \leq 0.05$) differences in the flux of ammonia and rate of nitrogen fixation between an impacted desert site and pristine site. The data also suggests that there are significant differences in the composition of the microbial population at the two sites. Lastly, the performance of the solid state ammonia sensor demonstrates its potential to monitor and collect real time data from large training areas. A field scale program based upon these results is the next step in developing the technology for monitoring and assessing training impacts on military ecosystems.

Table of Contents

	Page
INTRODUCTION.....	1
PROOF-OF-CONCEPT.....	3
Experiment 1.....	3
Experiment 2.....	3
Experiment 3.....	4
Experiment 4.....	4
Experiment 5.....	4
METHODOLOGY.....	4
Sites and Sampling.....	4
Nitrogen Fixation.....	6
Phospholipids.....	6
Microbial Enumeration.....	7
Ammonia Sensor.....	7
RESULTS AND DISCUSSIONS.....	9
CONCLUSIONS.....	15
REFERENCES.....	15

INTRODUCTION

Desert land Department of Defense (DOD) facilities encompass more than two million acres and provide a realistic training environment that is essential for maintaining military readiness. These include Navy and Marine Corps facilities at Marine Corps Air Ground Combat Center (MCAGCC) Twenty-Nine Palms, California, (595,367 acres), Marine Corps Base (MCB) Camp Pendleton, California, (125,000 acres), Naval Air Weapons Station (NAWS) China Lake, California, (608,190 acres), Yuma Marine Corps Air Station (MCAS), Arizona, (3,000 acres) and the associated Chocolate Mountain Aerial Gunnery Range at Niland, California, (460,000 acres), Marine Corps Logistics Base (MCLB) Barstow, California, (5,688 acres), Fallon Naval Air Station (NAS), Nevada, (7,982 acres), and the Army National Training Center, Fort Irwin, California, (640,000 acres).

However, desert ecosystems are extremely sensitive to disturbances, slow to recover, and they support unique organisms including threatened and endangered species. These concerns have resulted in a number of legislative initiatives aimed at protecting and preserving desert ecosystems. As a result, the DOD as stewards of these lands which also serve as de facto nature preserves has taken a proactive approach to the management of base ecosystems. Examples of this approach are DOD participation in the Mojave Ecosystem Database Program (MEDP) and the Strategic Environmental Research and Development Program (SERDP) sponsored land management research projects (e.g., the Army's Maneuver Impact Miles (MIM) and Integrated Dynamic Landscape Analysis and Modeling System (IDLAMS)). The objective of these programs is to develop tools that can be used to assess and manage training associated ecological impact.

The immediate and inevitable consequences of desert training impacts are the loss of plants, cryptogamic communities, and soil compaction. Since these impacts have been identified as the primary determinants of arid land degradation (Ray, 1995), the quantification of these events as land management tools is being pursued. However, the heterogeneity and dynamic nature of desert ecosystems including seasonal variability and successional changes brought about by fires, storms, and disease complicate the interpretation of these results (Schimel, 1995). Thus, there is a critical need for assessment tools that provide a real-time response to disturbance associated ecosystem impacts from which seasonal, spatial, and temporal variation can be filtered. An additional benefit would be the capability to provide ground-truth for approaches that use remote sensing.

Despite the seemingly impossible complexity that this requirement poses, Schlesinger (1991) and Schlesinger et al., (1990) have identified the interdependence of the biogeochemical cycles of carbon and nitrogen as a key element in assessing desert ecosystems. The biogeochemical cycles are the processes that shuttle elements between their inorganic (primarily CO₂ and N₂) and organic (e.g., proteins and nucleic acids) forms. The cycles are tightly coupled due to the carbon nitrogen stoichiometry which regulates the transfer of these elements within and between autotrophs (producers) and heterotrophs (consumers) and the overall regulation of these processes by water (Collatz et al., 1991).

Thus, selected aspects of the biogeochemical cycles of nitrogen and carbon and the structure of the bacterial community associated with them should be sensitive indicators of the ecological impact of training exercises. For example, Harrison et al., (1993) have shown that one of the consequences of agriculture induced disturbance of soil ecosystems is the loss of what they refer to as the fast turnover time pool of humic materials. Although the data are for

agricultural soils, the research demonstrates how large-scale disturbances are correlated with an essential soil component that is derived from biogeochemical cycling in the soil. Thinner and less robust desert soils appear to be even more susceptible to these types of disturbances.

Training associated destruction of plants and resultant decay releases carbon, phosphorus, and nitrogenous compounds to the soil. This influx of nutrients may initiate a shift in the microbial population of the soil, from nitrogen fixation and carbon storage to nitrogen release and carbon use. Klein et al., (1996) detected these types of responses in test plots that were amended with nitrogen. Additional studies have shown that adding nitrogen and carbon to an ecosystem affects succession and reestablishment of the plant community (McLendon and Redente, 1991; McLendon and Redente, 1992; Redente et al., 1992). Critical determinants of these processes are the nitrogen content (Mosier et al., 1991) and the phosphorus nitrogen ratio (Eisele et al., 1989). These processes are also sensitive to diffusive gas transport and the water content of the soil matrix (Adamsen and King, 1993), properties that are directly impacted by soil compaction associated with vehicle and troop traffic.

When military training impacts lead to plant loss, the rhizosphere bacterial communities are also greatly diminished or lost. The loss of root zone associated bacteria that are known to play an important role in nitrogen cycling and by association carbon cycling coupled with the spatial heterogeneity of desert ecosystems (Lajtha and Schlesinger, 1988; Skuji, 1984) may contribute to the slow recovery of desert ecosystems.

A particularly important aspect of the relationship between bacteria and plants is the biogeochemical cycling of nitrogen (Carter et al., 1995; Lajtha and Schlesinger, 1986; Herman et al., 1993; Hobbs et al., 1991). Plants also excrete compounds that promote the growth of beneficial bacteria and inhibit bacteria that may be detrimental (reviewed by Rovira, 1989). In all terrestrial ecosystems, numerous other bacteria essential to the health of the plants are closely associated with the plant root zone, (Newman, 1985; Parkin, 1993; Smith and Tiedje, 1979; Turner and Newman, 1984).

In training impacted desert soils, loss of plant cover and reduced percolation in compacted soils is associated with water stress (Pappendick and Campbell, 1981). In the saline alkaline desert soils typical of MCAGCC Twenty-Nine Palms, changes in the availability and quality of water may trigger changes in pH (Hall et al., 1995) and salinity (Galinski, 1995), both of which are known stressors.

As the effects of large-scale disturbances propagate through the ecosystem, they may lead to loss of species (biodiversity) and diversity within a species (genetic diversity). At the microbial level this includes groups of bacteria generically referred to as nitrogen fixing bacteria, methanotrophs, and nitrifiers, that perform essential functions related to the cycling of nitrogen and carbon. Loss of biodiversity and/or shifts in bacterial populations are linked to changes in the fluxes of nutrients and pollutants, and the stabilization of the physical environment. Consequently, disruption of the ecosystem at the microbial level is an unseen but insidious consequence of impacts on higher organisms. As a result, the overall health of the ecosystem may be compromised, higher plant and animal species are threatened, and the recovery of the ecosystem may be impeded. However, conducting a detailed biogeochemical assessment whenever training is scheduled is not practical.

The need for real-time data can be met by using sensors that measure biogeochemical associated processes directly (e.g., the concentration of ammonia and correlated with more detailed measurements derived from lab and field data). Alternatively, some parameters (e.g., soil compaction caused by vehicles and troops) might be measured with pressure transducers and

subsequently correlated with biogeochemical changes measured in the in the lab and field. Furthermore, sensors can be queried from remote locations, and the data incorporated directly into GIS based management tools supported by SERDP (e.g., IDLAMS) that provide ground truth for air-based remote sensing.

PROOF-OF-CONCEPT

In response to concerns raised by SERDP in response to the presentation of the full proposal at the SERDP SAB meeting held in September 1997, the following series of experiments (i.e., proof-of-concept) were proposed on 10 October 1997. This protocol was accepted and funded by SERDP with a start date of 01 January 1998. The experimental work was completed in April 1998 and presented to the SERDP at the In-progress Review held in May 1998.

Proposal: Investigate the potential for using biogeochemical characteristics of the Mojave Desert Ecosystem at MCAGCC Twenty-Nine Palms as one measure of the impact of military training exercises. Once developed, these measurements have the potential to be used in conjunction with a variety of tools (e.g., soil erosion modeling that are being developed in other laboratories for monitoring and assessing the impact of military training exercises on a variety of ecosystems).

Assumption: Training impacts are associated with detectable changes in biogeochemical parameters. To demonstrate this concept, laboratory experiments will focus on nitrogen fixation, ammonia flux, bacterial community structure, and chemical composition of soil collected at MCAGCC Twenty-Nine Palms.

Soil. Three soil samples (six total) will be collected from impacted and non-impacted areas at MCAGCC 29 Palms and the following experiments performed in triplicate on each soil sample. Since the intent is to demonstrate the validity of using biogeochemical assessments, adequate data will require this level of sampling and experimentation. The soil will be collected from sites identified by Ms. Sharon Jones and her colleagues at MCAGCC Twenty-Nine Palms.

Experiment 1. Bacterial community structure. Extract bacterial lipids from soil samples, convert to their methyl ester derivatives, chromatograph, and compare the fatty acid profiles using principle component analysis (PCA).

Rationale: Disturbance events cause the disruption and re-establishment of bacterial communities that can be distinguished by their fatty acid profiles.

Experiment 2. Rate of nitrogen fixation. Soil samples will be incubated with acetylene in closed containers. The headspace will be sampled at regular intervals and chromatographed to detect ethylene. The results will be expressed as millimoles of nitrogen fixed per day, per gram of soil (dry weight).

Rationale: Impacted and non-impacted lands exhibit nitrogen-dependent differences in biological community succession. Thus, impacted and non-impacted soils would be expected to exhibit differences in the rate and quantity of nitrogen fixed.

Experiment 3. Solid state ammonia sensor performance. Place the solid state ammonia electrode developed by American Research Corporation of Virginia (ARCOVA) in soil and determine the dynamic sensing range of the electrode. Various configurations of the sensor being developed by ARCOVA will be tested.

Rationale: Demonstrate that a solid state sensor can be used to measure the flux of ammonia when placed in the soil. This is a critical point since we propose to use these types of sensors to collect real-time data for ecosystem management.

Experiment 4. Rate of ammonia flux. Measure the rate of ammonia flux from soil samples by trapping with sulfuric acid, performing a Kjeldahl digestion and assay for ammonia.

Rationale: The Kjeldahl method is an accepted method for ammonia determinations and will be performed to confirm the results of the previous experiments and serve as an independent confirmation of the ammonia electrode results.

Experiment 5. Extract soil samples with chromatography grade water and use ion chromatography to analyze for major/minor cations and anions. Concentrations will be expressed as millimoles per gram of soil (dry weight).

Rationale: While the focus of the previous experiments is nitrogen, disturbance events may lead to changes in other chemical species. Exploratory analyses are necessary to investigate and identify biogeochemical parameters that can be used for ecosystem assessment.

The research reported was conducted in collaboration with ARCOVA who is developing sensors that are inexpensive and can withstand the rigors of the training environment. These sensors use a laser diode and cost between \$20 and \$50. By modifying the surface, they can be tailored to respond to a variety of chemical species. For this phase of the project, ARCOVA developed and tested an ammonia sensor.

METHODOLOGY

Sites and Sampling. An undisturbed pristine site that served as the control is shown in Figure 1 and a sampling point is shown in Figure 2. Figure 3 shows an area that is used for tank training exercises was used as a disturbed area. The mound in the background with PVC pipes inserted in it is a creosote revegetation project.




Figure 1. Pristine desert at MCAGCC 29 Palms from which soil representative of an undisturbed site was obtained.




Figure 2. Undisturbed soil sample location.




Figure 3. Heavily disturbed soil sampling. This disturbance is caused by tanks and other vehicles and is representative of environmental degradation associated with desert training activity. The PVC pipes in the background are an effort to revegetate adjacent areas with mesquite.

Since the focus of this phase of the project is the subsurface, samples were taken from a depth of 2 to 6 inches following removal of the top 1-inch of soil. Three samples were collected at each site and all analyses were run in triplicate on each sample. Average values were calculated and compared using the Student's T-Test to calculate a two-tailed "P" value. Soil collected from a third site that has not been used for training for several years was sent to ARCOVA to test the ammonia sensor.

Nitrogen Fixation. Soil samples were incubated in serum bottles containing acetylene, and the headspace sampled and chromatographed to quantify ethylene (Balderston et al., 1976; Bedard and Knowles, 1989; Fireston and Davidson, 1989; Tiedje, 1988).

Phospholipids. Gas chromatography of lipids extracted from soil samples was used to estimate microbial biomass (Vance, et al., 1987; Zelles et al., 1994; Zelles et al., 1995), the distribution of eukaryotes and prokaryotes, and as indicators of the stress experienced by the gram-negative bacteria (Balkwill, et al., 1988; Frostegard et al., 1991; Gillan and Hogg, 1984; Guckert et al., 1985; Peterson and King, 1994; Vestal and White, 1989). Atlas (1984) first described changes in microbial phospholipids in response to a variety of stressors. Numerous additional examples (Brown, 1990; Frostegard et al., 1993a; Frostegard et al., 1993b; McKinley and Vestal., 1984) have demonstrated the value of this method for assessing the status and

composition of the indigenous microbial community in response to environmental insults. This work was conducted in collaboration with Mr. Greg Davis at Microbial Insights, Inc.

Microbial Enumeration. Total heterotrophs were enumerated using the fifteen tube most probable number method in a basal salts medium developed specifically for growing and enumerating bacteria at Fallon Naval Air Station and MCAGCC Twenty-Nine Palms and used in previous bioremediation work (Table 1). To enumerate heterotrophs, the medium was amended with 0.025% (w/v) each of yeast extract and casamino acids.

Table 1. Basal Salts medium used to enumerate bacteria in alkaline desert soils characteristic of MCAGC 29 Palms. The composition of the medium is based on an analysis of desert soils found at Fallon Naval Air Station and MCAGCC 29 Palms.

COMPOUND	MOLECULAR WEIGHT	CONCENTRATION mM	GRAMS/LITER
CaCl ₂ ·2H ₂ O	147.02	0.5	0.0735
NaCl	58.44	100	5.844
KH ₂ PO ₄	136.1	2.5	0.3402
KNO ₃	101.1	5	0.5055
MgSO ₄ ·7H ₂ O	246.47	0.5	0.1232
Na ₂ SO ₄	142.04	50	7.102
Na ₂ CO ₃	105.99	5	0.530

The pH is adjusted to 8.8 to 8.9 with sulfuric acid and filter sterilized. To prepare Fallon Nutrient Broth the medium is amended with 0.025 g/L each of Difco Yeast Extract and Casamino Acids.

Ammonia Sensor. Construction and testing of the ammonia sensor was conducted by American Research Corporation of Virginia. The test system is illustrated in Figure 4 and the ammonia sensor is illustrated in Figure 5.




Figure 4. Experimental setup for the detection of ammonia using a solid state sensor. The sensor uses a photodiode to capture the total internal reflectance fluorescence (PD/TIRF)




Figure 5. Ammonia Sensor. The flow cell was placed on top of a column 3 cm wide by 11 cm deep packed with sand from MCAGCC Twenty-Nine Palms and gassed with ammonia mixed with nitrogen. The polymeric film is selectively permeable to ammonia and contains a dye that changes its absorption coefficient when exposed to ammonia. A laser is used to illuminate the dye and a photodiode detector is used to capture the absorption spectrum.

RESULTS AND DISCUSSION

The size of the microbial population in each soil sample was estimated from the total mass of extracted phospholipid fatty acids (PLFA). These data are shown in Figure 6 and suggest a larger microbial population in the soil samples from the disturbed site (i.e., 186,349 picomoles PLFA from the disturbed site vs 11,091 picomoles PLFA from the pristine site). When heterotrophs were enumerated in these soil samples using the MPN method, the average number of heterotrophs in the disturbed and pristine samples were $1.5 \times 10^7 \pm 2.1 \times 10^6$ and $1.04 \times 10^7 \pm 2.1 \times 10^6$ bacteria/g soil (dry weight), $P = 0.0478$, respectively. These data support the idea that the release of nutrients associated with the destruction of the vegetation supports an increase in the bacterial population.

An analysis of fatty acids characteristic of the growth phase of gram-negative bacteria also supports the proposal that the release of nutrients associated with the destruction of plants at the disturbed site promotes bacterial growth (Figure 7). However, in the samples from the disturbed site there is an increase in PLFAs characteristic of gram-negative bacteria exposed to environmental stressors (Figure 8). This observation is consistent with the proposal that soil compaction, destruction of plants, and other disturbance events stress the microbial ecosystem.




Figure 6. Biomass content of soil from impacted, samples 1, 3, and 5 and pristine, samples 2, 4, and 6.




Figure 7. Fatty acids characteristic of Gram-negative bacteria suggest that the bacteria are in stationary phase in all samples. However, the higher ratio of cy/ ω 7c in samples from the pristine site suggest that bacteria at this site turn over more slowly than the organisms at the impacted site.




Figure 8. Analysis of fatty acids associated with environmental stressors in gram-negative bacteria suggests the bacteria at the disturbed site are more highly stressed environment than the pristine site.

Characteristic phospholipids were used to estimate the microbial composition of each soil sample (Figure 9). These data suggest a fairly uniform distribution of prokaryotes and eukaryotes among all the samples but (see the legend on Figure 9) there are significantly fewer prokaryotes and eukaryotes at the pristine site. However, the dominant gram-negative bacteria belong to the genera *Actinomycetes* and *Desulfovibrio*. The former organisms are common in soil and sulfate-reducing bacteria would be expected in the saline sulfate rich desert soils. However, principle component analysis (Figure 10) of these data demonstrate that there is a distinctly unique distribution of bacteria at each site.

Additional differences between the soil samples were found in the flux of ammonia from the soil (Figure 11) and nitrogenase activity (Figure 12). The lower flux of ammonia from the impacted soil is consistent with the increase in biomass which would be expected to capture the ammonia for use as a molecular building block. The availability of ammonia in the disturbed site would also be expected to suppress nitrogen fixation which is supported by the data in Figure 12. Analysis of soil samples also supports the differences in the availability of nitrogen at the disturbed site compared to the pristine site (Table 2).

Performance of the solid state ammonia sensor when tested with 40 ppb and 20 ppb of ammonia are shown in Figures 13 and 14, respectively. As the ammonia passes through the column, the concentration may be reduced due to adsorption and chemical reactivity. Thus, the concentration at the sensor may be less than the input concentration. However, the response of the sensor to 20 ppb ammonia is within the range of ammonia flux expected from desert soils, this study is shown Figure 11 and Schlesinger et al., (1990).




Figure 9. Percent composition of the soil samples with respect to major groups of organisms as determined by phospholipid fatty acid analysis. In the disturbed soil samples the average picomoles of bacterial and eukaryote PLFAs are 4,538 and 1,705 respectively. In the pristine soil the values are 290 and 86 respectively. The abbreviations are; TerBrSats tertiary branched saturated fatty acids, Monos monounsaturated fatty acids, MidBrSats midbranched saturated fatty acids, Nsats, normal saturated fatty acids, and fatty acids characteristic of eukaryotes.




Figure 10. Principle component analysis of the PLFA of the samples from the impacted and non-impacted soil samples suggests that the organisms fall into separate groups.




Figure 11. Volatilization of ammonia from soil samples. A column 10 cm in diameter was filled with soil to a depth of approximately 20 cm and ammonia captured in 2N sulfuric acid for 24 h. Ammonium ion was analyzed following Kjeldahl digestion. The value for each soil sample is the average of three measurements. The averages are 21.9 ± 6.95 and 30.1 ± 6.47 for the impacted (samples 1, 3, and 5) and non-impacted (samples 2, 4, and 6) respectively. The two-tailed "P" value is 0.0194.




Figure 12. Average values of nitrogenase activity in impacted and non impacted soils are 3 ± 1.04 and 6 ± 2.05 respectively. The two-tailed "P" value is 0.0012. Approximately 50 g of soil was placed in a 120 ml serum bottle capped with a teflon faced rubber serum stopper. Acetylene (4 ml) generated from calcium carbide was injected and the vials incubated at room temperature. Ethylene was quantified by injecting 300 ul of headspace sample into a GC with an FID detector. Each soil sample was assayed in triplicate.

Table 2. Average values of selected parameters in pristine and disturbed soil samples. Each value is the average of nine analyses conducted on three soil samples from each site. The Student T Test was used to calculate the 2-tailed P value.

PARAMETER	AVERAGE \pm STANDARD DEVIATION ¹		2-TAILED P VALUE
	PRISTINE SITE	DISTURBED SITE	
% Moisture	2.0 \pm 0.81	2.0 \pm 0.20	0.875
pH	8.35 \pm 0.11	8.3 \pm 0.12	0.394
Phosphate	0.47 \pm 0.25	0.56 \pm 0.51	0.647
Nitrogen	45.6 \pm 4.88	64.6 \pm 8.70	<0.0001
Magnesium	1.3 \pm 0.15	1.3 \pm 0.21	0.8
Calcium	18.6 \pm 7.9	25.9 \pm 8.1	0.07
Sulfate	1290 \pm 256	1090 \pm 267	0.121
Sodium	1840 \pm 376	1854 \pm 322	0.937
Chloride	1250 \pm 135	1310 \pm 267	0.582
Nitrate	10.7 \pm 3.39	5.0 \pm 2.12	6.0
Manganese	10.3 \pm 4.4	10 \pm 1.6	0.832
Potassium	137 \pm 6.3	128 \pm 26.7	0.37

¹Concentrations expressed as mg/Kg soil dry weight




Figure 13. Response of the ammonia sensor to 41 ppb of ammonia.




Figure 14. Response of the ammonia sensor to 20 ppb of ammonia.

CONCLUSIONS

The data support the premise in the original proposal and the scaled down proof-of-concept which is that biogeochemical markers can be used to monitor the ecological impact of military training. Specifically, the data presented here confirm that there are measurable and statistically significant ($P \leq 0.05$) differences in the flux of ammonia and rate of nitrogen fixation between an impacted and pristine site.

These data can be interpreted in terms of the destruction of vegetation and release of nutrients. Thus, there is the potential for correlating these types of measurements with remote monitoring of vegetation cover. In addition, the data suggest significant differences in the composition of the microbial population at the two sites which may be associated with additional biogeochemical differences and/or correlated with compaction, soil moisture, and/or oxygen concentration in the soil as discussed in the original proposal. Furthermore, the performance of the solid state ammonia sensor demonstrates the potential for using this technology to monitor and collect real-time data from large areas.

REFERENCES

- Adamsen, A. P. S. and G. M. King. 1993. Methane consumption in temperate and subarctic forest soils: rates, vertical zonation, and responses to water and nitrogen. *Appl. Environ. Microbiol.* 59:485-490.
- Atlas, R. M. 1984. Use of microbial diversity measurements to assess an environmental stress. p. 540-545. *In* K. C. Marshall (ed.), *Current perspectives in microbial ecology*. American Society for Microbiology, Washington, D.C.
- Balderston, W. L., B. Sherr, and W. J. Ryan. 1976. Blockage by acetylene of nitrous oxide reduction in *Pseudomonas perfectomarinus*. *Appl. Environ. Microbiol.* 31:504-508.
- Balkwill, D. L., F. R. Leach, J. T. Wilson, J. F. McNabb, and D. C. White. 1988. Equivalence of microbial biomass measures based on membrane lipid and cell wall components, adenosine triphosphate, and direct counts in subsurface aquifer sediments. *Microb. Ecol.* 16:73-84.
- Bedard, C. and R. Knowles. 1989. Physiology, biochemistry, and specific inhibitors of CH_4 , NH_4^+ , and CO consumption by methanotrophs and nitrifiers. *Microbiol. Rev.* 53:68-84.
- Brown, A. D. 1990. *Microbial water stress physiology: principles and applications*. John Wiley and Sons Ltd., West Sussex, England.
- Carter, J. R., Y. H. Hsiao, S. Spiro, and D. J. Richardson. 1995. Soil and sediment bacteria capable of aerobic nitrate respiration. *Appl. Environ. Microbiol.* 61:2852-2858.

- Collatz, G. J., C. Griver, and J. T. Ball. 1991. Physiological and environmental regulation of stomatal conductance, photosynthesis, and transpiration: a model. *Agric. Forest Meteorol.* 54:107-136.
- Eisele, K. A., D. S. Schimel, L. A. Kapustka, and W. J. Parton. 1989. Effects of available P and N:P ratios on non-symbiotic dinitrogen fixation in tallgrass prairie soils. *Oecologia* 79:471-474.
- Firestone, M. K. and E. A. Davidson. 1989. Microbiological basis of NO and N₂O production and consumption in soil, p. 7-21. *In* M. O. Andreae and D. S. Schimel (ed.), Exchange of trace gases between terrestrial ecosystems and the atmosphere. John Wiley & Sons, Ltd., Chichester, England.
- Frostegard, A., E. Baathe, and A. Tunlid. 1991. Microbial biomass measured as a total lipid phosphate in soils of different organic content. *J. Microbiol. Methods* 14:151-163.
- Frostegard, A., E. Baathe, and A. Tunlid. 1993a. Shifts in the structure of soil microbial communities in limed forests as revealed by phospholipid fatty acid analysis. *Soil Biol. Biochem.* 25:723-730.
- Frostegard, A., A. Tunlid, and E. Baath. 1993b. Phospholipid fatty acid composition, biomass, and activity of microbial communities from two soil types experimentally exposed to different heavy metals. *Appl. Environ. Microbiol.* 59:3605-3617.
- Galinski, E. A. 1995. Osmoadaptation in bacteria. *Adv. Microbial Physiol.* 37:273-328.
- Gillan, F. T. and R. W. Hogg. 1984. A method for the estimation of bacterial biomass and community structure in mangrove-associated sediments. *J. Microbiol. Methods* 2:275-293.
- Guckert, J. B., C. P. Antworth, P. D. Nichols and D. C. White. 1985. Phospholipid ether-linked fatty acid profiles as reproducible assays for changes in prokaryotic community structure of estuarine sediments. *FEMS Microbiol. Ecol.* 31:794-801.
- Hall, H. K., K. L. Karem and J. W. Foster. 1995. Molecular responses of microbes to environmental pH stress. *Adv. Microbial Physiol.* 37:229-272.
- Harrison, K. G., W. S. Broecker, and G. Bonani. 1993. The effect of changing land use on soil radiocarbon. *Science* 262:725-726.
- Herman, R. P., K. R. Provencio, R. J. Torrez, and G. M. Seager. 1993. Effect of water and nitrogen additions on free-living nitrogen fixer populations in desert grass root zones. *Appl. Environ. Microbiol.* 59:3021-3026.
- Hobbs, N. T., D. S. Schimel, C. E. Owensby, and D. S. Ojima. 1991. Fire and grazing in the tallgrass prairie: contingent effects on nitrogen budgets. *Ecology* 72:1374-1382.

Klein, D. A., T. McLendon, M. W. Paschke, and E. F. Redente. 1996. Nitrogen availability and fungal-bacterial responses in successional semiarid steppe soils. *Arid Soil Research and Rehabilitation* **10**:321-332.

Lajtha, K. and W. H. Schlesinger. 1986. Plant response to variations in nitrogen availability in a desert shrubland community. *Biogeochemistry* **2**:29-37.

McKinley, V. L. and J. R. Vestal. 1984. Biokinetic analyses of adaptation and succession: microbial activity in composting municipal sewage sludge. *Appl. Environ. Microbiol.* **47**:933-941.

McLendon, T. and E. F. Redente. 1991. Nitrogen and phosphorous effects on secondary succession dynamics on a semi-arid sagebrush site. *Ecology* **72**:2016-2024.

McLendon, T. and E. F. Redente. 1992. Effects of nitrogen limitation on species replacement dynamics during early secondary succession on a semiarid sagebrush site. *Oecologia* **91**:312-317.

Mosier, A., D. Schimel, D. Valentine, K. Bronson, and W. Parton. 1991. Methane and nitrous oxide fluxes in native, fertilized and cultivated grasslands. *Nature* **350**:330-332.

Newman, E. I., 1985. The rhizosphere: carbon sources and microbial populations, p. 107-122. *In* A. H. Fitter (ed.), *Ecological interactions in soil-plants, microbes, and animals*. Blackwell Scientific Publications, Oxford.

Pappendick, R. I. and G. S. Campbell. 1981. Theory and measurement of water potential, p. 1-22. *In* J. F. Parr, W. R. Gardner, and L. F. Elliott (ed.), *Water potential relations in soil microbiology*. Soil Science Society of America, Madison, Wis.

Parkin, T. B. 1993. Spatial variability of microbial processes in soil - a review. *J. Environ. Qual.* **22**:409-417.

Peterson, S. O. and M. J. Klug. 1994. Effects of sieving, storage, and incubation temperature on the phospholipid fatty acid profile of a soil microbial community. *Appl. Environ. Microbiol.* **60**:2421-2430.

Ray, T. W. 1995. Remote monitoring of land degradation in arid/semiarid regions, Ph.D. thesis, California Institute of Technology, 415 pp. Available on-line at <http://www.planetary.caltech.edu>

Redente, E. F., J. E. Friedlander, and T. McLendon. 1992. Response of early and late semiarid seral species to nitrogen and phosphorus gradients. *Plant and Soil.* **140**:127-135.

Rovira, A. D. 1989. Plant root exudates. *Bot. Rev.* **35**:35-57.

Schimel, D. S. 1995. Terrestrial biogeochemical cycles: Global estimates with remote sensing. *Remote Sens. Environ.* **51**:49-56.

Schlesinger, W. H. 1991. Biogeochemistry: an analysis of global change. Academic Press, Inc. New York, NY.

Schlesinger, W. H., J. F. Reynolds, G. L. Cunningham, L. F. Huenneke, W. M. Jarrell, R. A. Virginia and W. G. Whitford. 1990. Biological feedbacks in global desertification. *Science* 247:1043-1048.

Skuji, J. 1984. Microbial ecology of desert soils, p. 49-91. *In* Marshall, C. C. (ed.), *Advances in microbial ecology*. Plenum Press, New York, NY.

Smith, M. S. and J. M. Tiedje. 1979. The effect of roots on soil denitrification. *Soil Sci. Am. J.* 43:951-955.

Tiedje, J. M. 1988. Ecology of denitrification and dissimilatory nitrate reduction to ammonium, p. 179-245. *In* A. J. B. Zehnder (ed.), *Biology of anaerobic microorganisms*. Wilder, New York, N. Y.

Turner, S. M. and E. I. Newman. 1984. Growth of bacteria on the roots of grasses: influence of mineral nutrient supply and interactions between species. *J. Gen. Microbiol.* 130:505-512.

Vance, E. D., P. C. Brookes and D. S. Jenkinson. 1987. An extraction method for measuring biomass C. *Soil Biol. Biochem.* 19:703-707.

Vestal, J. R. and D. C. White. 1989. Lipid analysis in microbial ecology. *BioScience* 39:535-541.

Zelles, L., Q. Y. Bai, R. M. Ma, R. Rackwitz, K. Winter and F. Beese. 1994. Microbial biomass, metabolic activity and nutritional status determined from fatty acids patterns and polyhydroxybutyrate in agriculturally managed soils. *Soil Biochem.* 19:115-123.

Zelles, L., Q. Y. Bai, R. Rackwitz, D. Chadwick, K. Winter and F. Beese. 1995. Determination of phospholipid- and lipopolysaccharide- derived fatty acids as an estimate of microbial biomass and community structures in soils. *Biol. Fertil. Soils* 14:87-98.