

5199

23

OTS: 60-41092

JPRS: 5199

3 August 1960

THE DYNAMICS OF LARGE-SCALE ATMOSPHERIC PROCESSES

By N. I. Bulevov and H. I. Martschuk

-US3R-

DISTRIBUTION STATEMENT A
Approved for Public Release
Distribution Unlimited

20000406 181

Distributed by:

OFFICE OF TECHNICAL SERVICES
U. S. DEPARTMENT OF COMMERCE
WASHINGTON 25, D. C.

REPRODUCED BY
**NATIONAL TECHNICAL
INFORMATION SERVICE**
U. S. DEPARTMENT OF COMMERCE
SPRINGFIELD, VA. 22161

U. S. JOINT PUBLICATIONS RESEARCH SERVICE
205 EAST 42nd STREET, SUITE 300
NEW YORK 17, N. Y.

Reproduced From
Best Available Copy

80P

JPRS: 5199
CSO: 4599-N

THE DYNAMICS OF LARGE-SCALE ATMOSPHERIC PROCESSES

(This analysis was made in 1951)

[This is a translation of an article by N.I. Bulayev and H.I. Martshuk, in Publication No 2, Institute for Physics of the Atmosphere, USSR Academy of Sciences; CSO: 4599-N]

Introduction

A complete conception of large-scale atmospheric processes determining the nature of the weather in many respects, can only be obtained by the inclusion of all basic factors in the baroclynic atmosphere.

In the analysis of large-scale atmospheric processes we find it necessary to formulate mathematical theories which show the nature and evolution of the real fields of meteorological elements as exactly as possible. In principle, the mathematical constructions based on the application of the laws of hydrothermodynamics in atmospheric conditions permit the pertinent system of nonlinear equations to determine the meteorological elements of interest to us. Therefore, in forecasting meteorological elements the solution of the complete system of nonlinear hydrothermodynamic equations appears to be the fundamental assignment for meteorological theory.

The attempt to obtain weather forecasts the theoretical way by studying the system of hydrodynamic equations was made for the first time by Richardson (9) in 1922. Richardson did not obtain sufficient results because he did not tackle the problem the right methodical way. Richardson took the initial wind field as the starting fact and hoped to obtain the evolution of atmospheric currents with time on the basis of the solution of the equation system. Inasmuch as the initial field of velocity generally includes manifold perturbations of all kinds of scales, Richardson should have taken into account the evolution of all types of small (sound, gravitational) perturbations together with the evolution of large-scale perturbations which are of meteorological interest, although the former are insignificant in meteorological respects. The description of the small perturbations require the utilization of correspondingly small scales and small time integrals. The length and time scales used by Richardson, were much larger than the degree of exactitude required; therefore, the sole forecast calculated by Richardson did not stand the test.

The first feasible solution of the equation system for atmospheric processes was found by I.A. Kibel (2) in 1940. He based his analysis on the assumption that large-scale atmospheric processes present themselves as quasi-geostrophic processes. This permitted I.A. Kibel to take the initial fields of pressure and temperature as starting factors and not the

initial wind field; his problem was the calculation of changing pressure and temperature fields in time.

However, in I.A. Kibel's analysis the possibilities of quasi-geostrophic samples were not completely exhausted. Therefore, the temperature derivative of time $\frac{\partial T}{\partial t}$ was only determined with a calculation of the horizontal mass transfer, and in the pressure change $\frac{\partial p}{\partial t}$ the convergence factor and the divergence of the air in the atmosphere were not considered.

In other meteorological analyses of the Friedmann-Kotshin School, the system of hydrothermodynamic equations was used more efficiently; the vortex velocity equation was examined as a promoting equation connecting the complete alteration of the vortex with the divergence of the air on the horizontal plane.

In 1941 M.E. Shvetz (6) obtained the expression for the vertical velocity in the shape of the integral according to altitude from the individual derivative of the vortex velocity.

In 1943 E.N. Blinova (1) managed to obtain the prognostic equations for pressure and temperature T on a certain "average" atmospheric level through linearization, taking into account the sphericity of the surface of the earth. E.N. Blinova started with the condition of maintenance of vorticity of ~~masses~~ ^{PARCELS} on the mean level and with the assumption that temperature changes in the atmosphere depend only upon the horizontal transfer of homogeneous masses of different temperatures. E.N. Blinova integrated the system of linear equations obtained in this manner, under general conditions of the elementary field:

Foreign meteorologists see the possibility of utilizing hydrodynamic equations for weather forecasts in a different way.

N. Ertel (8) analysed the equation ¹

$$-\frac{\partial \Delta p}{\partial t} = \frac{1}{L_p} (p, \Delta p)$$

obtained by him starting with the nature of the area of dependence of equation $\frac{\partial p}{\partial t}$ upon the field of function $(p, \Delta p)$; the result was a meteorologically incorrect conclusion according to which it is impossible to calculate the future pressure map in advance. The deficiency was clearly reflected in the results of A.M. Obuchov's (3) analysis published in 1949 in which the author obtained a more complete equation in the examination of the barotrope sample of the atmosphere:

$$\frac{1}{L_1^2} \frac{\partial p}{\partial t} - \Delta \frac{\partial p}{\partial t} = \frac{1}{L_p} (p, \Delta p),$$

$L_1 = \frac{\sqrt{gH_0}}{l}$; H_0 - altitude of homogeneous atmosphere; the solution of this equation is

$$\frac{\partial p}{\partial t} = \int_0^{\infty} \frac{1}{L_p} (p, \Delta p), K_0(r) r dr,$$

¹ We will keep to the symbols used in the first part of this analysis.

$$r = \frac{\sqrt{x^2 - y^2}}{L_1}, \quad K_0(r) - \text{Macdonald function of zero order.}$$

In 1950 N.I. Bulajev obtained ~~an equation~~ ^{the equation} for the change of pressure in the baroclinic atmosphere under ^{the condition of a} linear temperature ^{decrease with height}. The equation for the pressure change on the mean level of the troposphere was obtained as follows:

$$\frac{\partial p}{\partial t} - k_1 \Delta \frac{\partial p}{\partial t} = k_2(p, \Delta p) - k_3(\bar{T}, \Delta \bar{T}),$$

\bar{T} - average temperature of troposphere; k_1, k_2, k_3 - a certain constant quantity.

This analysis also brought us the formula for the calculation of vertical velocity at different levels of the atmosphere:

$$g\rho w = -a_1(z)\Delta(\bar{T}, p) - a_2(z) \left[(\bar{T}, \Delta p) + (p, \Delta \bar{T}) \right],$$

p - pressure at the mean level of atmosphere; $a_1(z), a_2(z)$ - the coefficient dependent on the altitude.

Charney's (7) analysis examined the barotropic sample of the atmosphere and the simple baroclinic sample. The author uses the equation for the winds in geostrophic approximation and the equation of the heat flux without considering the vertical current. Charney obtained Poisson's equation for the

derivative of pressure at different levels. The resulting equation in the barotropic case was integrated according to the method of finite difference with the aid of electronic equipment. To integrate the equation, Charney was forced to supplement artificial boundary conditions, pressure values and Laplacian pressure as time functions at the boundary of the area.

The baroclinic atmosphere sample was examined in the analysis published by I.A. Kibel (1953). The author started with the equation of the vortex in geostrophical approximation and with the equation of the heat influx without considering the vertical current. I.A. Kibel integrated Poisson's equation for the desired function $-\frac{\partial z}{\partial t}$ on surfaces $p = \text{const}$ and obtained the solution as follows:

$$\frac{\partial z}{\partial t} = \frac{1}{2\pi} \int_0^{2\pi} \int_0^{r_1} F(r, \phi, p) \ln \frac{r_1}{r} r dr d\phi - \overline{\frac{\partial z}{\partial t}},$$

$F(r, \phi, p)$ - the known function of the fields of meteorological elements in the entire atmosphere, $\overline{\frac{\partial z}{\partial t}}$ - the mean value of the desired value at level p up to the circumference of radius r_1 . Value $\overline{\frac{\partial z}{\partial t}}$ with sufficiently large r_1 (1,000km) appears small and may be eliminated. Excluding the average value of function $F(r, \phi, p)$ under the integral symbol and integrating the function of influence $\frac{r_1}{r}$ I.A. Kibel obtained the prognostic formula

$$\frac{\partial z}{\partial t} = m\bar{F}(r, \phi, p),$$

$m = \frac{r^2}{L}$. For purposes of simplification it was assumed that function $\bar{F}(r, \phi, p)$ equals $F(r, \phi, p)$ in regard to the examined point.

In this analysis the authors intended to obtain general equations for pressure and temperature changes and an equation for the vertical currents which takes all basic factors of the baroclinic atmosphere into account. Compared to previous analyses the complete inclusion of dynamic factors in the temperature changes appears to be a new factor.

In the integration of the obtained differential equations the authors found solutions which were expressed by space integrals ^{of the} determinate ^{which} expressions ^{ed} depending upon the distribution of pressure fields and temperature in space as well as the congruent influence functions.

1. Formulation of Problem

In the study of the dynamics of pressure and temperature changes and the formation of vertical motion in the baroclinic atmosphere we used the system of hydrothermodynamic equations in regard to atmospheric processes on a large scale. A chara

teristic feature of this system which distinguishes it from conventional hydrodynamic equations, appears to be the existence of the deflective force of the rotation of the earth in the equations of motion.

The local study of space regions with horizontal dimensions over several thousand kilometers permits us to consider the surface of the earth as a level surface subject to the limitations of the region under study and the use of the rectangular coordinate system in the initial equations.

The analysis excludes regions in the immediate proximity of the equator ($\phi = 0-30^\circ$) because the nature of the motion shows fundamental changes in these regions.

The initial system of hydrodynamic equations is assumed as follows:

Equations of Motion

$$\frac{du}{dt} = -\frac{1}{\rho} \frac{\partial p}{\partial x} + lv + \frac{1}{\rho} \frac{\partial}{\partial z} \mu \frac{\partial u}{\partial z}, \quad (1)$$

$$\frac{dv}{dt} = -\frac{1}{\rho} \frac{\partial p}{\partial y} - lv + \frac{1}{\rho} \frac{\partial}{\partial z} \mu \frac{\partial v}{\partial z}, \quad (1)$$

Equation of Statistics

$$\frac{\partial p}{\partial z} = -g\rho, \quad (1)$$

Equation of Continuity

$$\frac{\partial \rho}{\partial t} + \frac{\partial \rho u}{\partial x} + \frac{\partial \rho v}{\partial y} - \frac{\partial \rho w}{\partial z} = 0,$$

Equation of Heat Influx

$$\frac{dT}{dt} = \frac{\lambda_a}{g f} - \frac{dp}{dt} = \frac{\epsilon}{c_p}, \quad (1)$$

Equation of State

$$p = \rho RT. \quad (2)$$

The following symbols were used in the equations (1.1) and (1.6):

u, v, w - components of velocity vector \vec{V} to coordinate axis;
 p - pressure, ρ - density, T - air temperature, $l = 2\omega \cos \theta$
 ω - angular velocity of earth rotation, θ - supplement to local latitude, μ - coefficient of turbulence, g - acceleration due to gravity, R - gas constant, ϵ - flux of heat per mass unit dependent upon radiation and transition of water in atmosphere, from one phase state to the other, c_p - specific air under constant pressure, λ_a - adiabatic temperature gradient.

Boundary conditions and initial data are indispensable for the complete determination of the problem.

As boundary conditions with the earth surface we take the transformation of the vertical velocity to zero

$$w = 0 \quad \text{with} \quad z = 0$$

and the conditions of the free surface at the upper limit of the atmosphere

$$\frac{dp}{dt} \rightarrow 0 \quad \text{with} \quad z \rightarrow \infty. \quad (1)$$

Elementary data are pressure fields and temperatures at $t =$ $p(x, y, z)$ and $T(x, y, z)$.

It is advisable to go over to the coordinate system x', y', p, t' , where pressure p is assumed to be an independent variable; axes x' and y' are on the isobaric surfaces. The altitude of the isobaric surfaces z is now considered a function of coordinates x', y', p and time t' .

The transition from the elementary system of the coordinates x, y, z, t to the new system x', y', p, t' is realized through the following changes:

$$\left. \begin{aligned} \frac{\partial}{\partial x'} &= \frac{\partial}{\partial x} - \frac{\partial p}{\partial x} \frac{\partial}{\partial p}, & \frac{\partial}{\partial y'} &= \frac{\partial}{\partial y} - \frac{\partial p}{\partial y} \frac{\partial}{\partial p}, \\ \frac{\partial}{\partial p} &= -\frac{1}{g\rho} \frac{\partial}{\partial z}, & \frac{\partial}{\partial t'} &= \frac{\partial}{\partial t} - \frac{\partial p}{\partial t} \frac{\partial}{\partial p}, \end{aligned} \right\} \quad (2)$$

The result is that

$$\frac{\partial p}{\partial x} = g\rho \frac{\partial z}{\partial x'}, \quad \frac{\partial p}{\partial y} = g\rho \frac{\partial z}{\partial y'}, \quad \frac{\partial p}{\partial t} = g\rho \frac{\partial z}{\partial t'}. \quad (3)$$

In the new system the time derivative of any element has the following form:

$$\frac{d}{dt} = \frac{\partial}{\partial t} + u \frac{\partial}{\partial x} + v \frac{\partial}{\partial y} + w \frac{\partial}{\partial z} = \frac{\partial}{\partial t'} + u \frac{\partial}{\partial x'} + v \frac{\partial}{\partial y'} + \frac{dp}{dt} \frac{\partial}{\partial p} \quad (1.1)$$

This shows that in the system of coordinate x', y', p, t value $w = \frac{dp}{dt}$ plays the part of vertical velocity; this value gives the position of the air unit in regard to the isobaric surface. The transition from w to $\frac{dp}{dt}$ is realized through the correlation

$$w = \frac{\partial z}{\partial t} + u \frac{\partial z}{\partial x} + v \frac{\partial z}{\partial y} = \epsilon p w$$

or

$$w = \frac{1}{\epsilon p} \left(\frac{\partial z}{\partial t} + u \frac{\partial z}{\partial x'} + v \frac{\partial z}{\partial y'} \right) \quad (1.2)$$

Equations (1.1) and (1.2) in the new coordinate system are as follows:

$$\frac{\partial u}{\partial t'} + u \frac{\partial u}{\partial x'} + v \frac{\partial u}{\partial y'} + \epsilon \frac{\partial u}{\partial p} = - \epsilon \frac{\partial z}{\partial x'} + (v + \epsilon^2 \frac{\partial}{\partial p} p^2) \frac{\partial u}{\partial p} \quad (1.3)$$

$$\frac{\partial v}{\partial t'} + u \frac{\partial v}{\partial x'} + v \frac{\partial v}{\partial y'} + \epsilon \frac{\partial v}{\partial p} = - \epsilon \frac{\partial z}{\partial y'} + (u - \epsilon^2 \frac{\partial}{\partial p} p^2) \frac{\partial v}{\partial p} \quad (1.4)$$

$$\frac{\partial z}{\partial p} = -\frac{1}{g\rho} \quad (1.1)$$

In the system of the x', y', p, t' coordinate the equation of continuity (1.4) can be greatly simplified. To this end we use the equation of statics (1.3) and correlations (1.12) and (1.9). Considering that

$$\frac{\partial \rho}{\partial s} = -\frac{1}{g} \frac{\partial}{\partial z} \left(\frac{\partial p}{\partial s} \right) \quad (s = x, y, t),$$

$$\frac{\partial \rho}{\partial t} + \frac{\partial \rho v}{\partial y} = \frac{1}{g} \frac{\partial}{\partial z} \left(u \frac{\partial p}{\partial x} + v \frac{\partial p}{\partial y} \right) + \rho \frac{\partial \tau}{\partial p}$$

and

$$\begin{aligned} \frac{\partial \rho u}{\partial x} + \frac{\partial \rho v}{\partial y} &= \rho \left(\frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} \right) - \frac{1}{g} \left(u \frac{\partial}{\partial z} \frac{\partial p}{\partial x} + v \frac{\partial}{\partial z} \frac{\partial p}{\partial y} \right) = \\ &= \rho \left(\frac{\partial u}{\partial x'} + \frac{\partial v}{\partial y'} \right) + \rho \left(\frac{\partial u}{\partial p} \frac{\partial p}{\partial x} + \frac{\partial v}{\partial p} \frac{\partial p}{\partial y} \right) - \frac{1}{g} \left(u \frac{\partial}{\partial z} \frac{\partial p}{\partial x} + v \frac{\partial}{\partial z} \frac{\partial p}{\partial y} \right) = \\ &= \rho \left(\frac{\partial u}{\partial x'} + \frac{\partial v}{\partial y'} \right) - \frac{1}{g} \frac{\partial}{\partial z} \left(u \frac{\partial p}{\partial x} + v \frac{\partial p}{\partial y} \right), \end{aligned}$$

we obtain the following form (1.4)

$$\frac{\partial u}{\partial x'} + \frac{\partial v}{\partial y'} + \frac{\partial \tau}{\partial p} = 0. \quad (1)$$

The equation of heat influx (1.5) is presented in the following manner if we take the new variables into account:

$$\frac{\partial T}{\partial t'} + u \frac{\partial T}{\partial x'} - v \frac{\partial T}{\partial y'} = \frac{\alpha_0 - \gamma}{g\rho} \tau + \frac{\epsilon}{c_p}, \quad (1.17)$$

where

$$\gamma = - \frac{\partial T}{\partial z}.$$

Finally, we express the limit conditions (1.7) and (1.8) with the new variables as follows:

$$\tau = 0 \quad \text{with} \quad p = 0 \quad (1.18)$$

and

$$\tau = g\rho_0 \frac{\partial z_0}{\partial t'} \quad \text{with} \quad p = p_0. \quad (1.19)$$

It can be easily recognized that the coordinate system selected by us, permits the use of the hydrodynamic equations of the atmosphere in the same way as the equations for the incompressible fluid.

Now after the problem of ascertainment of meteorological elements was formulated, we approach the problem of change to obtain equations suitable for physical conclusions.

For this purpose we differentiate equation (1.13) by y' and equation (1.14) by x' ; we subtract the first equation from the second. We assume that ρ changes little in regard to the horizontal. Thus we obtain

$$\begin{aligned} & \frac{\partial}{\partial t'} \left(\frac{\partial v}{\partial x'} - \frac{\partial u}{\partial y'} \right) + u \frac{\partial}{\partial x'} \left(\frac{\partial v}{\partial x'} - \frac{\partial u}{\partial y'} \right) + v \frac{\partial}{\partial y'} \left(\frac{\partial v}{\partial x'} - \frac{\partial u}{\partial y'} \right) - \\ & + \left(\frac{\partial u}{\partial x'} + \frac{\partial v}{\partial y'} \right) \left(\frac{\partial v}{\partial x'} - \frac{\partial u}{\partial y'} \right) + \tau \frac{\partial}{\partial p} \left(\frac{\partial v}{\partial x'} - \frac{\partial u}{\partial y'} \right) + \frac{\partial v}{\partial p} \frac{\partial \tau}{\partial x'} - \\ & - \frac{\partial u}{\partial p} \frac{\partial \tau}{\partial y'} = - \left(\frac{\partial u}{\partial x'} + \frac{\partial v}{\partial y'} \right) - u \frac{\partial l}{\partial x'} - v \frac{\partial l}{\partial y'} + g^2 \frac{\partial}{\partial p} \rho \frac{\partial}{\partial p} \\ & \left(\frac{\partial v}{\partial x'} - \frac{\partial u}{\partial y'} \right). \quad (1.20) \end{aligned}$$

Compared to other terms of the equation, components of type $\frac{\partial v}{\partial p} \frac{\partial \tau}{\partial x'}$ are generally small and can be disregarded.

We insert the following designations:

$$\frac{\partial v}{\partial x'} - \frac{\partial u}{\partial y'} = \Omega, \quad \frac{\partial u}{\partial x'} + \frac{\partial v}{\partial y'} = \text{div}_p \vec{V}.$$

With the inclusion of new designations equation (1.20) take the following shape:

$$\frac{\partial(\Omega+l)}{\partial x'} + u \frac{\partial(\Omega+l)}{\partial x'} + v \frac{\partial(\Omega+l)}{\partial y'} - g^2 \frac{\partial}{\partial p} \rho^2 \frac{\partial \Omega}{\partial p} =$$

$$= -(\Omega+l) \operatorname{div}_p \vec{V} - \tau \frac{\partial \Omega}{\partial p} \quad (1.21)$$

With the equation of continuity (1.16) we exclude $\operatorname{div}_p \vec{V}$ from (1.21). We obtain:

$$\frac{\partial(\Omega+l)}{\partial x'} + u \frac{\partial(\Omega+l)}{\partial x'} + v \frac{\partial(\Omega+l)}{\partial y'} - g^2 \frac{\partial}{\partial p} \rho^2 \frac{\partial \Omega}{\partial p} =$$

$$= (\Omega+l) \frac{\partial \tau}{\partial p} - \tau \frac{\partial \Omega}{\partial p} \quad (1.22)$$

Inasmuch as value Ω is usually small compared to l , and inasmuch as the changes $\Omega+l$ which depend on the altitude, are small compared to the relative changes of τ according to altitude, we can simplify the equation to some extent as follows:

$$\frac{\partial(\Omega+l)}{\partial x'} + u \frac{\partial(\Omega+l)}{\partial x'} + v \frac{\partial(\Omega+l)}{\partial y'} - g^2 \frac{\partial}{\partial p} \rho^2 \frac{\partial \Omega}{\partial p} =$$

$$= l \frac{\partial \tau}{\partial p} \quad (1.22)$$

To simplify the analysis we check the dynamics of atmospheric motions first without the inclusion of frictional forces. The latter will be discussed at a later stage of the analysis.

For further simplification of equation (1.22) we assume that axis x' shows east and axis y' north. Then the following correlation is obtained:

$$\frac{\partial l}{\partial x'} = 0, \quad \frac{\partial l}{\partial y'} = \frac{2\omega \sin \theta}{a_0} = \beta$$

a_0 - radius of globe. In the following, parameter β will always be considered constant.

Taking into account all of the above conditions, equation (1.22) now takes the following form:

$$\frac{\partial \Omega}{\partial t'} + u \frac{\partial \Omega}{\partial x'} + v \frac{\partial \Omega}{\partial y'} + \beta v = l \frac{\partial \tau}{\partial p} \quad (1.23)$$

We put down the initial equations together with the limit conditions as follows:

$$\frac{du}{dt'} = -g \frac{\partial z}{\partial x'} + lv, \quad (1.24)$$

$$\frac{dv}{dt'} = -g \frac{\partial z}{\partial y'} - lu, \quad (1.25)$$

$$\frac{\partial \Omega}{\partial t'} + u \frac{\partial \Omega}{\partial x'} + v \frac{\partial \Omega}{\partial y'} + \beta v = \left(\frac{\partial \tau}{\partial p} \right), \quad (1.2)$$

$$\frac{\partial T}{\partial t'} + u \frac{\partial T}{\partial x'} + v \frac{\partial T}{\partial y'} = \frac{\kappa - \delta}{g p} z + \frac{e}{c_p}, \quad (1.2)$$

$$\frac{\partial z}{\partial p} = - \frac{1}{g p}, \quad (1.)$$

$$p = \rho R T, \quad (1.)$$

$$\tau = 0 \quad \text{with} \quad p = 0. \quad (1.)$$

$$\tau = g f_0 \frac{\partial z_0}{\partial t'} \quad \text{with} \quad p = p_0. \quad (1.)$$

In the study of atmospheric motion for which the characterizing dimensions in regard to the horizontal are in the order of 1,000 kilometers, and the characteristic horizontal velocities ^{is} 10 sec/m, the analysis of equations (1.24) and (1.25) shows that these motions may be considered geostrophic motion.

$$\begin{aligned} u &= U + u^*, \\ v &= V + v^*, \end{aligned} \quad (1.)$$

where

$$U = -\frac{g}{l} \frac{\partial z}{\partial y'}, \quad V = \frac{g}{l} \frac{\partial z}{\partial x'} \quad (1.33)$$

as component of geostrophic winds. u^* , v^* are small supplementing corrections.

The analysis of the expression for the vortex

$$\Omega = \frac{g}{l} \Delta z + \frac{\partial v^*}{\partial x'} - \frac{\partial u^*}{\partial y'}$$

where

$$\Delta z = \frac{\partial^2 z}{\partial x'^2} + \frac{\partial^2 z}{\partial y'^2}$$

shows that value $-\frac{g}{l} \Delta z$, i.e. the geostrophic approximation for the vorticity includes the main part of the vorticity.

We use the geostrophic approximation for the horizontal component of velocity u and v in equations (1.26) and (1.27); excluding ρ from (1.28) with (1.29) we obtain the following equations:

$$\frac{\partial \Delta z}{\partial t'} + \frac{g}{l} (z, \Delta z) + \beta \frac{\partial z}{\partial x'} = \frac{l^2}{g} \frac{\partial \tau}{\partial p}, \quad (1.34)$$

$$\frac{\partial T}{\partial t'} = \frac{g}{l} (T, z) + \frac{g}{c_p} + \frac{R(\gamma_a - \gamma)}{g} \frac{T}{p} \tau, \quad (1.35)$$

$$\frac{\partial z}{\partial p} = - \frac{R}{g} \frac{T}{p}, \quad (1.36)$$

$$(A, B) = \frac{\partial A}{\partial x'} \frac{\partial B}{\partial y'} - \frac{\partial A}{\partial y'} \frac{\partial B}{\partial x'}.$$

Together with limit conditions (1.30) and (1.31) and elementary data this will suffice to determine future pressure fields, temperature and vertical currents.

Today the ascertainment of the general solution of the non-linear system for the differential equations (1.34) to (1.36) meets difficulties which cannot be overcome. We have only taken up the problem to ascertain the first derivatives $\frac{\partial z}{\partial t}$, $\frac{\partial T}{\partial t}$ and function ρ according to the given distribution of the pressure fields and the temperature.

2. Equation for Pressure Change

We record the equation system (1.34) to (1.36) in the following manner:

$$\frac{\partial \Delta z}{\partial t} + \frac{g}{l} (z, \Delta z) + \rho \frac{\partial z}{\partial x} = \frac{l^2}{Pg} \frac{\partial \tau}{\partial z} \quad 1; \quad (2.1)$$

¹ Index (') in the coordinates x, y, t is left out.

$$\frac{\partial T}{\partial t} - \frac{g}{l} (T, z) - \frac{g}{c_p} = \frac{m^2 l^2}{PR} \frac{\tau}{\zeta} ; \quad (2.2)$$

$$T = - \frac{g}{R} \int \frac{\partial z}{\partial \zeta} , \quad (2.)$$

where $\zeta = \frac{p}{p_0}$ and P the mean pressure on the surface of the earth which was assumed with 1,000 millibar.

$$m^2 = \frac{R^2 T (\alpha_a - \gamma)}{g l^2} .$$

We assume that parameter m^2 changes ^{with respect to height} to a very small extent compared to the relative changes with the altitude of vertical velocity τ .

From equation (2.2) we obtain the derivative of temperature $\frac{\partial T}{\partial t}$ through the equation of statics. Thus we obtain:

$$- \frac{g}{l} \zeta \frac{\partial}{\partial \zeta} \left(\frac{\partial z}{\partial t} \right) - \frac{g}{l} (T, z) - \frac{g}{c_p} = \frac{m^2 l^2}{PR} \frac{\tau}{\zeta} . \quad (2.4)$$

Both parts of the equation are multiplied by ζ . Then we differentiate by ζ . Thus we obtain:

$$- \frac{g}{R} \frac{\partial}{\partial \zeta} \zeta^2 \frac{\partial}{\partial \zeta} \left(\frac{\partial z}{\partial t} \right) - \frac{\partial}{\partial \zeta} \zeta \left[\frac{g}{l} (T, z) + \frac{g}{c_p} \right] = \frac{m^2 l^2}{PR} \frac{\tau}{\zeta} .$$

Excluding derivative $\frac{\partial \sigma}{\partial \xi}$ from equations (2.1) and (2.5) we obtain

$$\left(\frac{\partial}{\partial \xi} \xi^2 \frac{\partial}{\partial \xi} + m^2 \Delta \right) \frac{\partial z}{\partial t} = f_1(x, y, \xi), \quad (2.6)$$

where

$$f_1(x, y, \xi) = -m^2 \left[\frac{g}{l}(z, \Delta z) + \beta \frac{\partial z}{\partial x} \right] - \frac{R}{g} \frac{\partial}{\partial \xi} \xi \left[\frac{g}{l}(T, z) + \frac{\epsilon}{c_p} \right]. \quad (2.7)$$

In the cylindrical coordinate system (r, ϕ, ξ) where $r = \frac{\sqrt{x^2 + y^2}}{m}$, ϕ - *polar* ~~azimuthal~~ angle and $\xi = \frac{p}{p_0}$ the reduced

altitude, equation (2.6) is changed in the following manner:

$$\left(\frac{\partial}{\partial \xi} \xi^2 \frac{\partial}{\partial \xi} + \frac{1}{r} \frac{\partial}{\partial r} r \frac{\partial}{\partial r} + \frac{1}{r^2} \frac{\partial^2}{\partial \phi^2} \right) \frac{\partial z}{\partial t} = f_1(r, \phi, \xi). \quad (2.8)$$

To formulate the problem completely, equation (2.8) should be supplemented by limit conditions.

As the first limit condition we assume the relation directly resulting from the heat influx equation (2.2) on the surface level of 1,000 millibar ^($\xi=1$) on the basis of condition (1.31) as follows:

$$\frac{\partial T_0}{\partial t} = \frac{g}{l}(T_0, z_0) + \frac{\epsilon_0}{c_p} + (\alpha - \gamma) \frac{\partial z_0}{\partial t}. \quad (2.9)$$

Considering that $T_0 = - \left(\frac{g}{R} \left(\frac{\partial z}{\partial \xi} \right) \right)_{\xi=1}$, we put (2.9) into the following shape:

$$\left(\xi \frac{\partial}{\partial \xi} + \alpha \right) \frac{\partial z}{\partial t} \Big|_{\xi=1} = - A(r, \phi, l), \quad (2.)$$

where

$$A(r, \phi, l) = \frac{R}{g} \left[\frac{g}{l} (T_0, z_0) + \frac{\xi_0}{c_p} \right], \quad (2.)$$

$$\alpha = \frac{R(\gamma_a - \gamma)}{g} \approx 0,1. \quad (2.)$$

The second limit condition is obtained from the heat influx equation for the upper limit of the atmosphere in regard to the volume unit of the air:

$$\lim_{\xi \rightarrow 0} \left(c_p \rho \frac{\partial T}{\partial t} \right) = \lim_{\xi \rightarrow 0} \left[c_p \rho \frac{g}{l} (T, z) + \xi' + \frac{c_p (\gamma_a - \gamma)}{g} \tau \right], \quad (2.)$$

ξ' - influx of heat at volume unit.

On the basis of (1.30) we have

$$\lim_{\xi \rightarrow 0} \left[\frac{c_p (\gamma_a - \gamma)}{g} \tau \right] = 0.$$

Moreover, the quantity of heat ξ' flowing at the volume unit, is very small in the upper atmospheric layer. Therefore

$$\lim_{\zeta \rightarrow 0} \varepsilon' = 0.$$

Finally, $\lim_{\zeta \rightarrow 0} \rho u = \lim_{\zeta \rightarrow 0} \rho v = 0$ and the derivatives of temperature $\frac{\partial T}{\partial x}$ and $\frac{\partial T}{\partial y}$ are *BOUNDED ON* the isobaric surfaces in regard to the large-scale motions; as a result

$$\lim_{\zeta \rightarrow 0} \left[c_p \rho \frac{g}{l} (T, z) \right] = 0.$$

In this way we obtain the relation

$$\lim_{\zeta \rightarrow 0} \left(c_p \rho \frac{\partial T}{\partial t} \right) = 0. \quad (2.14)$$

Condition (2.14) can be put down in the following satisfactory manner if we take (2.3) into account:

$$\zeta^2 \frac{\partial}{\partial \zeta} \left(\frac{\partial z}{\partial t} \right)_{\zeta \rightarrow 0} = 0. \quad (2.15)$$

In this manner the work done for the ascertainment of the first derivatives according to time led from the altitude of isobaric surfaces to the integration of the inhomogeneous, differential equation of the second order (2.8) under limit conditions (2.10) and (2.15).

Now we proceed to the problem (2.8), (2.10), (2.15). We assume that functions $f_1(r, \varphi, \zeta)$ and $A(r, \varphi, l)$ can be presented in the following form:

$$\left. \begin{aligned} f_1(r, \varphi, \zeta) &= \operatorname{Re} \sum_{n=-\infty}^{\infty} e^{in\varphi} \int_0^{\infty} F_n(\rho, \zeta) J_n(r\rho) \rho d\rho, \\ A(r, \varphi, l) &= \operatorname{Re} \sum_{n=-\infty}^{\infty} e^{in\varphi} \int_0^{\infty} G_n(\rho) J_n(r\rho) \rho d\rho, \end{aligned} \right\} (2.16)$$

where

$$\left. \begin{aligned} F_n &= \frac{1}{2\pi} \int_0^{2\pi} e^{-in\varphi} d\varphi \int_0^{\infty} f_1(r', \varphi, \zeta) J_n(\rho r') r' dr', \\ G_n &= \frac{1}{2\pi} \int_0^{2\pi} e^{-in\varphi} d\varphi \int_0^{\infty} A(r', \varphi, l) J_n(\rho r') r' dr'. \end{aligned} \right\} (2.17)$$

Here $J_n(x)$ is Bessel's function of the n -order, and the symbol Re means that only the ^{"Real"}(essential) parts of the corresponding expressions are examined.

We look for the solution of the problem in the following manner:

$$\frac{\partial z}{\partial t} = \operatorname{Re} \sum_{n=-\infty}^{\infty} e^{in\varphi} \int_0^{\infty} S_n(\rho, \zeta) J_n(r\rho) \rho d\rho. \quad (2.18)$$

We subject (2.18) and (2.16) under limit conditions (2.10) and (2.15) in equation (2.8). We combine the terms containing $e^{in\phi} J_n(\rho r)$ products with the same n , and adapt them to zero. We use correlation

$$\frac{1}{r} \frac{\partial}{\partial r} r \frac{\partial}{\partial r} J_n(\rho r) - \frac{n^2}{r^2} J_n(\rho r) = -\rho^2 J_n(\rho r),$$

As a result we obtain the following equation and the limit conditions for a new unknown function $S_n(\rho, \zeta)$: as follows:

$$\frac{\partial}{\partial \zeta} \zeta^2 \frac{\partial S_n}{\partial \zeta} - \rho^2 S_n = F_n(\rho, \zeta), \quad (2.19)$$

$$\left(\zeta \frac{\partial}{\partial \zeta} + \alpha \right) S_n \Big|_{\zeta=1} = -G_n(\rho), \quad (2.20)$$

$$\zeta^2 \frac{\partial S_n}{\partial \zeta} \Big|_{\zeta=0} = 0. \quad (2.21)$$

The solution of equation (2.19) is composed of the general solution of the homogeneous equation

$$\zeta^2 \frac{\partial^2 S_n}{\partial \zeta^2} + 2\zeta \frac{\partial S_n}{\partial \zeta} - \rho^2 S_n = 0 \quad (2.22)$$

and of the specific solution of inhomogenous equation (2.19).
 The general solution of homogenous equation (2.22) has the
 following form:

$$S_n^0(\rho, \zeta) = C_{n1} \zeta^{\nu_1} + C_{n2} \zeta^{\nu_2}, \quad (2.23)$$

where

$$\nu_1 = -\frac{1}{2} + \sqrt{\frac{1}{4} + \rho^2}, \quad \nu_2 = -\frac{1}{2} - \sqrt{\frac{1}{4} + \rho^2}, \quad (2.24)$$

C_{n1} and C_{n2} - constant quality depending upon ρ .

The particular solution $S_n^*(\rho, \zeta)$ of the inhomogenous equation (2.19) will be found through the method of variation of constants.

$$S_n^*(\rho, \zeta) = D_{n1}(\rho, \zeta) \zeta^{\nu_1} + D_{n2}(\rho, \zeta) \zeta^{\nu_2}. \quad (2.25)$$

As it is known functions D_{n1} and D_{n2} are determined from following equation system:

$$\left. \begin{aligned} D_{n1}' \zeta^{\nu_1} + D_{n2}' \zeta^{\nu_2} &= 0, \\ \nu_1 D_{n1}' \zeta^{\nu_1-1} + \nu_2 D_{n2}' \zeta^{\nu_2-1} &= \frac{F_n}{\zeta^2}, \end{aligned} \right\} \quad (2.26)$$

$$D'_{ni} = -\frac{D_{ni}}{\partial \xi}.$$

As a result of the solution of system (2.26) we obtain:

$$D'_{n1} = \frac{1}{\nu_1 - \nu_2} F_n \xi^{\nu_2}, \quad D'_{n2} = -\frac{1}{\nu_1 - \nu_2} F_n \xi^{\nu_1}. \quad (2.27)$$

We integrate (2.27) at the limits from zero to ξ . The arbitrarily chosen integration constants are excluded and we obtain:

$$D_{n1}(\rho, \xi) = \frac{1}{\nu_1 - \nu_2} \int_0^{\xi} F_n(\rho, \eta) \eta^{\nu_2} d\eta,$$

$$D_{n2}(\rho, \xi) = -\frac{1}{\nu_1 - \nu_2} \int_0^{\xi} F_n(\rho, \eta) \eta^{\nu_1} d\eta.$$

Consequently, the particular solution of the inhomogenous equation has the following form:

$$S_n^p = \frac{1}{\nu_1 - \nu_2} \int_0^{\xi} F_n(\rho, \eta) [\xi^{\nu_1} \eta^{\nu_2} - \xi^{\nu_2} \eta^{\nu_1}] d\eta. \quad (2.2)$$

In this manner we obtain the desired general solution of equation (2.19)

$$S_n(\rho, f) = C_{n1} f^{\nu_1} + C_{n2} f^{\nu_2} + \frac{1}{\nu_1 - \nu_2} \int_0^f F_n(\rho, \eta) \left[\eta^{\nu_1} - \eta^{\nu_2} \right] d\eta. \quad (2.2)$$

C_{n1} and C_{n2} are obtained from the limit conditions (2.20) and (2.21). The result of condition (2.21) is that

$$C_{n2} = 0. \quad (2.3)$$

Value C_{n1} can be determined from condition (2.20)

$$\nu_1 C_{n1} + \frac{1}{\nu_1 - \nu_2} \int_0^1 F_n(\rho, \eta) \left[\eta^{\nu_1} - \eta^{\nu_2} \right] d\eta + \alpha C_{n1} + \frac{\alpha}{\nu_1 - \nu_2} \int_0^1 F_n(\rho, \eta) \left[\eta^{\nu_2} - \eta^{\nu_1} \right] d\eta = -G_n(\rho),$$

resulting in

$$C_{n1} = \frac{1}{\nu_1 - \nu_2} \int_0^1 F_n(\rho, \eta) \left[\eta^{\nu_1} - \eta^{\nu_2} \right] d\eta - \frac{1}{\alpha + \nu_1} \int_0^1 F_n(\rho, \eta) \eta^{\nu_1} d\eta - \frac{G_n(\rho)}{\alpha + \nu_1}. \quad (2.3)$$

On the basis of (2.30) and (2.31) in (2.29) we obtain

$$S_n(\rho, \zeta) = -\frac{1}{\nu_1 - \nu_2} \int_0^1 F_n(\rho, \eta) \left[\delta_1 \zeta^{\nu_2} \eta^{\nu_1} + \delta_2 \zeta^{\nu_1} \eta^{\nu_2} - (\zeta \eta)^{\nu_1} \right] d\eta - \frac{1}{\alpha + \nu_1} \int_0^1 F_n(\rho, \eta) \zeta^{\nu_1} \eta^{\nu_1} d\eta - \frac{G_n(\rho)}{\alpha + \nu_1} \zeta^{\nu_1}, \quad (2)$$

where

$$\delta_1 = \begin{cases} 1 & \text{with } \eta \leq \zeta, \\ 0 & \text{with } \eta > \zeta, \end{cases} \quad \delta_2 = \begin{cases} 0 & \text{with } \eta \leq \zeta, \\ 1 & \text{with } \eta > \zeta. \end{cases}$$

Now we take (2.32) as a basis for (2.18). We obtain:

$$\frac{\partial z}{\partial t} = -\operatorname{Re} \sum_{n=-\infty}^{\infty} e^{in\phi} \int_0^{\infty} \left\{ \frac{1}{2\mu} \int_0^1 F_n(\rho, \eta) \frac{1}{\sqrt{\zeta\eta}} \left[\delta_1 \left(\frac{\eta}{\zeta}\right)^{\mu} + \delta_2 \left(\frac{\zeta}{\eta}\right)^{\mu} - (\zeta\eta)^{\mu} \right] d\eta + \frac{1}{\alpha - \frac{1}{2} + \mu} \int_0^1 F_n(\rho, \eta) \frac{1}{\sqrt{\zeta\eta}} (\zeta\eta)^{\mu} d\eta - \frac{G_n(\rho)}{\alpha - \frac{1}{2} + \mu} \frac{\zeta^{\mu}}{\sqrt{\zeta}} \right\} J_n(r\rho) \rho d\rho, \quad (2.33)$$

where $\mu = \sqrt{\frac{1}{4} + \rho^2}$.

We also take the values of $F_n(\rho, \eta)$ and $G_n(\rho)$ from (2.1) as a basis for (2.33) and change the sequence of integration

The result is as follows:

$$\frac{\partial z}{\partial t} = -\frac{1}{2\pi} \operatorname{Re} \sum_{n=-\infty}^{\infty} e^{in\phi} \int_0^{\infty} \int_0^{2\pi} e^{-in\psi} \left\{ \int_0^1 f_1(r', \psi, \eta) M_1^{(n)}(r, \xi, r', \eta) d\eta + A(r', \psi, 1) M^{*(n)}(r, \xi, r') \right\} r' d\psi dr', \quad (1)$$

where

$$M_1^{(n)}(r, \xi, r', \eta) = \frac{1}{2\sqrt{\xi\eta}} \int_0^{\infty} \left[\delta_1 \left(\frac{r}{\xi} \right)^\mu + \delta_2 \left(\frac{r'}{\eta} \right)^\mu - (\xi\eta)^\mu \right] \frac{J_n(r\rho) J_n(r'\rho)}{\rho} \rho d\rho + \frac{1}{\sqrt{\xi\eta}} \int_0^{\infty} (\xi\eta)^\mu \frac{J_n(r\rho) J_n(r'\rho)}{\alpha - \frac{1}{2} + \mu} \rho d\rho,$$

$$M^{*(n)}(r, \xi, r') = \frac{1}{\sqrt{\xi}} \int_0^{\infty} \xi^\mu \frac{J_n(r\rho) J_n(r'\rho)}{\alpha - \frac{1}{2} + \mu} \rho d\rho. \quad (2)$$

If we take the vertical line crossing the zero of the coordinate system $(r = 0)^1$ as a testing point, we get $n \neq 0$ all values if $J_n(0) = 0$ with $n \neq 0$ and $J_0(0) = 1$ are taken into account.

$$M_1^{(n)} = 0 \quad \text{and} \quad M^{*(n)} = 0$$

¹ We can always combine the vertical axis of the coordinate the studied vertical.

Only functions $M_1^{(0)}$ and $M^{*(0)}$ differ from zero.

In this manner, the sumtotal (2.34) is transformed to a composed solution, with $r = 0$.

$$\left. \begin{aligned} \frac{\partial z}{\partial t} \Big|_{r=0} &= - \int_0^{\infty} \int_0^1 F_1(r', \eta) M_1(\zeta, r', \eta) r' d\eta dr' - \\ &- \int_0^{\infty} A(r', 1) M^*(\zeta, r') r' dr'. \end{aligned} \right\} \quad (2.37)$$

$$\left. \begin{aligned} F_1(r', \eta) &= \frac{1}{2\pi} \int_0^{2\pi} f_1(r', \psi, \eta) d\psi, \\ A(r', 1) &= \frac{1}{2\pi} \int_0^{2\pi} A(r', \psi, 1) d\psi, \end{aligned} \right\} \quad (2.38)$$

$$M^*(x, r) = \frac{1}{\sqrt{x}} \int_0^{\infty} x^\mu \frac{J_0(r' \rho)}{\alpha - \frac{1}{2} + \mu} \rho d\rho, \quad (2.39)$$

$$\left. \begin{aligned} M_1(\zeta, r', \eta) &= M^*(\int \eta, r') + M_3(\zeta, r', \eta), \\ M_3(\zeta, r', \eta) &= \frac{1}{2\sqrt{\zeta \eta}} \int_0^{\infty} \left[\delta_1 \left(\frac{\eta}{\zeta} \right)^\mu + \delta_2 \left(\frac{\zeta}{\eta} \right)^\mu - (\zeta \eta)^\mu \right] \\ &\quad \frac{J_0(r' \rho)}{\mu} \rho d\rho. \end{aligned} \right\} \quad (2.40)$$

The result of (2.37) is that the pressure change at the points which are situated on any vertical line, is determined by the mean value of the exact functions of the pressure and temperature fields, in the circumferences on the isobaric surfaces, the center being on the examined vertical.

This result can be obtained through a simpler mathematical process. We integrate equation (2.8) with φ , from zero to 2π . We find the solution for the derivatives $\frac{\partial z}{\partial t}$, obtained with the circumference of radius r ; thus we assume that in the obtained solution $r = 0$.

In the following this method will be applied for the obtainment of the solution for $\frac{\partial T}{\partial t}$ and τ .

Now we concentrate on the fact that according to (2.7) the "thermal coefficients" $-\frac{g}{l}(T, z) + \frac{g}{c_p}$ enter the solution (2.37) in a very complicated manner as

$$f_1(r', \psi, \eta) = -B(r', \psi, \eta) - \frac{\partial}{\partial \eta} \eta A(r', \psi, \eta),$$

where

$$\left. \begin{aligned} B(r', \psi, \eta) &= m^2 \left[\frac{g}{l}(z, \Delta z) + \beta \frac{\partial z}{\partial x} \right] \\ A(r', \psi, \eta) &= \frac{R}{g} \left[\frac{g}{l}(T, z) + \frac{g}{c_p} \right] \end{aligned} \right\} \quad (2.37)$$

To simplify solution (2.37) we try to find the function of influence referring directly to $A(r', \eta)$. For this purpose we use expression $-\frac{\partial}{\partial \eta} \eta A(r', \eta)$ instead of $F_1(r', \eta)$ in (2.37). We integrate in stages and obtain:

$$\begin{aligned} & \int_0^1 \frac{\partial}{\partial \eta} \eta A(r', \eta) M_1(\zeta, r', \eta) d\eta - A(r', 1) M^*(\zeta, r') = \\ & = \eta A(r', \eta) M_1(\zeta, r', \eta) \Big|_0^1 - \int_0^1 A(r', \eta) \eta \frac{\partial}{\partial \eta} M_1(\zeta, r', \eta) d\eta \\ & - A(r', 1) M^*(\zeta, r') = - \int_0^1 A(r', \eta) \eta \frac{\partial}{\partial \eta} M_1(\zeta, r', \eta) d\eta = \\ & = \int_0^1 A(r', \eta) M_2(\zeta, r', \eta) d\eta, \end{aligned} \quad (2.)$$

where

$$\left. \begin{aligned} M_2(\zeta, r', \eta) &= -\eta \frac{\partial}{\partial \eta} M_1 = N(\zeta, r', \eta) + \\ &- \frac{1}{2} M_3(\zeta, r', \eta) + \alpha M^*(\zeta, r'), \\ N(\zeta, r', \eta) &= \frac{1}{2\sqrt{\zeta\eta}} \int_0^\infty \left[\delta_2 \left(\frac{\zeta}{\eta}\right)^\mu - \delta_1 \left(\frac{\eta}{\zeta}\right)^\mu - (\zeta\eta)^\mu \right] \\ & J_0(r' \rho) \rho d\rho. \end{aligned} \right\} (2..)$$

Considering (2.41) and (2.42) we, finally, present the solution in the following form:

$$\begin{aligned} \frac{\partial z}{\partial t} = & \frac{1}{2\pi} \int_0^1 \int_0^\infty \int_0^{2\pi} m^2 \left[\frac{g}{l}(z, \Delta z) + \beta \frac{\partial z}{\partial x} \right] M_1(\zeta, r, \eta) r d\varphi dr d\eta + \\ & + \frac{1}{2\pi} \int_0^1 \int_0^\infty \int_0^{2\pi} \frac{R}{g} \left[\frac{g}{l}(T, z) + \frac{\epsilon}{c_p} \right] M_2(\zeta, r, \eta) r d\varphi dr d\eta. \end{aligned} \quad (2.44)$$

The influence function $M_1(\zeta, r, \eta)$ and $M_2(\zeta, r, \eta)$ is characterized by the dependence area of the solution of the meteorological element fields in the surrounding space.

To calculate functions of M^* , M_1 and M_2 it is appropriate to make certain changes in (2.39), (2.41) and (2.43). The values $(\zeta\eta)^\mu$, $(\frac{\eta}{\zeta})^\mu$ and $(\frac{\zeta}{\eta})^\mu$ are presented as instructive functions of type e^{-x} :

$$(\zeta\eta)^\mu = e^{-\mu \ln \frac{1}{\zeta\eta}}, \quad \left(\frac{\eta}{\zeta}\right)^\mu = e^{-\mu \ln \frac{\zeta}{\eta}}, \quad \left(\frac{\zeta}{\eta}\right)^\mu = e^{-\mu \ln \frac{\eta}{\zeta}}$$

With the inequalities

$$\zeta\eta \leq 1, \quad \frac{\eta}{\zeta} \leq 1, \quad \frac{\zeta}{\eta} \leq 1$$

the functions $\ln \frac{1}{\zeta\eta}$, $\ln \frac{\zeta}{\eta}$ and $\ln \frac{\eta}{\zeta}$ are always positive.

As a result function $M_3(\zeta, r, \cdot)$ is converted to the algebraic sumtotal of three integrals of type (4):

$$\sigma(x, r) = \int_0^{\infty} e^{-\mu \ln \frac{1}{x}} \frac{J_0(rp)}{\mu} \rho d\rho = \frac{e^{-\frac{1}{2} \sqrt{\ln^2 \frac{1}{x} + r^2}}}{\sqrt{\ln^2 \frac{1}{x} + r^2}}, \quad (2.45)$$

where $\mu = \sqrt{\frac{1}{4} + \rho^2}$, and x takes value $\{\eta, \frac{1}{\eta}, \frac{1}{\eta}\}$.

If we use the symbols of (2.45), function $M_3(\zeta, r, \eta)$ is presented in the following manner:

$$M_3(\zeta, r, \eta) = \frac{1}{2\sqrt{\zeta\eta}} \left[\delta_1 \sigma\left(\frac{\eta}{\zeta}, r\right) + \delta_2 \sigma\left(\frac{\zeta}{\eta}, r\right) - \sigma(\zeta\eta, r) \right]. \quad (2.46)$$

To calculate $M^*(x, r)$ we use the transformation

$$\int_0^{\infty} x^{\mu - \frac{1}{2}} \frac{J_0(rp)}{\mu - \frac{1}{2} + \mu} \rho d\rho = x^{-\alpha} \int_0^x \lambda^{\alpha - \frac{1}{2}} \frac{\partial}{\partial \lambda} \left\{ \int_0^{\infty} \lambda^{\mu} \frac{J_0(rp)}{\mu} \rho d\rho \right\} d\lambda.$$

Considering (2.45) we obtain:

$$\begin{aligned} M^*(x, r) &= x^{-\alpha} \int_0^x \lambda^{\alpha - \frac{1}{2}} \frac{\partial \sigma(\lambda, r)}{\partial \lambda} d\lambda = \\ &= \frac{\sigma(\lambda, r)}{\sqrt{x}} + \left(\frac{1}{2} - \alpha \right) x^{-\alpha} \int_0^x \lambda^{\alpha - \frac{3}{2}} \sigma(\lambda, r) d\lambda. \quad (2.47) \end{aligned}$$

We introduce

$$\frac{\sigma(x, r)}{\sqrt{x}} = U(x, r). \quad (2.4)$$

As a result we can present function $M_1(\zeta, r, \eta)$ through the following expression:

$$M_1(\zeta, r, \eta) = \frac{1}{2} \left[\frac{\delta_1}{\zeta} U\left(\frac{x}{\zeta}, r\right) + \frac{\delta_2}{\eta} U\left(\frac{x}{\eta}, r\right) + U(\zeta\eta, r) \right] + \left(\frac{1}{2} - \alpha\right) (\zeta\eta)^{-\alpha} \int_0^{\zeta\eta} x^{\alpha-1} U(x, r) dx, \quad (2.49)$$

ζ - the level for which $\frac{\partial \xi}{\partial t}$ is determined, r, η - the variable quantities of integration.

To calculate function $N(\zeta, r, \eta)$ we use the transformati

$$\int_0^{\infty} x^{\mu - \frac{1}{2}} J_0(rp) p dp = \sqrt{x} \frac{\partial}{\partial x} \int_0^{\infty} x^{\mu} \frac{J_0(rp)}{p} p dp \quad (2.50)$$

and then also correlations (2.45) and (2.48). Then $N(\zeta, r, \eta)$ becomes the sumtotal of three integrals of type

$$I(x, r) = \int_0^{\infty} x^{\mu - \frac{1}{2}} J_0(rp) p dp = \sqrt{x} \frac{\partial \sigma(x, r)}{\partial x} = \frac{1}{2} \frac{\ln \frac{1}{x} \left(2 + \sqrt{\ln^2 \frac{1}{x} + r^2} \right)}{\ln^2 \frac{1}{x} + r^2} U(x, r). \quad (2.51)$$

Taking note of (2.46), (2.48), and (2.51) function M_2 can now be described in the following manner:

$$M_2(\zeta, r, \eta) = \frac{1}{2} \left\{ \frac{\delta_2}{\eta} I \left(\frac{\zeta}{\eta}, r \right) - \frac{\delta_1}{\zeta} I \left(\frac{\eta}{\zeta}, r \right) - I(\zeta, \eta, r) \right\} + \frac{1}{4} \left\{ \frac{\delta_1}{\zeta} U \left(\frac{\eta}{\zeta}, r \right) + \frac{\delta_2}{\eta} U \left(\frac{\zeta}{\eta}, r \right) - U(\zeta, \eta, r) \right\} + \alpha M^*(\zeta, \eta, r). \quad (2.52)$$

We note particularly that with $\zeta = 1$, the influence function M_1 and M_2 has a usual form as follows:

$$M_1(1, r, \eta) = M^*(\eta, r), \quad (2.53)$$

$$M_2(1, r, \eta) = -I(\eta, r) + \alpha M^*(\eta, r). \quad (2.54)$$

Functions $U(x, r)$, $I(x, r)$, $M^*(x, r)$ and the fields of functions $M_1(\zeta, r, \eta)$ and $M_2(\zeta, r, \eta)$ for different cutting through the vertical plane, are graphically shown in Fig. 1 - 12.

We will analyse the reaction of these functions with different values of the independent variables x, r, η .

Function $U(x, r)$ (Fig. 1) is $U(1, r) = \frac{1}{r} e^{-\frac{r}{2}}$ with $x = 1$, and $U(0, r) = 0$ with $x = 0$; with a small x the function

tends toward zero such as $\frac{1}{\ln \frac{1}{x}}$. With $r \neq 0$ $U(x, 0) = \frac{1}{\ln x}$.

with growing r and a fixed value x function $U(x, r)$ decrease in a regular manner; with $r \rightarrow \infty$ it tends toward zero, as $\frac{1}{r} e^{-\frac{r}{2}}$. The function shows its characteristic trait at point $(x=1, r=0)$.

With $x=1$ and $r \neq 0$ function $I(x, r)$ (Fig. 2) equals zero; with $x=0$ $I(0, r) = 0$; with a small x the function

tends toward zero, as $\frac{1}{2 \ln \frac{1}{x}}$. With $r=0$ $I(x, 0) = \frac{2 + \ln \frac{1}{x}}{2 \ln^2 \frac{1}{x}}$. With growing r and a fixed value x function

$I(x, r)$ decreases in a regular manner; with $r \rightarrow \infty$ it

tends toward zero, as $\frac{1}{2r^2} e^{-\frac{r}{2}}$. The function is character-

istic at point $(x=1, r=0)$.

The characteristics of function $M^*(x, r)$ (Fig. 3) are similar to function $U(x, r)$. Values $M^*(x, r)$ are 1.5 to two times higher than values $U(x, r)$.

With $x=0$ $M^*(0, r) = 0$; with $r \rightarrow \infty$ $M^*(x, r)$ appears

as $\frac{1}{r} e^{-\frac{r}{2}}$ (Fig. 4). The tendency of the function toward

zero is governed by the rule $\frac{x}{\ln \frac{1}{x}}$.

Function $M_1(l, r, \eta)$ (Fig. 5) describes the integral act of "the dynamic factors" $\frac{g}{l}(z, \Delta z) + \beta \frac{\partial z}{\partial x}$ in regard to the pressure change at the point on the earth's surface. The act distance $R = \sqrt{m^2} \approx 750$ km corresponds to the relative length




Fig. 1
Graphic Presentation
of Function $U(x, r)$
with Different r
Values




Fig. 2
Graphic Presentation
of Function $I(x, r)$
with Different r
Values




Fig. 3. Graphic Presentation of Function $M^*(x, r)$ with Different r Values




Fig. 4. Graphic Presentation of Function $M^*(x, r)$ with Varying x .

Attention must be given to the vast expansion of the action area of the dynamic factors in horizontal direction. For better illustration and analysis of vanishing of action on the part of the dynamic factors with increasing r values, we will show the influence of $rM_1(1, r, \eta)$, in a graphical manner, describing the importance of average values of dynamic factors $\frac{g}{T}(z, \Delta z) + \beta \frac{\partial z}{\partial x}$ relative to circumferences of radius r (Fig. 6). We see that function rM_1 vanishes very slowly with increasing r . With $r \rightarrow \infty$ $rM_1(1, r, \eta)$ decreases as $e^{-\frac{r}{2}}$.

In actual practice, the influence of dynamic factors on the change of surface pressure will always be limited by radius R of the 2,000 km order, because the average values of $\frac{g}{T}(z, \Delta z) + \beta \frac{\partial z}{\partial x}$ relative to the circumferences 2,000 - 3,000 km are actually small.

The limited effect on the change of surface pressure and dynamic factors in the upper layers, can be seen in a clearer manner in Fig. 6. With an equal distribution of the dynamic factors according to altitude, their effect from the surface limited by radius $r = 1$ (300 millibar) is reduced twofold compared to the effect of these factors from the 900 millibar surface which is limited by the same radius.

Function $M_2 (1, r, \eta)$ (Fig. 7) gives an idea of the effect of thermal factors $-\frac{E}{t} (T, z) + \frac{E}{c_p}$ or, to be more precise, of the effect of local heat flux in the atmosphere and the change of surface pressure. The negative values of function $M_2 (1, r, \eta)$ in the surrounding area of the atmosphere show that at any level of the atmosphere in this area the local heat flux causes a pressure drop at the earth, ^{while for} heat emission ~~and temperature~~ ^{PRESSURE} increases.




Fig. 5. Field of Function $M_1 (\zeta, r, \eta)$ with $\zeta = 1.0$




Fig. 6. Field of Function $rM_1 (1, r, \eta)$

Compared to the action area of dynamic factors, the region of activity of thermal factors appears to be more limited. Indeed, the maximum value of function $M_2 (1, r, \eta)$ along the vertical line with $r = 1$ is more than twenty times smaller than the maximum value of $M_2 (1, r, \eta)$ along the vertical line

with $r = 0,2$, during the same period in which, for instance, the function of influence $M_1 (1, r, \eta)$ decreases 4 times.

The difference between the areas of influence of thermic and dynamic factors can be recognized by the asymptotic reaction of functions $M_2 (1, r, \eta)$ and $M_1 (1, r, \eta)$.

Function $I (\eta, r)$ which appears as the main part of function $M_2 (1, r, \eta)$, tends toward zero, with higher r values, as

$$\frac{1}{r^2} e^{-\frac{r}{2}}, \text{ but } M_1 (1, r, \eta) \text{ shows the same reaction as } \frac{1}{r} e^{-\frac{r}{2}}.$$

We also want to show function $r M_2 (1, r, \eta)$ which represents the summary influence of thermal factors on the circumference of radius r , as well as the change of surface pressure. The analysis of function $r M_2 (1, r, \eta)$ shows that the influence of thermal factors in the upper layers of the atmosphere on the change of surface pressure, is usually very small. Inasmuch as the average values of the thermal factors $\frac{\epsilon}{T} (T, z) + \frac{\epsilon}{c_p}$ on circumference r with a sufficiently high r , can be considered important only in the case of greater thermobarometric disturbances in the atmosphere. On the basis of the graph shown in Fig. 8 it can be maintained that the relatively important influence of thermic factors of the upper layers on the change of surface pressure can exist only in the case of great thermobarometric disturbances.




Fig. 7. Field of Function $M_2(\xi, r, \eta)$ with $\xi = 1.0$




Fig. 8. Field of Function $rM_2(1, r, \eta)$

Now we examine the nature of the functions of influence of $M_2(\xi, r, \eta)$ and $M_2(\xi, r, \eta)$ with $\xi < 1$, i.e. the effect of the above mentioned dynamic and thermic factors on the change of pressure at a certain level above the surface of the earth.

Fig. 9 and 10 show functions $M_1(0.7, r, \eta)$ and $M_1(0.5, r, \eta)$. In principle, they do not differ from functions which are similar to them, i.e. from functions for $\xi = 1$, which were studied above. The difference is only that the area of maximum values of the function differed here from levels $\xi = 0.7$ and $\xi = 0.5$, that is to say from levels of examined points.




Fig. 9. Field of Function $M_1(\zeta, r, \eta)$ with $\zeta = 0.7$




Fig. 10. Field of Function $M_1(\zeta, r, \eta)$ with $\zeta = 0.5$

A completely different type is function $M_2(\zeta, r, \eta)$ with $\zeta < 1$ compared to $M_2(1, r, \eta)$. Fig. 11 and 12 show functions $M_2(0.7, r, \eta)$ and $M_2(0.5, r, \eta)$. We can see that the local flux of heat in the upper part of the atmosphere relative to level ζ causes a drop of temperature; on the other hand, the heat flux in the lower part relative to level ζ leads to a rise in temperature. The ^{EMISSION} of heat at different levels ζ of the atmosphere causes the reverse effect.

At a certain level ζ^* the influences of thermic factors of the upper and lower layers on the pressure change compensate each other so that the pressure change depends only upon the dynamic factors. This level will be called "mean level" of the atmosphere.




Fig. 11. Field of Function $M_2(\xi, r, \eta)$ with $\xi = 0.7$




Fig. 12. Field of Function $M_2(\xi, r, \eta)$ with $\xi = 0.5$

Theoretically, the existence of such a level was already mentioned in I.A. Kibel's (2) analysis. In the first place the isobaric field at this level, as it was shown by I.A. Kibel, determines the motion of barometric and thermal disturbances in the vicinity of the earth. However, empirically the existence of the steering current was shown already at an earlier date by S.I. Troitzky (5).

In each concrete synoptic situation function $M_2(\xi, r, \eta)$ permits us to determine the position of such a level. In general, this level does not remain constant neither relative to the vast spaces nor the time, because in each practical case its position will be dependent upon the distribution of the heat flux $E = c_p \frac{\partial}{\partial t} (T, z) + \mathcal{E}$.

If we admit that the local heat flux is constant relative to the vertical line along the entire thickness of the atmosphere, the mean level, in this case, would be somewhere between the surfaces of 700 and 500 millibar.

The position of the mean level remains dependent upon the values of thermic and barometric distribution in the atmosphere. With increasing r values the negative values of function $rM_2(\xi, r, \eta)$ in area $\eta < \xi$ disappear at a slower rate than the positive values in area $\eta > \xi$; the conclusion is that with the same relative distribution of the heat flux $E(r, \phi, \eta)$ along the vertical line, the mean level will be higher in the case of great disturbances than in the case of small disturbances.

With $\xi \rightarrow 0$ solution (2.44) for $\frac{\partial z}{\partial t}$ is converted to the following expression:

$$\left. \frac{\partial z}{\partial t} \right|_{\xi=0} = \frac{1}{2\pi} \int_0^{\infty} \int_0^{2\pi} m^2 \left[\frac{g}{l}(z, \Delta z) - \beta \frac{\partial z}{\partial x} \right]_{\xi=0} \frac{U(1, r)}{2} r d\phi dr ;$$

$$+ \frac{1}{2\pi} \int_0^{\infty} \int_0^2 \frac{E}{g} \left[\frac{g}{l}(T, z) + \frac{\xi}{c_p} \right]_{\xi=0} \frac{U'(1, r)}{4} r d\phi dr, \quad (2.45)$$

where

$$U'(1, r) = U(1, r) - 2I(1, r),$$

i.e.

$$U^1(l, r) = \begin{cases} U(l, r) & \text{with } r > 0 \\ 0 & \text{with } r = 0. \end{cases} \quad (2.56)$$

The functions of influence $M_1(\xi, r, \eta)$ and $M_2(\zeta, r, \eta)$ characterize the region of dependence of the studied value $\frac{\partial z}{\partial t}$ at the respective point upon the environmental setting of the fields of meteorological elements; these functions can also be interpreted on the basis of the principle of reversibility in the following manner.

At a certain point of space (x, y, ζ) we place a single "source of substance".

$$B = \alpha^2 \left[\frac{B}{l} (z, \Delta z) + \beta \frac{\partial z}{\partial x} \right] \quad \text{or} \quad A = \frac{R}{g} \left[\frac{g}{l} (T, z) + \frac{L}{c_p} \right].$$

Accordingly, function $M_1(\eta, r, \xi)$ or $M_2(\eta, r, \zeta)$ causes the change of pressure in the surrounding space through the influence of this source. The positive source of the dynamic substance B at the level results in an increase of pressure in the surrounding area at all levels, reflected by space function $M_1(\eta, r, \xi)$. It can be easily seen that function $M_2(\eta, r, \zeta)$ is identical with $M_1(\xi, r, \eta)$.

The single "thermal source A " at level ζ results in decreased pressure in the upper part of the atmosphere according to function $M_2(\eta, r, \zeta)$ as far as level ξ is concerned.

as well as a rise of pressure in the lower part of the atmosphere. Negative sources of the above mentioned dynamic and thermic substances cause, in an analogous manner, a reverse tendency.

Such an interpretation of the influence function gives us a clear idea of the radius of influence of thermic-barometric disturbances in the atmosphere on the change of pressure changes in neighboring regions.

An analogous interpretation will also be given by the solutions for $\frac{\partial T}{\partial t}$ and τ , which will be described in the following paragraphs.

3. Equations for Temperature Changes.

The equations for temperature changes at different levels will be obtained through the separation from the equation system (2.1) - (2.3) of the vertical velocity τ and derivative $\frac{\partial z}{\partial t}$.

To this end we differentiate (2.1) with ξ . We obtain:

$$\frac{\partial}{\partial \xi} \left(\Delta \frac{\partial z}{\partial t} \right) + \frac{\partial}{\partial \xi} \left[\frac{g}{L} (z, \Delta z) + \beta \frac{\partial z}{\partial x} \right] = \frac{L}{P_g} \frac{\partial^2 z}{\partial \xi^2} .$$

The equation of state^{ics} (2.3) results as follows: (3.1)

$$\frac{\partial}{\partial \xi} \left(\Delta \frac{\partial z}{\partial t} \right) = - \frac{R}{g} \frac{1}{\xi} \Delta \frac{\partial T}{\partial t} . \quad (3.2)$$

$$\frac{\partial}{\partial \xi} \left[\frac{g}{l} (z, \Delta z) + \beta \frac{\partial z}{\partial x} \right] = - \frac{R}{g} \frac{1}{f} \left[\frac{g}{l} (T, \Delta z) + \frac{g}{e} (z, \Delta T) - \right. \\ \left. + \beta \frac{\partial T}{\partial x} \right]. \quad (3.3)$$

We incorporate (3.2) and (3.3) into (3.1) and obtain:

$$- \frac{1}{f} \Delta \frac{\partial T}{\partial t} - \frac{1}{f} \left[\frac{g}{l} (T, \Delta z) + \frac{g}{e} (z, \Delta T) + \beta \frac{\partial T}{\partial x} \right] = \frac{l^2 \partial^2 \tau}{PR \partial \xi^2}. \quad (3.4)$$

To exclude $\frac{\partial^2 \tau}{\partial \xi^2}$ from (3.4) we dissolve the equation in regard to τ and differentiate same twice with ξ assuming, as previously, that parameter m^2 changes according to altitude. We obtain:

$$\left(\xi \frac{\partial^2}{\partial \xi^2} + 2 \frac{\partial}{\partial \xi} \right) \frac{\partial T}{\partial t} - \left(\xi \frac{\partial^2}{\partial \xi^2} + 2 \frac{\partial}{\partial \xi} \right) \left[\frac{g}{l} (T, z) + \frac{g}{c_p} \right] = \\ = \frac{m^2 l^2 \partial^2 \tau}{PR \partial \xi^2}. \quad (3.5)$$

Now we multiply (3.4) with m^2 and subtract from (3.5).

The result is as follows:

$$\left(\frac{\partial}{\partial \xi} \xi^2 \frac{\partial}{\partial \xi} + m^2 \Delta \right) \frac{\partial T}{\partial t} = f_2 (x, y, \xi), \quad (3.6)$$

$$f_2(x, y, \zeta) = \frac{\partial}{\partial \zeta} \zeta^2 \frac{\partial}{\partial \zeta} \left[\frac{g}{l} (T, z) + \frac{\epsilon}{c_p} \right] -$$

$$- m^2 \left[\frac{g}{l} (T, \Delta z) + \frac{g}{l} (z, \Delta T) + \beta \frac{\partial T}{\partial x} \right]. \quad (3.7)$$

In regard to the nature of limit conditions for equation (3.6) we assume the following (see conditions (2.9) and (2.14

$$\left. \begin{aligned} \frac{\partial T}{\partial t} \Big|_{\zeta=1} &= Q_0(x, y), \\ \frac{\partial T}{\partial t} \Big|_{\zeta=0} &= 0, \end{aligned} \right\} \quad (3.8)$$

$$Q_0(x, y) = \frac{g}{l} (T_0, z_0) + \frac{\epsilon_0}{c_p}. \quad (3.9)$$

The term $(\alpha - \gamma) \frac{\partial z_0}{\partial t}$ contained in $Q_0(x, y)$ is disregarded in this analysis because this term is of a higher order¹.

In view of the fact that equation (3.6) has a structure which is analogous to the structure of equation (2.6) for $\frac{\partial z}{\partial t}$

¹ The exclusion of term $(\alpha - \gamma) \frac{\partial z_0}{\partial t}$ contained in $Q(x, y)$, is not of basic nature. It could also be retained as $\frac{\partial z_0}{\partial t}$ can be determined according to formula (2.44) in the previous paragraph.

the solution for $\frac{\partial T}{\partial t}$ as well as for $\frac{\partial z}{\partial t}$ will be determined by the mean value of exact functions on the peripheries with the center being at the examined point. Therefore, we apply a simpler mathematical method for the solution of equation (3.6). The possibility of applying this method was already mentioned above.

We note equation (3.6) in the cylindrical coordinate system (r, ϕ, ξ) , $r = \frac{\sqrt{x^2 + y^2}}{m}$, ϕ - polar angle, $\xi = p^z$ - mentioned altitude. We integrate this equation with ϕ from zero to 2π .

The result is as follows:

$$\left(\frac{\partial}{\partial \xi} \int^2 \frac{\partial}{\partial \xi} + \frac{1}{r} \frac{\partial}{\partial r} r \frac{\partial}{\partial r} \right) \frac{\partial T}{\partial t} = f_2(r, \xi), \quad (3.10)$$

$$\frac{\partial T}{\partial t} = \frac{1}{2\pi} \int_0^{2\pi} \frac{\partial T}{\partial t} d\phi, \quad f_2(r, \xi) = \frac{1}{2\pi} \int_0^{2\pi} f_2(r, \phi, \xi) d\phi \quad (3.11)$$

These are values $\frac{\partial T}{\partial t}$ and f_2 ascertained on the circumference of radius r .

The limit conditions (3.8) are reflected in the following manner:

$$\left. \begin{aligned} \frac{\partial T}{\partial t} \Big|_{\xi=1} &= Q_0(r), \\ \xi \frac{\partial T}{\partial t} \Big|_{\xi=0} &= 0. \end{aligned} \right\} \quad (3.12)$$

The solution for $\frac{\partial T}{\partial t}$ will be tried in the form of the Fourier-Bessel integral

$$\frac{\partial T}{\partial t} = \int_0^{\infty} S(\rho, \xi) J_0(\rho r) \rho d\rho. \quad (3.13)$$

We assume that function $f_2(r, \xi)$ and $Q_0(r)$ can be presented in the form of the Fourier-Bessel integral:

$$\left. \begin{aligned} f_2(r, \xi) &= \int_0^{\infty} J_0(\rho r) \rho d\rho \int_0^{\infty} f_2(r', \xi) J_0(\rho r') r' dr', \\ Q_0(r) &= \int_0^{\infty} J_0(\rho r) \rho d\rho \int_0^{\infty} Q_0(r') J_0(\rho r') r' dr'. \end{aligned} \right\} \quad (3.14)$$

We take (3.13) and (3.14) as a basis for equation (3.10) and limit conditions (3.12) and use the known correlation

$$\frac{1}{r} \frac{\partial}{\partial r} \left\{ r \frac{\partial J_0(\rho r)}{\partial r} \right\} = -\rho^2 J_0(\rho r),$$

In this manner we obtain the following equation and limit conditions for $S(\rho, \xi)$:

$$\frac{\partial}{\partial \xi} \xi^2 \frac{\partial S}{\partial \xi} - \rho^2 S = F_2(\rho, \xi), \quad (3.1)$$

$$\left. \begin{aligned} S &= G_2(\rho), \\ \int S \Big|_{\xi=0} &= 0, \end{aligned} \right\} \quad (3.1)$$

$$\left. \begin{aligned} F_2(\rho, \xi) &= \int_0^{\infty} f_2(r', \xi) J_0(\rho r') r' dr', \\ G_2(\rho) &= \int_0^{\infty} Q_0(r') J_0(\rho r') r' dr'. \end{aligned} \right\} \quad (3.1)$$

In view of the fact that equation (3.15), as far as its structure is concerned, is analogous to equation (2.19) examined in the last paragraph, the general solution of equation (3.15) will have a form which is analogous to (2.29).

$$S(\rho, \xi) = C_1 \xi^{\nu_1} + C_2 \xi^{\nu_2} + \frac{1}{\nu_1 - \nu_2} \int_0^{\xi} F_2(\rho, \xi') \left[\xi^{\nu_1} \eta^{\nu_2} - \xi^{\nu_2} \eta^{\nu_1} \right] d\xi' \quad (3.1)$$

$$v_1 = -\frac{1}{2} + \sqrt{\frac{1}{4} + \rho^2}, \quad v_2 = -\frac{1}{2} - \sqrt{\frac{1}{4} + \rho^2}.$$

We take arbitrarily constant quantities C_1 and C_2 from the limit conditions and obtain:

$$S(\rho, \zeta) = -\frac{1}{2\mu} \int_0^1 F_2(\rho, \eta) \frac{1}{\sqrt{\zeta\eta}} \left[\delta_1 \left(\frac{\eta}{\zeta} \right)^\mu + \delta_2 \left(\frac{\zeta}{\eta} \right)^\mu - (\zeta\eta)^\mu \right] d\eta + G(\rho) \zeta^{\mu - \frac{1}{2}}, \quad (3.19)$$

$$\mu = \sqrt{\frac{1}{4} + \rho^2}.$$

We take (3.19) and (3.13) as a basis and replace $F_2(\rho, \eta)$ and $G(\rho)$ according to formula (3.17). If we also change the order of integration we obtain:

$$\begin{aligned} \frac{\partial T}{\partial t} = & - \int_0^1 \int_0^\infty f_2(r', \eta) \left\{ \frac{1}{2\sqrt{\zeta\eta}} \int_0^\infty \left[\delta_1 \left(\frac{\eta}{\zeta} \right)^\mu + \right. \right. \\ & \left. \left. + \delta_2 \left(\frac{\zeta}{\eta} \right)^\mu - (\zeta\eta)^\mu \right] \frac{J_0(r\rho) J_0(r'\rho)}{\mu} \right\} r' dr' d\eta + \\ & + \int_0^\infty Q_0(r') \left\{ \frac{1}{\sqrt{\zeta}} \int_0^\infty \zeta^\mu J_0(r\rho) J_0(r'\rho) \rho d\rho \right\} r' dr'. \quad (3.) \end{aligned}$$

Now we assume that in (3.20) $r = 0$. As a result we obtain the solution for $\frac{\partial T}{\partial t}$ at the points which are located at various levels ζ , along the coordinate axis.

$$\frac{\partial T}{\partial t} = - \int_0^1 \int_0^{\infty} f_2(r_1, \eta) M_3(\zeta, r', \eta) r' dr' d\eta + \int_0^{\infty} Q_0(r') I(\zeta, r') r' dr', \quad (3.2)$$

$$M_3(\zeta, r', \eta) = \frac{1}{2\sqrt{\zeta\eta}} \int_0^{\infty} \left[\delta_1 \left(\frac{\eta}{\zeta}\right)^{\mu} + \delta_2 \left(\frac{\zeta}{\eta}\right)^{\mu} - (\zeta\eta)^{\mu} \right] \frac{J_0(r'\rho)}{\mu} \rho d\rho \quad (3.2)$$

and

$$I(\zeta, r') = \int_0^{\infty} \zeta^{\mu} - \frac{1}{2} J_0(r'\rho) \rho d\rho \quad (3.2)$$

These are functions which had already been used previously

We take $f_2(r', \eta)$ from (3.11) and (3.7) as a basis for (3.21) and partially integrate thermic factor $\frac{E}{l}(T, z) + \frac{E}{c_p}$

Thus we can finally give a description of solution (3.21) for $\frac{\partial T}{\partial t}$ as follows¹:

¹ Derivative $\frac{\partial T}{\partial t}$ can be directly obtained from solution (2.3) for $\frac{\partial^2 T}{\partial t^2}$ by differentiating with ζ and utilizing correlation

$$\begin{aligned}
\frac{\partial T}{\partial t} &= \frac{1}{2\pi} \int_0^{\infty} \int_0^{2\pi} \left[\frac{g}{l} (T_0, z_0) + \frac{\epsilon_0}{c_p} \right] I(\xi, r) r d\varphi dr + \\
&+ \frac{1}{2\pi} \int_0^1 \int_0^{\infty} \int_0^{2\pi} \eta \frac{\partial}{\partial \eta} \left[\frac{g}{l} (T, z) + \frac{\epsilon}{c_p} \right] M_4(\xi, r, \eta) r d\varphi dr d\eta + \\
&+ \frac{1}{2\pi} \int_0^1 \int_0^{\infty} \int_0^{2\pi} m^2 \left[\frac{g}{l} (T, \Delta z) + \frac{g}{l} (z, \Delta T) + \beta \frac{\partial T}{\partial z} \right] M_3(\xi, r, \eta) \\
&rd\varphi dr d\eta, \tag{3.2}
\end{aligned}$$

$$M_4(\xi, r, \eta) = \eta \frac{\partial M_3}{\partial \eta}.$$

(continued)

$$\frac{\partial T}{\partial t} = -\frac{g}{R} \int \frac{\partial}{\partial \xi} \left(\frac{\partial z}{\partial t} \right).$$

Then we obtain:

$$\begin{aligned}
\frac{\partial T}{\partial t} &= \frac{1}{2\pi} \int_0^{\infty} \int_0^{2\pi} Q_0(r, \varphi) G_1(\xi, r) r d\varphi dr + \frac{1}{2\pi} \int_0^1 \int_0^{\infty} \int_0^{2\pi} f(r, \varphi, \eta) \\
&G_2(\xi, r, \eta) r d\varphi dr d\eta,
\end{aligned}$$

$$f(r, \varphi, \eta) = \frac{gm^2}{R} \left[\frac{g}{l} (z, \Delta z) + \beta \frac{\partial z}{\partial \eta} \right] + \frac{\partial}{\partial \eta} \eta \left[\frac{g}{l} (T, z) + \frac{\epsilon}{c_p} \right]$$

The functions of influence M_3 and M_4 are expressed by functions of type $U(x, r)$ and $I(x, r)$ which were used in the preceding paragraph.

$$\begin{aligned}
 M_3(\xi, r, \eta) &= \frac{1}{2} \left[\frac{\delta_1}{\xi} U\left(\frac{\eta}{\xi}, r\right) + \frac{\delta_2}{\eta} U\left(\frac{\xi}{\eta}, r\right) - U(\xi\eta, r) \right] \\
 M_4(\xi, r, \eta) &= \frac{1}{2} \left\{ \frac{\delta_1}{\xi} I\left(\frac{\eta}{\xi}, r\right) - \frac{\delta_2}{\eta} I\left(\frac{\xi}{\eta}, r\right) - I(\xi\eta, r) \right\} - \left. \begin{aligned} & \\ & - \frac{1}{2} M_3(\xi, r, \eta). \end{aligned} \right\} \quad (3.2)
 \end{aligned}$$

Fig. 13-17 show the function of $I(\xi, r)$ and the sections in the vertical field ^{value of} of functions $M_4(\xi, r, \eta)$ and $M_3(\xi, r, \eta)$.

The analysis of the values of function $I(\xi, r)$ (Fig. 13) shows that the first term on the right (3.24) calculated according to altitude, steadily vanishes with the vaning ξ .

(continued)

$$\begin{aligned}
 G_2(\xi, r, \eta) &= -\xi \frac{\partial M_1}{\partial \xi} = \frac{1}{2} \left\{ \frac{\delta_1}{\xi} I\left(\frac{\eta}{\xi}, r\right) - \frac{\delta_2}{\eta} I\left(\frac{\xi}{\eta}, r\right) + I(\xi\eta, r) \right\} \\
 &+ \frac{1}{2} M_3(\xi, r, \eta) + \alpha M^*(\xi, r),
 \end{aligned}$$

$$G_1(\xi, r) = \xi \frac{\partial M^*(\xi, r)}{\partial \xi} = I(\xi, r) - \alpha M^*(\xi, r).$$

Function $M_4(\zeta, r, \eta)$ (Fig. 14 and 15) is used for calculating the irregular distribution of the local heat flux along the vertical. The first two integrals on the right (3.24) give complete data on the influence of local heat flux distributed over the entire atmosphere and on the temperature change at a fixed point.

Analytically, function $M_4(\zeta, r, \eta)$ consists of two parts:

$$M_4(\zeta, r, \eta) = X(\zeta, r, \eta) - \frac{1}{2} M_3(\zeta, r, \eta), \quad (3.26)$$

$$X(\zeta, r, \eta) = \frac{1}{2} \left\{ \frac{\delta_1}{\zeta} I\left(\frac{\eta}{\zeta}, r\right) - \frac{\delta_2}{\eta} I\left(\frac{\zeta}{\eta}, r\right) - I(\zeta, \eta, r) \right\}. \quad (3.2)$$




Fig. 13. Graph of Function $I(\zeta, r)$ for various ζ .

In the environment of radius $r \approx 0.5$ $X(\zeta, r, \eta)$ forms the main part of values $M_4(\zeta, r, \eta)$. On the other hand, $\frac{1}{2} M_3(\zeta, r, \eta)$ appears to be relatively small. With higher values of r ($r \gg 1$), the corresponding equilibrium of both components contained in $M_4(\zeta, r, \eta)$, is outbalanced; but each of these components is small.

Function $M_3(\zeta, r, \eta)$ (Fig. 16 and 17) gives an idea of the sphere of influence of the dynamic factors $m_2 \left[\frac{E}{T}(T, \Delta z) + \frac{E}{T}(z, \Delta T) + \beta \frac{\partial T}{\partial \chi} \right]$ on the change of temperature at the respective point on level ζ . This function has its maximum values in the immediate vicinity of the examined point and vanishes with the increasing r but also in the direction of the upper and lower boundaries of the atmosphere. $\eta = 0$ and $\eta = 1$ $M_3(\zeta, r, \eta) = 0$, if $\zeta \neq 0$.

It can be easily recognized that with higher r values function $M_3(\zeta, r, \eta)$ is presented in an asymptotic manner as follows: $\frac{1}{r^2} e^{-\frac{r}{\lambda}}$. According to the increase of r values, the area of maximum values of function $M_3(\zeta, r, \eta)$ with level ζ is formed at the upper part.

The result of the analysis of the properties of function $M_3(\zeta, r, \eta)$ is that the temperature change at any level ζ , expressed by the last term in (3.24), is determined by the dynamic factors of the atmospheric layers bordering level ζ to a greater extent. In regard to level ζ the dynamic factors of the upper layers have a higher ^{weight} ~~importance~~ than similar factors in lower layers.




Fig. 14. Field of Function $M_4(\xi, r, \eta)$ with $\xi = 0.7$




Fig. 15. Field of Function $M_4(\xi, r, \eta)$ with $\xi = 0.5$

Now we assume in what manner the solution for $\frac{\partial T}{\partial t}$ describes the change of temperature at the upper border of the atmosphere.




Fig. 16. Field of Function $M_3(\xi, r, \eta)$ with $\xi = 0.7$




Fig. 17. Field of Function $M_3(\xi, r, \eta)$ with $\xi = 0.5$

With $\xi \rightarrow 0$ function $I(\xi, r) \rightarrow 0$, $M_4(\xi, r, \eta)$ is converted to $\frac{\delta_1}{2\xi} \left[I(1, r) - \frac{1}{2} U(1, r) \right]$, $M_3(\xi, r, \eta)$ and $\frac{\delta_1}{2\xi} U(1, r)$. Then solution (3.24) takes the following shape

with $\xi \rightarrow 0$:

$$\begin{aligned} \frac{\partial T}{\partial t} \Big|_{\xi=0} &= -\frac{1}{2\pi} \int_0^{\infty} \int_0^{2\pi} \eta \frac{\partial}{\partial \eta} \left[\frac{g}{l}(T, z) + \frac{g}{c_p} \right]_{\xi=0} \frac{U'(1, r)}{4} r d\varphi dr + \\ &+ \frac{1}{2\pi} \int_0^{\infty} \int_0^{2\pi} m^2 \left[\frac{g}{l}(T, \Delta z) + \frac{g}{l}(z, \Delta T) + \beta \frac{\partial T}{\partial x} \right]_{\xi=0} \frac{U(1, r)}{2} r d\varphi dr, \end{aligned} \quad (3.28)$$

Analogous to (2.56)

$$U'(1, r) = \begin{cases} U(1, r) & \text{with } r > 0, \\ 0 & \text{with } r = 0. \end{cases}$$

Now we recall correlation (3.3)

$$\xi \frac{\partial}{\partial \xi} \left[\frac{g}{l}(z, \Delta z) + \beta \frac{\partial z}{\partial x} \right] = -\frac{R}{g} \left[\frac{g}{l}(T, \Delta z) + \frac{g}{l}(z, \Delta T) + \beta \frac{\partial T}{\partial x} \right]$$

Assuming that

$$\lim_{\xi \rightarrow 0} \left\{ \xi \frac{\partial}{\partial \xi} \left[\frac{g}{l}(z, \Delta z) + \beta \frac{\partial z}{\partial x} \right] \right\} = 0^1, \quad (3.29)$$

¹ Qualification (3.29) means that the kinetic energy of the u of mass $\frac{v^2}{2}$ according to altitude does not increase faster than the increase of $\frac{1}{\xi}$.

we obtain

$$\left[\frac{\varepsilon}{l}(T, \Delta z) + \frac{\varepsilon}{l}(z, \Delta T) + \beta \frac{\partial T}{\partial x} \right]_{\zeta=0} = 0. \quad (3.30)$$

The result of (3.30) is that derivatives $\frac{\partial T}{\partial x}$ and $\frac{\partial T}{\partial y}$ at the upper border of the atmosphere equal zero; this means that the temperature on the surface $\zeta = \text{const}$, is constant with $\zeta \rightarrow 0$. In other words,

$$(T, z)_{\zeta=0} = 0. \quad (3.31)$$

We further assume that $\zeta \rightarrow 0$

$$\zeta \frac{\partial \varepsilon}{\partial \zeta} \Big|_{\zeta=0} = 0. \quad (3.32)$$

Meeting conditions (3.30) to (3.32) from (3.28) we obtain the result:

$$\frac{\partial T}{\partial t} \Big|_{\zeta=0} = 0. \quad (3.33)$$

4. Equations for Vertical Velocity.

Vertical motion appears as a component of the mechanism of atmospheric circulation. The redistribution of kinetic, potential and internal energies of air from one level to the

other is realized through vertical currents. Therefore, in the study of atmospheric processes the examination of vertical motion should be given careful attention.

At the same time, vertical motions are of interest as such because they are the main factor in the process of formation of cloud conditions and precipitation.

To obtain an equation for vertical motion ζ in the atmosphere, it is indispensable to exclude derivatives $\frac{\partial^2 \zeta}{\partial t^2}$ and $\frac{\partial T}{\partial t}$ from system (2.1) and (2.3).

For this purpose we differentiate the equation (2.1) with ζ . We obtain:

$$\frac{l^2}{Pg} \frac{\partial^2 \zeta}{\partial \zeta^2} = \frac{\partial}{\partial \zeta} \left(\Delta \frac{\partial z}{\partial t} \right) + \frac{\partial}{\partial \zeta} \left[\frac{g}{l}(z, \Delta z) + \beta \frac{\partial z}{\partial x} \right]. \quad (4.1)$$

The equation of statics gives us correlations

$$\frac{\partial}{\partial \zeta} \left(\Delta \frac{\partial z}{\partial t} \right) = - \frac{R}{g} \frac{1}{\zeta} \Delta \frac{\partial T}{\partial t}, \quad (4.2)$$

$$\frac{\partial}{\partial \zeta} \left[\frac{g}{l}(z, \Delta z) + \beta \frac{\partial z}{\partial x} \right] = - \frac{R}{g} \frac{1}{\zeta} \left[\frac{g}{l}(T, \Delta z) + \frac{g}{l}(z, \Delta T) + \beta \frac{\partial T}{\partial x} \right]. \quad (4.3)$$

We take (4.2) and (4.3) as a basis for (4.1). Then we obtain

$$\frac{l^2}{PR} \frac{\partial^2 \tau}{\partial \xi^2} = - \frac{1}{f} \Delta \frac{\partial T}{\partial t} - \frac{1}{f} \left[\frac{g}{l} (T, \Delta z) + \frac{g}{l} (z, \Delta T) + \beta \frac{\partial T}{\partial x} \right]. \quad (4.4)$$

We exclude the derivative $\frac{\partial T}{\partial t}$ of (4.4) through the equation of heat flux (2.2) and obtain:

$$\zeta^2 \frac{\partial^2 \tau}{\partial \xi^2} + \frac{R^2}{gl^2} \Delta [T(\alpha - \gamma) \tau] = f_3(x, y, \zeta), \quad (4.5)$$

$$f_3(x, y, \zeta) = -P \frac{R}{l^2} \left\{ \frac{g}{l} \Delta (T, z) + \frac{\Delta \epsilon}{c_p} + \frac{g}{l} (T, \Delta z) + \frac{g}{l} (z, \Delta T) + \beta \frac{\partial T}{\partial x} \right\}. \quad (4.6)$$

We assume that the changes of value $T(\alpha - \gamma)$ in regard the vertical and horizontal are small compared to the relative changes of vertical velocity. Thus equation (4.5) can be formulated in the following manner:

$$\zeta^2 \frac{\partial^2 \tau}{\partial \xi^2} + m^2 \Delta \tau = f_3(x, y, \zeta), \quad (4.7)$$

m^2 - parameter introduced above.

The limit conditions for equation (4.7) are as follows:

$$\begin{aligned} \tau|_{\xi=0} &= 0, \\ \tau|_{\xi=1} &= \tau_0, \end{aligned} \quad (4.8)$$

$$\tau_0 = g \rho_0 \frac{\partial z_0}{\partial t} \quad - \text{small quantity.}$$

We now come to the cylindrical coordinate system r, φ, ξ , already used above.

In this coordinate system equation (4.7) assumes the following shape:

$$\xi^2 \frac{\partial^2 \bar{\tau}}{\partial \xi^2} + \frac{1}{r} \frac{\partial}{\partial r} r \frac{\partial \bar{\tau}}{\partial r} + \frac{1}{r^2} \frac{\partial^2 \bar{\tau}}{\partial \varphi^2} = f_3(r, \varphi, \xi). \quad (4.9)$$

Integrating this equation with φ from zero to 2π , we obtain the following equation for function $\bar{\tau} = \frac{1}{2\pi} \int_0^{2\pi} \tau d\varphi$:

$$\xi^2 \frac{\partial^2 \bar{\tau}}{\partial \xi^2} + \frac{1}{r} \frac{\partial}{\partial r} r \frac{\partial \bar{\tau}}{\partial r} = F_3(r, \xi), \quad (4.10)$$

$$F_3(r, \xi) = \frac{1}{2\pi} \int_0^{2\pi} f_3(r, \varphi, \xi) d\varphi. \quad (4.11)$$

The limit conditions of equation (4.10) are as follows:

$$\left. \begin{aligned} \bar{\tau}|_{\xi=0} &= 0, \\ \bar{\tau}|_{\xi=1} &= \bar{\tau}_0(r). \end{aligned} \right\} \quad (4.12)$$

The solution of equation (4.10) is obtained through

$$\bar{\tau}(r, \xi) = \int_0^{\infty} S(\rho, \xi) J_0(r\rho) \rho d\rho. \quad (4.13)$$

As we have done above, we assume that function $F_3(r, \zeta)$ and $\bar{F}_0(r)$ can be presented as intervals according to Fourier-Bessel

$$\left. \begin{aligned} F_3(r, \zeta) &= \int_0^{\infty} J_0(r\rho) \rho d\rho \int_0^{\infty} F_3(r', \zeta) J_0(\rho r') r' dr', \\ \bar{F}_0(r) &= \int_0^{\infty} J_0(r\rho) \rho d\rho \int_0^{\infty} \bar{F}_0(r') J_0(\rho r') r' dr'. \end{aligned} \right\} (4.14)$$

We take (4.13) and (4.14) as basis for (4.10) and (4.12). Analogous to the above we obtain the following differential equations and limit conditions for the new unknown function $S(\rho, \zeta)$:

$$\zeta^2 \frac{\partial^2 S}{\partial \zeta^2} - \rho^2 S = G_3(\rho, \zeta), \quad (4.15)$$

$$\left. \begin{aligned} S|_{\zeta=0} &= 0, \\ S|_{\zeta=1} &= G_3(\rho), \end{aligned} \right\} (4.16)$$

$$\left. \begin{aligned} F_3(\rho, \xi) &= \int_0^{\infty} F_3(r', \xi) J_0(\rho r') r' dr', \\ G_3(\rho) &= \int_0^{\infty} \bar{v}_0(r') J_0(\rho r') r' dr'. \end{aligned} \right\} (4.17)$$

Following is the general solution of the equation (4.15)

$$s(\rho, \xi) = c_1 \xi^{\frac{1}{2} + \mu} + c_2 \xi^{\frac{1}{2} - \mu} + \frac{1}{2\mu} \int_0^{\xi} F_3(\rho, \eta) \sqrt{\frac{\xi}{\eta}} \left[\left(\frac{\xi}{\eta} \right)^{\mu} - \left(\frac{\eta}{\xi} \right)^{\mu} \right] d\eta, \quad (4.18)$$

as above

$$\mu = \sqrt{\frac{1}{4} + \rho^2}.$$

In the selection of C_1 and C_2 we will see to it that solution (4.18) meets limit conditions (4.16). The result is as follows

$$s(\rho, \xi) = -\frac{1}{2\mu} \int_0^1 F_3(\rho, \eta) \sqrt{\frac{\xi}{\eta}} - \left[\delta_1 \left(\frac{\eta}{\xi} \right)^{\mu} + \delta_2 \left(\frac{\xi}{\eta} \right)^{\mu} - (\xi \eta)^{\mu} \right] d\eta + G_3(\rho) \xi^{\frac{1}{2} + \mu}. \quad (4.19)$$

Bearing in mind that $F_3(\rho, \xi)$ and $G_3(\rho)$ can be expressed the integral way (4.17), we will use it as a basis for (4.19). Moreover, the obtained expression is taken as a bases for (4.13). Now we change the order of integration and assume th $r = 0$.




Fig. 18. Graph of Function $\xi I(\xi, r)$ for various ξ .

Then we obtain:

$$\tau(\xi) = - \int_0^1 \int_0^{\infty} F_3(r', \eta) M_3(\xi, r', \eta) r' dr' d\eta + \int_0^{\infty} \tau_0(r') \xi I(\xi, r') r' dr', \quad (4.2)$$

$M_3(\xi, r, \eta)$ and $I(\xi, r)$ - Functions (3.22) and (3.23) introduced previously.

Substituting $F_3(r, \eta)$ and $\tau_0(r)$ in (4.20) according to (4.11) and (4.12) through (4.6), we formulate solution (4.20) in the final form as follows:

$$\begin{aligned} \tau(\xi) = & \frac{1}{2\pi} \int_0^1 \int_0^\infty \int_0^{2\pi} P \frac{R}{l^2} \left\{ \frac{g}{l} \Delta(T, z) + (T, \Delta z) + (z, \Delta T) \right. \\ & + \frac{\Delta E}{c_p} + \beta \frac{\partial T}{\partial x} \left. \right\} \xi M_3(\xi, r, \eta) r d\varphi dr d\eta + \\ & + \frac{1}{2\pi} \int_0^\infty \int_0^{2\pi} g \rho_0 \frac{\partial z_0}{\partial t} \xi I(\xi, r) r d\varphi dr. \end{aligned} \quad (4.2)$$

The graphic presentation of function $\xi I(\xi, r)$ with various ξ (Fig. 18) shows that the second term in the right part (4.2) quickly vanishes at the respective altitude.

Considering that $\tau_0 = g \rho_0 \frac{\partial z_0}{\partial t}$ usually does not exceed 10 millibar (12 hours), we may disregard the second component in (4.21).

Fig. 19 and 20 show the fields of functions $\xi M_3(\xi, r, \eta)$ for different ξ .




Fig. 19. Field of Function $\xi M_3(\xi, r, \eta)$ with $\xi = 0.9$




Fig. 20. Field of Function $\xi M_3(\xi, r, \eta)$ with $\xi = 0.7$

The properties of function $M_3(\xi, r, \eta)$ show that in the formation of vertical motions in the intermediary troposphere the dynamic processes in the intermediary and upper parts of the troposphere play the decisive part. The vertical motions at the level of 3-5 km are mainly determined by particularities of the pressure fields, temperature and heat flux at the 3-8 levels.

Equation (4.21) shows that in the source region of the heat flux the terms of type $\Delta \left[-\frac{g}{T} (T, z) + \frac{E}{c_p} \right]$ will always provide the anabatic motions. But in the heat discharge regions they will provide the catabatic motions. Consequently, part of the heat influx entering any important region of the atmosphere is transferred, together with the vertical currents to higher layers.

On the other hand, the heat discharge is partly compensated by the transfer of heat through vertical currents from higher layers.

This points out the important role played by vertical motions of the atmosphere in the redistribution of thermic energy.

Factor $\beta \frac{\partial T}{\partial x}$ in the solution for τ which appears in the initial equations of the parameter change of the coriolis according to latitude, is important only in the presence of great thermic disturbances of the atmosphere. In the eastern part of the thermal crest this factor becomes an anabatic component of vertical velocity, in the western part of the crest however, a catabatic component.

Literature.

1. E.N. Elinova. The Hydrodynamic Theory of Pressure Waves, Temperature Waves, and Action Centers of the Atmosphere. State Academy of Sciences of the USSR, 39, No. 7, 1943.
2. I.A. Kibel. Contribution on the Meteorology of Mechanical Equations of Baroclyne Fluid. Bull., Ac.Sc. of the USSR, Geogr. and Geophys. Series, No.r, 1940.
3. A.M. Obuchov. On the Problem of Geostrophic Winds. Bull. Ac.Sc. USSR, Geogr. and Geophys. Series, No.4, 1949
4. I.M. Ryshik and I.S. Gradstein. Tables on Integrals, Sum totals, Series and Products. State Techn. Publ., M.-L., 1951.
5. S.N. Troitzkyj. Determination of Aerosymptotic Criteria for Weather Forecasts, 1933.
6. M.E. Shvetz. Determination of Vertical Velocities in Mob Masses Through Hydrodynamic Equations, Bull., Ac.Sc. USSR, Geogr. and Geophys. Section, No.4, 1950.
7. Charney, Fjoertoft, Neumann. Numerical Integration of Ba tropic Vorticity Equations. Tellus, Vol.2, No.4, 195

8. Ertel. On New Atmospheric Equations of Motion. Meteorol.
Mag., Vol. 58, No.3, 1941.

9. L. Richardson. Weather Forecasting by Numerical Processes.
Cambridge, 1922.

- END -