

PHOTONIC CRYSTALS AND LIGHT LOCALIZATION

NATO Advanced Study Institute
June 19-30, 2000
Crete, Greece

DNIS QUALITY INSPECTED 4

20000816 045

DISTRIBUTION STATEMENT A
Approved for Public Release
Distribution Unlimited

Organizing Committee Members:

| NAME: | INSTITUTION: |
|----------------|--|
| E. N. Economou | Professor, Physics Department, University of Crete, Heraklion, Crete, Greece |
| S. John | Professor, Physics Department, University of Toronto, Ontario, Canada |
| A. Lagendijk | Professor, Physics Department, University of Amsterdam , Amsterdam, The Netherlands |
| E. Ozbay | Professor, Physics Department, Bilkent University, Ankara, Turkey |

ASI SPONSOR

Scientific Affairs Division of the North Atlantic Treaty Organization

ASI CO-SPONSORS

Ames Laboratory - U.S. Department of Energy
University of Crete and Forth
U.S. Army and Research Laboratory--European Research Office
U.S. Army Research Office
General Secretariat for Research and Technology, Greece

CONFERENCE SECRETARY

Rebecca A. Shivers
Ames Laboratory
Iowa State University
Ames, Iowa 50011 U.S.A.
Telephone: (515) 294 -3481
FAX: (515) 294-0689
e-mail: shivers@ameslab.gov

111440

Number of visitors since 15 April 1999

e-mail: shivers@ameslab.gov
Last updated: 0830 CDST 22 MAY 2000

**NATO Advanced Study Institute
Photonic Crystals and Light Localization
Conference Program**

**June 18-30, 2000
Creta Maris Hotel
Limin Hersonnisou, Crete, Greece**

All sessions will be held in the Creta Maris Hotel Conference Room. The Lectures will have 50 minutes for presentations and 10 minutes for discussion. The Talks will have 25 minutes for presentation and 5 minutes for discussion. The Posters will be displayed during two evening sessions. The poster presenter will be allowed to show one (1) view-graph, which is most representative of the poster to pique the interest of the audience.

Sunday, June 18, 2000

19:30-21:30

Welcoming Party at the Creta Maris Hotel

Monday, June 19, 2000

- 8:30 Registration
9:30 Opening and Welcoming Remarks

Morning Session

Chairperson: E. N. Economou

- 10:00 **Lecture:** J. D. Joannopoulos, Massachusetts Institute of Technology, *The Magical World of Photonic Crystals: I*
- 11:00 Coffee Break
- 11:30 **Lecture:** A. Lagendijk, University of Amsterdam, *Localization of Light I*
- 12:30 **Lecture:** J. Pendry, Imperial College, *Computing the Response of Structured Metals to Light*
- 13:30 Lunch and Discussions

Evening Session

Chairperson: S. John

- 17:30 **Lecture:** S. Lin, Sandia National Laboratory, *Microfabrication of 2d and 3d Photonic Crystals: I*
- 18:30 **Panel Discussion:** *Photonic Band Gaps*
- 19:30 End of Session

Tuesday, June 20, 2000

Morning Session

Chairperson: A. Lagendijk

9:00

Lecture: J. D. Joannopoulos, Massachusetts Institute of Technology, *The Magical World of Photonic Crystals: II*

10:00

Lecture: S. John, Toronto, *A Theoretical Roadmap for 3-d Photonic Band Gap Materials and Their Uses: I*

11:00

Coffee Break

11:30

Lecture: E. Ozbay, Bilkent University, *Physics and Applications of Photonic Crystals*

12:30

Lecture: S. Noda, Kyoto, *Semiconductor Photonic Crystals*

13:30

Lunch and Discussions

Evening Session

Chairperson: Kurt Busch

17:30

Session I: **Oral Presentation of the Posters**

18:00 - 19:30

Posters

Wednesday, June 21, 2000

Morning Session

Chairperson: J. D. Joannopoulos

- 9:00 **Lecture:** J. Pendry, Imperial College, *Intense Focusing of Light Using Metals*
- 10:00 **Lecture:** S. John, Toronto, *A Theoretical Roadmap for 3-D Photonic Band Gap Materials and Their Uses: II*
- 11:00 Coffee Break
- 11:30 **Lecture:** E. N. Economou, Research Center of Crete, *Elastic Waves in Periodic Composite Material: I*
- 12:30 **Talk:** C. T. Chan, Hong Kong, *Photonic Crystals From Metal-coated Spheres*
- 13:30 Lunch and Discussions

Evening Session

Chairperson: D. A. Papaconstantopoulos

- 17:30 **Lecture:** E. N. Economou, Research Center of Crete, *Elastic Waves in Periodic Composite Materials: II*
- 18:00 **Talk:** J. Page, University of Manitoba, *Acoustic Band Gap Materials*
- 18:30 **Talk:** F. J. Garcia-Vidal, University of Madrid, *Surface Plasmons of Lamella Metallic Gratings*
- 19:00 **Talk:** D. R. Smith, University of California, San Diego, *Left-handed Materials*
- 19:45 End of the Session

Thursday, June 22, 2000

Morning Session

Chairperson: J. Pendry

- 9:00 **Lecture:** Ad Lagendijk, University of Amsterdam, *Random Lasers*
- 10:00 **Talk:** H. Cao, Northwestern University, *Localization of Laser Light in Active Random Media*
- 11:00 Coffee Break
- 11:30 **Talk:** Z. Vardeny, University of Utah, *Random and Distributed Feedback Lasing in Opal Photonic Crystals*
- 12:00 **Lecture:** C. M. Soukoulis, Ames Laboratory, Iowa State University, *Theory and Simulation of Random Lasers*
- 13:00 **Talk:** J. P. Woerdman, Leiden University, *Special Resonators for Random and Fractal Lasers*
- 13:30 Lunch and Discussions

Evening Session

Chairperson: K. Asakawa

- 17:30 **Talk:** S. Lin, Sandia National Laboratory, *Microfabrication of 2d and 3d Photonic Crystals: II*
- 18:00 **Talk:** T. Baba, Yokohama National University, *Light Propagation Characteristics in Defect Waveguides in a Photonic Crystal Slab*
- 18:30 **Talk:** D. C. Allan, Corning Incorporated, *Photonic Band Gap Fibers for Real World Applications*
- 19:00 **Talk:** R. E. Slusher, Lucent Technology—Bell Laboratories, *Symmetry, Laser Action, and Output Coupling in 2d Photonic Crystals*
- 19:30 End of the Session

Friday, June 23, 2000

Morning Session

Chairperson: D. J. Norris

- 9:00 **Lecture:** W. Vos, University of Amsterdam, *Strong Interaction Between Light and Photonic Crystals Made Using Self-Assembly*
- 10:00 **Lecture:** K. Busch, University of Karlsruhe, *Tunable Photonic Crystals*
- 11:00 Coffee Break
- 11:30 **Talk:** F. Meseguer, University of Valencia, *Inverse Opals Fabrication*
- 12:00 **Talk:** A. van Blaaderen, Utrecht University, *Manipulating Colloidal Crystallization for Photonic Applications*
- 12:30 **Talk:** S. G. Romanov, University of Wuppertal, *Light Manipulation of 3d Templated Photonic Crystals*
- 13:00 **Talk:** A. Polman, FOM, Amsterdam, *Fabrication and Characterization of Silicon-based Photonic Band Gap Materials*
- 13:30 Lunch and Discussions

Evening Session

Chairperson: A. Polman

- 17:30 **Talk:** K. Asakawa, The Femtosecond Technology Research Association (FESTA), *Photonic Crystal at FESTA in Japan - A Step Toward an Extremely Miniaturized Optical Pulse Control Device/circuit*
- 18:00 **Talk:** D. J. Norris, NEC Research Institute, *Self-organized 3D Photonic Crystals*
- 18:30 **Talk:** R. Biswas, Ames Laboratory, Iowa State University, *Structure of Colloidal Photonic Crystals*
- 19:00 **Talk:** H. M. van Driel, University of Toronto, *Optical Reflectivity of Strongly Photonic Crystals: Multiple Diffraction and Bragg Peaks That Do Not Move*
- 19:30 End of the Session

Saturday, June 24, 2000

Morning Session

Chairperson: R. Sprik

- 9:00 **Lecture:** W. Vos, University of Amsterdam, *Strong Interaction between Light and Photonic Crystals Made Using Self-Assembly: II*
- 10:00 **Talk:** Y. Fink, Massachusetts Institute of Technology, *A Comprehensive Approach to Photonic Crystal Formation in One, Two and Three Dimensions Using Self-assembled Block Copolymer Structures*
- 10:30 **Talk:** J. Martorell, University of Catalunya, *Quadratic Nonlinear Interactions in Photonic Crystals*
- 11:00 Coffee Break
- 11:30 **Talk:** V. Shalaev, New Mexico State University and Univ. P. and M. Curie, *Anderson Localization of Surface Plasmons and Nonlinear Optics of Percolating Composites.*
- 12:00 **Talk:** J. J. Saenz, University of Madrid, *Statistical Properties of Wave transport through Surface Corrugated Waveguides*
- 12:30 **Talk:** D. Cassagne, University of Montpellier, *Tight-binding Wannier Function Method for Photonic Band Gap Materials*
- 13:00 **Talk:** B. van Tiggelen, Grenoble, *Wave Transport Around the Localization Threshold*
- 13:30 Lunch and Discussions

Sunday, June 25, 2000

Field Trip

Monday, June 26, 2000

Morning Session

Chairperson: J. Page

- 9:00 **Lecture:** E. Yablonovitch, University of California, Los Angeles, *Electromagnetic Band Gaps at Photonic and Radio Frequencies: I*
- 10:00 **Lecture:** T. F. Krauss, University of St. Andrews, *Patterned Photonic Crystal Waveguides: I*
- 11:00 Coffee Break
- 11:30 **Lecture:** A. Lagendijk, University of Amsterdam, *Localization of Light: II*
- 12:30 **Talk:** D. Wiersma, University of Florence, *Propagation of Light in Partially Disordered Systems*
- 13:00 **Talk:** P. Halevi, Puebla, Mexico, *Homogenization of 2d Periodic Composites and Photonic Crystal Optics*
- 13:30 Lunch and Discussions

Evening Session

Chairperson: A. Moroz

- 17:30 **Lecture:** C. Beenakker, Leiden University, *Random Matrix Theory of Random Lasers and Photonic Localization: I*
- 18:30 Session II: **Oral Presentation of Posters**
- 19:00 **Posters**
- 20:00 End of the Session

Tuesday, June 27, 2000

Morning Session

Chairperson: M. Sigalas

9:00

Lecture: E. Yablonovitch, University of California, Los Angeles, *Electromagnetic Band Gaps at Photonic and Radio Frequencies: II*

10:00

Lecture: H. Benisty, Ecole Polytechnique, Palaiseau, *Applications of 2d Photonic Crystals to Optoelectronics: I*

11:00

Coffee Break

11:30

Talk: A. Forchel, Wuerzburg University, *Integration of Photonic Crystals with Active Optoelectronic Devices*

12:00

Talk: R. B. Wehrspohn, Max-Planck Institute of Microstructure, Halle, *Electrochemically Prepared High-Aspect-ratio Pore Arrays for 2d Photonic Crystal Applications*

12:30

13:00

Lunch and Discussions

Evening Session

Chairperson: E. Yablonovitch

17:30

Lecture: T. F. Krauss, University of St. Andrews, *Patterned Photonic Crystal Waveguides: II*

18.30

Panel Discussion: *Applications of Photonic Band Gap*

19:30

End of the Session

Wednesday, June 28, 2000

Morning Session

Chairperson: H. Benisty

- 9:00 **Lecture:** C. Beenakker, Leiden University, *Random Matrix Theory of Random Lasers and Photonic Localization: II*
- 10:00 **Talk:** Zhao-Qing Zhang, Hong Kong, *Photonic Crystals and Perturbation Theory*
- 10:30 **Talk:** Y. Vlasov, Ioffe Institute, *Influence of Disorder in Photonic Crystals*
- 11:00 Coffee Break
- 11:30 **Talk:** C. M. Soukoulis, Ames Laboratory, Iowa State University, *The Effects of Disorder in Photonic Band Gap Structures*
- 12:00 **Talk:** V. Kuzmiak, Chez Republic, *The Localization of Light in a Random 2d Dielectric Medium*
- 12:30 **Talk:** G. Cwilich, Yeshiva University, *Photonic Crystals and Light Localization*
- 12:45 **Talk:** S. Venakides Duke University, *Computational and Theoretical Results in Photonic Crystals*
- 13:00 **Panel Discussion:** *Photonic Band Gaps and Disorder*
- 13:30 Lunch and Discussions

Evening Session

Chairperson: P. Halevi

- 17:30 **Talk:** R. Biswas, Ames Laboratory, Iowa State University, *Antenna Applications of Photonic Crystals*
- 18:00 **Talk:** M. Sigalas, Agilent Laboratories, *Waveguide Bends in 3D Layer-by-Layer Photonic Band Gap Materials*
- 18:30 **Talk:** N. A. Nicorovici, University of Sydney, *Localization and Homogenization in Photonic Crystals*
- 19:00 **Talk:** A. Orłowski, Warsaw, *From Proximity Resonances to Anderson Localization: Universality Effects in Multiple Scattering*
- 19:30 End of the Session

Thursday, June 29, 2000

Morning Session

Chairperson: T. F. Krauss

- 9:00 **Lecture:** H. Benisty, Ecole Polytechnique, Palaiseau,
Applications of 2d Photonic Crystals to Optoelectronics: II
- 10:00 **Talk:** D. Chigrin, University of Wuppertal, *One-*
Dimensional Photonic Crystals: Omnidirectional Reflection
and Superprism Phenomena
- 10:30 **Talk:** A. Moroz, Utrecht University, *Complete Photonic*
Band Gap Structures Below Infrared Wavelengths
- 11:00 Coffee Break
- 11:30 **Lecture:** E. Yablonovitch, University of California, Los
Angeles, *Electromagnetic Band Gaps at Photonic and*
Radio Frequencies: III
- 12:30 **Panel Discussion:** *Photonic Crystals and Optoelectronic*
Devices
- 13:30 Lunch and Discussions

Evening Session

Chairperson: C. M. Soukoulis

- 17:30 **Remarks and Discussion:** *Open and Future Directions*
- 18:00 **Panel Discussion:** *Open and Future Directions*
- 19:30 **Closing Session:** End of the Institute

Abstracts of Lectures and Talks

Monday, June 19, 2000

- | | |
|------|-------------------------------|
| 8:30 | Registration |
| 9:30 | Opening and Welcoming Remarks |

Morning Session **Chairperson: E. N. Economou**

10:00 **Lecture:** J. D. Joannopoulos, Massachusetts Institute of Technology, *The Magical World of Photonic Crystals: I*

Within the past several years we have witnessed the emergence of a new class of materials which provide capabilities along a new dimension in the control and manipulation of light. These materials, known as "photonic crystals," are viewed ideally as a composite of a periodic array of macroscopic dielectric scatterers in a homogeneous dielectric matrix. A photonic crystal affects the properties of a photon in much the same way that a semiconductor affects the properties of an electron. Consequently, photons in photonic crystals can have band structures, localized defect modes, surface modes, etc. This new ability to mold and guide light leads naturally to many novel applications of these materials as optoelectronic components. An introductory survey including recent exciting developments in the field of photonic crystals is presented.

11:30 **Lecture:** A. Lagendijk, University of Amsterdam, *Localization of Light: I*

We will introduce and review the concept of localization of light. Ideas will be illustrated using some new experimental results. Experiments that will be discussed are transmission and enhanced backscattering. We will show that a recently introduced interferometric technique allows us to obtain phase information as well as to perform measurements with very short time scales.

12:30 **Lecture: J. Pendry**, Imperial College, *Computing the Response of Structured Metals to Light*

Metals present special challenges to computation and the problems are directly related to the plasmon modes they support. In a dielectric insulator details of the electromagnetic fields are confined to the scale of the wavelength of light in the dielectric. This limit is very helpful when planning a computation particularly if we use a real space description of the fields. Insulators are easy targets for computation in another respect: their dielectric functions tend to be independent of frequency and therefore we are free to use a variety of techniques including time domain methods.

Metals belong to the awkward squad when it come to calculation. The length scale of the wavefield can vary from the wavelength of light in free space down to the smallest detail of the metallic structure. Extreme concentrations of radiation into a few cubic nanometres are not uncommon and must be described in detail if an accurate overall response is to be calculated.

To handle these difficulties the transfer matrix method is commonly used as it computes at a fixed frequency. Hence dispersion of the dielectric function is not a problem. Another technique that is employed for metals is the adaptive mesh whereby the grid of points on which the field is specified expands or contracts according to the details expected in the fields. As it happens there are particularly easy ways of implementing this potentially tricky technique without rewriting computer code. Finally it is possible to adapt the 'FDTD', finite difference time domain, methods to work for dispersive materials whilst retaining the advantageous 'order N' scaling of this methods with system size.

Evening Session

Chairperson: S. John

17:30 **Lecture: S. Lin**, Sandia National Laboratory, *Microfabrication of 2d and 3d Photonic Crystals: I*

In this talk, the challenges and promises of three-dimensional (3D) photonic crystals in the infrared and optical wavelengths will be described. Emphasis will be placed on historical development, nano-fabrication techniques, construction of high-Q cavities, and the alternation of emission process by 3D photonic crystals. Additionally, the realization of a new class of 2D photonic crystal slab structures will be reported. The 2D crystal is capable of fully confining light and operates at 1.5mm communication wavelengths.

18:30

Panel Discussion: Photonic Band Gaps

19:30

End of Session

Tuesday, June 20, 2000

Morning Session

Chairperson: A. Lagendijk

9:00 **Lecture:** J. D. Joannopoulos, Massachusetts Institute of Technology, *The Magical World of Photonic Crystals: II*

10:00 **Lecture:** S. John, Toronto, *A Theoretical Roadmap for 3-d Photonic Band Gap Materials and Their Uses: I*

Unlike semiconductors which facilitate the coherent propagation of electrons, photonic band gap (PBG) materials execute their novel functions through the coherent localization of photons. I review and discuss our recent synthesis of a large scale three-dimensional silicon photonic crystal with a complete photonic band gap near 1.5 microns. When a PBG material is doped with impurity atoms which have an electronic transition that lies within the gap, spontaneous emission of light from the atom is inhibited. Inside the gap, the photon forms a bound state to the atom. Outside the gap, radiative dynamics in the colored vacuum is highly non-Markovian. I discuss the influence of these memory effects on laser action. When spontaneous emission is absent, the next order radiative effect (resonance dipole-dipole interaction between atoms) must be incorporated, leading to anomalous nonlinear optical effects which occur at a much lower threshold than in ordinary vacuum. I describe the collective switching of two-level atoms near a photonic band edge, by external laser field, from a passive state to one exhibiting population inversion. This effect is forbidden in ordinary vacuum. Finally, I discuss the prospects for a phase sensitive, single (3-level) atom quantum memory device, onto which information may be written by an external laser pulse.

11:30 **Lecture:** E. Ozbay, Bilkent University, *Physics and Applications of Photonic Crystals*

Photonic crystals are three dimensional periodic structures having the property of reflecting the electromagnetic (EM) waves in all dimensions, for a certain range of frequencies. Defects or cavities around the same geometry can also be built by means of adding or removing material. The electrical fields in such cavities are usually enhanced, and by placing active devices in such cavities, one can make the device benefit from the wavelength selectivity and the large enhancement of the resonant EM field within the cavity. We used three-dimensional photonic crystals to demonstrate resonant cavity enhanced detectors and antennas, and waveguiding.

We have demonstrated the resonant cavity enhanced (RCE) effect by placing microwave detectors in defect structures built around dielectric and metallic based photonic crystals. A power enhancement factor of 3450 was measured for planar cavity structures built around dielectric based photonic crystals. A resonant antenna was also built around the same structure. We measured a maximum directivity of 310, and a power enhancement of 180 at the resonant frequency of the cavity. The measured radiation patterns agree well with our theoretical results.

We used the defect structures to demonstrate waveguiding around layer-by-layer photonic crystals. An air gap introduced between two photonic crystal walls was used as the waveguide. We observed full (100%) transmission of the electromagnetic (EM) waves through these planar waveguide structures within the frequency range of the photonic band gap. The dispersion relations obtained from the experiment were in good agreement with the predictions of our waveguide model. By using coupled periodic defects, we have also experimentally observed a new type of waveguiding in a photonic crystal. A complete transmission was achieved throughout the entire waveguiding band.

12:30 **Lecture:** S. Noda, Kyoto, *Semiconductor Photonic Crystals*

In this talk, I would like to review the present status of 3D and 2D photonic crystals based on III-V semiconductors. On the 3D photonic crystals, the development of complete photonic crystals in optical wavelengths, and their applications to 3D sharp bend waveguide and control of light-emission will be discussed. On the 2D photonic crystals, a very unique coherent oscillation of 2D photonic crystal laser and some other functional devices will be described.

Evening Session

Chairperson: Kurt Busch

17:30 Session I: **Oral Presentation of the Posters**

18:00 - 19:30 **Posters**

Wednesday, June 21, 2000

Morning Session

Chairperson: J. D. Joannopoulos

9:00 **Lecture:** J. Pendry, Imperial College, *Intense Focusing of Light Using Metals*

We are familiar with the fact that in vacuum light may only be focused to an area of the same order as the wavelength. However metal surfaces support plasma modes that couple to incident light, and the focusing of surface plasma modes is not restricted by the free space wavelength. Such nanofocusing offers the possibility of huge concentrations of radiative energy in very small volumes, impossible to achieve with conventional focusing with lenses. For very modest power input, local concentrations of energy may be great enough to excite nonlinear effects that depend on the intensity.

In fact most nanostructured metal surfaces will produce such focusing as evidenced by the Surface Enhanced Raman effect which is known to be due to intense local concentrations of electric field. Further evidence of intense local field concentrations is dramatically demonstrated in recent experiments by Ebbesen et al., who studied transmission of light at a wavelength of 1500 nanometres, through a 200-nanometre-thick silver film into which cylindrical holes with a diameter of 150 nanometres had been cut. In this system light plays this Houdini-like trick by coupling to surface modes of the metal that are very closely confined to the surface and therefore find it much easier to filter through the tiny holes, as demonstrated in a recent theoretical paper.

10:00 **Lecture:** S. John, Toronto, *A Theoretical Roadmap for 3-D Photonic Band Gap Materials and Their Uses: II*

11:30 **Lecture:** E. N. Economou, Research Center of Crete, *Acoustic waves in Ordered and Disordered Systems: Recent Advances: I*

A brief review of the equations governing wave propagation in composite fluid-fluid, fluid-solid and solid-solid ordered and disordered systems will be presented together with relevant experimental and theoretical results. The role of multiscattering will be emphasized and demonstrated by considering acoustic propagation in bubbly water. Finally, preliminary recent results in exciting by light scattering optical-like vibrational modes in solutions containing copolymers will be presented.

12:30 **Talk:** C. T. Chan, Hong Kong, *Photonic Crystals From Metal-coated Spheres*

We show theoretically and experimentally that photonic band gaps can be realized using metal or metal-coated spheres as building blocks. Robust photonic gaps exist in any periodic structure built from such spheres when the filling ratio of the spheres exceeds a threshold. The frequency and the size of the photonic gap are essentially determined by the local order and they are not sensitive to the symmetry or the global long range order. These photonic crystals can withstand a large amount of disorder. For example, stacking faults cause almost no degradation. The gaps persist even in a random packing of such spheres. Good agreement between theory and experiment is obtained in the microwave regime. Calculations show that the approach can be scaled up to IR and optical frequencies.

Work done in collaboration with Z.L. Wang, W.Y. Zhang, X.Y. Lei, D.G. Zheng, W.Y. Tam and P. Sheng

Evening Session **Chairperson: D. A. Papaconstantopoulos**

17:30 **Lecture:** E. N. Economou, Research Center of Crete, *Acoustic Waves in Ordered and Disordered systems: Recent Advances: II*

18:00 **Talk:** J. Page, University of Manitoba, *Acoustic Band Gap Materials*

We use ultrasonic experiments to study the acoustic analogue of photonic crystals. As an illustrative example, we present recent pulsed transmission measurements on hexagonal close packed (hcp) arrays of 0.80-mm-diameter stainless steel beads immersed in water. High quality crystals are prepared, starting with extremely monodisperse steel beads, by using a manual assembly technique. We directly measure the ultrasonic wave field transmitted through slab-shaped samples of different thicknesses, allowing us to determine both the dispersion curve and amplitude transmission coefficient. We compare our experimental results with theoretical calculations based on multiple scattering theory (MST) for elastic waves, which is ideally suited to the spherical scatterer geometry of the present experiments. We find that the low-frequency band structure and transmission coefficient calculated using this approach are in very good agreement with our experiments.

Work performed in collaboration with A.L. Goertzen, Zhengyou Liu, C.T. Chan and Ping Sheng

18:30 **Talk: F. J. Garcia-Vidal**, University of Madrid, *Surface Plasmons of Lamella Metallic Gratings*

Surface electromagnetic modes (surface plasmons) of lamellar metallic gratings with very narrow and deep grooves or slits are reviewed from the theoretical point of view. In the reflection configuration [1] we will show how there are basically two kind of modes: surface plasmon polaritons (SPPs) and waveguide resonances in which light is strongly localized inside the grooves. Experimental evidence of the excitation of these two kind of plasmons in the optical regime is also given. In the transmission configuration [2], transmission resonances associated to the excitation of surface plasmons of the grating can appear in the spectrum. We will show that there are two possible ways of transferring light from the upper surface to the lower one: by the excitation of coupled SPPs on both surfaces or by the coupling of the incident light with waveguide resonances located in the slits. Both mechanisms can lead to almost perfect transmittance for those particular wavelengths.

19:00 **Talk: D. R. Smith**, University of California, San Diego, *Left-handed Materials*

We have recently introduced a new periodic metallic structure¹ based on two interlaced lattices of straight wires² and Split Ring Resonators³. The size of the scattering elements and the lattice constant are significantly less than the incident wavelength (at frequencies of interest), so that the system can be considered as an effective medium characterized by frequency-dependent material constants. By simulations and experimental measurements, we demonstrate a frequency band where both material constants—the permittivity (ϵ) and permeability (μ)—are simultaneously negative. When both material constants are negative, Maxwell's Equations allow for plane wave solutions, but with a twist: $\mathbf{E} \times \mathbf{B}$ is in the direction of $-\mathbf{k}$ rather than $+\mathbf{k}$ for plane waves, and thus a "left-hand" rule relates the three vectors. The result of this simple reversal is that many common electromagnetic phenomena, such as the Doppler shift, Cerenkov radiation, and even Snell's Law, are also reversed⁴, leading to the potential of novel electromagnetically active materials and surface coatings based on LH media.

Thursday, June 22, 2000

Morning Session

Chairperson: J. Pendry

- 9:00 **Lecture:** Ad Lagendijk, University of Amsterdam, *Random Lasers*
Disordered systems that can both scatter and amplify light have emerged as an exciting field. We will discuss in some detail the threshold behaviour while presenting some new experimental and theoretical results.

All our papers are available on line. This collection includes review papers, tutorials, and a collection of high-quality Dutch theses on multiple scattering, localization, and random lasers:

- 10:00 **Talk:** H. Cao, Northwestern University, *Localization of Laser Light in Active Random Media*
We have observed random laser action in highly disordered semiconductor powder and polycrystalline films. Since the scattering mean free path is less than the emission wavelength, recurrent light scattering arises and provides coherent feedback for lasing. Above the lasing threshold, discrete lasing modes appear in the spectrum in addition to a drastic increase of emission intensity. Our measurement of spatial distribution of laser light intensity indicates that photons are localized in spatial regions of the size on the order of optical wavelength. The laser emission from the random media can be observed in all directions. In addition to macroscopic-scale random media, we have observed lasing action in micrometer-sized random media. This discovery leads to the fabrication of microlaser with disordered medium. Anderson localization of light provides a new physical mechanism of optical confinement for microlaser.

- 11:30 **Talk:** Z. Vardeny, University of Utah, *Random and Distributed Feedback Lasing in Opal Photonic Crystals*
Stimulated emission and lasing have been studied in various scattering gain media, including luminescent B-conjugated polymer films and opal micro-crystals infiltrated with polymer or dye solutions. In addition to amplified spontaneous emission that is usually observed in high gain media upon intense photoexcitation, a lasing process was revealed using stripe excitation geometry. The new emission regime is characterized by a finely structured spectrum with narrow lines, of the order of 0.1nm. We interpret this laser like emission as due to random lasing where the optical feedback is provided by weak light scattering inside the gain medium. Coherent back-scattering measurements of the same media systems were found to be in agreement with this interpretation.

In addition we demonstrate a three-dimensional photonic crystal laser based on infiltrated dyes in a single crystal opal, operating simultaneously in the blue (435nm) and red (670-720nm) spectral regions. The feedback mechanism underlying this type of laser is the Bragg scattering off different sets of crystalline (hkl) silica planes, via a distributed feedback process. The laser emission wavelength was selected by proper orientation of the lasing axis with respect to the [111] and [220] directions for the red and blue lasing, respectively. Defect modes caused by stacking faults in the opal crystal were identified in the laser spectrum as sharp lines, which were found to depend on the specific illuminated area on the opal surface.

12:00 **Lecture:** C. M. Soukoulis, Ames Laboratory, Iowa State University, *Theory and Simulation of Random Lasers*

Disordered systems that both scatter and amplify light (the so-called random lasers) have been a fascinating subject to study. Over the last five years, there have been substantial theoretical and experimental efforts to unravel the mechanism that gives rise to this amazing behavior.

A model to simulate the phenomenon of random lasing is presented. It couples Maxwell's equations with the rate equations of electronic population in a disordered system. Finite difference time domain methods are used to obtain the field pattern and the spectra of localized lasing modes inside the system. A critical pumping rate exists for the appearance of the lasing peaks. The dependence of on the length of the system and the strength of disorders is obtained. The number of lasing modes increases with the pumping rate and the length of the system. There is a lasing mode repulsion. This property leads to a saturation of the number of modes for a given size system and a relation between the localization length and average mode length . Similar behavior is expected to be seen in photonic crystals, too.

13:00 **Talk:** J. P. Woerdman, Leiden University, *Special Resonators for Random and Fractal Lasers*

Random lasers do not necessarily require strong microscopic scattering. A suitably designed multi-mode laser cavity can act as a nonresonant system if it supports a manifold of spectrally overlapping modes. This manifold must have equal mode losses and strong mixing. We will discuss various cavity configurations within this category. Particularly interesting is the geometrically-unstable cavity, which has chaotic ray characteristics. The mode patterns of this cavity are self-similar and are characterized by a fractal dimension (Nature 402, 138 (1999)).

Evening Session Chairperson: K. Asakawa

17:30 **Talk:** S. Lin, Sandia National Laboratory, *Microfabrication of 2d and 3d Photonic Crystals: II*

18:00 **Talk:** T. Baba, Yokohama National University, *Light Propagation Characteristics in Defect Waveguides in a Photonic Crystal Slab*

Channel waveguides composed of a series of defects in an air hole photonic crystal slab was characterized theoretically and experimentally. They were similar to GaInAsP/SiO₂/InP waveguides described in the first reports [1,2]. We employed the SOI wafer for stable fabrication of long guides. The light transmission through the waveguides was confirmed in > 100-um-long waveguides at wavelengths of 1.47 to 1.60 um. The propagation loss was measured by the Fabry-Perot resonance method to be less than 3 dB for a waveguide length of 56 um. The dependence of the propagation loss on wavelength, slab thickness, channel width, etc., will be discussed with 3-D FDTD simulations. The possibility of various lightwave devices[3], i.e. sharp bends, Y blanches and directional couplers will also be discussed.

18:30 **Talk:** D. C. Allan, Corning Incorporated, *Photonic Band Gap Fibers for Real World Applications*

Conventional telecommunication optical waveguide glass fiber is the backbone of the internet revolution. This highly optimized and highly transparent material consists of a higher index core glass inside a lower index clad glass. Light is localized in the core by total internal reflection at the core/clad boundary. The transmission range between amplifiers of today's fibers, about 70km, is limited in part by the small but nonzero optical nonlinearity of silica. It is also limited in part by the small but nonzero amount of intrinsic scattering. It has recently been demonstrated, both theoretically[1] and experimentally[2], that photonic band gap confinement can be used to form a new kind of optical waveguide fiber. This photonic band gap fiber (PBGF) guides light in a LOW index core, in contrast to conventional fiber waveguides. This low index core can be air, allowing for dramatic reduction in nonlinearity and scattering. This presentation describes the fabrication of photonic band gap fibers, their experimental characterization, their predicted properties, and their significance in the context of optical telecommunication.

19:00 **Talk:** R. E. Slusher, Lucent Technology—Bell Laboratories, *Symmetry, Laser Action, and Output Coupling in 2d Photonic Crystals*

We have investigated the lasing and output coupling characteristics of two-dimensional photonic crystals. The lasers/couplers are experimentally realized with organic semiconductors and involve photo-excitation of patterned combinations of organic semiconductor and dielectrics (e.g., SiO₂). The patterning is accomplished by advanced optical lithography or electron-beam lithography followed by dry etching and the subsequent deposition of the organic gain medium as a thin coating. We show that by employing advanced optical lithography, which is driven by silicon electronics, feature sizes required for photonic crystals that operate in the visible can be attained. Organic media are also very convenient in investigating the basic physics of novel resonators and optical structures [1]. Most of the photonic crystals employed in this work do not possess a complete gap; however theoretical extensions of this work to systems with a complete bandgap will be discussed along with some experimental data.

The basic principle employed in our designs is based on our recent discovery that the output coupling characteristics of two-dimensional gratings (or PC's) are significantly different from one-dimensional gratings. Since we do not employ the lowest energy bandstructure features to provide the feedback necessary for laser oscillation, phase matching conditions result in diffractive coupling of some of the laser emission out of the plane of the waveguide. In two-dimensional PC's, phase matching conditions result in the coupling of light to one or a discrete number of directions instead of a cylindrical wave. We have designed numerous combinations of lasers and couplers which have the potential to couple to a single spot normal to the plane of the waveguide.

One design that has been successfully implemented is a square two-dimensional PC, which functions as a laser and output coupler. Detailed theoretical calculation show that a careful study of band-structure features is required to attain an accurate understanding of mechanism of laser action in two-dimensional photonic crystals. Laser action occurs at points in the Brillouin zone where the density of photon states is peaked and where the photon group velocity goes to zero. Laser action does not necessarily take place at band extrema; occurring at saddle points where the density of states is sometimes higher. We have also found a variety of ways to combine lasers and couplers resulting in composite devices with interesting and potentially useful optical properties.

Friday, June 23, 2000

Morning Session

Chairperson: D. J. Norris

9:00 Lecture: W. Vos, University of Amsterdam, *Strong Interaction Between Light and Photonic Crystals Made Using Self-Assembly: I*

The following issues will likely be discussed: - the fabrication of 3D photonic crystals that strongly interact with light and that are made using self-organizing systems as templates, and new developments. - the detailed characterization of the crystals, by electron microscopy and by synchrotron small-angle x-ray scattering. - optical experiments such as reflectivity, transmission, enhanced backscattering, phase-sensitive ultrashort-pulse interferometry. - studies of the emission by light sources (excited molecules or atoms) located *inside* photonic crystals, both cw and time-resolved. - the interpretation of the experiments.

10:00 Lecture: K. Busch, University of Karlsruhe, *Tunable Photonic Crystals*

We demonstrate that complete three-dimensional Photonic Band Gaps (PBG), spanning roughly 10% and 15% of the gap center frequency are attainable by incomplete infiltration of an opal with silicon and germanium, respectively. In addition, we show that when an optically birefringent nematic liquid crystal is infiltrated into the void regions of an inverse opal PBG material, the resulting composite material exhibits a completely tunable PBG. Measurements on analogous two-dimensional Photonic Crystals show that the photonic band edge may be shifted by more than 100 nm. Furthermore, we present results of bandstructure calculations for Photonic Crystals whose constituent materials exhibit Faraday or natural optical activity. This provides additional flexibility for tuning the optical properties of Photonic Crystals.

11:30 Talk: F. Meseguer, University of Valencia, *Inverse Opals Fabrication*

Here we report on different approaches to the fabrication of inverse opals. We start from silica opal templates with sphere size in the range between 200 nm and 1300 nm. The opal porous lattice is fully (above 90 % of the opal void volume) infiltrated with semiconductors (CdS, Ge) as well as polymers by several methods. Afterwards, the template is removed, from the composite, by a soft chemical etching method. Thus, an inverse opal is obtained. The periodicity of the template can be chosen to guarantee that photonic gaps or pseudogaps are in the transparency region of the bulk infiltrated material.

Optical properties of CdS, Ge, and polymer inverse opals, as well as the comparison with theoretical calculations of the transmission and reflectance are reported. Ge inverse opal has a huge refractive index contrast (4) well above the onset value to present a sizeable full band gap.

12:00 **Talk:** A. van Blaaderen, Utrecht University, *Manipulating Colloidal Crystallization for Photonic Applications*

Colloidal particles are the ideal building blocks for photonic crystals because their properties can be tuned by chemical modification and their self-organization or crystallization into 3D periodic lattices can be manipulated by changing the interaction potentials between the particles and by using external fields. In this contribution we focus 1) on how large colloidal crystals with known crystal orientation and symmetry can be grown by colloidal epitaxy. And 2) how colloidal crystallization can be manipulated by electric fields; both in symmetry, size and orientation of the crystals.

12:30 **Talk:** S. G. Romanov, University of Wuppertal, *Light Manipulation of 3d Templated Photonic Crystals*

The photonic energy band structure, resulting from a suitable periodic modulation of dielectric properties in a solid, makes it possible to control the propagation and emission of light without dissipation. To realise the full extent of these possibilities, electromagnetic waves have to be confined in 3-dimensions. Three-dimensional crystalline packages of silica or polymeric balls (synthetic opals and similar structures) as well as their replicas, i.e. inverted opal, have been recognised as suitable models of 3-dimensional photonic crystals. In our lecture the following topics related to experimental studies of templated opal-based photonic crystals operating in the visible range of spectrum will be discussed: Technological aspects of the fabrication of 3D opal-like and opal-templated photonic crystals. Examination techniques and results. Bragg diffraction from silica- and polymeric-based photonic crystals, mapping of corresponding anisotropic photonic bandgap structures. Enhancement of the photonic stop-band (bandgapwidth enlargement, bandgap dispersion smoothening) by means of infilling and coating opals with high refractive index materials (S, TiO₂, CdS, InP, GaP) and inversion of opals. Comparison of stop-band characteristics of bulk and thin-film opal-like photonic crystals Design and testing of organic-inorganic opal-based photonic crystals (e.g. opal-TiO₂-polymer-dye/rare earth ion structures). Peculiarities of light propagation in photonic crystals with double PBG structure (GaP-opal/Air-opal). Probing the density of photonic states in opal-like photonic crystals. Self-focusing of the emission from opal-like photonic crystals, focusing enhancement with the increase of the refractive index contrast. Modelling of this effect using the superprism approach. Probing the changes in the spontaneous emission rate of emitters embedded in photonic crystals. Comparison of emission from wide (laser dyes) and narrow (rare earth ions) band emitters embedded in photonic crystals. Prospective opal-based device structures.

13:00 **Talk:** A. Polman, FOM, Amsterdam, *Fabrication and Characterization of Silicon-based Photonic Band Gap Materials*

Silicon is one of the best known materials in the world. It is available in high purity and the processing technologies for silicon are by now very well established. While the properties of silicon as a semiconductor are well explored, it is not much used in photonic technology. However, Si is an excellent optical waveguide material, as it is completely transparent at 1.5 μm . In this work we have exploited silicon as a substrate and optical waveguide material in a variety of novel photonic structures. Our experiments focussed on a) the design of two-dimensional photonic bandgap structures, b) measurements on the spontaneous emission rate of atoms to probe the local optical density of states. 2-D photonic crystal waveguides are fabricated in Si by reactive ion etching of Si pillars that are 5 μm tall, 205 nm in diameter, and arranged in a square lattice with a 570 nm pitch. Bandstructure calculations predict a gap for TM modes at a wavelength of 1.5 μm with a gap width equal to 37 % of the center energy. In order to meet the severe nano-tolerance requirements in such a device, the SF₆/O₂ electron cyclotron resonance plasma process conditions at reduced temperature are optimized. By removing an array of Si pillars, the waveguide is defined. In order to achieve confinement in the third dimension, a 2 μm thick region of the Si structure was amorphized before processing, using 4 MeV Xe irradiation. Input and output waveguides are integrated with the structure. In order to measure the local density of states in such photonic structures, optical probe ions must be incorporated. We have studied Er ion implantation into Si and found the optimum implantation and annealing conditions to obtain optically active Er, luminescing at 1.535 μm in Si. The luminescence shows strong quenching above 200 K, but lifetimes as long as 1 ms are achieved at 77 K. We have also studied the coating of the Si pillar structures with a novel wet chemical process, in which we have used a reaction between tetraethoxysilane, ammonia, and water to nucleate and grow well-controlled layers of SiO₂ on surfaces with extreme aspect ratio. Using confocal microscopy we show the uniform coating of such layers doped with an eosin dye. Finally, we show how we can strongly affect the radiative lifetime of Er probe ions implanted into silica colloidal particles with a diameter of 300 nm, by comparing three geometries, 1) placed on a Si substrate, 2) arranged in a 3-D crystal, or 3) surrounded by an index-matching fluid.

Evening Session

Chairperson: A. Polman

17:30 **Talk:** K. Asakawa, The Femtosecond Technology Research Association (FESTA), *Photonic Crystal at FESTA in Japan - A Step Toward an Extremely Miniaturized Optical Pulse Control Device/Circuit*

AlGaAs-based two-dimensional photonic band-gap crystal (PBC) slab test devices have been fabricated using electron beam lithography and a reactive ion beam etching technology. Air hole-based triangular lattices with lattice constants ranging from 0.32 to 0.45 μm have been fabricated on both semiconductor-clad (SC) and air-bridge (AB) structures. In a transmission/reflection spectrum with wavelength ranging from 0.85 to 1.1 μm on an SC-type device, a band-gap with large attenuation more than 30 dB has been observed for both E- and H-polarization. Quantitatively more precise characteristics are now under measurement using a sub-ps pulse propagation technique. Prospect for achievement of a PBC-based optical pulse control device involving a branching/interferometer waveguide with optically nonlinear (NL) medium will be addressed. A site-controlled quantum dot assisted with a nano-probe technique will also be shown as a representative of NL medium. An AB-type device will be presented at a poster session in the meeting. This work was supported by NEDO within the framework of the Femtosecond Technology Project.

18:00 **Talk:** D. J. Norris, NEC Research Institute, *Self-organized 3D Photonic Crystals*

Recently colloidal self-organization has been used to make 3D photonic crystals from a variety of materials. A typical approach is to induce sub-micron silica spheres to organize on an fcc lattice. This template, which is periodic on an optical length scale, then serves as a three-dimensional scaffolding into which another material may be infiltrated. Subsequent removal of the template by selective etching then yields a 3D photonic crystal, also called an inverted opal. These structures are particularly interesting since they have been predicted to exhibit an omnidirectional photonic band gap and enhanced optical nonlinearities. Here we show how to use colloidal self-organization to make three different materials: 1) semiconductors to explore high refractive index contrast, 2) metallo-dielectrics to explore coupling to plasmons, and 3) conjugated polymers to explore optical nonlinearities. We then discuss the optical properties of our photonic crystals. In particular, we use optical microscopy to probe a single crystalline domain in each of our samples. By measuring spatially resolved reflection and emission spectra, inhomogeneities due to averaging over inherent disorder (always present in self-assembled photonic crystals) can be eliminated. "Defect-free" spectra are obtained, from which the intrinsic photonic band structure can be extracted. Furthermore, this technique allows us to probe the specific band structure at higher frequencies, up to the "4th-order" stop band. This capability is important since the omnidirectional photonic band gap should be at the "2nd-order" stop band in inverted opals (i.e. between the 8th and 9th bands).

18:30 **Talk:** R. Biswas, Ames Laboratory, Iowa State University, *Structure of Colloidal Photonic Crystals*

I will survey theoretical calculations and experimental measurements on colloidal photonic crystals synthesized at Iowa State University. These crystals show reflectivity peaks at optical wavelengths. Optical measurements on these photonic crystals indicate a geometry that is considerably distorted from ideal. Band structure calculations of the positions of the stop bands will be utilized to determine the structure of the crystal and refractive index of the background material. The dependence of photonic properties with stacking and disorder will also be discussed. Alternative photonic lattices with photonic gaps will be discussed. (In collaboration with G. Subramania, K. Constant, M. M. Sigalas, K.-M. Ho, C. M. Soukoulis.) Supported by the U.S. Dept. of Energy/Office of Basic Energy Sciences.

19:00 **Talk:** H. M. van Driel, University of Toronto, *Optical Reflectivity of Strongly Photonic Crystals: Multiple Diffraction and Bragg Peaks That Do Not Move*

We report results of angle, polarization and frequency resolved optical reflectivity measurements on fcc crystals consisting of air spheres in titania. The carefully grown crystals show excellent structural order, as confirmed by x-ray scattering and electron microscopy. The reflectivity spectra of these strongly photonic crystals show unusual features in comparison to simple Bragg diffraction of weakly photonic crystals. Over a wide range of reflection angles, the (111) Bragg peak splits into at least two peaks that show an avoided crossing. Calculated photon dispersion curves show that the multiple peaks result from band repulsions of Bloch states due to simultaneous (200) and (111) Bragg resonances. Such multiple Bragg diffraction results in flat dispersion relations, and thus plays an essential role in photonic band gap formation. In the range of the 2nd order Bragg diffraction, where a complete photonic band gap is predicted for fcc crystals with higher dielectric contrast than our materials, at least three peaks are observed with central frequencies independent of reflection angle over a wide range. Theoretical analysis allows us to assign these peaks to stop-gaps or characteristics of dispersionless bands. A significant conclusion is that reflectivity spectra can mimic the signature of a complete photonic band-gap without such a band-gap actually being present. We thank Judith Wijnhoven for sample preparation, Henry Schriemer and Rudolf Sprik for discussions, and Ad Lagendijk especially for organizing HMvD's visit.

Saturday, June 24, 2000

Morning Session

Chairperson: R. Sprik

9:00 **Lecture:** W. Vos, University of Amsterdam, *Strong Interaction between Light and Photonic Crystals Made Using Self-Assembly: II*

10:00 **Talk:** Y. Fink, Massachusetts Institute of Technology, *A Comprehensive Approach to Photonic Crystal Formation in One, Two and Three Dimensions Using Self-assembled Block Copolymer Structures*

A conceptual framework for creating photonic crystals from self-assembling block copolymers has been formulated. In order to form useful band gaps in the visible regime, periodic dielectric structures made of typical block copolymers need to be modified to obtain appropriate characteristic distances and dielectric constants. Moreover, the absorption and defect concentration must also be controlled. This affords the opportunity to tap into the large structural repertoire, the flexibility and intrinsic tunability that these self-assembled block copolymer systems offer. A symmetric poly(styrene-*b*-isoprene) (~400k M) block copolymer was used to achieve a photonic band gap in the visible regime. By swelling the diblock copolymer with lower molecular weight constituents, we have been able to control the location of the stop band across the visible regime. One and three-dimensional crystals have been formed by changing the volume fraction of the swelling media. Methods for incorporating defects of prescribed dimensions into the self-assembled structures have been explored leading to the construction of a microcavity light-emitting device.

10:30 **Talk:** J. Martorell, University of Catalunya, *Quadratic Nonlinear Interactions in Photonic Crystals*

Soon after the birth of the field of photonic crystals it became clear that the intrinsic properties of such materials made them very suitable to control the nonlinear interaction between the radiation and matter. In this presentation, I will show that some of these properties are particularly useful when the interaction that we consider is the second order nonlinear interaction. For an efficient generation of second harmonic light or for any other type of efficient parametric interaction, one must combine, within the same material, a lack of inversion symmetry and a large nonlinearity with a phase matching mechanism. I will present experimental evidence that 3-dimensional photonic crystals offer the ideal frame to combine in the same material these three requirements. On the other hand, I will show that 1-dimensional truncated periodic structures can be used to enhance cascaded second order nonlinear interactions and at the same time eliminate the phase sensitivity of this type of quadratic interaction.

11:30 **Talk:** V. Shalaev, New Mexico State University and Univ. P. and M. Curie, *Anderson Localization of Surface Plasmons and Nonlinear Optics of Percolating Composites*.

Anderson localization of surface plasmons in random metal-dielectric composites near the percolation threshold has been predicted and observed using scanning near-field optical microscopy. A scaling theory of local-field distributions and optical nonlinearities is developed. The theory predicts that the local fields are very inhomogeneous and consist of sharp peaks representing localized surface plasmons (sp). The localization maps the Anderson localization problem described by the random Hamiltonian with both on- and off-diagonal disorder. The local fields exceed the applied field by several orders of magnitudes resulting in giant enhancements of various optical phenomena. The white-light generation dramatically enhanced by the localized sp has been experimentally obtained. A feasibility of nonlinear single-molecule spectroscopy in percolation films has been shown.

12:00 **Talk:** J. J. Saenz, University of Madrid, *Statistical Properties of Wave transport through Surface Corrugated Waveguides*

Statistical properties of waves transmitted [1] and reflected [2] from random media are discussed in terms of Random Matrix Theory (RMT). Exact numerical calculations in surface randomly corrugated waveguides are presented. By increasing the length of the disordered region, we are able to analyze the transition from ballistic to diffusive regimes as well as from diffusion to localization in detail. Although the transport in these systems is strongly nonisotropic, the analysis of the probability distributions of both the transport coefficients and the spatial intensity [3] remarkably agrees with the analytical predictions of RMT. Our results predict unexpected strong finite size effects when the number of propagating channels is small [3].

12:30 **Talk:** D. Cassagne, University of Montpellier, *Tight-binding Wannier Function Method for Photonic Band Gap Materials*

Using the concept of generalized Wannier functions, adapted from the electronic theory of solids, we demonstrate for two-dimensional photonic crystals the existence of a localized state basis and we establish an efficient computational method allowing a tight-binding-like parameter free modelization of any dielectric structure deviating from periodicity. Examples of numerical simulations using this formalism, including modal analysis of microcavities and waveguides and calculations of the transmission coefficients are presented to prove the ability of this approach to deal accurately with large scale systems and complex structures.

13:00 **Talk:** B. van Tiggelen, Grenoble, *Wave Transport Around the Localization Threshold*

I present a new theory that is capable of describing coherent backscattering and transmission of waves very close to the mobility edge of localization. Theory is a generalization of the self-consistent theory of localization, devised 20 years ago by Götze, Vollhardt and Wölfle. It is able to deal with boundary condition, an important requirement for a quantitative description of coherent backscattering. A comparison will be presented to experiments conducted in Florence and Amsterdam.

Monday, June 26, 2000

Morning Session

Chairperson: J. Page

- 9:00 **Lecture:** E. Yablonovitch, University of California, Los Angeles, *Electromagnetic Band Gaps at Photonic and Radio Frequencies: I*
Engineering design is sometimes inspired by Nature. The natural world is filled with crystals, periodic structures that interact with Schrodinger Waves. Drawing on this analogy, we are designing artificial crystal structures which are intended for Electromagnetic Waves instead. This has now unleashed the collective scientific imagination, engendering a profusion of synthetic electromagnetic crystal structures. In correspondence to semiconductor crystals these usually have an electromagnetic bandgap, a band of frequencies in which electromagnetic waves are forbidden. We will present a pictorial portfolio of various 2 and 3 dimensional crystal structures which have been conceived, and indicate the applications, such as opto-electronic light emitters, radio antennas, and color pigments, for which they are intended. In addition we will emphasize some of the new ideas that will be important for the future, including the further development of radio frequency bandgap structures, and of surface plasmon band structure.
- 10:00 **Lecture:** T. F. Krauss, University of St. Andrews, *Patterned Photonic Crystal Waveguides: I*
I will review the development of semiconductor-based planar photonic crystals, highlight the underlying concepts, materials and fabrication issues and point towards important issues for device applications. Examples of successful photonic crystal-based devices include photonic crystal waveguides and photonic integrated circuit elements as well as lasers and LEDs.
- 11:30 **Lecture:** A. Lagendijk, University of Amsterdam, *Localization of Light: II*
- 12:30 **Talk:** D. Wiersma, University of Florence, *Propagation of Light in Partially Disordered Systems*
The nematic phase of a liquid crystal is characterized by a global alignment of the molecules in a direction called the nematic director and an otherwise translational disorder. The strong opacity of the nematic phase comes about from local fluctuations of the nematic director that elastically scatter light. The strong scattering combined with the partial ordering of the nematic phase leads to an anisotropic diffusion process. There are several ways of combining nematics with either ordered (photonic crystals) or disordered (porous solids) dielectric structures. We have studied experimentally various aspects of anisotropic light diffusion, amongst which the anisotropy in the diffusion constant by means of time-resolved transmission experiments.

13:00 **Talk:** P. Halevi, Puebla, Mexico, *Homogenization of 2d Periodic Composites and Photonic Crystal Optics*

This talk concerns composite materials that have periodicity in a plane. They are formed by identical, long cylinders that constitute a two-dimensional, arbitrary Bravais lattice in the plane perpendicular to the cylinders. The cross-section of the cylinders is also of arbitrary form. The cylinder material and the interstitial material are assumed to be homogeneous and isotropic. In the last decade, such structures came to be known as photonic or phononic crystals - depending whether the materials involved are of transparent or of elastic nature. The study of these periodic composites usually focuses on a high-frequency band gap - the phononic or photonic gap in which, respectively, light or sound cannot propagate. Here, on the contrary, we are interested in the low-frequency regime, where the wavelength (corresponding to the Bloch vector) is much greater than the lattice spacing. Thus, we present a theory of homogenization applicable to a very general class of composites with two-dimensional periodicity. Our method is applicable to a wide class of materials: dielectric, magnetic, electrically conducting, thermally conducting, etc. For simplicity, the material properties are given in terms of the dielectric constants of the cylinder and host substances.

The homogenization is performed by taking the quasistatic limit of Maxwell's equations. In this limit, we prove that the composite behaves like either a uniaxial or a biaxial (homogeneous) optical medium. That is, it can be described by means of a hermitian dielectric tensor that can be diagonalized. It has three eigenvalues - the Principal Dielectric Constants - that, in general, are all different. We have derived exact and compact formulas for these dielectric constants, given in terms of summations over the two-dimensional reciprocal lattice and a matrix inversion. These formulas have been applied to a variety of Bravais lattices, cylinder cross-sectional shapes, and materials. In each case, the Principal Dielectric Constants are computed as a function of the cylinder filling fraction. Convergence is very rapid, and some of our results are of unprecedented precision.

Evening Session

Chairperson: A. Moroz

17:30 **Lecture:** C. Beenakker, Leiden University, *Random Matrix Theory of Random Lasers and Photonic Localization: I*

A random laser differs from a conventional laser in that the feedback is provided by multiple scattering from disorder rather than by confinement from mirrors. The statistical properties of the radiation are strikingly different because the scattering is chaotic (in contrast to integrable scattering in a conventional laser). Random-matrix theory provides a powerful tool for the study of the quantum optics of such chaotic resonators. Because it is a non-perturbative approach, it can deal with the strongly disordered regime in which the photons are localized by the interference of multiply scattered waves. The interplay of localization and stimulated or spontaneous emission is a fascinating new field of research, that is just beginning to be explored.

Tuesday, June 27, 2000

Morning Session

Chairperson: M. Sigalas

9:00 **Lecture:** E. Yablonovitch, University of California, Los Angeles,
Electromagnetic Band Gaps at Photonic and Radio Frequencies: II

10:00 **Lecture:** H. Benisty, Ecole Polytechnique, Palaiseau, *Applications of 2d Photonic Crystals to Optoelectronics: I*

In these lectures, we will start with a brief reminder of basics of semiconductor optoelectronics by addressing the role of heterostructures (not only as diodes but also as guides or Bragg mirrors) in edge-emitting laser, light-emitting diode (LED) and VCSELs. We will classify the different modes in heterostructures through the effective index. We will also introduce the more complex "photonic integrated circuits" which are able to carry out sophisticated treatments of light beams, ideally on a monolithic system. We will then turn to the impact of lower photon dimensionality on spontaneous emission in optoelectronics. The planar microcavity case is interesting to start with, as it is a case of modulation of the DOS in different directions, without much difference of the overall DOS and it is well documented for application to LEDs. Then, the feasibility of single mode emission in a cavity defined in a waveguide and/or in a micropillar defined in a microcavity will be examined in the limit case of negligible " Purcell effect ."

We will next discuss specifically the modelling tools and the properties of 2D PC etched through waveguides. One issue of paramount importance for applications in optoelectronic devices is how far these structures can be regarded as ideal 2D ones, i.e. as infinite air rods in a homogeneous matrix. The light line is a basic concept to examine the role of the guide index contrast (3.3:1 in the case of InP-based self-supported membranes and 3.5:3, or less, in the case of conventional substrates). However, the light line is not adapted to discuss the defect modes, which are the only to be used in devices. A complementary perturbative approach based on a separable solution of the etched waveguide will be introduced. It will be shown to provide quite realistic estimates of light losses in various kind of experiments with a single parameter added to the 2D problem. Cavities, which are an essential building-block of various devices, will be used as examples to illustrate possible treatments of these losses.

The last part of these lectures will be devoted to issues in integrated optics : What are the approaches to design a basic PC-based straight guide? What are the technological issues towards large scale? How could coupling be engineered in all-PC structures, e.g. between a PC microcavity and a guide ? How far do the existing tool box and concepts extend? and also what remains to be set up towards optoelectronics applications?

11:30 **Talk:** A. Forchel, Wuerzburg University, *Integration of Photonic Crystals with Active Optoelectronic Devices*

We have investigated the integration of photonic crystal structures with active optoelectronic devices based on ridge waveguide lasers. The photonic crystal is realized by a triangular lattice of air holes etched into the semiconductor. Confinement of the light parallel to the holes is provided by the waveguide of the laser.

An InGaAs/AlGaAs structure with a single InGaAs quantum well emitting at 990 nm in TE polarization (\parallel QW) is used for laser fabrication. The ridge waveguide is etched to a distance of 150 nm from the GaAs waveguide layer. Electron beam lithography is then used to define a triangular array of holes with lattice constants between 160 and 400 nm in PMMA resist. The pattern is etched in an intermediate metal mask and finally transferred into the semiconductor.

12:00 **Talk:** R. B. Wehrspohn, Max-Planck Institute of Microstructure, Halle, *Electrochemically Prepared High-aspect-ratio Pore Arrays for 2d Photonic Crystal Applications*

Formation and characterization of two-dimensional arrays of pores with very high aspect ratios based on anodic etching of silicon and aluminum will be described. Regular pattern formation by self-organization and by lithographic techniques will be dealt with. In both cases, these pore arrays can be used as two-dimensional photonic crystals, which exhibit an optical bandgap from the infrared to the visible depending on the anodization conditions. Diverse defect structures can be easily introduced by lithographic prepatterning. For example wave-guides as well as microcavities have been fabricated and characterized optically.

Evening Session Chairperson: E. Yablonovitch

17:30 **Lecture:** T. F. Krauss, University of St. Andrews, *Patterned Photonic Crystal Waveguides: II*

I will review the development of semiconductor-based planar photonic crystals, highlight the underlying concepts, materials and fabrication issues and point towards important issues for device applications. Examples of successful photonic crystal-based devices include photonic crystal waveguides and photonic integrated circuit elements as well as lasers and LEDs.

18.30 **Panel Discussion:** *Applications of Photonic Band Gap*

Wednesday, June 28, 2000

Morning Session

Chairperson: H. Benistry

9:00 **Lecture:** C. Beenakker, Leiden University, *Random Matrix Theory of Random Lasers and Photonic Localization: II*

10:00 **Talk:** Zhao-Qing Zhang, Hong Kong, *Photonic Crystals and Perturbation Theory*

We show here, with three examples, that a simple perturbative approach is very useful to understand certain photonic band gap problems. In the first example, we will demonstrate that the perturbation analysis can provide us a simple, systematic and efficient way to engineer a photonic band gap. The change of a gap size caused by the variation in the microstructure can be easily determined from the knowledge of eigenfield distribution at two band edge states. Thus, by knowing the field distribution at some symmetry points, one should be able to manipulate an existing gap by changing its microstructure. Explicit examples are given for certain 2D photonic crystals where full/absolute gaps are enlarged significantly. By extending this perturbation analysis to a disordered photonic crystal, in the second example, we explain why a gap is more robust against positional disorder of the scatterers and more sensitive to the randomness in the size of the scatterers. Specific examples are presented for some 2D and 3D photonic crystals. In the final example, we will use the various eigenfield distributions to explain why an incomplete infiltration has a larger photonic gap in an inverse opal photonic crystal. By careful mode analysis, we see that the enlargement of the photonic band gap due to incomplete infiltration in inverse opal is the result of subtle changes in the photonic bands at two particular k-points in the Brillouin zone due to the depletion of high dielectric material.

10:30 **Talk:** Y. Vlasov, Ioffe Institute, *Influence of Disorder in Photonic Crystals*

Self-organized synthetic opals possessing a face centered cubic fcc lattice are promising for fabrication of three-dimensional photonic crystal with a full photonic band gap in the visible. The fundamental limiting factor of this method is the large concentration of lattice defects and, especially, planar stacking faults, which are intrinsic to self-assembling growth of colloidal crystal. On the other hand this makes synthetic opals an ideal model system for the studies of the effects of disorder on photonic band structure. We have numerically analyzed the mechanisms of light localization in such a periodic-on-average system with variable amount of disorder. It is known that photonic Bloch states can become strongly localized near the band edges in a disordered photonic crystal. We show that Bloch states are disrupted and the new localization regime establishes when local fluctuations of the band edge frequency caused by randomization of refractive index profile becomes as large as the bandgap width. This results in strong inhomogeneous broadening of the photonic stopbands in experimental reflection and transmission spectra. Transmission experiments performed on opal photonic crystal have shown the exponential decay of light throughout the gap region, which is ascribed to building up of this second regime of light localization.

11:30 **Talk:** C. M. Soukoulis, Ames Laboratory, Iowa State University, *The Effects of Disorder in Photonic Band Gap Structures*

By using two ab initio numerical methods we study the effects that disorder has on the spectral gaps and on wave localization in two-dimensional photonic band gap materials. We find that there are basically two different responses depending on the lattice realization (solid dielectric cylinders in air or vice versa), the wave polarization, and the particular form under which disorder is introduced. Two different pictures for the photonic states are employed, the "nearly free" photon and the "strongly localized" photon. These originate from the two different mechanisms responsible for the formation of the spectral gaps, ie. multiple scattering and single scatterer resonances, and they qualitatively explain our results.

12:00 **Talk:** V. Kuzmiak, Chez Republic, *The Localization of Light in a Random 2d Dielectric Medium*

We systematically study the strong localization of the light in a two-dimensional, randomly disordered, dielectric medium, that is periodic in average. We consider a simple scattering geometry consisting of an infinite array of infinitely long, parallel, dielectric rods of square and circular cross section embedded in a vacuum. In the case of the parallel rods of square cross sections the dielectric constant of each rod is independent random variable whose values are uniformly distributed about an average value ϵ_a . In systems formed by the rods of circular cross section the disorder is introduced by statistically uncorrelated dielectric constant of each rod, positions of the scatterers and radius of each rod all of which are small deviations satisfying a gaussian probability distribution. We calculate photonic band structure of average periodic system. and by using finite-difference time-domain methods based on the numerical excitation of the mode by virtual oscillating dipole embedded in the center of the computational domain we evaluate the electromagnetic energy stored in consecutive pairs of cylinders centered at the position of the dipole. By inspecting the time-averaged energy associated with the modes with the frequencies close to the band gap edge we observe the strong localization characterized by an exponential decay. We simultaneously calculate the optical transmission of a slab of disordered 2D systems to show the correspondence between the Anderson localized wave functions and the resonances in energy transfer to the transmitted Bragg waves.

12:30 **Talk:** G. Cwilich, Yeshiva University, *Photonic Crystals and Light Localization*

The propagation of a signal in a disordered or a periodic structure is considered in a simple model that contains, as limiting cases, both the propagation of a classical persistent random walker AND the coherent propagation of a wave in the presence of elastic scatterers of mesoscopic physics. A complete statistical analysis of the problem is performed by using a loop-expansion of all the possible paths in a one-dimensional system; the statistics of the distribution of the length of the paths is presented and the transition from the ballistic to the diffusive regime is discussed. A combination of "strip-techniques" allows to generalize this analysis to higher order dimensional systems, for different boundary conditions, and to obtain reflection and transmission coefficients. In particular results for propagation in small $N \times M$ structures will be presented, showing the presence of band gaps. The possible application to simple imaging problems will be discussed. The different roles of the "randomness" in the underlying statistical model and the disorder of the medium and their interaction will be discussed, both analytically and through numerical simulations.

12:45 **Talk: S. Venakides Duke University, *Computational and Theoretical Results in Photonic Crystals***

(a) We compute the transmission properties of 2-d electromagnetic TM waves that are normally incident on a Fabry-Perot structure with mirrors consisting of photonic crystals. We use a boundary integral formulation with quadratic boundary elements and utilize the Ewald representation for the Green's functions. We trace the frequencies of the Fabry-Perot cavity modes traversing the photonic bandgap as the cavity length increases and calculate corresponding Q-values. We do the computation with both lossy and lossless materials. Our results are in very good agreement with experiments performed by H. Everitt's group at Duke.

(b) We derive theoretically Second Harmonic Generation in an LC circuit chain with nonlinear inductors and with capacitors that alternate in value along the chain. We use this as a discrete model of a one-dimensional photonic crystal. The periodicity in the chain is used to achieve phase matching.

Evening Session

Chairperson: P. Halevi

17:30 **Talk: R. Biswas, Ames Laboratory, Iowa State University, *Antenna Applications of Photonic Crystals***

Three-dimensional photonic band gap crystals are utilized to tailor the radiation patterns of antennas. By utilizing the resonances of a Fabry Perot cavity formed with photonic band gap crystals, we were able to create uniquely directional antennas by placing antennas within such cavity. Very good agreement was obtained between finite difference time domain (FDTD) simulations and measurements. Exceptionally directional patterns over narrow frequency ranges were obtained, with half power beam widths less than 10 degrees were obtained. We will compare the theoretical performance of 3-D and 1-D photonic crystals.

These recent results will be compared with past configurations where dipole antennas were placed on top of photonic crystals and the reflecting properties of the PBG crystals were used to improve antenna performance. Such directional promise several new applications.

18:00 **Talk: M. Sigalas, Agilent Laboratories, *Waveguide Bends in 3D Layer-by-Layer Photonic Band Gap Materials***

Using finite difference time domain (FDTD) simulations, we study waveguide configurations in a three dimensional layer-by-layer photonic crystal. We study waveguides lying in the plane of the rods. These can be created by removing one or more rods from one layer and the wave is propagating along the removed rods. An L-shaped waveguide is created by removing part of a rod in one layer and part of a rod in the next layer. The removed rods are perpendicular to each other. We found that 100 % transmission efficiency can be achieved through the 90° bend. Also, waveguides created by removing a whole or a part of a rod from each unit cell along the stacking direction have been studied. In that case the wave is mainly concentrated around the defects. The overlap of the wavefunction of neighboring defects allows the propagation of the wave.

18:30 **Talk:** N. A. Nicorovici, University of Sydney, *Localization and Homogenization in Photonic Crystals*

We describe a new method which allows us to investigate photonic crystals modelled as stacks of gratings consisting of metallic, circular cylinders. Our formulation enable us to compute photonic band diagrams of two- and three-dimensional arrays of inclusions and to estimate accurately the edges of the band gaps in the case of lossless materials. In turn, this provides useful estimates of the gap edges for lossy materials. The technique, which is applicable to structures whose layers do not interpenetrate (thereby permitting the use of plane wave expansions for fields between layers) has, at its heart, an eigenvalue problem which is derived by applying a quasi-periodicity condition. The resulting dispersion relation (which involves the scattering matrices of the single representative layer) can then be reduced to an algebraic eigenvalue problem. The solutions of the eigenvalue equation also provide a generalised technique to determine the field modes of the structure. We note that our technique is intrinsically more stable than similar methods as it avoids the inversion of particular scattering matrices, the condition or stability of which cannot be guaranteed. Also, the accuracy of the method is not affected by the high contrast between the cylinders and the surrounding medium. We use the reflection and transmission scattering matrices of a single grating, to generate the corresponding scattering matrices for a finite stack of gratings, and demonstrate closed form expressions for solutions of the recurrence relations. By considering the limit as the stack length increases without bound, we calculate the reflection scattering matrix R^∞ , which is the fixed point of the recurrence relation for the reflection scattering matrices. By means of R^∞ it should be possible to define an effective permittivity at general points in the Brillouin zone. In the quasistatic limit, we obtain an effective permittivity identical to that obtained using the electrostatic theory.

The method can be applied to the study of both ordered and disordered photonic crystals, and can yield analytic insights into problems such as localization and homogenisation. At long wavelengths, the individual grating layers homogenise and the structure behaves as a one-dimensional stack in which Anderson localization of waves is evident, until the wavelength is sufficiently large so as to render the crystal homogeneous. This method can also be generalized to coated cylinders and to crossed stacks with rotation of layers (one layer to the next). A particular choice of crossing angles of 0° and 90° produces approximate polarization insensitivity.

19:00 **Talk:** A. Orłowski, Warsaw, *From Proximity Resonances to Anderson Localization: Universality Effects in Multiple Scattering*

Multiple elastic scattering of both scalar and electromagnetic waves from a collection of randomly distributed objects is studied. Novel universal properties of the spectra of random Green matrices involved in the description are discovered. A striking physical interpretation within various models of the single scatterer is elaborated. Proximity resonances and Anderson localization are considered as two illustrative examples. Generalizations of 2D numerical transmission experiments are also given.

Thursday, June 29, 2000

Morning Session Chairperson: T. F. Krauss

9:00 **Lecture:** H. Benisty, Ecole Polytechnique, Palaiseau, *Applications of 2d Photonic Crystals to Optoelectronics: II*

10:00 **Talk:** D. Chigrin, University of Wuppertal, *One-Dimensional Photonic Crystals: Omnidirectional Reflection and Superprism Phenomena*
A review on photonic bandgap related phenomena in the simplest type of photonic crystals, i.e., one-dimensional (1D) periodic structures, will be presented. The main focus will be on the omnidirectional reflection and superprism phenomena displayed by 1D photonic crystals. Possibilities to design optoelectronic devices based on these phenomena, e.g., laser microcavities, hollow fibers, collimators, and prisms, will be specifically addressed.

10:30 **Talk:** A. Moroz, Utrecht University, *Complete Photonic Band Gap Structures Below Infrared Wavelengths*

Till recently, no two- and three-dimensional structures have been known to yield a practical complete photonic bandgap (CPBG) for refractive indices available below infrared wavelengths, mainly because of the simultaneous requirements on the dielectric contrast and the modulation (the total number and the length of periodicity steps). We show a promising new route to achieve a CPBG below the infrared wavelengths, using a periodic arrangements of scatterers of material with Drude-like behavior of the dielectric function. In two-dimensions, for a simple square lattice several CPBG's open and a relative gap width g_w (gap width to the midgap frequency ratio) can be as large as 34 even if the host dielectric constant $\epsilon_h=1$. In three-dimensions, despite of recent claims, simple face-centered-cubic (fcc) structures of spheres are shown to yield a tunable CPBG in the region from GHz down to optical wavelengths using currently available experimental techniques. Up to four CPBG's can open in the frequency region $0.6 \omega_0 \leq \omega_1 \leq \omega_p$ with ω_p being the plasma frequency, where the sphere material bulk absorption is assumed to be small (a nonabsorptive window). The relative gap width g_w can be as large as 10 even if the host refractive index $n_h=1$. These properties are rather robust against coating of such spheres with a semiconductor or an insulator. Using different coatings and supporting liquids, the width and midgap frequency of a CPBG can be tuned considerably. These results open a door to fabrication of CPBG structures in the visible using colloidal crystals. In the latter case, CPBS's can be switched on and off by applying an electric field, since the latter allows to switch in ms from an fcc colloidal crystal to a body centered tetragonal (bct) crystal: a so-called martensitic transition.

11:30 **Lecture:** E. Yablonovitch, University of California, Los Angeles, *Electromagnetic Band Gaps at Photonic and Radio Frequencies: III*

Evening Session Chairperson: C. M. Soukoulis

17:30 **Remarks and Discussion:** *Open and Future Directions*

18:00 **Panel Discussion:** *Open and Future Directions*

19:30 **Closing Session:** End of the Institute

Poster Presentations

| | | |
|-----|---|--|
| P1 | Mario Agio, University of Pavia, Pavia, Italy | Impurity modes in a photonic crystal: coupling efficiency and quality factor |
| P2 | Ivan Alvarado-Rodriguez, UCLA Optoelectronics, Los Angeles, CA | Photonic crystal defect cavities in InGaAs |
| P3 | Richard Amos, DERA, United Kingdom | Fabrication of large-area colloidal crystals using one- and two-dimensional shear |
| P4 | Dimitris Angelakis, Imperial College, London, United Kingdom | Coherent effects in two band photonic crystals |
| P5 | Vasily Astratov, University of Sheffield, United Kingdom | Direct study of heavy photon dispersions in photonic crystal waveguides |
| P6 | Mark Auslender, Ben-Gurion University of the Negev, Israel | On the correct use of plane-wave discretization in photonic-structure simulation: Grating diffraction example |
| P7 | Mehmet Bayindir, Bilkent University, Ankara, Turkey | Propagation of photons through localized coupled-cavity modes in photonic band gap structures |
| P8 | Mona Berciu, University of Toronto, Canada | Theory of fluorescence in three-dimensional photonic crystals |
| P9 | Navin Bhat, University of Toronto, Canada | Optical pulse propagation in nonlinear photonic crystals |
| P10 | Alvaro Blanco, Instituto de Ciencia de Materiales de Madrid, Madrid, Spain | Photonic properties of CdS inverted opals |
| P11 | Muriel Botey Cumella, Universitat Politecnica de Catalunya, barcelona, Spain | Momentum nonconserving interaction within a layer of nonlinear material localized in a subwavelength region of the space |
| P12 | Niclas Carlsson, FESTA, Ibaraki, Japan | Formation of 2D photonic crystal airbridge structures in AlGaAs based semiconductor heterostructures |
| P13 | Martin Charlton, University of Southampton, United Kingdom | Analysis of photonic crystal waveguide devices |
| P14 | Crina Maria Cojocaru, Universitat Politecnica de Catalunya, Barcelona, Spain | Dispersive properties of a truncated 1-dimensional photonic crystal |
| P15 | M. Cornelia Cuisin, Institut d'Electronique fondamentale, Orsay, France | PMMA resist templates for dielectric and metallic "Yablonovite" photonic crystals at submicrometer scales |
| P16 | Michiel de Dood, FOM, The Netherlands | Fabrication and optical doping of 1, 2, and 3-dimensional photonic materials using ion beams |
| P17 | George Deligeorgis, FORTH, Crete, Greece | Application of reactive ion etching for GaAs-based integrated optoelectronics |
| P18 | Ihab El-Kady, Iowa State University, United States | Waveguides in two-dimensional photonic band gap structures |
| P19 | | Metallic photonic crystals at optical wavelengths |
| P20 | Stefan Enoch, Faculté des Sciences et Techniques de St. Jérôme, Marseille, France | Anomalous refractive properties of photonic crystals at the band edges |
| P21 | Nayer Eradat, University of Utah, United States | Optical studies of metal infiltrated opal photonic crystals |
| P23 | Alexei Erchak, Massachusetts Institute of Technology, Cambridge, MA, U.S.A. | Increased light extraction from a light-emitting diode using a two-dimensional photonic crystal |
| P23 | Jan Fagerstrom, FOA, Sweden | Characterization of a three-dimensional microwave photonic crystal |
| P24 | Lucia Florescu, University of Toronto, Canada | Time-dependent diffusion model of lasing in a random amplifying medium |
| P25 | Marian Florescu, University of Toronto, Canada | Non-Markovian atomic switching in photonic band gap materials |
| P26 | Eliane Flück, University of Twente, The Netherlands | Light interference in waveguides containing periodic one-dimensional air rod arrays |
| P27 | Stavroula Foteinopoulou, Iowa State University, United States | Tight Binding Parametrization of Photonic Crystals |
| P28 | Valentin Freilikher, Bar-Ilan University, Israel | Transport and localizations in quasi-one-dimensional systems |
| P29 | Antonio Garcia-Martin, Universidad Autonoma de Madrid, Spain | Conductance distributions and transport eigenchannels: From conductance quantization to Anderson localization |
| P30 | Florencio Garcia Santamaria, Universidad Autonoma de Madrid, Spain | Photonic properties of CdS inverted opals |
| P31 | Jacek Generowicz, University of Southampton, United Kingdom | Modelling photonic crystals using finite elements |
| P32 | Cécile Goffaux, Laboratoire de Physique du Solide, FUNDP, Namur, Belgium | Acoustic and elastic waves propagation in two and three dimensional periodic heterostructures |
| P33 | Jaime Gomez Rivas, Van der Waals-Zeeman Instituut, Amsterdam, The Netherlands | Optical experiments on germanium powders close to the Anderson localization transition |
| P34 | Raquel Gomez-Medina, Universidad Autonoma de Madrid, Spain | Enhanced electromagnetic field-particle interactions in a waveguide |

| | | |
|-----|--|--|
| P35 | Serguei Grabtchak, University of Toronto, Canada | Self-assembly of 3-D PBG materials |
| P36 | Boris Gralak, Faculté des Sciences et Techniques Centre de Saint Jérôme, France | Transmission by photonic crystals: unbounded transfer operators approach |
| P37 | Zhong-Ze Gu, Kanagawa Academy of Science and Technology, Japan | Control of photonic band structure by photo-irradiation |
| P38 | Christian Hermann, Institute of Technical Physics, Germany | Lightfield propagation in 2D photonic crystal slab waveguide structures |
| P39 | Reinald Hillebrand, Max Planck Institute of Microstructure Physics, Halle, Germany | Theoretical study of complete photonic band gaps for varying 2D filling patterns of oval-shaped type |
| P40 | Marta Ibsate, Instituto de Ciencia de Materiales de Madrid (CSIC), Madrid, Spain | Photonic properties of CdS inverted opals |
| P41 | Arnout Imhof, Van der Waals-Zeeman Instituut, University of Amsterdam, The Netherlands | Diffusive and coherent transport of light in strongly scattering porous GaP |
| P42 | Steven Johnson, Massachusetts Institute of Technology, United States | Photonic crystals in 3D planar structures and integrated optics |
| P43 | Maria Kafesaki, University of Crete, and FORTH, Crete, Greece | Band gaps phenomena in twinned periodic elastic composites |
| P44 | Robin Kaiser, INLN, France | Coherent backscattering of light by laser cooled atoms |
| P45 | Erik Knudsen, Technical University of Denmark, Kgs. Lyngby, Denmark | Waveguiding properties of photonic crystal fibres |
| P46 | Femius Koenderink, University of Amsterdam, The Netherlands | Enhanced backscattering from photonic crystals |
| P47 | Patrick Kramper, Universitaet Konstanz, Germany | Spectroscopy of defect modes in photonic crystals |
| P48 | Sissy Kyriazidou, University of California-Los Angeles, United States | Fundamental electromagnetic excitation of PBG materials: A formal analogy of PBG and natural crystals |
| P49 | Miguel Laso, Publi University of Navarre, Spain | Photonic crystals in microstrip technology |
| P50 | Ralk Lenke, University of Konstanz, Germany | Magnetic Field Effects on Coherent Backscattering in case of Mie Scattering |
| P51 | | Comparison between the 'The Glory' and Coherent Backscattering of Light in turbid Media |
| P52 | Stephen Leonard, University of Toronto, Canada | Two-dimensional single-mode photonic crystal waveguides fabricated in silicon |
| P53 | Marine Levassor d'Yerville, Université Montpellier, France | Influence of defect positions on the diffraction properties of photonic crystals |
| P54 | Txema Lopetegi, Universidad Autonoma de Madrid, Spain | Vacancy lattices in two-dimensional acoustic crystals |
| P55 | Fernando López-Tejiera, Universidad Autonoma de Madrid, Spain | Diffraction effects in artificial opals |
| P56 | Virginie Lousse, Facultés Universitaires Notre Dame de la Paix, Namur Belgium | Influence of illumination on the density of modes of photorefractive nonlinear photonic crystals |
| P57 | Torsten Maka | Thin film photonic crystals |
| P58 | Vinothan Manoharan, University of California--Santa Barbara, United States | Ordered macroporous ritle titanium dioxide by emulsion templating |
| P59 | Jesus Manzanares Martinez, Université Montpellier II, France | Photonic defect states in imperfect coated opals |
| P60 | Luis Martin-Moreno, Universidad de Zaragoza, Spain | Self-Assembled triply periodic minimal surfaces as moulds for photonic band gap materials |
| P61 | Beatriz Martinez, Public University of Navarra, Spain | Patch antenna design with a photonic bandgap substrate |
| P62 | Hernan Miguez, la Universidad Politecnica de Valencia, Valencia, Spain | Photonic properties of CdS inverted opals |
| P63 | Daniel Mittleman, Rice University, United States | Using colloidal crystals as templates for the formation of novel photonic materials |
| P64 | Jessica Mondia, University of Toronto, United States | Tunable two-dimensional photonic crystals using liquid crystal infiltration |
| P65 | Maria Caterina Netti, University of Southampton, United Kingdom | 2D photonic crystal waveguides: Band gaps and light cones |
| P66 | Bianca Nelson, University of Stanford | A periodic dielectric stack as a one-dimensional photonic crystal for wavelength demultiplexing by beam shifting |
| P67 | Georgios Nikolopoulos, University of Crete, Greece | Beyond single-photon localization at the edge of a photonic band gap |
| P68 | Ségolène Olivier, Ecole Polytechnique, Palaiseau Cedex, France | Optical properties of photonic crystal straight waveguides |
| P69 | Raluca Penciu, University of Crete, Greece | Vibrational modes in solution of copolymers |

| | | |
|-----|---|--|
| P70 | Suresh Pereira, University of Toronto, Canada | Canonical quantization of envelope function equations in a PBG material |
| P71 | David Peyrade, Université Montpellier, France | GaN 2D photonic crystals |
| P72 | Felipe Pinheiro, Centro Brasileiro de Pesquisas Fisicas, Brasil | Multiple scattering of light in disordered magnetic media: Localization parameter, transport velocity and diffusion constant |
| P73 | Luca Plattner, University of Southampton, United Kingdom | Advances in modelling of <i>Morpho</i> -type diffractive structures |
| P74 | Vladimir Poborchii, National Institute for Advanced Interdisciplinary Research, Japan | Photonic-band-gap properties of 2-dimensional lattices of Si nanopillars |
| P75 | Iounnis Psarobas, University of Athens, Greece | A layer Korringa-Kohn-Rostoker method for phononic crystals |
| P76 | Min Qiu, Royal Institute of Technology, Sweden | Studying photonic band-gap structures by finite-difference time-domain method |
| P77 | Rajesh Rengarajan, Rice University, United States | Controlling the optical band gap properties of macroporous polymers by the application of mechanical compression |
| P78 | Andrew Reynolds, Glasgow, Scotland, United Kingdom | Analysis of membrane support structures for integrated antenna usage on 2-dimensional photonic band gap structures |
| P79 | Jorge Ripoll, ICMM, Spain | Multiple layered diffusive media: Theory and experiments |
| P80 | Andrey Rogach, University of Hamburg, Germany | Chemical route to three-dimensional colloidal photonic crystals doped with semiconductor nanoparticles |
| P81 | José Sanchez-Dehesa, Universidad Autonoma de Madrid, Spain | Vacancy lattices in two dimensional acoustic crystals |
| P82 | Juan Vicente Sanchez-Perez, Universidad Politecnica de Valencia, Spain | Experimental and theoretical determination of acoustic dispersion relations in two-dimensional systems |
| P83 | Jorg Schilling, Max-Planck-Institut for Microstructure Physics, Halle, Germany | Photonic crystals made of macroporous silicon: Down to near IR, up to 3 dimensions |
| P84 | Serguei Skipetrov | Temporal fluctuations and instabilities of waves in nonlinear disordered media |
| P85 | David Taylor, DERA, United Kingdom | Colloidal crystals in a liquid crystal host |
| P86 | Burak Temelkuran, Bilkent University, Turkey | Silicon micromachined quasi-metallic photonic crystals |
| P87 | Victor Tikhomirov, University of Leeds, United Kingdom | High refractive index chalcogenide glasses for photonic band-gap devices |
| P88 | Adriaan Tipl, FOM, The Netherlands | Band structure for absorptive photonic crystals |
| P89 | Ovidiu Toader, University of Toronto, Canada | Photonic crystals exhibiting full photonic band gaps |
| P90 | Han van der Lem, FOM, AMOLF, The Netherlands | Two-dimensional Complete Photonic-Bandgap Structures in the Visible |
| P91 | Jerome Vasseur, Universite de Lille, France | Defect mode in one-dimensional combined photonic waveguides--application to the resonant tunneling between two continua |
| P92 | Krassimir Velikov, Utrecht University, The Netherlands | Photonic crystals from core-shell colloidal particles |
| P93 | Jean-Pol Vigneron, Facultes Universitaires Notre Dame De La Paix, Belgium | Tight-binding description of polarization waves in periodic arrays of metal dots |
| P94 | Joachim Wagner, Universität des Saarlandes, Germany | Preparation of Mesoscale Ordered Structures by Means of Immobilized Colloidal Crystals |
| P95 | Vassilos Yannopoulos, NTVA, Greece | Effect of moderate disorder on the absorbance of plasma spheres distributed in a host dielectric medium |
| P96 | Ji Zhou, Nanyang Technological University, Singapore | Three-dimensional photonic bandgap structure of a polymer-metal composite |
| P97 | Majd Ellas Zoorob, University of Southampton, United Kingdom | Investigation into very highly symmetric photonic quasicrystals based on the pinwheel quasicrystal lattice structure |

NATO ASI PCLL Participants and Observers

Mario Agio

Department of Physics "A. Volta"
University of Pavia
Via Bassi 6 1-27100
Pavia
Italy

Telephone 39-0523-975540

Fax 39-0523-975288

agio@bloch.pv.infn.it

Douglas Allan

Fundamental Research
SP FR 5
Corning Incorporated
Corning, NY 14831
U.S.A.

Telephone 1-607-974-3498

Fax 1-607-974-2410

allandc@corning.com

Ivan Alvarado-Rodriguez

UCLA Optoelectronics
Room 64-144 Eng IV
420 Westwood Plaza
Los Angeles, CA 90095
U.S.A.

Telephone 1-310-206-3724

Fax 1-310-206-2596

ialva@ee.ucla.edu

Richard Amos

PD104 DERA
St. Andrews road
Malvern Worcs
WR143PS
United Kingdom

Telephone 44 1684 896693

Fax 44 1684 896270

ramos@dera.gov.uk

Dimitris G. Angelakis

Optics Section
Blackett Laboratory
Imperial College
London SW7 2BW
United Kingdom

Telephone 441715947728

Fax 441718238376

d.angelakis@ic.ac.uk

Kiyoshi Asakawa

FESTA
5-5, Tokodai
Tsukuba, 300-2635

Japan

Telephone 81-298-47-5181

Fax 81-298-47-4417

asakawa@festa.or.jp

Vasily Astratov

Dept. of Physics and Astronomy
University of Sheffield
Hounsfield Rd.
Hicks Building
United Kingdom

Telephone 44(0)114 2224522

Fax 44(0)114 2728079

V.Astratov@sheffield.ac.uk

Mark Auslender

Ben-Gurion University of the Negev
Dept. of Electrical and Computer Eng.
POB 653
Beer-Sheva 84105
Israel

Telephone 972 7 6461583

Fax 972 7 642949

marka@ee.bgu.ac.il

NATO ASI PCLL Participants and Observers

Toshihiko Baba

Yokohama National University
Div. of Electrical & Computer Eng.
79-5 Tokiwadai, Hodogayaku,
Yokohama 240-8501
Japan

Telephone 81 45 339 4258

Fax 81 45 339 4258

baba@dnj.ynu.ac.jp

Mehmet Bayindir

Dept. of Physics
Bilkent University
Bilkent, 06544 Ankara

Turkey

Telephone 90 312 290 2305

Fax 90 312 266 4579

bayindir@fen.bilkent.edu.tr

Carlo Beenakker

Instituut-Lorentz
Leiden University
P.O. Box 95006
2300 RA Leiden
The Netherlands

Telephone 31 71 5275532

Fax 31 71 5275511

beenakker@lorentz.leidenuniv.nl

Henri Benisty

Laboratoire de Physique de la Matière
Condensée
Ecole Polytechnique
F-91128, Palaiseau cedex
France

Telephone 01 69 33 39 59

Fax 01 69 33 30 04

hb@pmc.polytechnique.fr

Monica Berciu

University of Toronto
Department of Physics
60 St. George Street
Toronto, ON M5S 1A7
Canada

Telephone 416-946-5858

Fax 416-978-2537

berciu@physics.utoronto.ca

Navin Bhat

Department of Physics
University of Toronto
60 St. George St.
Toronto, ON M5S 1A7

Canada

Telephone 416 978 5444

Fax 416 978 2537

nbhat@physics.utoronto.ca

Rana Biswas

Microelectronics Research Center
Iowa State University
Ames, Iowa 50011

United States

Telephone 1-515-294-6987

Fax 1-515-294-0689

biswasr@ameslab.gov

Alvaro Blanco

Instituto de Ciencia de Materiales de Madrid
(CSIC)
Campus de Cantoblanco
28049 Madrid
Spain

Telephone 34 91 334 90 00

Fax 34 91 372 06 23

ablanco@icmm.csic.es

NATO ASI PCLL Participants and Observers

Muriel Botey Cumella

Universitat Politecnica de Catalunya
Departament de Fisica i Enginyeria Nuclear
C/Colom 11
08222 Terrassa (Barcelona)
Spain

Telephone 93 739 82 86

Fax 93 739 81 01

muriel.botey@upc.es

Niclas Carlsson

The Femtosecond Technology Research
Association
FESTA
5-5 Tokodai
Japan

Telephone 81 298 47 5181

Fax 81 298 47 4417

carlsson@festa.or.jp

Donald Brown

School of Physics and Astronomy
JF Allen Laboratories
North Haugh
St. Andrews, Fife, KY16 9SS
United Kingdom

Telephone 44 1334 463158

Fax 44 1334 463104

rab16@st.and.ac.uk

David Cassagne

Groupe d'Etude des Semiconducteurs
Universite Montpellier II, CC074
Place E. Bataillon
34095 Montpellier Cedex 05
France

Telephone 33 4 67 14 48 26

Fax 33 4 67 14 37 60

cassagne@ges.univ-montp2.fr

Kurt Busch

Institute for Theory of Condensed Matter
University of Karlsruhe
P.O. Box 6980
76128 Karlsruhe
Germany

Telephone 49 721-608-3366

Fax 49-721-698-150

kurt@tkm.physik.uni-karlsruhe.de

Che Ting Chan

Physics Department
Hong Kong University of Science and
Technology
Clear Water Bay
Hong Kong

Telephone 852 2358-7487

Fax 852 2358-1652

phchan@ust.hk

Hui Cao

Department of Physics and Astronomy
Northwestern University
2145 Sheridan Road
Evanston, IL 60208-3112
U.S.A.

Telephone 1-847-467-5452

Fax 1-847-491-9982

h-cao@nwu.edu

Martin Charlton

The University of Southampton
Dept. of Electronics & Computer Science
Mountbatten 2021
Highfield
United Kingdom SO171BJ

Telephone 02380593127

Fax 02380593029

mdc1@soton.ac.uk

NATO ASI PCLL Participants and Observers

Dmitry Chigrin

Institute of Materials Science & Dept. of Elec.
Eng.
University of Wuppertal
D-42097 Wuppertal
Germany

Telephone 49 202 439 3727

Fax 49 202 439 3037

chigrin@materials.uni-wuppertal.de

Crina Maria Cojocaru

Univeersitat Politecnica de Catalunya
Departament De Fisica 1 Enginyeria Nuclear
Colom 11
08222 Terrassa (Barcelona)
Spain

Telephone 34 93 739 8286

Fax 34 93 739 8101

crina.maria.cojocaru@upc.es

Maria Cornelia Cuisin

Institut d'Electronique Fondamentale
Bat. 220
91405 Orsay

France

Telephone 33 1 69 15 41 08

Fax 33 1 69 15 40 90

cuisin@ief.u-psud.fr

Gabriel Cwilich

Department of Physics
Yeshiva University
500 W. 185th Street
New York, NY 10033
U.S.A.

Telephone 1-212-960-5342

Fax 1-212-960-5245

cwilich@ymail.yu.edu

Michiel J.A. de Dood

FOM
AMOLF
Kruislaan 407
1098 SJ Amsterdarm
The Netherlands

Telephone (31)20-6081234

Fax (31)20-6684106

dood@amolf.nl

George Deligeorgis

FORTH-IESL Microelectronics Group
Vassilika Vouton
P. O. Box 1527
Heraklion-Crete 71110
Greece

Telephone 30 81 394129

Fax 30 81 394106

deligeo@fawn.physics.uoc.gr

Marcel Doosje

Institute for Theoretical Physics
University of Groningen
Nijenborgh 4
9747 AG Groningen
The Netherlands

Telephone 315-03634965

Fax 31503634947

M.Doosje@phys.rug.nl

Eleftherios Economou

Department of Physics
F.O.R.T.H.
Research Center of Crete
Iraklion, 71110 Crete
Greece

Telephone (30) 81 391560

Fax (30) 81 391569

economou@admin.forth.gr

NATO ASI PCLL Participants and Observers

Inigo Ederra

ESA-ESTEC (TOS-EEA)
Keplerlaan, 1 -P. O. Box 299
2200 AG Noorwijk ZH

The Netherlands

Telephone 31 71 565 4832
Fax 31 71 565 4999

iederrau@estec.esa.nl

Alexei Erchak

Massachusetts Institute of Technology
77 Massachusetts Ave., Room 13-3073
Cambridge, MA 02139

United States

Telephone 1-617 253 4346
Fax 1-617 258 6640

aaerchak@mit.edu

Ihab El-Kady

Department of Physics and Astronomy
Iowa State University
Ames, IA 50010

USA

Telephone 1-515-294-2985
Fax 1-515-294-0689

ihab@iastate.edu

Jan Fagerstrom

FOA 31
P. O. Box 1165
SE-581 11 Linkoping

Sweden

Telephone 46 13 378340
Fax 46 13 378519

fager@lin.foa.se

Stefan Enoch

Laboratoire d'Optique Electromagnétique
UPRES-A 6079
Faculté des Sciences et Techniques de St.
Jérôme, case 262 Av,
France

Telephone 33-4-91 28 87 09
Fax 33-4 91 67 44 28

enoch@loe.u-3mrs.fr

Rolando Ferrini

Institut de micro et Optoelectronique
Departement de Physique
EPFL
CH-1015 Lausanne
Switzerland

Telephone 41 21 693 54 87
Fax 41 21693 54 90

ferrini@dpmail.epfl.ch

Nayer Eradat

Department of Physics
115 S 1400E, Room 201
University of Utah
Salt lake City, UT 84112
U.S.A.

Telephone 1-801-581-4402
Fax 1-801-581-4801

nayer@physics.utah.edu

Anne-Marie Filip

Ryerson Polytechnic University
Dept. of Mathematics
Physics and Computer Science
350 Victoria St., Toronto, Ontario M5B 2K3
Canada

Telephone 416 979 5000
Fax 416 979 5343

amfilip@acs.ryerson.ca

NATO ASI PCLL Participants and Observers

Yoel Fink

MIT Bldg. 12-127A
77 Massachusetts Ave.
Cambridge, MA 02139

USA

Telephone 1-617-253-4407

Fax

yoel@mit.edu

Marian Florescu

Mc Lennan Physical labs
60 St. George St.
University of Toronto
Toronto, Ontario, M5S 1A7
Canada

Telephone

Fax

marian@physics.utoronto.ca

Lucia Florescu

Mc Lennan Physical labs
60 St. George St.
University of Toronto
Toronto, Ontario, M5S 1A7
Canada

Telephone

Fax

lucia@physics.utoronto.ca

Eliane Flück

Applied Optics Group
Dept. of Applied Physics
University of Twente
P. O. Box 217
The Netherlands

Telephone 31534893871

Fax 31534891105

e.flueck@tn.utwente.nl

Stavroula Foteinopoulou

Dept. of Physics and Astronomy
Iowa State University
Ames, IA 50011

U.S.A.

Telephone 1-515-663-0508

Fax 1-515-294-0689

sfoteino@iastate.edu

Valentin Freilikher

Department of Physics
Bar-Ilan University
Ramat-Gan 52 900

Israel

Telephone 972 3 531 8647

Fax 972 3 353 3298

freiliv@mail.biu.ac.il

Antonio Garcia-Martin

Dep. Fisica de la Materia Condensada
Universidad Autonoma de Madrid
E-28049 Madrid

Spain

Telephone 34 91 397 47 57

Fax 34 91 397 39 61

a.garcia@uam.es

Florencio Garcia-Santamaria

CSIC
Campus de Catoblanco 28049
Madrid

Spain

Telephone

Fax

foren@icmm.csic.es

NATO ASI PCLL Participants and Observers

F. J. Garcia-Vidal

Departamento de Fisica Teorica de la materia
Condensada (C-V)
Universidad Autonoma de Madrid
E-28049 Madrid
Spain

Telephone 34913978628

Fax 34913974950

fj@uamca4.fmc.uam.es

Jamie Gomez Rivas

Van der Waals-Zeeman Instituut
Valckenierstraat 65
1018 XE Amsterdam

The Netherlands

Telephone 31 20 5255650

Fax 31 20 5255788

rivas@wins.uva.nl

Jacek Generowicz

Microelectronics
Dept. of Electronics & Computer Science
University of Southampton
Highfield, Southampton SO17 1BJ
United Kingdom

Telephone 44 2380592950

Fax 44 2380593903

jmg@ecs.soton.ac.uk

Raquel Gomez-Medina

Dept. Fisica de la Materia Condensada
Universidad Autonoma de Madrid
28049-Madrid

Spain

Telephone 34 91 397 4757

Fax 34 91 397 3961

raquel@tejon.fmc.uam.es

Martina Gerken

Ginzton Laboratory
Stanford University
Stanford, CA 94305

USA

Telephone 1-650- 497-9262

Fax 1-650-725-7509

gerken@stanford.edu

Serge Grabtchak

Department of Physics
University of Toronto
60 St. George St.
Toronto, Ontario M6A 1X3
Canada

Telephone (416) 978-4395

Fax (416) 978-5848

serge@physics.utoronto.ca

Cecile Goffaux

Laboratoire de Physique de Solide
Facultés Universitaires Nore Dame de la paix
FUNDP
Rue de Brux 61
Belgium

Telephone 32 81 72 47 02

Fax 32 81 72 47 07

cecile.goffaux@scf.fundp.ac.be

Boris Gralak

Laboratoire D'Optique Electromagnétique
ESA 6079, Case 262
Faculté des Sciences et Techniques Centre
de Saint Jérôme
France

Telephone 33 291288175

Fax 33 491 674 428

gralak@loe.u-3mrs.fr

NATO ASI PCLL Participants and Observers

Zhong-Ze Gu

Special Research Laboratory for Optical
Science
Kanagawa Academy of Science and
Technology
Japan

Telephone 81-44-819-2073

Fax 81-44-819-2070

gu@fchem.chem.t.u-tokyo.ac.jp

Peter Halevi

INAOE
Apdo.Post. 51
Puebla, Puebla 72000

Mexico

Telephone 52 2 284-5779

Fax 52 2 247-0517

halevi@inaoep.mx

Christian Hermann

Theoretical Quantum electronics
Institute of Technical Physics, DLR
Pfaffenwaldring 38-40
D-70569 Stuttgart
Germany

Telephone 49-711-6862-794

Fax 49-711-6862-349

Christian.Hermann@dlr.de

Reinold Hillebrand

Max Planck Institute of Microstructure
Physics
Weinberg 2
D-06120 Halle
Germany

Telephone 49-245-5582 911

Fax 49-345-5511 223

hi@mpi-halle.mpg.de

Marta Ibisate

Universidad Politcnica de Valencia
Centro Tecnol6gico de Ondas
Unidad Asociada CSIC-UPV,
Avd de los Naranjos s/n, 46022 Valencia
Spain

Telephone 3463879840

Fax 3463879849

ibisate@upv.es

Arnout Imhof

Van der Waals-Zeeman instituut
University of Amsterdam
Valckenierstraat 65
1018 XE Amsterdam
The Netherlands

Telephone 31 20 525 5646

Fax 31 20 525 5788

imhof@wins.uva.nl

J. Joannopoulos

Department of Physics, Bldg. 12-116
Massachusetts Institute of Technology
77 Massachusetts Ave.
Cambridge, MA 02139
USA

Telephone 1-617-253-4806

Fax

Joannop@mit.edu

Sajeev John

Department of Physics
University of Toronto
60 St. George Street
Toronto, ON M5S 1A7
Canada

Telephone (416) 978-3459

Fax (416) 978-2537

john@physics.utoronto.ca

NATO ASI PCLL Participants and Observers

Steven Johnson

305 Memorial Drive #303B
Cambridge, MA 02139

U.S.A.

Telephone 1-617-253-4780

Fax 1-617-253-262

stevenj@alum.mit.edu

Maria Kafesaki

Laboratorio de Nanotecnologia
CSIC
Serrano 144
28006 Madrid
Spain

Telephone 34 600468205

Fax 34 91 5631560

kafesaki@fsp.csic.es

Robin Kaiser

INLN
1361 Route des Lucioles
F-06560 Valbonne

France

Telephone 33 492067391

Fax 33 493652517

kaiser@inln.cnrs.fr

Martin Kamp

Technische Physik
Am Hubland
97074 Wuerzburg

Germany

Telephone 49 931 888 5174

Fax 49 931 888 5143

mkamp@physik.uni-wuerzburg.de

Kihong Kim

Department of Physics
College of Natural Sciences
Ajou University
Suwon 442-749
Korea

Telephone 823 312192584

Fax 823312191615

khkim@madang.ajou.ac.kr

Barry Klein

Office of the Provost
University of California
One Shields Avenue
Davis, CA 95616
U.S.A.

Telephone 1-530-752-2072

Fax 1-530-752-6359

bmklein@ucdavis.edu

Erik Knudsen

Bygn. 345v/181
DK-2800 Kgs. Lyngby

Denmark

Telephone 45 4525 3811

Fax 45 4593 6581

ek@com.dtu.dk

Femius Koenderink

Van der Waals-Zeeman instituut
University of Amsterdam
Valckenierstraat 65
1018 XE Amsterdam

The Netherlands

Telephone 31 20 525 5792

Fax 31 20 525 5788

fkoender@wins.uva.nl

NATO ASI PCLL Participants and Observers

Patrick Kramper
Universitaet Konstanz
FB Physik, LS Mlynek
Universitaet Str. 10
D-78457
Germany

Telephone 49 7531883830
Fax 49 7531 883072

patrick.kramper@uni-konstanz.de

T. F. Krauss
School of Physics
University of St. Andrews
North Haugh
St. Andrews, Fife, KY16 9SS
Scotland

Telephone 44 1334 463107
Fax 44 1334 463104

tfk@st-andrews.ac.uk

Peter Kuchment
Mathematics Department
Wichita State University
Wichita, KS 67260-0033

USA

Telephone 1-316-978-3939
Fax 1-316-978-3748

kuchment@twsuvm.uc.twsu.edu

Vladimir Kuzmiak
University of California—Irvine
Department of Physics and Astronomy
4129 Fredeerick Reines Hall
Irvine, CA 92697-4575
U.S.A.

Telephone 1-949-824-2587
Fax 1-949-824-2174

vkuzmiak@luey.ps.uci.edu

Sissy Kyriazidou
Electrical Eng. Dept.
University of California—Los Angeles
53-145 Eng. IV Building
Los Angeles, CA 90095
USA

Telephone 1-310 825-1027
Fax 1-310 206-5306

kyria@ee.ucla.edu

Ad Lagendijk
Van-der-Waals-Zeeman
University of Amsterdam
Valckenierstraat 65-67
1018 XE Amsterdam
The Netherlands

Telephone (31) 20 5255725
Fax (31)20 5255788

adlag@phys.uva.nl

Miguel Laso
Electrical and Electronic Eng. Dept.
Public University of Navarre
Campus Arrosadia s/n
31006 Pamplona
Spain

Telephone 34 948 169 328
Fax 34 948 169 720

mangel.gomez@unavarra.es

Christoph Lellig
Universität des Saarlandes
Physikalische Chemie
Geb. 9.2
Im Stadtwald
Germany

Telephone 49 681 302 4041
Fax 49 681 302 4759

lellig@pc.uni-sb.de

NATO ASI PCLL Participants and Observers

Ralf Lenke

University of Konstanz
Department of Physics
Lehrstuhl Maret
D-78457 Konstanz
Germany

Telephone 07531/883862
Fax 07531/883090

ralf.lenk@uni-konstanz.de

Stephen W. Leonard

Department of Physics
University of Toronto
60 St. George Street
Toronto, Ontario M5S 1A7
Canada

Telephone 4169784395
Fax 416978 5848

leonard@physics.utoronto.ca

David Leuenberger

Blancherie 5
Rue de la
CH-1022 Chavannes

Switzerland

Telephone 41 78 682 48 72
Fax 41 21 693 43 43

leuenber@dpmail.epfl.ch

Marine Levassor d'Yerville

Groupe d'Etude des Semiconducteurs
Universite Montpellier II, CC074
Place E. Bataillon 34095 Montpellier Cedex
05
France

Telephone 33 4 67 14 38 29
Fax 33 4 67 14 37 60

levassor@ges.univ-montp2.fr

Christian Liguda

Technische Universität Hamburg-Harburg
Materialien der Mikroelektronik
Eissendorfer Str. 38
21071 Hamburg
Germany

Telephone 49 40 42878 3853
Fax 49 40 42878 2229

liguda@tu-harburg.de

Shawn-Yu Lin

Sandia National Laboratories
P. O. Box 5800
Albuquerque, NM 87185

U.S.A.

Telephone 1-505-844-8097
Fax 1-505-844-8985

slin@sandia.gov

Txema Lopetegi

Electrical and Electronic Eng. Dept.
Public University of Navarre
Campus Arrosadia s/n
31006 Pamplona
Spain

Telephone 34948 169 328
Fax 34 948 169 720

txema.lopetegi@unavarra.es

Fernando López-Tejeira

Departamento de Fisica Teorica de la Materia
Condensada
Facultad de Ciencias
Universidad Autonoma de Madrid
Spain

Telephone 34 913 978 647
Fax 34 913 974 950

flt@uamca3.fmc.uam.es

NATO ASI PCLL Participants and Observers

Virginie Lousse

Bld de la Dodaine, 15
B-1400 Nivelles

Belgium

Telephone 32 67/210217

Fax 32 81/724707

virginie.lousse@scf.fundp.ac.be

Torsten Maka

Institut für Materialwissenschaften
Bergische Universität Wuppertal
GauBstr. 20
42097 Wuppertal
Germany

Telephone 49 202 439 3727

Fax 49 202 439 3037

maka@uni-wuppertal.de

Vinothan Manoharan

Dept. of Chemical Eng.
University of California—Santa Barbara
Santa Barbara, CA 93106

USA

Telephone 1-805-893-7862

Fax 1-805-893-4731

vinny@alumni.princeton.edu

Jesus Manzanares Martinez

Groupe d'Etude des Semiconducteurs
Universite Montpellier II, CC074
Place E. Bataillon 34095
Montpellier Cedex 05
France

Telephone 33 4 67 14 47 34

Fax 33 4 67 14 37 60

manzanares@ges.univ-montp2.fr

Luis Martin Moreno

Departamento de Fisica de la Materia
Condensada
universidad de Zaragoza
C/ Pedro Cerbuna
Spain

Telephone 976-762457

Fax

lmm@posta.unizar.es

Beatriz Martinez

Electric and Electronic Engineering Dept.
Public University of Navarra
Campus Arrosadia
31006 Pamplona
Spain

Telephone 34 948 16 92 26

Fax 34 948 16 97 20

beatriz.martinez@unavarra.es

Jordi Martorell

Departament de Fisica i Enginyeria Nuclear
Universitat Politecnica de Catalunya
C/Colom 11
08222 Terrassa (Barcelona)
Spain

Telephone 34-93-739 8076

Fax 34-92-739 8138

jordi.martorell@upc.es

Reinhard März

Infineon Technologies
CPR PH
Otto-Han Ring 6
D-81739 Munich
Germany

Telephone 49 89 234 44695

Fax 49 89 234 44695

Reinhard.Maerz@infineon.com

NATO ASI PCLL Participants and Observers

Francisco Meseguer

Centro Tecnológico de Ondas
Unidad Asociada CSIC-UPV
Edificio de Institutos II
Universidad Politécnica de Valencia
Spain

Telephone 34 96 3879841

Fax 34 96 3879849

fmese@fis.upv.es

Alexander Moroz

Van'Hoff laboratory for Physical & Colloid
chemistry
Debye institute
Utrecht University
The Netherlands

Telephone 49-40-42848-2914

Fax 49 40 42838-6798

moroz@amolf.nl

Hernan Miguez

Centro Tecnológico de Ondas
(CSIC-UPV)
Edificio Nuevos Institutos II
Campus de la Universidad Politécnica de
Spain

Telephone 34 96 387 98 40

Fax 34 96 387 9849

hmiguez@icmm.csic.es

Bianca Nelson

Ginzton Laboratory
Stanford University
Stanford, CA 94305

U.S.A.

Telephone 1-650-725-2291

Fax 1-650-725-7509

bnelson@leland.stanford.edu

Daniel Mittleman

Electrical & Computer Engineering Dept.
MS-366
Rice University
6100 Main Street
U.S.A.

Telephone 1-713-348-5452

Fax 1-713-348-5686

daniel@rice.edu

Maria Caterina Nettì

Physics and Astronomy Department
University of Southampton
Southampton SO17 1BJ

United Kingdom

Telephone 44 1703 596695

Fax 44 1703 593910

mcnetti@soton.ac.uk

Jessica Mondia

Department of Physics
60 St. George St.
University of Toronto
Toronto, Ontario, M5S 1A7
Canada

Telephone 416-978-4395

Fax 416-978-5848

mondia@physics.utoronto.ca

Nicolae-Alexandru Nicorovici

Dept. of Theoretical Physics
School of Physics, A29
University of Sydney
NSW 2006
Australia

Telephone 61 2 9351 5897

Fax 61 2 9351 7726

nicolae@physics.usyd.edu.au

NATO ASI PCLL Participants and Observers

Georgios Nikolopoulos

Dept. of Physics
University of Crete
P. O. Box 2208
G4 710 03 Heraklion
Greece

Telephone 30-81-394163
Fax 30-81-394-101

nikolg@physics.uoc.gr

Susumu Noda

Department of Electronic Science and
Engineering
Kyoto University
Kyoto 606-8501
Japan

Telephone 81 75 753-5297
Fax 81 75 751-1576

snoda@kuee.kyoto-u.ac.jp

David Norris

NEC Research Institute
4 Independence Way
Princeton, NJ 08540

U.S.A.

Telephone 1-609-951-2640
Fax 1-609-951-2482

dnorris@research.nj.nec.com

Stephen O'Brien

Room 816
Condensed Matter Theory Group
Blackett Lab
Imperial College
United Kingdom

Telephone
Fax

s.obhriain@ic.ac.uk

Segolene Olivier

Laboratoire PMC
Ecole polytechnique
91128 Palaiseau Cedex

France

Telephone 33 1 69 33 39 59
Fax 33 1 69 33 30 04

so@pmc.polytechnique.fr

Arkadiusz Orlowski

Instytut Fizyki PAN
Aleja Lotnikow 32/46
02-668 Warszawa

Poland

Telephone 48 22 8437001
Fax 48 22 8430926

orlow@punio.ifpan.edu.pl

Ekmel Ozbay

Department of Physics
Bilkent University
Bilkent 06533 Ankara

Turkey

Telephone (90) 312 2664397
Fax (90) 312 2664579

ozbay@fen.bilkent.edu.tr

John Page

Department of Pphysics
University of Manitoba
Winnipeg, MB R3T 2N2

Canada

Telephone 204-474-9852
Fax 204-474-7622

jhpage@cc.UManitoba.ca

NATO ASI PCLL Participants and Observers

Dimitris Papaconstantopoulos

Complex Systems Theory Branch
Naval Research Laboratory
Code 6690
Washington, DC 20375-5345
U.S.A.

Telephone 1-202-767-6886

Fax 1-202-404-7546

papacon@dave.nrl.navy.mil

Michael Patra

Instituut-Lorentz for Theoretical Physics
Universiteit Leiden
Postbus 9506
2300 RA Leiden
The Netherlands

Telephone 31 71 527 5504

Fax 31 71 527 5511

patra@lorentz.leidenuniv.nl

Raluca Penciu

Dept. of Physics
University of Crete
P. O. Box 2208
710 03 Heraklion, Crete
Greece

Telephone

Fax

raluca@physics.uoc.gr

John Pendry

The Blackett lab
Imperial College
London SW7 2BZ

England

Telephone 44 171 594 7500

Fax 44 171 594 7504

j.pendry@ic.ac.uk

Suresh Pereira

Department of Physics
60 St. George St.
University of Toronto
Toronto, Ontario, M5S 1A7
Canada

Telephone 416 978 5208

Fax 416 978 2537

Pereira@physics.utoronto.ca

Alexandros Pertsinidis

Physics Dept.
Brown University
Box 1843
Providence, RI 02912
U.S.A.

Telephone 1-401 863 9337

Fax 1-401 863 2024

pertsin@barus.physics.brown.edu

David Peyrade

196 Ave. H. Ravera
92225 Bagneux

France

Telephone 1 42 31 73 74

Fax 1 42 31 73 78

David.Peyrade@L2M.cnrs.fr

Felipe Pinheiro

Centro Brasileiro de Pesquisas Fisicas,
CBPF-DE
Rua Dr. Xavier Sigaud, 150 Urca
CEP 22290-180
Brasil

Telephone 55 21 586 7372

Fax 55 21 586 7400

felipe@cbpf.br

NATO ASI PCLL Participants and Observers

Luca Plattner

Microelectronics
Dept. of Electronics & Computer Science
University of Southampton
Highfield
United Kingdom

Telephone 44 2380 593127

Fax 44 2380 593127

lpl@soton.ac.uk

Vladimir Poborchii

JRCAT
c/o National Institute for Advanced
Interdisciplinary Research
1-1-4 Higashi, Tsukuba, Ibaraki 305-0046
Japan

Telephone 81-298-61-2627

Fax +81-298-61-2576

vpob@jrkat.or.jp

Albert Polman

Head, Optoelectronic Materials Program
FOM-Institute for Atomic and Molecular
Physics
Kruislaan 407, 1098, SJ Amsterdam
The Netherlands

Telephone 31-20-6081234

Fax 31-20-6684106

polman@amolf.nl

Sergey Polyakov

School of Optics/CREOL
University of Central Florida
4000 Central florida Blvd.
Orlando, FL 32816
U.S.A.

Telephone 1-407-823-6944

Fax 1-407-823-6955

serge@creol.ucf.edu

Ioannis Psarobas

The University of Athens
Solid State Physics
Panepistimioupolis Zografou
GR-157 84 Athens
Greece

Telephone 30 1 7274758

Fax 30 1 7257689

ipsarob@cc.uoa.gr

Min Qiu

Department of Electromagnetic Theory
Royal Institute of Technology
100 44 Stockholm

Sweden

Telephone 46 8 7908371

Fax 46 9 108327

min@tet.kth.se

Rajesh Rengarajan

6100 Main St.
Electrical and Computer Engineering Dept.
Rice University
Houston, TX 77005
U.S.A.

Telephone

Fax

rajeshr@rice.edu

Antti Renko

Nokia Research Center
P. O. Box 407 FIN-00045
Nokia Group

Finland

Telephone 358405470806

Fax 358943766213

antti.renko@nokia.com

NATO ASI PCLL Participants and Observers

Andrew Reynolds

Dept. of Electronics & Elec. Eng.
Rankine Building
Oakfield Avenue
University of Glasgow
Scotland

Telephone 44 141-330-6022

Fax 44 141-330-6002

areynolds@elec.gla.ac.uk

Jorge Ripoll

Dept. Teoria de la Materia Condensada
Instituto de Ciencia de Materiales
ICMM
Campus de Cantoblanco
Spain

Telephone 34 91 3349000

Fax 34 91 3720623

jripoll@icmm.csic.es

Andrey Rogach

Institute of Physical Chemistry
University of Hamburg
Bundesstr. 45
D-20146 Hamburg
Germany

Telephone

Fax

andrey.rogach@mailcity.com

Sergei Romanov

Institute of material Science and Dept. of
Electrical Eng.
University of Wuppertal
Gauss-strasse 20
Germany

Telephone 49 202 439 2232

Fax 49 202 439 9037

romanov@uni-wuppertal.de

Juan Jose Saenz Gutierrez

Dept. Fisica de la Materia Condensada
Universidad Autonoma de Madrid
28049 Madrid

Spain

Telephone 34 91 397 3804

Fax 34 91 397 3961

juanjo.saenz@uam.es

Birsen Saka

Hacettepe University
Dept. of Electrical and Electronics Eng.
06532 Beytepe
Ankara
Turkey

Telephone 90 312 2977045

Fax 90 312 2992125

birsen@hacettepe.edu.tr

Jose Sanchez-Dehesa

Departamento de Fisica Teorica de la Materia
Condensada
Facultad de Ciencias (C-5)
Universidad Autonoma de Madrid
Spain

Telephone 91 397 4905

Fax 91 397 4950

jsdehesa@uamca3.fmc.uam.es

Juan Vicente Sanchez-Perez

Universidad Politcnica de Valencia
Centro Tecnologico de Ondas
Avd de los Naranjos s/n
46022 Valencia
Spain

Telephone 00 34 96 3879840

Fax 00 34 96 3877975

jusanc@fis.upv.es

NATO ASI PCLL Participants and Observers

Lorenzo Sanchis Martinez

Pizarro 26-8a
46004 Valencia

Spain

Telephone 963518400

Fax

lsanchis@fis.upv.es

Jorg Schilling

Max-Planck-institute for Microstructure
Physics
Weinberg 2
D-06120 Halle
Germany

Telephone 49 345 5582 901

Fax 49 345 55 11 22 3

schill@mpi-halle.de

Vladimir Shalaev

Physics Dept.
MSC 3D
New Mexico State University
Las Cruces, NM 88003
U.S.A.

Telephone 1-505-646-1932

Fax 1-505-646-1934

vshalaev@nmsu.edu

Rebecca Shivvers

Condensed Matter Physics
Ames Laboratory
Iowa State University
Ames, IA 50011
U.S.A.

Telephone 1-515-294-3481

Fax 1-515-294-0689

shivvers@ameslab.gov

Mihail Sigalas

Communications and Optics Lab
Agilent Laboratories, Agilent Technologies
3500 Deer Creek Road, MS26M7
Palo Alto, CA 94304
U.S.A.

Telephone 1-650 857 4819

Fax 1-650 857 6241

msigalas@santorin.labs.agilent.com

Serguei Skipetrov

Laboratoire de Physique et Modelisation
des Milieux Condenses, Universite
Joseph Fourier, Maison des Magisteres
CNRS, B.P. 166
France

Telephone 33 2 76 88 74 97

Fax 33 4 76 88 79 81

sergey@belledonne.polycnrs-gre.fr

Richart Slusher

Lucent Technologies--Bell laboratories
600 Mountain Avenue
Room 1D-227
Murray Hill, NJ 07974
U.S.A.

Telephone 1-908-582-4094

Fax 1-908-582-3260

res@lucent.com

Christopher Smith

Optoelectronics Research Group
Rankine Building
Oakfield Avenue
University of Glasgow
Scotland

Telephone 44 141 330 4108

Fax 44 141 330 6002

C.Smith@elec.gla.ac.uk

NATO ASI PCLL Participants and Observers

David R. Smith

Physics Dept.
9500 Gilman Drive
University of California—San Diego
La Jolla, CA 92093-0319
U.S.A.

Telephone 1-858 534-1510
Fax 1-858 534 8059

drs@sdss.ucsd.edu

Costas Soukoulis

Dept. of Physics
Iowa State University
Ames, IA 50011
U.S.A.

Telephone 1-515-294-2816
Fax 1-515-294-0689

soukoulis@ameslab.gov

Rudolf Sprik

van der Waals-Zeeman instituut
Valckenierstraat 65
1018 XE Amsterdam
The Netherlands

Telephone 31 20 525 5645
Fax 31 20 525 5788

sprik@phys.uva.nl

Adam Stelmach

c/o Bohlin
Solberga Angsv. 13 1 tr
S-125 44 Alvsjo
Sweden

Telephone 46 8 749 64 16
Fax 46 8 727 38 30

adam.stelmach@uab.ericsson.se

David Taylor

DERA
PD104
St. Andrews Rd., Malvern
WR14 3PS
United Kingdom

Telephone 01684 894847
Fax 01684 896720

dtaylor@falcor.dera.gov.uk

Burak Temelkuran

Dept. of Physics
Bilkent University
Bilkent 06533 Ankara
Turkey

Telephone 90 312 290 1963
Fax 90 312 266 4579

burak@fen.bilkent.edu.tr

Victor Tikhomirov

Department of Materials
University of Leeds
Leeds LS2 9JT
United Kingdom

Telephone 44-113-2332567
Fax 44-113-2422531

V.Tikhomirov@leeds.ac.uk

Adriaan Tip

FOM
AMOLF
Kruislaan 407
1098SF Amsterdam
The Netherlands

Telephone 31 20 6081353
Fax 31 20 6684106

tip@amolf.nl

NATO ASI PCLL Participants and Observers

Ovidiu Toader

60 St. George Street
University of Toronto
Toronto, ON, M5S-1A7

Canada

Telephone 416-978-5444

Fax 416-978-2537

ovi@physics.utoronto.ca

Alfons van Blaaderen

Condensed Matter Dept.
Debye Institute,
Utrecht University
P. O. Box 80000
The Netherlands

Telephone 31 30 2532414

Fax 31 30 2532403

A.vanBlaaderen@phys.uu.nl

Han van der Lem

FOM
AMOLF
Kruislaan 407
1098 SJ Amsterdam
The Netherlands

Telephone 31 20 6081348

Fax 31 20 6684106

han@amolf.nl

Henry van Driel

Department of Physics
University of Toronto
Toronto, Canada
M5S 1A7
Canada

Telephone 416-978-4200

Fax 416-978-2537

vandriel@physics.utoronto.ca

Bart Van Tiggelen

CNRS/Laboratoire de Physique et
Modelisation des Systemes Condenses
Universite Joseph Fourier
Maison de Magisteres
France

Telephone 33-4 76 88 12 76

Fax 33-4 76 88 79 81

tiggelen@belledonne.polycnrs-gre.fr

Valy Vardeny

Physics Department
115 South 1400 East
201 JFB
University of Utah
U.S.A.

Telephone 1-801-581-8372

Fax 1-801-581-4801

val@physics.utah.edu

Karri Varis

Helsinki University of Technology
Optoelectronics Laboratory
PO Box 3000
FIN-02015 HUT
Finland

Telephone 358 9 451 3129

Fax 358 9 451 3128

Karri.Varis@hut.fi

Mikhail Vasilevskiy

Departamento de Fisica
Universidade do Minho
Campus de Gualtar
4710 - 057 Braga
Portugal

Telephone 351-253-604327

Fax 351-253-678981

mikhail@fisica.uminho.pt

NATO ASI PCLL Participants and Observers

Panagiotis Vasilopoulos

Concordia University
Dept. of Physics
1455 de Maisonneuve Ouest
Montreal, Quebec
Canada

Telephone 514 84803290
Fax 514-848-2828

takis@boltzmann.concordia.ca

Jean-Pol Vigneron

Facultes Universitaires Notre Dme De La Paix
Laboratoire de Physique du Solide
rue de Bruxelles, 61
B-5000 Namur
Belgium

Telephone 32 81/724711
Fax 32 81/724707

jean-pol.vigneron@scf.fundp.ac.be

Jérôme Vasseur

Equipe de Dynamique des Interfaces
LDSMM-UPRESA CNRS 8024 Bat P5bis
UFR de Physique - Universite de Lille I
59655 Vileneuve d'Ascq Cedex
France

Telephone
Fax

jerome.vasseur@univ-lille1.fr

Yuri Vlasov

loffe Institute of Physics and Technology
St. Petersburg

Russia

Telephone 609-951-2836
Fax 1-609-951-2482

vlasov@research.nj.nec.com

Krassimir Velikov

Chemistry and Physics of Condensed Matter
Ornstein Laboratory
Debye Institute
Utrecht University
The Netherlands

Telephone
Fax

K.P.Velikov@chem.uu.nl

Willem Vos

van der Waals-Zeeman Instituut
Universiteit van Amsterdam
Valckenierstraat 65
NL-1018 XE Amsterdam
The Netherlands

Telephone 31-20-525-5663
Fax 31-20-525-5788

wvos@wins.uva.nl

Stephanos Venakides

Duke University
Department of Mathematics
Durham, NC 27708

USA

Telephone 1-919-660-2815
Fax 1919-660-2821

ven@math.duke.edu

Joachim Wagner

Universität des Saarlandes
Physikalische Chemie
Geb. 9.2
Im Stadtwald
Germany

Telephone 49 681 302 3229
Fax 49 681 302 4759

wagnerjch@rz.uni-sb.de

NATO ASI PCLL Participants and Observers

Ralf Wehrspohn

Max-Planck-Institute for Microstructure
Physics
Weinberg 2
06108
Germany

Telephone 49 345 5582726

Fax 49 345 5511223

wehrspoh@mpi-halle.de

Diederik Wiersma

Istituto Nazionale di Fisica della Materia
Largo E. Fermi 2
50125 Florence

Italy

Telephone 390 55 2307830

Fax 390 55 224072

wiersma@lens.unifi.it

Carol Wilson

Agilent Laboratories
3500 Deer Creek Road
Palo Alto, CA 94304

U.S.;.A.

Telephone 1-650-857-8067

Fax 1-650-857-6241

carol_wilson@agilent.com

J. P. Woerdman

Huygens Laboratory
Leiden University
P. O. Box 9504
2300 RA Leiden
The Netherlands

Telephone

Fax 31 71 5275819

qo@molphys.leidenuniv.nl

Eli Yablonovitch

Electrical Engineering Department, Room
66-147K
Eng IV, Box 951594, 420 Westwood Plaza
University of California-Los Angeles
USA

Telephone 1-310-206-2240

Fax 1-310-206-4685

elij@ee.ucla.edu

Vassilios Yannopoulos

Dept. of Applied Mathematics and Physics
National Technical University of Athens
Zografou Campus
GR-157 80 Athens
Greece

Telephone 30 1 7274787

Fax 30 1 7257689

vyannop@cc.uoa.gr

Fumiyo Yoshino

CREOL
University of Central Florida
4000 Central Florida Blvd.
Orlando, FL 32816
U.S.A.

Telephone 1-407-823-6944

Fax 1-407-823-6955

fumy@creol.ucf.edu

Zhao-Qing Zhang

Physics Department
Hong Kong University of Science &
Technology

Hong Kong

Telephone

Fax 852-2358-7487

phzzhang@ust.hk

NATO ASI PCLL Participants and Observers

Ji Zhou

Photonics Lab, S-1
School of Electrical and Electronic Eng.
Nanyang Technological University
Singapore 639798
Singapore

Telephone 65 7906319-24

Fax 657904101

ejzhou@ntu.edu.sg

Majd Elias Zoorob

Room 2043 Mountbatten Building
Department of Electronics & Computer
Science
University of Southampton
United Kingdom

Telephone

Fax

mez@soton.ac.uk

| AD NUMBER | DATE | DTIC ACCESSION |
|---|------|---|
| 1. REPORT IDENTIFYING INFORMATION | | REQUE |
| A. ORIGINATING AGENCY F.O.R.T.H., GREECE | | 1. Put you on reve |
| B. REPORT TITLE AND/OR NUMBER PHOTONIC CRYSTALS AND LIGHT LOCALIZATION | | 2. Complet |
| C. MONITOR REPORT NUMBER R+D 8919-PH-02 | | 3. Attach fi mailed |
| D. PREPARED UNDER CONTRACT NUMBER N68171-00-M-5733 | | 4. Use unc Informu |
| 2. DISTRIBUTION STATEMENT | | 5. Do not o for 6 to |
| APPROVED FOR PUBLIC RELEASE DISTRIBUTION UNLIMITED | | DTIC: 1. Assign. 2. Return |
| PROCEEDINGS | | TIONS ARE OBSOLETE |

2
 3
 4
 5
 6
 7
 8
 9
 10
 11
 12
 13
 14
 15
 16
 17
 18
 19
 20
 21
 22
 23
 24
 25
 26
 27
 28
 29
 30
 31
 32
 33
 34
 35
 36
 37
 38
 39
 40
 41
 42
 43
 44
 45
 46
 47
 48
 49
 50
 51
 52
 53
 54
 55
 56
 57
 58
 59
 60
 61
 62
 63
 64
 65
 66
 67
 68
 69
 70
 71
 72
 73
 74
 75
 76
 77
 78
 79
 80
 81
 82
 83
 84
 85
 86
 87
 88
 89
 90
 91
 92
 93
 94
 95
 96
 97
 98
 99
 100