

REPORT DOCUMENTATION PAGE

AFRL-SR-BL-TR-00-

0452

Source, collection of Reports shall be

The public reporting burden for this collection of information is estimated to average 1 hour per response, including gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments of information, including suggestions for reducing the burden, to Department of Defense, Washington Headquarters (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.

| | | | | | |
|---|------------------------|---|----------------------------|--|---|
| 1. REPORT DATE (DD-MM-YYYY) 8/31/00 | | 2. REPORT TYPE AFOSR/AASERT Final Technical Report | | 3. DATES COVERED (From - To) 6/1/97 - 5/31/00 | |
| 4. TITLE AND SUBTITLE State-Resolved Thermal/Hyperthermal Collision Dynamics of Atmospheric Species | | | | 5a. CONTRACT NUMBER F49620-97-1-0458 | |
| | | | | 5b. GRANT NUMBER 153-1108 | |
| | | | | 5c. PROGRAM ELEMENT NUMBER | |
| 6. AUTHOR(S) David J. Nesbitt | | | | 5d. PROJECT NUMBER | |
| | | | | 5e. TASK NUMBER | |
| | | | | 5f. WORK UNIT NUMBER | |
| 7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) JILA, Campus Box 440 University of Colorado Boulder, CO 80309-0440 | | | | 8. PERFORMING ORGANIZATION REPORT NUMBER | |
| 9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Air Force Office of Scientific Research 801 North Randolph Street Room 732 Arlington, Virginia 22203-1977 | | | | 10. SPONSOR/MONITOR'S ACRONYM(S) | |
| | | | | 11. SPONSOR/MONITOR'S REPORT NUMBER(S) | |
| 12. DISTRIBUTION/AVAILABILITY STATEMENT APPROVED FOR PUBLIC RELEASE: DISTRIBUTION UNLIMITED | | | | | |
| 13. SUPPLEMENTARY NOTES 20000922 075 | | | | | |
| 14. ABSTRACT Direct absorption IR laser methods developed under AFOSR support have been used to study state-to-state reactive scattering dynamics in collisions systems under crossed supersonic jet, single collision conditions. This past year has focused on reaction dynamics in three collision systems, i) $F + HD \rightarrow HF(v,J) + D$, ii) $F + CH_4 \rightarrow HF(v,J) + CH_3$, iii) $F + NH_3 \rightarrow HF(v,J) + NH_2$. Time resolved flash kinetic studies of the $OH/HO_2/O_3$ chemical chain reaction has been performed from 300K down to 190K, providing first access to temperature conditions relevant to accurate modeling of the lower stratosphere. Most recently, these methods have been extended to study "airglow" dynamics of highly rotationally excited $OH(v,N)$ radicals formed from $H+O_3$ reactions. | | | | | |
| 15. SUBJECT TERMS state-to-state collision dynamics, atmospheric radical kinetics | | | | | |
| 16. SECURITY CLASSIFICATION OF: | | | 17. LIMITATION OF ABSTRACT | 18. NUMBER OF PAGES 5 | 19a. NAME OF RESPONSIBLE PERSON |
| a. REPORT Unclass | b. ABSTRACT Unclass | c. THIS PAGE Unclass | | | 19b. TELEPHONE NUMBER (Include area code) |

AFOSR/AASERT Final Report: 9/1/00

State-to-state thermal/hyperthermal collision dynamics of atmospheric species

Abstract:

Direct absorption IR laser methods developed under AFOSR support have been used to study state-to-state reactive scattering dynamics in collisions systems under crossed supersonic jet, single collision conditions. This past year has focused on reaction dynamics in three collision systems, i) $F + HD \rightarrow HF(v,J) + D$, ii) $F + CH_4 \rightarrow HF(v,J) + CH_3$, iii) $F + NH_3 \rightarrow HF(v,J) + NH_2$. Time resolved flash kinetic studies of the $OH/HO_2/O_3$ chemical chain reaction has been performed from 300K down to 190K, providing first access to temperature conditions relevant to accurate modeling of the lower stratosphere. Most recently, these methods have been extended to study "airglow" dynamics of highly rotationally excited $OH(v,N)$ radicals formed from $H+O_3$ reactions.

Brief overview of work:

Direct absorption IR laser methods developed under AFOSR support have been used to study state-to-state reactive scattering dynamics in collisions systems under crossed supersonic jet, single collision conditions. This past year has focused on reaction dynamics in three collision systems, i) $F + HD \rightarrow HF(v,J) + D$, ii) $F + CH_4 \rightarrow HF(v,J) + CH_3$, iii) $F + NH_3 \rightarrow HF(v,J) + NH_2$, corresponding to our proposed plan to move from atom + diatom systems onto more complicated atom + polyatom reactive scattering.

In the $F + HD$ system, the key interest is in the probing of long lived "quantum scattering resonances" at the transition state. These resonances correspond to well developed vibrational nodal structure in wave packet calculations, and recently predicted by Skodje et al to occur at $E_{com} \approx 0.5$ kcal/mole. Theory predicts a dramatic shift in nascent state populations in the $HF(v=2,J)$ manifold in the scattering resonance regime, essentially from a more smooth Boltzmann-like J distribution to a relatively "flat-top" albeit structured distribution which drops off rapidly at the energetic limit. The results are shown in Fig. 1, which are in remarkably good agreement with wave packet theoretical calculations. This is particularly interesting in light of the immense effort that has been




Fig 1. $F+HD$ reactive scattering into $HF(v=2,J)$ manifold near the transition state resonance predicted at 0.5 kcal.mole: experiment vs theory

invested in attempts to verify similar resonance scattering behavior in the $F + H_2$ system, where in fact it appears that all resonance structure in FH_2^- photodetachment can now be theoretically assigned to van der Waals angular momentum barriers.

In a second direction, efforts toward atom + polyatom reactive scattering have been pursued. State-to-state reactive scattering of $F + CH_4 \rightarrow HF(v,J) + CH_3$ is studied using crossed supersonic jets and high-resolution ($\Delta v \approx 0.0001$ cm^{-1}) IR laser direct absorption




Fig 2. Nascent populations from F + CH₄ reactive scattering

energetically accessible J values in both the $v=2$ and $v=3$ vibrational manifolds, which provides experimental support for a bent F-H-C transition state structure.

In addition, product recoil data from F + CH₄ → HF(v,J) + CH₃ reactive scattering are obtained using crossed supersonic jets and narrow band ($\Delta\nu \approx 0.0001 \text{ cm}^{-1}$) IR laser direct absorption techniques. The high resolution IR profiles of HF(v,J) product exhibit extensive Doppler broadening that directly reflects quantum state-resolved *translational* distributions in the laboratory frame under collision free conditions. Analysis of Doppler profiles via singular value decomposition (SVD) methods yield




Fig 3. HF(v,J) z axis recoil translational energy from F+CH₄ reactive events, indicating small propensity for forward/backward scattering

information on state-resolved differential cross sections into the HF ($v=1$) and HF($v=2$) vibrational manifolds, which identify a small propensity for forward/backward scattering ($|\cos(\theta)| \approx 1$) vs. side scattering ($|\cos(\theta)| \approx 0$) in the center of mass frame. This can be nicely seen in terms of a plot of experimentally observed recoil kinetic energy along the IR probe (z) direction vs total available energy (Fig 3). When compared with the theoretical predictions for purely sideways or isotropic scattering, the data clearly indicates a systematic shift toward more forward/backward than isotropic behavior.

techniques. Rovibrational state-resolved HF column-integrated absorption profiles are obtained under single collision conditions and converted to populations via appropriate density-to-flux transformation. Nascent rovibrational distributions in each HF(v,J) state are now available (Fig. 2) and reported in the literature. Summed over all product rotational levels, the nascent *vibrational* quantum state populations for HF(v) [$v=3$: 0.106(3); $v=2$: 0.667(14); $v=1$: 0.189(27); $v=0$: 0.038(78); 2σ error bars] are in agreement with previous flow cell studies by Setser, Heydtmann, and co-workers [Chem. Phys. **94**, 109 (1985)]. At the *rotational* state level, however, the current studies indicate nascent distributions for HF(v,J) that are significantly hotter than previously reported, ostensibly due to reduced collisional relaxation effects under supersonic jet conditions. Final HF rotational states from F + CH₄ are observed near the maximum

We have most recently been studying the $F + NH_3$ system, which has now yielded first rigorously collision free $HF(v,J)$ populations for this reaction. There is an interesting bimodality in the $HF(v=1,2)$ rotational distributions, specifically a long tail toward higher J levels that may indicate two different channels in the reaction mechanism.

Time resolved flash kinetic studies of the $OH/HO_2/O_3$ chemical chain reaction has

been performed from 300K down to 190K, providing first access to temperature conditions relevant to accurate modeling of the lower stratosphere (Fig. 4). Most recently, these methods have been extended to study "airglow" dynamics of highly rotationally excited $OH(v,N)$ radicals formed from $H+O_3$ reactions. Preliminary work has indicated the presence of high rotational levels up to $N \approx 33$, as indeed directly observed in the upper atmosphere by efforts in the Air Force laboratories. Further work in progress will be necessary to distinguish between direct formation into high N states versus anomalously rapid and efficient V-R transfer from the highest vibrational manifold.

Induction Decay Rate vs. $[O_3]$




Fig 4. Temperature dependent rate constant data for $OH/HO_2/O_3$ chemical chain reaction kinetics

Awards or Fellowships received 1999-2000

William F. Meggers Award, Optical Society of America 1999

Alexander von Humboldt Fellowship, Humboldt Foundation, 1999-2000

Graduate students supported by AASERT grant:

Scott Davis

William Chapman

Bradley Blackmon

Joanna Fair

Evan Jochnowitz

Papers published, in press or submitted since last progress report acknowledging AFOSR support:

- 1) S. A. Nizkorodov, W. W. Harper and D. J. Nesbitt, "State-to-state reaction dynamics in crossed supersonic jets: threshold evidence for non-adiabatic channels in $F + H_2$ ", *Faraday Disc.* 113, 107 (1999).
- 2) J. R. Fair and D. J. Nesbitt, "Dynamics of Collisional Alignment in Supersonic Expansions: Trajectory Studies of $He + CO$, O_2 and CO_2 ", *J. Chem. Phys.* 111, 6821 (1999).
- 3) S. A. Nizkorodov, W. W. Harper, W. B. Chapman, B. W. Blackmon and D. J. Nesbitt, "Energy dependent cross sections and non-adiabatic reaction dynamics in $F(^2P_{3/2}, ^2P_{1/2}) + n-H_2 \rightarrow HF(v,J) + H$ ", *J. Chem. Phys.* 111, 8404 (1999).
- 4) S. Davis, D. Uy, and D. J. Nesbitt, "Laser spectroscopy of jet-cooled ethyl radical: Infrared studies in the CH_2 stretch manifold", *J. Chem. Phys.* 112, 1823 (2000).
- 5) S. A. Nizkorodov, W. W. Harper, B. W. Blackmon and D. J. Nesbitt, "Temperature dependent kinetic studies of the $OH/HO_2/O_3$ chain reaction by time resolved high resolution laser absorption spectroscopy", *J. Phys. Chem* 104, 3964 (2000).
- 6) W. W. Harper, S. A. Nizkorodov, and D. J. Nesbitt, "Quantum state-resolved reactive scattering of $F + CH_4 \rightarrow HF(v,J) + CH_3$: Nascent $HF(v,J)$ product state distributions", *J. Chem. Phys.* (in press).

Invited talks since last progress report acknowledging AFOSR support:

"Non-adiabatic reaction dynamics in $F + n-H_2$ from IR laser based state-to-state reactive scattering studies", American Chemical Society (218th National Meeting), New Orleans, LA, August 23, 1999.

"Single particle microscopy above and below the diffraction limit", Optical Society of America, Santa Clara, CA, September 28, 1999.

"Chemical physics with lasers: From slit jet discharges to single molecule spectroscopy", Department of Chemistry, University of Wisconsin, Madison, WI, October 26, 1999.

"Where Chemistry meets Physics", CU Wizards Science Outreach Program, Department of Chemistry, University Colorado, Boulder, CO, October 30, 1999.

"From state-to-state reaction dynamics to single molecule microscopy", Department of Chemistry, University of Maryland, College Park, MD, November 11, 1999.

"Chemical dynamics with a twist: From state-resolved reactions in supersonic jets to single molecule microscopy", Department of Chemistry, University of Southern California, Los Angeles, CA, January 10, 2000.

"Chemical kinetics with a twist: From state-to-state reaction dynamics to single molecule microscopy", Department of Chemistry, University of Arizona, Tucson, AZ, January 24, 2000.

"Microscopy at and below the diffraction limit via resonant scattering and laser induced fluorescence: Recent progress from apertureless NSOM", American Physical Society, Minneapolis, MN, March 21, 2000.

"Probing quantum state to state dynamics: From clusters to chemical reactions", American Chemical Society (219th national Meeting), San Francisco, CA, March 26, 2000.

"From Single Collisions to Single Molecules", Institute for Physical Chemistry, University of Goettingen, Goettingen, Germany, April 13, 2000

"Spectroscopy above and below the diffraction limit", Max Planck Institute for Biophysical Chemistry, Goettingen, Germany, April 28, 2000.

"State-resolved IR Laser Studies of Fundamental Reaction Dynamics", Atomic and Molecular Interactions Gordon Conference, New London, NH, July 4, 2000.

"Spectroscopy, kinetics and single collision dynamics via high resolution IR laser studies of radicals", 16th International Symposium on Gas Kinetics, Cambridge, UK, July 23, 2000.