

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE Nov 2000	3. REPORT TYPE AND DATES COVERED Proceedings on Invironmental Ergonomics VIII	
4. TITLE AND SUBTITLE EFFECTS OF WEARING IMPERMEABLE AND PERMEABLE PROTECTIVE CLOTHING ON THERMOREGULATORY RESPONSES WHILE SEDENTARY		5. FUNDING NUMBERS	
6. AUTHOR(S) THOMAS L. ENDRUSICK AND RICHARD R. GONZALEZ		8. PERFORMING ORGANIZATION REPORT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) U.S. ARMY RESEARCH INSTITUTE OF ENVIRONMENTAL MEDICINE NATICK, MA 01760-5007		10. SPONSORING / MONITORING AGENCY REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. ARMY MEDICAL RESEARCH & MATERIEL COMMAND FORT DETRICK, MD 21702-5012		11. SUPPLEMENTARY NOTES	
12a. DISTRIBUTION / AVAILABILITY STATEMENT APPROVED FOR PUBLIC RELEASE; DISTRIBUTION UNLIMITED		12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) The purpose of this study was to investigate if protective overgarments made from the same material but with different moisture vapor transmission rates would influence thermoregulatory responses in volunteers exposed to typical indoor workplace environments. Eight volunteers wore an impermeable overgarment (IO) and a permeable overgarment (PO) during a 4 h sedentary exposure at 18.3°C/50% rh (COOL) and 29.7°C/52% rh (WARM). During both COOL and WARM, skin and core temperatures were lower when wearing the PO. Skin wettedness was significantly higher during both COOL and WARM when wearing the IO. The IO had the highest weight increase due to absorption of non-evaporated moisture vapor during the 4 h test. These results showed that a moisture vapor permeable overgarment reduced overall thermal strain, minimized underclothing absorption of sweat and increased evaporation of moisture vapor during an extended sedentary exposure to simulated workplace environments.			
14. SUBJECT TERMS protective clothing, moisture vapor transmission rate, indoor workplace environments, cleanroom manufacturing.		15. NUMBER OF PAGES 4	
17. SECURITY CLASSIFICATION OF REPORT U		16. PRICE CODE	
18. SECURITY CLASSIFICATION OF THIS PAGE U		20. LIMITATION OF ABSTRACT U	
19. SECURITY CLASSIFICATION OF ABSTRACT U		20. LIMITATION OF ABSTRACT U	

DTIC QUALITY IMPROVED 4

20001124 055

Environmental Ergonomics VIII

International Series on Environmental Ergonomics, Volume 1.

Editors: James A. Hodgdon
Jay H. Heaney
Michael J. Buono

Naval Health Research Center
and
San Diego State University
San Diego, California, USA



Selected papers from the
8th International Conference on Environmental Ergonomics

San Diego, California, USA

18-23 October 1998

EFFECTS OF WEARING IMPERMEABLE AND PERMEABLE PROTECTIVE CLOTHING ON THERMOREGULATORY RESPONSES WHILE SEDENTARY

Thomas L. Endrusick and Richard R. Gonzalez

U.S. Army Research Institute of Environmental Medicine
Natick, MA 01760-5007 USA

INTRODUCTION

The use of protective clothing as a barrier against occupational and environmental hazards has increased dramatically in recent years. Certain types of protective overgarments are also being utilized in cleanroom manufacturing environments where contamination of the work site by personnel is a major concern. In the computer semiconductor manufacturing business, there is a reported industry-wide perception that the sedentary nature of the work does not justify the wearing of more costly, "breathable" protective clothing versus inexpensive, disposable, non-permeable garments. It is understandable that some managers of large-scale, industrial protective clothing and equipment programs would purchase specific garments based solely on a minimal cost per unit basis. Nevertheless, a recent study suggests that the use of higher-cost, vapor-permeable, and reusable protective clothing can actually be more economical when analyzed on a cost per use basis (1). The purpose of this present study was to investigate if protective overgarments manufactured from the same basic materials but with different levels of permeability would have an influence on thermoregulatory responses in volunteers who were sedentary and exposed to two typical indoor workplace environments.

MATERIALS AND METHODS

Eight healthy males (age= 21.0 ± 1.9 yrs, height=173.3 ± 5.6 cm, weight=72.5 ± 6.3 kg, body surface area=1.86 ± 0.10 m²) volunteered for the study. They were informed of the purpose, procedures and risks of the study. All volunteers expressed an understanding of the study by signing a statement of informed consent. All test overgarments were manufactured from material containing a waterproof/ breathable, protective membrane. The material was made by W.L. Gore and Associates. The protective membrane was composed of a thin layer of microporous polytetrafluoroethylene (PTFE). The PTFE membrane can be manufactured with varying levels of permeability. Test overgarment materials were evaluated by the manufacturer for permeability according to ASTM Standard E96-80 (2) which is used to calculate a moisture vapor transmission rate (MVTR, g·m⁻²·24 h⁻¹). All volunteers wore both an impermeable overgarment (IO, MVTR=5) and a permeable overgarment (PO, MVTR=864) during a 4-hour sedentary exposure to two different

environments: 18.3°C/50% rh (COOL); 29.7°C/52% rh (WARM). There was a constant air velocity of 1.1 m·s⁻¹ directed at the volunteers as they sat in the climatic chamber. All volunteers also wore lightweight 100% polyester underwear, gloves, socks, and leather boots. Mean weighted skin temperature (\bar{T}_{sk} , 8 sites, °C), rectal temperature (T_{re} , °C), skin wettedness (w, %) calculated from dew point temperature within the overgarment, and heart rate (HR, bpm) were measured. Total body mass loss (\dot{m}_b , g·h⁻¹) and moisture absorption (g) by the various garments were determined by pre- and post-experiment weights of all clothing items.

RESULTS

Table 1. Initial and final values (Mean \pm 1 SD) of \bar{T}_{sk} and T_{re} of volunteers (n=8) when wearing the IO and PO during COOL and WARM.

GARMENT		COOL	WARM
IO	Initial \bar{T}_{sk}	29.5 (1.0)	30.8 (1.6)
	Final	27.8 (0.9)	32.9 (1.5)
	Initial T_{re}	36.9 (0.2)	37.0 (0.3)
	Final	36.5 (0.5)	37.2 (0.2)
PO	Initial \bar{T}_{sk}	29.2 (1.3)	30.7 (1.3)
	Final	27.7 (0.7)	32.1 (1.3)
	Initial T_{re}	36.9 (0.3)	37.0 (0.3)
	Final	36.3 (0.7)	36.9 (0.3)

Figure 1. shows w of the volunteers while wearing both overgarments during COOL and WARM. There were significant increases in w during both environmental conditions when wearing the IO. At 4 hours exposure, w approached 0.9 when wearing the IO during WARM. Excessive skin wettedness has been shown to exacerbate the rate of body heat storage (3, 4).




Figure 1. Local skin wettedness (Mean \pm 1 SD) of volunteers (n=8) while wearing the IO and PO during COOL and WARM as a function of time of exposure.

Although there were no significant differences in HR when wearing either overgarment, HR during WARM was elevated an average of 24% and 19% above COOL values with the IO and PO, respectively. Mean m_b was lower during COOL (IO=77.5, PO=78.5) and higher during WARM (IO=92.3, PO=103). The IO had the highest mean weight increase (11 g, COOL and 44 g, WARM) due to absorption and/or condensation of moisture vapor within the overgarment during the 4-hour test. Absorption of moisture vapor also caused higher mean underclothing/footwear weight increases (22 g, COOL and 43 g, WARM) when worn with the IO. These weight increases were lower (20 g, COOL and 13 g, WARM) when wearing the PO.

CONCLUSIONS

These results showed that a moisture-vapor-permeable overgarment reduced overall thermal strain, reduced underclothing absorption of sweat and increased evaporation of moisture vapor when compared with a non-permeable overgarment during an extended sedentary exposure to simulated workplace environments. Cleanroom personnel can be required to wear completely-encapsulating protective clothing ensembles for up to 12 hours during an extended work shift. The use of protective clothing ensembles with sufficient thermal resistance and increased levels of moisture vapor transmission can improve overall thermal comfort that could lead to subsequent improvements in task performance and workforce morale.

DISCLAIMER

The views, opinions, and/or findings contained in this paper are those of the authors and should not be construed as an official Department of the Army position, policy, or decision unless so designated by other official documentation. Citation of

commercial organizations and registered trade names of products in this paper do not constitute an official Department of the Army endorsement or approval of the products or services of the organization. Approved for public release; Distribution is unlimited.

REFERENCES

1. Schwope, A.D. and Renard, E.P. 1992. Estimation of the cost of using chemical protective clothing. In J.P. McBriarty and N.W. Henry (eds.), *Performance of Protective Clothing: Fourth Volume*, ASTM STP 1133 (American Society of Testing and Materials, Philadelphia). 972-981.
2. American Society for Testing and Materials 1984. *Standard Test Methods for Water Vapor Transmission of Materials, Standard Test Method E96-80*. Philadelphia, PA.
3. Gagge, A.P. and Gonzalez, R.R. 1974. Physiological and physical factors associated with warm discomfort in sedentary man. *Environmental Research*, 7, 230-242.
4. Gonzalez, R.R. 1988. Biophysics of heat transfer and clothing considerations. In K.B. Pandolf, M.N. Sawka, and R.R. Gonzalez (eds.), *Human Performance Physiology and Environmental Medicine at Terrestrial Extremes* (Benchmark Press, Inc., IN). 45-94.