

AD _____

Award Number: DAMD17-99-1-9328

TITLE: STAT5A Regulates the Survival of Mammary Epithelial Cells
and the Development of Mammary Cancer

PRINCIPAL INVESTIGATOR: Robin Humphreys, Ph.D.

CONTRACTING ORGANIZATION: National Institutes of Health
Bethesda, Maryland 20892

REPORT DATE: July 2000

TYPE OF REPORT: Annual Summary

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

DTIC QUALITY INSPECTED 3

20010108 158

REPORT DOCUMENTATION PAGE*Form Approved*
OMB No. 074-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE July 2000	3. REPORT TYPE AND DATES COVERED Annual Summary (1 Jul 99 - 30 Jun 00)	
4. TITLE AND SUBTITLE STAT5A Regulates the Survival of Mammary Epithelial Cells and the Development of Mammary Cancer			5. FUNDING NUMBERS DAMD17-99-1-9328	
6. AUTHOR(S) Robin Humphreys, Ph.D.				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) National Institutes of Health Bethesda, Maryland 20892 E-MAIL: robinh@box-r.nih.gov			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012			10. SPONSORING / MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES Report contains color photos				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited				12b. DISTRIBUTION CODE
13. ABSTRACT (Maximum 200 Words) The <i>in vivo</i> relationship between epidermal growth factor (EGF) and prolactin/Jak/Stat signaling pathways in mammary gland development and tumorigenesis was explored in transgenic mice overexpressing the TGF alpha gene (TGF α TG) and lacking the prolactin-activated transcription factor, Stat5a (Stat5aKO/TGF α TG). We present evidence that Stat5a has a crucial role regulating programmed cell death (PCD) during mammary gland involution and the development of EGF-dependent tumorigenesis. TGF α TG mice exhibit delayed mammary gland involution and the absence of Stat5a facilitated involution-associated changes in morphology and the extent and timing of PCD. Stat5a remained phosphorylated through involution in TGF α TG suggesting that Stat5a is constitutively "activated" by the EGF signaling pathway. These results demonstrated TGF α epithelium is prevented from progressing into the normal pattern of involution and apoptosis by the "activated" form of Stat5a. This activated Stat5a affected EGF-receptor initiated mammary gland tumorigenesis. Overexpression of TGF α in mammary epithelium generates mammary hyperplasia and tumors. In the presence of an activated EGF receptor, deletion of Stat5a delayed initial hyperplasia and mammary tumor development by 6 weeks. These observations demonstrate that Stat5a is a survival factor, is required for mammary gland epithelium to resist regression and involution-mediated apoptosis and can influence the development of mammary gland tumorigenesis.				
14. SUBJECT TERMS Breast Cancer, prolactin, STAT, EGF, mouse model, tumor				15. NUMBER OF PAGES 28
				16. PRICE CODE
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited	

FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the U.S. Army.

RCH Where copyrighted material is quoted, permission has been obtained to use such material.

RCH Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

RCH Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.

X In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and use of Laboratory Animals of the Institute of Laboratory Resources, national Research Council (NIH Publication No. 86-23, Revised 1985).

N/A For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.

N/A In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

N/A In the conduct of research utilizing recombinant DNA, the investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules.

N/A In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories.


PI - Signature 7.24.00.
Date

Table of Contents

1. Cover.....	1
2. SF298.....	2
3. Foreword.....	3
4. Table of Contents.....	4
5. Introduction.....	5
6. Body.....	6
7. Key Research Accomplishments.....	8
8. Reportable Outcomes.....	9
9. Appendices	10
Manuscripts and Abstracts	

Introduction

The objective of this research is to elucidate the relationship between the action of the transcription factor, Stat5a, and the regulation of programmed cell death in the mammary gland in order to establish the role of Stat5a in the development of breast cancer. This will be achieved through the development of a unique mouse model of breast cancer and the molecular analysis of the signaling pathways critical for the activation of this factor. This mouse model will permit evaluation of specific molecules involved in the Stat5a signaling pathway as modulators of involution and tumorigenesis.

Body

This research project began with interbreeding of the WAP-TGF α transgenic and Stat5a knockout mouse lines and to provide a detailed study of the effects of Stat5a deletion on WAP TGF α -induced tumor progression. This combination transgenic/knockout mouse line (TGF α /Stat5aKO) was established and was expanded through standard breeding protocols to generate sufficient experimental and control mice for the tumorigenesis studies. After 8 months of breeding and data collection it was evident that the absence of Stat5a did have an effect on the development of tumors in the WAP-TGF α transgenic tumor model. In the presence of TGF α overexpression, complete genetic deletion of Stat5a delayed initial mammary tumor development by 6 weeks. This preliminary data was presented in a poster format to the Breast Cancer Think Tank at Chantilly, Virginia. The alteration in the progression of involution was confirmed by histological analysis. WAP-TGF α transgenic mice have a significant delay in the onset of involution at the end of lactation. This delay is suggested to be one of the mechanisms for mammary tumor formation in this mouse model. It was observed that TGF α /Stat5aKO mammary glands underwent a significant reorganization and deletion of cell structure during involution. These changes replicated those observed during involution of a normal mammary gland. This observation suggested that the absence of Stat5a in the TGF α /Stat5aKO mice was allowing involution to occur and possibly reducing the tumorigenic capacity of the epithelium. Stat5a requires phosphorylation in order to be activated and is normally rapidly dephosphorylated at the beginning of involution (within 12 hours). Phosphorylation analyses of the tissues from WAP-TGF α transgenic mice revealed that Stat5a was inappropriately phosphorylated during involution. This provided evidence that the EGF signaling pathway was activating Stat5a during involution and preventing the regression of the epithelium. The EGF pathway was simultaneously activated implicating the EGF-dependent signaling kinase, MAPK, as the molecule initiating this phosphorylation event. This confirmed our earlier suspicions about the potential action of TGF α on Stat5a signaling in the mammary gland. Detailed molecular analysis of proliferation and programmed cell death levels revealed that there was an increase in apoptosis in the TGF α /Stat5aKO mouse when compared to the WAP-TGF α transgenic control gland. Surprisingly, apoptosis was increased during late pregnancy in the TGF α /Stat5aKO glands when normally cell death is low or absent in the wildtype gland. This was the first evidence that Stat5a can act as a survival factor for mammary epithelium. These experiments demonstrated that the absence of Stat5a affected the initiation of involution and the levels of apoptosis in the WAP-TGF α mammary gland. This data lead us to conclude that Stat5a can mediate signaling from the EGF receptor and can influence the progression of EGF-initiated breast cancer.

The absence of Stat5a delayed the initial appearance of tumor formation by 6-8 weeks. It did not completely abrogate the appearance of tumors and the rate of tumor formation was the same as the control group. This suggested that the process of involution can remove significant numbers of cells that could potentially contribute to the development of tumors. But it also suggested that there remained sufficient cells after the process of involution, that could respond to EGF-activated signaling, to initiate transformation of the epithelium. The data generated was sufficient to write and submit a manuscript "*Signal Transducer and Activator of Transcription 5a Influences Mammary Epithelial Cell Survival and Tumorigenesis*". The manuscript was accepted and published in Cell Growth and Differentiation in October, 1999.

The effect of Stat5a on the level and expression of cell death regulating molecules was examined through RNase protection assays. No significant change in the expression patterns of the Bcl-2 family members was observed. The activation of caspases 3, 6 and 9 molecules is currently under examination with protein and enzymatic assays. Initial experiments have been performed with specific kinase inhibitors to block the selected pathways that may phosphorylate and activate Stat5a protein. These experiments are aimed at determining the level and duration of exposure required for significant inhibition of Stat5a phosphorylation and possibly induction of Stat5a-dependent apoptosis. Implants were generated from plastic pellets combined with inhibitors for the MAPK, JAK and cAMP-dependent kinases. Results from these studies have demonstrated initial levels of kinase inhibitor required to effect minor morphological changes in the mammary epithelium immediately around the implant. No change in Stat5a status has been detected to date. This may require an increase in the dose and or duration of the kinase inhibitor. This data and the data published in the manuscript in October 1999, was presented in a platform talk at the Keystone Conference "*Advances in Breast and Prostate Cancer*" in Lake Tahoe in March of 2000.

A review article "*Transforming growth factor alpha and mouse models of human breast cancer*" was submitted and published in Oncogene in January 2000. This manuscript reviewed the known mouse models of human disease generated with the TGF α gene including my work with the TGF α /Stat5aKO.

This project is also interested in the possible mechanisms of compensation or competition for Stat5a action by related family members. Based on conclusions from our work with the TGF α /Stat5aKO mouse model a collaboration was setup with Dr. David Levys lab at New York Medical Center, New York to study the role of Stat3 in the development and transformation of the mammary gland. I have developed another combination transgenic-knockout mouse model to examine the role of this molecule in mouse mammary cancer and development. This mouse model is currently being developed.

Key Research Accomplishments

1. **Paper:** *Signal Transducer and Activator of Transcription 5a Influences Mammary Epithelial Cell Survival and Tumorigenesis* published in *Cell Growth and Differentiation* **10**:685-694, October, 1999.
2. **Paper:** *Transforming growth factor alpha and mouse models of human breast cancer* published in *Oncogene* (review article) **19**: 1085-1091, January ,2000.
3. **Platform talk** :Presented at Keystone Meeting "Advances in Breast and Prostate Cancer" , March , 2000.
4. **Poster:** Presented at Breast Cancer Think Tank Meeting, July, 1999. Received poster award to attend meeting.
5. Generated the combined WAP-TGF α transgenic Stat5a knockout mouse model of breast cancer.
6. Demonstrated Stat5a has a cell survival function in the mammary epithelium.
7. Demonstrated a connection between Stat5a activity and the development of mammary cancer.
8. Established that Stat5a has a regulatory role in the involution of the mammary gland .
9. Used an *in vivo* model of breast cancer to demonstrated that the regulation of the activity a transcription factor can affect the development and progression of transformation in the mammary gland.
10. Applied for several faculty positions based on research from this work. University of Washington, University of Arizona , University of Oregon, University of Washington (St. Louis).
11. Collected tissue from all animal models to generate a tissue bank of various stages of WAP-TGF α transgenic and TGF α /Stat5aKO mammary glands. These tissues were used for various mammary gland analyses within and outside our lab. Including microarray analysis of gene expression of tumor types in the mammary gland. The data garnered from this analysis was used to prepare a manuscript from the Hennighausen lab on the development of a microarray analysis program(submitted).
12. Developed a collaboration to study the role of a related Stat molecule in the same breast cancer mouse model.

Reportable Outcomes

1. **Paper:** *Signal Transducer and Activator of Transcription 5a Influences Mammary Epithelial Cell Survival and Tumorigenesis* published in *Cell Growth and Differentiation* 10:685-694, October, 1999.
2. **Paper:** *Transforming growth factor alpha and mouse models of human breast cancer* published in *Oncogene* (review article) 19: 1085-1091, January ,2000.
3. **Platform talk:** Presented at Keystone Meeting "Advances in Breast and Prostate Cancer" , March , 2000.
4. **Poster:** Presented at Breast Cancer Think Tank Meeting, July, 1999. Received poster award to attend meeting.
5. Generated the combined WAP-TGF α transgenic/Stat5a knockout mouse model of breast cancer.

Signal Transducer and Activator of Transcription 5a Influences Mammary Epithelial Cell Survival and Tumorigenesis¹

Robin C. Humphreys² and Lothar Hennighausen

Laboratory of Genetics and Physiology, National Institute of Digestive, Diabetes and Kidney Disease, NIH, Bethesda, Maryland 20892

Abstract

The mammary gland undergoes extensive tissue remodeling and cell death at the end of lactation in a process known as involution. We present evidence that the prolactin-activated transcription factor signal transducer and activator of transcription 5a (Stat5a) has a crucial role in the regulation of cell death during mammary gland involution. In a transforming growth factor- α transgenic mouse model that exhibited delayed mammary gland involution, the absence of Stat5a facilitated involution-associated changes in morphology of the gland and the extent and timing of programmed cell death. These Stat5a-dependent changes also affected epidermal growth factor receptor-initiated mammary gland tumorigenesis. Overexpression of the transforming growth factor α transgene in the mammary epithelium reproducibly generated mammary hyperplasia and tumors. In the presence of the activated epidermal growth factor receptor, deletion of Stat5a delayed initial hyperplasia and mammary tumor development by 6 weeks. These observations demonstrate that Stat5a is a survival factor, and its presence is required for the epithelium of the mammary gland to resist regression and involution-mediated apoptosis. We also suggest that Stat5a is one of the antecedent, locally acting molecules that initiate the process of epithelial regression and reorganization during involution.

Introduction

Involution is the phase of mammary gland development that remodels the molecular and morphological characteristics of the gland after lactation. The systemic levels of prolactin, one of the dominant hormones in the stimulation and maintenance of lactational competence of the gland, decreases at the onset of involution (1). The characteristics of the biolog-

ical activity and regulation of prolactin argue for prominent involvement of this systemic hormone in the induction of mammary gland involution. Prolactin secretion is tightly regulated and is stimulated by suckling. The half-life of prolactin is very short in the blood, and there is evidence that exogenous treatment with prolactin can delay involution at certain stages of development (2, 3). Despite the clear role of this systemic hormone in the initiation of involution, the role of its subservient signal transduction pathways in the regulation of the morphological and transcriptional changes of involution is unknown. The prolactin-activated transcription factor Stat5a³ is essential for the development and differentiation of the mammary gland (4). Stat5a phosphorylation sharply declines within 12 h after weaning, suggesting a role in the onset of involution. In fact, Stat5a has been implicated in the regulation of apoptotic processes in other tissues (5, 6).

Mammary gland involution can be separated into two molecularly and morphologically distinct phases. The first phase is defined by changes in the expression patterns of genes involved in milk synthesis (WAP and ornithine decarboxylase), cell survival and death (Bax, Bcl-2, Bcl-x_s, SGP-2, interleukin-converting enzyme 1, and p53), and proteins that act to regulate potential survival factors (insulin-like growth factor-1, insulin-like growth factor-binding protein 5, and TGF- β) and modulate interactions with the extracellular matrix (tissue inhibitor of metalloproteinase 1; Refs. 3 and 7–11). Importantly, dephosphorylation of the prolactin-dependent transcription factors Stat5a and Stat5b can be detected within 12 h of the initiation of involution (12). Despite the high levels of apoptosis early in involution, the integrity of the alveolar basement membrane is not disrupted. The second phase of involution is characterized by increases in the expression of proteins and genes involved in reorganizing the extracellular matrix (gelatinase 1, stromelysin 1, and urokinase plasminogen activator) and abrogation of milk gene expression. These molecular changes precede the dramatic degradation of the lobuloalveolar acini and the absorption of apoptotic cells. It is presumed that changes in the concentrations of the systemic hormones prolactin, oxytocin, and progesterone, in concert with changes in the activity of locally acting factors, initiate and stimulate the process of involution.

Received 5/11/99; revised 8/4/99; accepted 8/10/99.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked advertisement in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

¹ Supported by Department of Defense, Medical Research and Material Command Grant DAMD17-99-1-9328 (to R. C. H.).

² To whom requests for reprints should be addressed, at Laboratory of Genetics and Physiology, National Institute of Digestive, Diabetes and Kidney Disease, NIH, Building 8, Room 111, Bethesda MD 20892. Phone: (301) 435-6634; Fax: (301) 480-7312; E-mail: robinh@box-r.nih.gov.

³ The abbreviations used are: Stat, signal transducer and activator of transcription; TGF, transforming growth factor; Jak, janus kinase; MAPK, mitogen-activated protein kinase; WAP, whey acidic protein; EGFR, epidermal growth factor receptor; BrdUrd, bromodeoxy uridine; WT, wild-type; PCD, programmed cell death; EGF, epidermal growth factor; TBST, Tris buffered saline Tween; TUNEL, terminal deoxynucleotidyl transferase-mediated uridine nick end labeling; ERK, extracellular signal-regulated kinase.

The importance of the Stat proteins in the growth and differentiation of many cell types is exemplified by their ubiquitous expression and widespread utilization in the signal transduction pathways of multiple cytokines (13). Stats are capable of transmitting both differentiative and proliferative signals, depending on the cellular and receptor context (14, 15). In the mammary gland, Stat5a activation by prolactin is required for functional differentiation of the epithelium during pregnancy and lactation (4, 16). Singular tyrosine phosphorylation by the prolactin receptor-associated intracellular kinase Jak-2 is obligatory to maintain Stat5a transcriptional activity (17). However, novel alternative mechanisms of regulating Stat5a activity have been identified. Prolactin-activated signal transduction can tyrosine- and serine-phosphorylate multiple isoforms of the Stat5a molecules (18). Furthermore, truncated forms of Stat5a have transcriptional dominant negative activity (19, 20) and have been linked to the regulation of apoptosis (5). In the liver, EGF can stimulate phosphorylation and nuclear translocation of the Stat5a isoform, Stat5b (21). Importantly, serine phosphorylation of Stat5a by the prolactin-independent, MAPK pathway has been identified, (22) including a direct association between Stat5a and the serine kinase ERK-1/2(MAPK) (23). Serine phosphorylation by the MAPK pathway is required for transcriptional activity of Stat1 and Stat3, but serine phosphorylation is not obligatory for prolactin-dependent transcriptional activity of Stat5 proteins (24). Taken together, these data imply that Stat5a activity can be regulated by the prolactin and EGF signaling pathways and possesses diverse functional roles within a single cell that are dependent on its phosphorylation state and expressed isoforms.

Initiation and progression of breast cancer rely on the inappropriate temporal and qualitative activity of normal cellular signaling pathways (25). These alterations in signaling activity often disrupt the normal mechanisms of mammary gland development. In one mouse model (WAP-TGF- α), overexpression of TGF- α , which binds to and activates the EGFR, delays mammary involution and transforms the mammary epithelium (26). The prolactin signaling pathway has been implicated in loss of normal epithelial differentiation (4, 16) and transformation (27, 28). The role of Stat5a in tumorigenesis is implicated by the appearance of Stat5a overexpression in leukemic (29, 30) and oncogenic (31) transformed cells. Additionally, constitutive activation of the Jak/Stat pathway occurs in multiple leukemia-, v-abl-, and human T-cell lymphotropic virus-transformed T cells (32, 33), and specific inhibition of constitutive Jak-2 activity in acute lymphoblastic leukemia can block cell growth and induce cell death (34). We used the WAP-TGF- α transgenic and Stat5a-null mouse models to explore the *in vivo* role of Stat5a in mammary gland involution and tumorigenesis.

Results

Delayed Mammary Involution in the TGF α TG Mice Is Curtailed in the Absence of Stat5a. The contribution of Stat5a in regulating involution and the establishment of mammary tumors was investigated in mice that carried the TGF- α transgene (TGF α TG mice) or were null for the Stat5a gene (Stat5aKO mice) and in Stat5a-null mice that expressed the

transgene (Stat5aKOTGF α mice). H&E-stained mammary glands from TGF α TG, Stat5aKO, Stat5aKOTGF α , and WT nontransgenic control mice at day 18 of pregnancy and days 1, 3, and 7 of involution were examined by light microscopy (Fig. 1). Morphological differences were observed between the TGF α TG and Stat5aKOTGF α mammary glands within the first pregnancy and involution. TGF α TG mammary glands contained hyperproliferative alveoli and ducts. Alveolar lumina contained heterogeneous eosin-positive staining material. A significant increase in the number of stromal cells was evident, in addition to increased deposition of stromal collagen (Fig. 1*l*). Involution was delayed in the TGF α TG mice, as reported previously (26). Some cell loss and condensation of lobuloalveolar structures was observed during involution, but the gland lacked the dramatic epithelial regression of the WT gland (Fig. 1, *n* versus *p*). Mammary tissue from pregnant Stat5aKOTGF α mice was histologically similar to pregnant tissue from TGF α TG mice but contained more epithelium than observed in the glands of Stat5aKO mice (Fig. 1, *a*–*d*). Secretory structures appeared to be more uniform and lacked the observed heterogeneous precipitations in the alveolar lumens. During involution, the Stat5aKOTGF α gland remodeled more rapidly and extensively than the TGF α TG gland (Fig. 1, *o* versus *p*). This effect was enhanced after subsequent rounds of pregnancy and involution (data not shown). The primary and secondary ducts of the Stat5aKOTGF α mice lacked concentrations of hyperplastic cells like those found in the TGF α TG mice. On rare occasions, hyperplastic regions were evident after three or more rounds of pregnancy in Stat5aKOTGF α mice (data not shown).

Apoptosis and Proliferation Levels Are Altered in the Stat5aKOTGF α and TGF α TG Mice. High levels of apoptosis, decreased proliferation, and the collapse of lobuloalveolar structures in the mammary epithelium are hallmarks of involution in the mammary gland (7). To establish and define the characteristics of mammary involution in the Stat5aKOTGF α and TGF α TG mice, the levels of apoptosis and proliferation were analyzed by TUNEL and BrdUrd immunohistochemistry, respectively (Tables 1 and 2). WT glands exhibited increased levels of apoptosis within 1 day of the initiation of weaning and exhibited maximal levels at day 3, as expected (9, 35). Consistent with an inhibition or delay in involution in the TGF α TG gland, the number of apoptotic cells was slightly reduced at day 1 of involution compared with WT and Stat5aKOTGF α glands (Fig. 2A). Apoptosis was reduced in the TGF α TG gland at day 7 when compared with WT gland at day 7. Whereas Stat5aKOTGF α mammary glands at day 18 of pregnancy ($P = 0.02$) and at day 1 of involution ($P = 0.015$) had significantly higher levels of apoptosis than TGF α TG and WT mammary glands over the same time period, the levels were similar at days 3 and 7. Interestingly, Stat5aKO mammary glands also displayed significantly high levels of apoptosis at day 18 of pregnancy and day 1 of involution ($P = 0.03$).

Levels of proliferation, as measured by BrdUrd incorporation, were equivalent in mammary tissue of pregnant Stat5aKOTGF α , TGF α TG, Stat5aKO, and WT mice (Fig. 2B). Generally, 6–10% of the cells in the pregnant gland were




Fig. 1. Involution is enhanced in the absence of *Stat5a*. H&E staining of inguinal mammary glands from *Stat5a*-null nontransgenic (NTGKO), WT, *Stat5a*-null TGF- α transgenic (KOTG), and TGF- α transgenic (TG) mice at 18 days of pregnancy (p18, a–d) and days 1 (i1, e–h), 3 (i3, i–l), and 7 (i7, m–p) of involution. Samples were collected from mice during or immediately after their first pregnancy. Compare epithelial condensation at day 3 and day 7 of involution in the WT (j) versus the KOTG (k) and TG (l). Bar in a, 200 μ m.

Table 1 Percentage of apoptotic cells detected with TUNEL

Values are the mean value $\pm$ SE. Superscript symbols denote mean values that were significantly different from the WT mean value of the same day.

	KOTG	TG	WT	NTGKO
Preg. day 18	4.21 $\pm$ 0.65 ^a	0.86 $\pm$ 0.31	1.47 $\pm$ 0.56	2.61 $\pm$ 0.52
Involution day 1	9.01 $\pm$ 1.46 ^b	2.01 $\pm$ 0.49	3.21 $\pm$ 0.96	7.65 $\pm$ 1.19 ^c
Involution day 3	6.96 $\pm$ 0.41	4.92 $\pm$ 1.92	5.97 $\pm$ 0.75	4.35 $\pm$ 0.14
Involution day 7	3.32 $\pm$ 0.74	1.75 $\pm$ 0.56	4.56 $\pm$ 1.42	4.16 $\pm$ 1.89

^a $P = 0.02$.

^b $P = 0.015$.

^c $P = 0.03$.

Table 2 Percentage of proliferating cells detected with BrdUrd labeling

Values are the mean $\pm$ SE. Superscript symbols denote mean values that were significantly different from the WT mean value of the same day.

	KOTG	TG	WT	NTGKO
Preg. day 18	10.30 $\pm$ 5.16	10.03 $\pm$ 1.77	6.97 $\pm$ 1.18	4.09 $\pm$ 1.72
Involution day 1	7.40 $\pm$ 2.12 ^a	3.63 $\pm$ 1.33	1.38 $\pm$ 0.49	1.53 $\pm$ 0.23
Involution day 3	6.43 $\pm$ 0.64 ^b	4.00 $\pm$ 1.84	0.67 $\pm$ 0.27	0.96 $\pm$ 0.15
Involution day 7	6.87 $\pm$ 0.80 ^c	5.62 $\pm$ 1.05 ^d	0.07 $\pm$ 0.07	0.65 $\pm$ 0.28

^a $P = 0.05$.

^b $P = 0.001$.

^c $P = 0.001$.

^d $P = 0.007$.

labeled with BrdUrd (Fig. 2B). Proliferation was low in involuting tissue from *Stat5a*KO and WT mice (<1%). In contrast, the TGF- α TG and *Stat5a*KOTGF- α glands had detectable, persistent proliferation levels of 4–7% throughout involution. Apoptotic cells were observed in *Stat5a*KOTGF- α mammary glands at day 18 of pregnancy and day 1 of involution in animals labeled exclusively with BrdUrd (Fig. 2C, arrows). The appearance of apoptotic, BrdUrd-labeled cells demon-

strated that DNA synthesis in these cells preceded the initiation of PCD.

WAP-TGF- α Transgene Expression Is Present throughout Mammary Gland Development in *Stat5a*KOTGF- α and TGF- α TG Mice. The WAP gene promoter has been exploited to target high-level expression of transgenes, including TGF- α , to the mammary epithelium of pregnant mice (36–40). To assure that the hormonal and morphological changes




Fig. 2. A, KOTG mammary glands display elevated levels of apoptotic cells during pregnancy and early involution. Apoptosis levels in Stat5a-null TGF- α transgenic (KOTG), TGF- α transgenic (TG), Stat5a-null nontransgenic (NTGKO), and WT mammary glands after analysis by the TUNEL technique. Values represent the percentage of cells (means $\pm$ SE) displaying the label in a minimum population of 1000 cells. Analysis was performed on mice during or immediately after their first pregnancy. Serial sections of tissue from the same mouse were used for the determination of proliferation and apoptosis. A minimum of three mice were analyzed from each genotype (see "Materials and Methods."). Total number of cells counted, 7.5×10^5 . a, b, and c denote significant differences between the KOTG and NTGKO values when compared with apoptosis levels in WT glands at day 18 of pregnancy and day 1 of involution. Specific values for Fig. 2, A and B, are shown in Tables 1 and 2. B, proliferation is higher in TG and KOTG mammary glands through involution. Proliferation levels in Stat5a-null TGF- α transgenic (KOTG), TGF- α transgenic (TG), Stat5a-null nontransgenic (NTGKO), and WT mammary glands after analysis by BrdUrd labeling. Values represent the percentage of cells (means $\pm$ SE) displaying the label in a minimum population of 1000 cells. Analysis was performed on mice during or immediately after their first pregnancy. Serial sections of tissue from the same mouse were used for determination of proliferation and apoptosis. A minimum of three mice were analyzed from each genotype. Total number of cells counted, 6.5×10^5 . a, b, c, and d significant differences between TG and KOTG samples when compared with WT at day 1, 3, and 7 of involution. C, proliferation analysis of mammary gland in Stat5a-null TGF- α transgenic mammary glands. This figure demonstrates the appearance of BrdUrd-labeled apoptotic cells in late pregnancy (p18) and days 1 and 7 of involution (i1 and i7). Mice were injected 2 h before sacrifice with $20 \mu\text{g/g}$ body weight BrdUrd, as described in "Materials and Methods." Gray arrows, BrdUrd-labeled apoptotic cells. Black arrowheads, intact BrdUrd-labeled nuclei. Bar in p18, $100 \mu\text{m}$.

associated with involution and the absence of Stat5a did not disrupt transgene expression, TGF- α expression was evaluated in mammary tissue of pregnant and involuting TGF α TG, WT, and Stat5aKOTGF α mice. Northern blot analysis of TGF- α transgene expression in TGF α TG and Stat5aKOTGF α mice revealed expression of an appropriately sized transcript at day 18 of pregnancy. Transgene expression persisted through days 1, 3, and 7 of involution in multiple animals (Fig. 3). This expression pattern deviates from the expression of the endogenous WAP gene, which declines at the initiation of involution (41). Endogenous TGF- α expression was only detected in an overloaded lane containing wildtype tissue at day 1 of involution. TGF- α transgene expression persisted through multiple rounds of pregnancy (three to nine rounds) in TGF α TG and Stat5aKOTGF α mice (data not shown).

The EGFR-activated Serine Threonine Kinase MAPK Remains Active throughout Involution. Binding of TGF- α to the EGFR initiates receptor dimerization and inherent kinase activation, which stimulates the Ras/Raf/Mek signaling cascade. To establish that the EGFR-dependent pathway was being activated by the expression of the TGF- α transgene, Western blot analysis was performed on protein extracts from Stat5aKOTGF α , WT, Stat5aKO, and TGF α TG animals with an antibody that only recognizes the active (phosphorylated) form of MAPK. Active MAPK was detected in TGF α TG and Stat5aKOTGF α mammary glands at day 18 of pregnancy and days 1, 3, and 7 of involution (Fig. 4, A and B). Increased levels of the active form of this enzyme were detectable after multiple rounds of pregnancy in the TGF α TG and Stat5aKOTGF α mammary glands. Very low levels of




Fig. 3. TGF- α transgene expression persists through involution in KOTG and TG mammary glands. Northern analysis of TGF- α expression in TG, KOTG, and WT mammary glands. Total RNA (20 μ g) was separated and analyzed on a single Northern blot. Each lane represents RNA from a single animal. Samples were collected from mice during or immediately after their first pregnancy. The TGF- α blot was exposed for 4 h at -70°C . EtBr, ethidium bromide staining of RNA.

active MAPK were detected in the nontransgenic Stat5aKO gland (Fig. 4C). Levels were also very low in pregnant and involuting WT mammary glands (Fig. 4D). A significant level of MAPK is present in the lactating, WT mammary gland.

Stat5a Proteins Are Phosphorylated during Involution in TGF α TG and Stat5aKOTGF α Mice. Stat5a is phosphorylated on tyrosine 694 by Jak-2, which is associated with the dimerized prolactin receptor (13). Stat5a must remain phosphorylated to maintain its transcriptional activity. One of the earliest molecular events after the onset of involution is the dephosphorylation of Stat5a and Stat5b (42). Immunoprecipitation and Western blot analysis of protein extracts from TGF α TG mice detected high levels of tyrosine-phosphorylated Stat5a and Stat5b at days 1, 3, and 7 of involution (Fig. 5). Conversely, the phosphorylation of Stat5a and Stat5b was decreased at day 1 and remained low to day 7 of involution in WT controls, as reported previously and shown in Fig. 5B (42). Additionally, tyrosine-phosphorylated Stat5a and Stat5b were also detected by immunoprecipitation and Western blot analysis in multiple individual TGF α TG tumors (data not shown). Stat5aKOTGF α mammary glands, lacking any Stat5a protein, were assayed for the phosphorylation of Stat5b to determine whether there was a compensation for the absence of Stat5a phosphorylation. A slight increase in Stat5b phosphorylation was detected on days 1, 3, and 7 of involution relative to WT controls.

TGF- α -initiated Mammary Tumorigenesis Is Delayed in Stat5aKOTGF α Mice. The rat TGF- α transgene under control of the WAP gene promoter induces mammary tumors in mice bred *ad lib* after three pregnancies (26). The contribution of Stat5a to the establishment of mammary hyperplasias and tumors was investigated in TGF α TG and Stat5aKOTGF α mice. Stat5aKOTGF α and TGF α TG mice were bred *ad lib* and examined after each round of pregnancy for palpable hyperplasias and tumors. All 48 TGF α TG mice contained palpable hyperplasia in multiple glands by the third pregnancy. TGF α TG mammary glands consistently displayed a swollen appearance by the second pregnancy, emblematic of hypertrophic glands and hyperplasias (data not shown). Gross histological examination revealed the presence of multiple hyperplastic alveolar nodules in multiple glands. TGF α TG glands also contained fluid-filled cysts and stromal proliferation. Some removed tumors were characterized by




Fig. 4. MAPK is active during pregnancy and involution in TG and KOTG mammary glands. Phosphorylation of MAPK was examined by Western blot analysis with an antibody specific for the active phosphorylated form of MAPK and is shown in the top panel of each section. 18p, i1, i3, and i7 represent samples collected from TG (A) KOTG (B), NTGKO (C), and WT (D) mammary glands. 4L, a control sample from a WT, 4 day lactating mammary gland. α , a sample from a KOTG mammary gland after three pregnancies. β , a TG mammary gland after three pregnancies. Each immunoblot was stripped and reblotted with an antibody to MAPK to determine loading, and the result is shown directly under the phosphorylated MAPK immunoblot in each panel. Arrows in A indicate the two subunits of MAPK, ERK-1 (top) and ERK-2 (bottom).

abnormal hyperproliferation of glandular structures with relatively little hyperplasia. The latency of mammary hyperplasia and tumor appearance and the histology of the hyperplasias and tumors were consistent with the results described previously (26). Hyperplasia and hypertrophy appeared less frequently in the Stat5aKOTGF α mammary gland. There was a significant difference in the rate of initial hyperplasia and tumor formation between the TGF α TG and Stat5aKOTGF α mice, ($P = 0.003$; Fig. 6). The initial appearance in the Stat5aKOTGF α mice was delayed 1.5 months compared with the TGF α TG mice. In Stat5aKOTGF α mice




Fig. 5. A, Stat5a remains phosphorylated during involution in TG mammary glands. Stat5a and Stat5b phosphorylation was examined in KOTG and TG mice mammary glands. Protein extracts were immunoprecipitated with anti-Stat5a antibodies (Stat5a, 1.2×10^5) or anti-Stat5b antibodies (Stat5b, 1.1×10^5) overnight. Immunoprecipitates were separated by PAGE and immunoblotted with anti-phosphotyrosine (p-Tyr, 1.1×10^3) antibodies. Samples were collected from mice during or immediately after their first pregnancy. Immunoblots probed initially with p-Tyr were stripped and reblotted with anti-Stat5a antibodies in the TG samples (5a/5b; 5a/5b) or anti-Stat5b antibodies in the KOTG samples (5a/5b; 5a/5b), respectively, to determine protein loading. i1, i3, and i7, days 1, 3, and 7 of involution, respectively. B, Stat5a and Stat5b are dephosphorylated during involution. Stat5a and Stat5b phosphorylation was examined in NTGKO and WT mice mammary glands. Protein extracts were immunoprecipitated with anti-Stat5a antibodies (Stat5a, 1.2×10^5) or anti-Stat5b antibodies (Stat5b, 1.1×10^5) overnight. Immunoprecipitates were separated by PAGE and immunoblotted with anti-phosphotyrosine (p-Tyr, 1.1×10^3) antibodies. Samples were collected from mice during or immediately after their first pregnancy. Immunoblots probed initially with p-Tyr were stripped and reblotted with anti-Stat5a antibodies or anti-Stat5b antibodies to determine protein loading. p18, i1, i3, and i7, day 18 of pregnancy and days 1, 3, and 7 of involution, respectively.

that did develop hyperplasias or tumors, usually only a single gland was involved, as compared with multiple gland involvement in the TGF α TG mice.

Discussion

Loss of Stat5a Activity Is an Early Event in Mouse Mammary Gland Involution. Dephosphorylation and hence inactivation of the transcription factors Stat5a and Stat5b and *de novo* phosphorylation of Stat3 occur in the first phase within several hours of pup removal (42), implicating the Stats and prolactin as early regulatory molecules in the re-




Fig. 6. Absence of Stat5a delays EGFR-mediated hyperplasia and tumor formation. Kaplan-Meier plot of the percentage of hyperplasia and tumor-free KOTG and TG mice over a 9-month time period. All mice were examined at each pregnancy for palpable tumors. There is a significant difference between the two curves ($P = 0.0357$). TG, $n = 48$. KOTG, $n = 34$.

gression of the gland. Using a combination of transgenic and gene deletion mouse models, we have now demonstrated a role for Stat5a in involution that is dependent on its phosphorylation and can be regulated by constitutive EGFR signal transduction. Involution is abrogated on continued activation of the EGFR (and Stat5a) through the expression of a TGF α transgene. In mammary tissue of TGF α TG mice, Stat5a remained phosphorylated, as did Stat5b. However, on deletion of the Stat5a gene, involution proceeded unabated, despite expression of the TGF α transgene. Rapid involution in the absence of Stat5a was accompanied by increased levels of apoptotic cells. Our experiments not only demonstrate that Stat5a is a survival factor *in vivo* but also show that its presence is mandatory to avoid the initiation of PCD in mammary epithelial cells.

Despite the lack of known genetic targets for Stat5a in the involuting gland, there is evidence to suggest that prolactin and Stat5a can control cell survival. Activation of Stat5a has been implicated in the inhibition of apoptosis in T cells through interleukin 2 (6) and interleukin 9 (43) signaling. In addition, selection of apoptosis-resistant immature T cells leads to the activation of a dominant negative form of Stat5a (5). Prolactin can also stimulate cell survival in mammary epithelial cells *in vitro* (44), and prolactin and growth hormone can promote cell survival in the involuting rat mammary gland (2, 3). The presence of apoptotic cells in the mammary tissue of pregnant mice that lack the Stat5a gene and express the TGF α transgene suggests that cell dependence on the presence of phosphorylated Stat5a precedes the actual systemic hormonal switch into involution. This observation also implies that the Stat5a-dependent mechanism of apoptosis induction may be hormonally independent. We did not observe this apoptotic phenomena in the WT and TGF α transgenic mice. This increased number of apoptotic cells during pregnancy was also detected in BrdUrd-labeled Stat5aKOTGTG mice. Analysis of these glands revealed the presence of BrdUrd-labeled apoptotic cells at day 18 of pregnancy and at days 1, 3, and 7 of involution. These cells replicated their DNA and then proceeded directly into PCD.

This association between the cell's ability to replicate its DNA and the progression into the apoptotic program has been reported previously (45). Cells may be programmed to initiate PCD before they enter the cell cycle; consequently, BrdUrd-labeled cells undergoing apoptosis are detected. Interestingly, not all the cells in the Stat5aKOTGF α pregnant gland undergo PCD. This suggests that some cells have obviated the requirement for Stat5a, possibly through an unknown compensatory mechanism. Alternatively, these cells may have a unique differentiation state that precludes them from entering the apoptotic pathway.

Despite persistent phosphorylation of Stat5a in involuting mammary tissue, TGF α TG mice did undergo involution-dependent epithelial reorganization and activation of PCD. This morphological change was characterized by moderate condensation of alveolar structures and loss of cells through the induction of PCD. We propose that the dephosphorylation of Stat5a is not the only guiding factor for apoptosis progression in the involuting mammary gland and that other parallel signals may be required for the complete induction of both phases of involution (12).

Transgenic TGF- α Activates the EGFR and Ensures Persistent Phosphorylation of Stat5a. TGF- α stimulates the EGFR and activates MAPK through the Ras/Raf/Mek signaling cascade. This is exemplified by the presence of the activated form of MAPK during involution in the TGF α TG and Stat5aKOTGF α mice. Persistent activation and phosphorylation of Stat5a in the TGF α TG mice could occur through three distinct mechanisms. The EGFR can directly phosphorylate Stat1 and Stat3 in the absence of Jak activation (46–48). Although direct phosphorylation of Stat5a by the EGFR cannot be ruled out, it was not possible to coimmunoprecipitate this receptor with anti-Stat5a antibodies in the TGF α TG mammary gland (data not shown). Secondly, stimulation of MAPK by EGF can lead to phosphorylation and nuclear translocation of the Stat5a isoform Stat5b in the liver (21). Support for this mechanism comes from results describing that Stat5a and its shorter isoforms can be tyrosine- and serine-phosphorylated by prolactin (18), and Stat5a and Stat5b can be tyrosine- and serine-phosphorylated by prolactin-independent, MAPK-dependent mechanisms (22, 49). A direct interaction was recently demonstrated between MAPK(ERK1/2) and Stat5a in growth hormone-dependent signaling (23). Growth hormone is known to influence mammary gland involution, although this effect is not as dominant as that observed for prolactin (2). Thirdly, EGFR could activate Stat5a through Jak-2. The EGFR is known to use Jak-1 to activate Stat1 and Stat3 (14, 50) and can modulate Stat5 expression in mammary epithelial cells (51). Theoretically, the constitutively active EGFR in the TGF α TG and Stat5aKOTGF α mice could stimulate Jak-2, although this was not formally proven. Alternatively, EGFR could stimulate a secondary Jak, which inappropriately uses Stat5a as a substrate that has been demonstrated in IFN- γ signaling (52, 53). Presumably, the stimulation of the EGFR kinase in the presence of the TGF- α transgene leads to phosphorylation and activation of a downstream kinase, possibly including a Jak, that maintains Stat5a in its phosphorylated state. We suggest that one or more of these mechanisms regulate the

phosphorylation of Stat5a and control apoptosis in the mouse mammary gland.

Persistent Stat5a Activity Contributes to Mouse Mammary Tumorigenesis, and Its Absence Acts to Delay Hyperplastic Epithelial Formation. One of the dominant pathways commonly altered in breast cancer is the EGF signal transduction pathway (25, 54). Gene deletion studies in mice have demonstrated that the EGFR is not dispensable for normal mammary development (55, 56), and that aberrant signaling through this receptor induces neoplastic growth and frank tumors (26, 57–59). Using an EGF-dependent transgenic mouse model, we can now show that the persistent activation of Stat5a is linked to the transformation of the mammary gland. We suggest that the transformation of the mammary epithelium occurs at the expense of a loss of cell survival regulation, which is contingent on the persistent phosphorylation of Stat5a and permits the inappropriate survival of epithelial cells. The appearance of tumors and hyperplasias in Stat5aKOTGF α mice suggests that the absence of Stat5a will not completely block mammary epithelial transformation, and that the TGF- α transgene can activate alternative (non-Stat5a) pathways to bypass PCD and involution and support tumor formation. Such pathways may include Stat5b, which, under specific circumstances, substitutes for Stat5a (60). The involvement of a single gland in a majority of the Stat5aKOTGF α tumors suggests that significant numbers of potentially transformed cells are destroyed during involution. This loss of cells probably contributes to the observed lower rate of tumor formation and the delay in initial tumor appearance. In spite of single gland involvement, survival of the Stat5aKOTGF α mice was no greater than that observed for the TGF α TG mice after initial tumor appearance.

Our findings that Stat5a is a survival factor and that its absence reduces mammary tumorigenesis support findings from other settings. Prolactin and other type 1 cytokines do possess mitogenic activity, implicating the associated Stat molecules with contributing to the mitogenesis of these organs (13). Constitutive Stat activity has been observed in cells transformed by oncogenic viruses (32) and oncogenes (61) and in leukemia cells (29). Moreover, constitutive activation of Jak expression can lead to transformation (27, 62).

Stat5a Is a Survival Factor for the Mammary Epithelium. The absence of Stat5a disrupts the prolactin-mediated survival of the mammary epithelium. Whereas overexpression of TGF- α in the presence of an intact Stat5a results in delayed involution, apoptosis and involution in the absence of Stat5a lead to elimination of the majority of the epithelium. The dephosphorylation or lack of *de novo* phosphorylation and inactivation of Stat5a that occur during involution in the WT gland are mimicked by the absence of Stat5a in the Stat5aKOTGF α mice. Therefore, we propose that the presence of the activated form of this transcription factor prolongs cell survival, and its inactivation or deletion permits PCD to occur. These experiments demonstrate that Stat5a can act as a survival factor for the epithelium of the mammary gland. In addition, they illustrate that the inhibition and/or disruption of the mechanisms that regulate this involution process are critical in mammary gland tumorigenesis.

The conclusions from this study lead us to hypothesize that the inhibition or disruption of Stat5a phosphorylation can lead to protection from transformation in the mammary gland. The application of Stat and/or Jak-specific tyrosine or serine kinase inhibitors may be an approach to blunt the development of mammary tumors originating from EGFR deregulation.

Materials and Methods

Materials. The TGF- α cDNA probe was a kind gift from Dr. David Lee (Lineberger Comprehensive Cancer Center, University of North Carolina, Chapel Hill, NC). The generation of the Stat5a antibodies has been described previously (63). Phosphorylated MAPK antibody was purchased from Promega (Madison, WI). Anti-phosphotyrosine antibody was purchased from Upstate Biotechnology, Inc. (Lake Placid, NY). Goat anti-rabbit and rabbit antimouse secondary antibodies were purchased from Transduction Laboratories (Lexington, KY).

Generation of TGF α TG and Stat5aKOTGF α Mice. WAP-TGF α TG mice were a gift of Dr. Eric Sandgren (University of Wisconsin, Madison, WI Ref. 26). The generation of the Stat5a-null mice has been described previously (4). TGF α TG mice were interbred with Stat5a-null mice (SvEv129/C57B6) to generate F₁ founders that were hemizygous for Stat5a deletion (Stat5aHT) and transgenic for TGF- α expression. Female Stat5aHTTGF α mice were bred with Stat5aHTTGF α males to generate offspring that were transgenic for TGF- α and homozygous for deletion of the Stat5a gene. These mice were backcrossed for five generations to generate a pure inbred Stat5aKOTGF α strain of mice. The same generation TGF α TG and control WT littermates was used for molecular analyses. Confirmation of the presence of the TGF- α transgene was performed by PCR analysis with TGF- α forward primer 5'-TGTCAGGCTCTGGAGAA-CAGC-3' and reverse primer 5'-CACAGCGAACACCCACGTACC-3'. Stat5a PCR was performed with two sets of primers to knockout (null) and WT alleles: (a) Stat5a forward (5'-CTGGATTGACGTTCTTACCTG-3') and Stat5a reverse (5'-TGGAGTCACTAGTCTGTCTCT-3') and (b) Neo forward (5'-AGAGGCTATTCGGCTATGACTG-3') and Neo reverse 5'-TTCGTCCAGATCATCTGATC-3'. PCR for all primers was performed with a denaturing step of 94°C for 3 min, followed by 30 cycles of 94°C for 40 s, 56°C for 40 s, and 68°C for 40 s, followed by 10 min at 68°C. Genotype of the mice was confirmed after tissue collection by Northern and Western blot analysis for TGF- α gene expression and absence of Stat5a protein expression, respectively. All animals were housed and handled according to the approved protocol established by the Institutional Animal Care and Use Committee and NIH guidelines.

Mammary Gland Collection. Mammary glands were surgically removed from anesthetized and cervically dislocated mice at day 18 of pregnancy and days 1, 3, and 7 of involution. Day 1 of involution was designated as 24 h after the morning that the pups were born. Pups were immediately removed from the dam after birth and fostered onto a WT mother. The mammary lymph node was removed before homogenization of all glands. Tissues were prepared immediately for RNA and protein extraction, as described previously (42). Whole (number 4) inguinal or (number 3) thoracic mammary glands were surgically excised from mice, spread on Omniset tissue cages (Fisher Scientific, Pittsburgh, PA), fixed for 5 h in Tellyzinckys fixative, and stored in 70% ethanol until processed by standard embedding and sectioning techniques onto Probe-On Plus slides (Fisher Scientific). Sections were stained with H&E.

Immunoprecipitations and Western Blot Analysis. Preparation of protein extracts and immunoprecipitations have been described previously (42). Briefly, 2 mg of fresh and frozen tissue were homogenized in 2 ml of lysis buffer with protease inhibitors; phenylmethylsulfonyl fluoride, leupeptin, and aprotinin at 50 μ g/ml on ice. Protein lysates were rocked for 1 h at 4°C and then cleared by centrifugation at 14,000 $\times$ g for 15 min. The supernatants were removed, mixed with 2 $\times$ loading buffer, and heated to 90°C for 3 min. Samples were spun briefly, electrophoresed under denaturing conditions on 8% precast tris-glycine gels, and transferred to polyvinylidene difluoride membranes according to manufacturer's protocol (Novex, San Diego, CA). Western blot analysis was performed essentially as described with the following exceptions; primary antibody (Stat5a, 1:20,000 dilution; Stat5b, 1:10,000 dilution; anti-phosphotyrosine, 1:5,000 dilution; anti-MAPK, 1:5,000 dilution) was incubated

overnight at 4°C with gentle rocking, and all incubations with antibodies and initial blocking were performed with 3% nonfat dried milk in 1 $\times$ TBST. Detection was performed with the enhanced chemiluminescence kit according to manufacturer's protocol (Amersham) and exposed to Kodak (Rochester, NY) MR autoradiography film. Exposure times are between 1 s and 2 min. Immunoprecipitations with anti-Stat5a antibodies were carried out as described previously (4). Stripping was performed by incubating blots at 56°C in 6.25 mM Tris-HCl (pH 6.8), 2% SDS, and 1% β -mercaptoethanol for 30 min. Blots were washed extensively in TBST and then blocked with 3% nonfat dried milk in TBST.

Northern Blot Analysis. Total RNA was isolated from fresh tissue by homogenization in lysis buffer as described previously (63). RNA was quantified by spectrophotometry and prepared for Northern blot analysis by heating in loading buffer at 65°C. Total RNA was separated by 1.5% agarose gel electrophoresis and transferred to Hybond N+ (Amersham) nylon membrane by capillary transfer with 10 $\times$ SSC. After overnight transfer, the membrane was UV-irradiated and hybridized to each cDNA probe in Quikhyb (Stratagene, La Jolla, CA). Hybridization for the TGF- α cDNA probe was performed in Quikhyb (Stratagene) at 65°C for 16 h, followed by two washes in 1 $\times$ SSC and 0.5% SDS for 30 min, followed by one wash in 0.1 $\times$ SSC and 0.5% SDS for 30 min at 56°C. Blots were hybridized for 16 h and then washed twice in 1 $\times$ SSC and 0.5% SDS for 30 min, followed by one wash in 0.1 $\times$ SSC and 0.5% SDS for 30 min. cDNA and oligo probes were random-primed and Klenow-labeled with [α -³²P]dCTP. After hybridization and washing, blots were exposed from 30 min to 24 h, as specifically described in the figure legends, to Kodak MR autoradiography film at -70°C.

BrdUrd and TUNEL Assays. Protocols for BrdUrd and TUNEL analysis have been described elsewhere (64). Mice were injected 2 h before sacrifice with 20 μ g/g body weight BrdUrd labeling reagent, as described by the manufacturer (Amersham). Each proliferation and apoptosis sample counted represents a minimum of three random fields (at $\times$ 200) and a minimum of 1000 total cells/section for each mouse. A minimum of three mice per timepoint were collected and analyzed. The total number of cells counted for proliferation assay is 6.5×10^5 . The total number of cells counted for the apoptosis assay is 7.5×10^5 . The number of mice collected and the number of mice analyzed are listed by genotype and by timepoints (pregnancy day 18/involution day 1/involution day 3/involution day 7) for apoptosis and proliferation, respectively: KOTG, 4/4/4/3 and 4/3/3/3; TG, 3/5/3/5 and 3/3/3/5; WT, 3/4/5/3 and 3/3/3/3; NTGKO, 3/3/2/3 and 3/3/3/3.

Hyperplasia and Tumor Analysis. Mice were palpated at the 18th day of pregnancy for mammary hyperplasia, hypertrophy, and tumors. Animals were scored for the presence of either hyperplasia or tumor at every pregnancy. Most animals were sacrificed at the third pregnancy by anesthesia followed by cervical dislocation. In studies that required long-term breeding for tumor development, dams were rebred within 2 days after birth and removal of the pups.

Acknowledgments

We thank Dr. Eric Sandgren for the kind gift of TGF- α transgenic mice. We thank Drs. J. Shillingford and E. Rucker for critical reading of the manuscript and Drs. P. Furth and B. Gusterson for essential discussions. We thank Ashton Garrett, Bill Kemp, and Sandra Price for animal husbandry and technical support and Dr. J. Shillingford for developing the Stat5a PCR assay.

References

- Vonderhaar, B. K. Prolactin: transport, function and receptors in mammary gland development and differentiation. In: M. C. Neville and C. W. Daniel (eds.), *The Mammary Gland: Development, Regulation and Function*, pp. 383-438. New York: Plenum Publishing, 1985.
- Travers, M. T., Barber, M. C., Tonner, E., Quarrie, L., Wilde, C. J., and Flint, D. J. The role of prolactin and growth hormone in the regulation of casein gene expression and mammary cell survival: relationships to milk synthesis and secretion. *Endocrinology*, 137: 1530-1539, 1996.
- Tonner, E., Quarrie, L., Travers, M., Barber, M., Logan, A., Wilde, C., and Flint, D. Does an IGF-binding protein (IGFBP) present in involuting rat mammary gland regulate apoptosis? *Prog. Growth Factor Res.*, 6: 409-414, 1995.

4. Liu, X., Robinson, G. W., Wagner, K. U., Garrett, L., Wynshaw-Boris, A., and Hennighausen, L. Stat5a is mandatory for adult mammary gland development and lactogenesis. *Genes Dev.*, **11**: 179–186, 1997.
5. Bovolenta, C., Testolin, L., Benussi, L., Lievens, P. M., and Liboi, E. Positive selection of apoptosis-resistant cells correlates with activation of dominant-negative STAT5. *J. Biol. Chem.*, **273**: 20779–20784, 1998.
6. Zamorano, J., Wang, H. Y., Wang, R., Shi, Y., Longmore, G. D., and Keegan, A. D. Regulation of cell growth by IL-2: role of STAT5 in protection from apoptosis but not in cell cycle progression. *J. Immunol.*, **160**: 3502–3512, 1998.
7. Strange, R., Friis, R. R., Bemis, L. T., and Geske, F. J. Programmed cell death during mammary gland involution. *Methods Cell Biol.*, **46**: 355–368, 1995.
8. Marti, A., Feng, Z., Altermatt, H. J., and Jaggi, R. Milk accumulation triggers apoptosis of mammary epithelial cells. *Eur. J. Cell Biol.*, **73**: 158–165, 1997.
9. Lund, L. R., Romer, J., Thomasset, N., Solberg, H., Pyke, C., Bissell, M. J., Dano, K., and Werb, Z. Two distinct phases of apoptosis in mammary gland involution: proteinase-independent and -dependent pathways. *Development (Camb.)*, **122**: 181–193, 1996.
10. Hadsell, D. L., Greenberg, N. M., Fligger, J. M., Baumrucker, C. R., and Rosen, J. M. Targeted expression of des(1–3) human insulin-like growth factor I in transgenic mice influences mammary gland development and IGF-binding protein expression. *Endocrinology*, **137**: 321–330, 1996.
11. Li, F., Strange, R., Friis, R. R., Djonov, V., Altermatt, H. J., Saurer, S., Niemann, H., and Andres, A. C. Expression of stromelysin-1 and TIMP-1 in the involuting mammary gland and in early invasive tumors of the mouse. *Int. J. Cancer*, **59**: 560–568, 1994.
12. Li, M., Liu, X., Robinson, G., Bar-Peled, U., Wagner, K. U., Young, W. S., Hennighausen, L., and Furth, P. A. Mammary-derived signals activate programmed cell death during the first stage of mammary gland involution. *Proc. Natl. Acad. Sci. USA*, **94**: 3425–3430, 1997.
13. Leonard, W. J., and O'Shea, J. J. Jak and STATs: biological implications. *Annu. Rev. Immunol.*, **16**: 293–322, 1998.
14. Zhong, Z., Wen, Z., and Darnell, J. E., Jr. Stat3: a STAT family member activated by tyrosine phosphorylation in response to epidermal growth factor and interleukin-6. *Science (Washington DC)*, **264**: 95–98, 1994.
15. Iwamoto, Y., Chin, Y. E., Peng, X., and Fu, X. Y. Identification of a membrane-associated inhibitor(s) of epidermal growth factor-induced signal transducer and activator of transcription activation. *J. Biol. Chem.*, **273**: 18198–18204, 1998.
16. Teglund, S., McKay, C., Schuetz, E., van Deursen, J. M., Stravopodis, D., Wang, D., Brown, M., Bodner, S., Grosveld, G., and Ihle, J. N. Stat5a and Stat5b proteins have essential and nonessential, or redundant, roles in cytokine responses. *Cell*, **93**: 841–850, 1998.
17. Darnell, J. E., Jr. STATs and gene regulation. *Science (Washington DC)*, **277**: 1630–1635, 1997.
18. Kirken, R. A., Malabarba, M. G., Xu, J., Liu, X., Farrar, W. L., Hennighausen, L., Larner, A. C., Grimley, P. M., and Rui, H. Prolactin stimulates serine/tyrosine phosphorylation and formation of heterocomplexes of multiple Stat5 isoforms in Nb2 lymphocytes. *J. Biol. Chem.*, **272**: 14098–14103, 1997.
19. Moriggi, R., Gouilleux-Gruart, V., Jahne, R., Berchtold, S., Gartmann, C., Liu, X., Hennighausen, L., Sotiropoulos, A., Groner, B., and Gouilleux, F. Deletion of the carboxyl-terminal transactivation domain of MGF-Stat5 results in sustained DNA binding and a dominant negative phenotype. *Mol. Cell Biol.*, **16**: 5691–5700, 1996.
20. Wang, D., Stravopodis, D., Teglund, S., Kitazawa, J., and Ihle, J. N. Naturally occurring dominant negative variants of Stat5. *Mol. Cell Biol.*, **16**: 6141–6148, 1996.
21. Ruff-Jamison, S., Chen, K., and Cohen, S. Epidermal growth factor induces the tyrosine phosphorylation and nuclear translocation of Stat 5 in mouse liver. *Proc. Natl. Acad. Sci. USA*, **92**: 4215–4218, 1995.
22. Yamashita, H., Xu, J., Erwin, R. A., Farrar, W. L., Kirken, R. A., and Rui, H. Differential control of the phosphorylation state of proline-juxtaposed serine residues Ser⁷²⁵ of Stat5a and Ser⁷³⁰ of Stat5b in prolactin-sensitive cells. *J. Biol. Chem.*, **273**: 30218–30224, 1998.
23. Pircher, T. J., Petersen, H., Gustafsson, J. A., and Haldosen, L. A. Extracellular signal-regulated kinase (ERK) interacts with signal transducer and activator of transcription (STAT) 5a. *Mol. Endocrinol.*, **13**: 555–565, 1999.
24. Wartmann, M., Cella, N., Hofer, P., Groner, B., Liu, X., Hennighausen, L., and Hynes, N. E. Lactogenic hormone activation of Stat5 and transcription of the β -casein gene in mammary epithelial cells is independent of p42 ERK2 mitogen-activated protein kinase activity. *J. Biol. Chem.*, **271**: 31863–31868, 1996.
25. Dickson, R. B., and Lippman, M. E. Growth factors in breast cancer. *Endocr. Rev.*, **16**: 559–589, 1995.
26. Sandgren, E. P., Schroeder, J. A., Qui, T. H., Palmiter, R. D., Brinster, R. L., and Lee, D. C. Inhibition of mammary gland involution is associated with transforming growth factor α but not c-myc-induced tumorigenesis in transgenic mice. *Cancer Res.*, **55**: 3915–3927, 1995.
27. Schwaller, J., Frantsve, J., Aster, J., Williams, I. R., Tomasson, M. H., Ross, T. S., Peeters, P., Van Rompaey, L., Van Etten, R. A., Ilaria, R., Jr., Marynen, P., and Gilliland, D. G. Transformation of hematopoietic cell lines to growth-factor independence and induction of a fatal myelo- and lymphoproliferative disease in mice by retrovirally transduced TEL/JAK2 fusion genes. *EMBO J.*, **17**: 5321–5333, 1998.
28. Wennbo, H., Gebre-Medhin, M., Gritli-Linde, A., Ohlsson, C., Isaksson, O. G., and Tornell, J. Activation of the prolactin receptor but not the growth hormone receptor is important for induction of mammary tumors in transgenic mice. *J. Clin. Invest.*, **100**: 2744–2751, 1997.
29. Zhang, Q., Nowak, I., Vonderheid, E. C., Rook, A. H., Kadin, M. E., Nowell, P. C., Shaw, L. M., and Wasik, M. A. Activation of Jak/STAT proteins involved in signal transduction pathway mediated by receptor for interleukin 2 in malignant T lymphocytes derived from cutaneous anaplastic large T-cell lymphoma and Sezary syndrome. *Proc. Natl. Acad. Sci. USA*, **93**: 9148–9153, 1996.
30. Hayakawa, F., Towatari, M., Iida, H., Wakao, H., Kiyoi, H., Naoe, T., and Saito, H. Differential constitutive activation between STAT-related proteins and MAP kinase in primary acute myelogenous leukaemia. *Br. J. Haematol.*, **101**: 521–528, 1998.
31. Yu, C. L., Jove, R., and Burakoff, S. J. Constitutive activation of the Janus kinase-STAT pathway in T lymphoma overexpressing the Lck protein tyrosine kinase. *J. Immunol.*, **159**: 5206–5210, 1997.
32. Migone, T. S., Lin, J. X., Cereseto, A., Mulloy, J. C., O'Shea, J. J., Franchini, G., and Leonard, W. J. Constitutively activated Jak-STAT pathway in T cells transformed with HTLV-I. *Science (Washington DC)*, **269**: 79–81, 1995.
33. Danial, N. N., Pernis, A., and Rothman, P. B. Jak-STAT signaling induced by the *v-ab1* oncogene. *Science (Washington DC)*, **269**: 1875–1877, 1995.
34. Meydan, N., Grunberger, T., Dadi, H., Shahar, M., Arpaia, E., Lapidot, Z., Leeder, J. S., Freedman, M., Cohen, A., Gazit, A., Levitzki, A., and Roifman, C. M. Inhibition of acute lymphoblastic leukaemia by a Jak-2 inhibitor. *Nature (Lond.)*, **379**: 645–648, 1996.
35. Quarrie, L. H., Addey, C. V., and Wilde, C. J. Apoptosis in lactating and involuting mouse mammary tissue demonstrated by nick-end DNA labelling. *Cell Tissue Res.*, **281**: 413–419, 1995.
36. Wagner, K. U., Young, W. S., III, Liu, X., Ginns, E. I., Li, M., Furth, P. A., and Hennighausen, L. Oxytocin and milk removal are required for postpartum mammary gland development. *Genes Funct.*, **1**: 233–244, 1997.
37. Tzeng, Y. J., Zimmermann, C., Guhl, E., Berg, B., Avantiaggiati, M. L., and Graessmann, A. SV40 T/t-antigen induces premature mammary gland involution by apoptosis and selects for p53 missense mutation in mammary tumors. *Oncogene*, **16**: 2103–2114, 1998.
38. Li, B., Murphy, K. L., Laucirica, R., Kittrell, F., Medina, D., and Rosen, J. M. A transgenic mouse model for mammary carcinogenesis. *Oncogene*, **16**: 997–1007, 1998.
39. Jhappan, C., Geiser, A. G., Kordon, E. C., Bagheri, D., Hennighausen, L., Roberts, A. B., Smith, G. H., and Merlino, G. Targeting expression of a transforming growth factor β 1 transgene to the pregnant mammary gland inhibits alveolar development and lactation. *EMBO J.*, **12**: 1835–1845, 1993.

40. Jager, R., Herzer, U., Schenkel, J., and Weiher, H. Overexpression of Bcl-2 inhibits alveolar cell apoptosis during involution and accelerates c-myc-induced tumorigenesis of the mammary gland in transgenic mice. *Oncogene*, 15: 1787-1795, 1997.
41. Pittius, C. W., Sankaran, L., Topper, Y. J., and Hennighausen, L. Comparison of the regulation of the whey acidic protein gene with that of a hybrid gene containing the whey acidic protein gene promoter in transgenic mice. *Mol. Endocrinol.*, 2: 1027-1032, 1988.
42. Liu, X., Robinson, G. W., and Hennighausen, L. Activation of Stat5a and Stat5b by tyrosine phosphorylation is tightly linked to mammary gland differentiation. *Mol. Endocrinol.*, 10: 1496-1506, 1996.
43. Bauer, J. H., Liu, K. D., You, Y., Lai, S. Y., and Goldsmith, M. A. Heteromerization of the γ c chain with the interleukin-9 receptor α subunit leads to STAT activation and prevention of apoptosis. *J. Biol. Chem.*, 273: 9255-9260, 1998.
44. Leff, M. A., Buckley, D. J., Krumenacker, J. S., Reed, J. C., Miyashita, T., and Buckley, A. R. Rapid modulation of the apoptosis regulatory genes bcl-2 and bax by prolactin in rat Nb2 lymphoma cells. *Endocrinology*, 137: 5456-5462, 1996.
45. Evan, G. I., Brown, L., Whyte, M., and Harrington, E. Apoptosis and the cell cycle. *Curr. Opin. Cell Biol.*, 7: 825-834, 1995.
46. Sartor, C. I., Dziubinski, M. L., Yu, C. L., Jove, R., and Ethier, S. P. Role of epidermal growth factor receptor and STAT-3 activation in autonomic proliferation of SUM-102PT human breast cancer cells. *Cancer Res.*, 57: 978-987, 1997.
47. David, M., Wong, L., Flavell, R., Thompson, S. A., Wells, A., Lerner, A. C., and Johnson, G. R. STAT activation by epidermal growth factor (EGF) and amphiregulin. Requirement for the EGF receptor kinase but not for tyrosine phosphorylation sites or JAK1. *J. Biol. Chem.*, 271: 9185-9188, 1996.
48. Leaman, D. W., Pisharody, S., Flickinger, T. W., Commane, M. A., Schlessinger, J., Kerr, I. M., Levy, D. E., and Stark, G. R. Roles of JAKs in activation of STATs and stimulation of c-fos gene expression by epidermal growth factor. *Mol. Cell. Biol.*, 16: 369-375, 1996.
49. Kirken, R. A., Malabarba, M. G., Xu, J., DaSilva, L., Erwin, R. A., Liu, X., Hennighausen, L., Rui, H., and Farrar, W. L. Two discrete regions of interleukin-2 (IL2) receptor β independently mediate IL2 activation of a PD98059/rapamycin/wortmannin-insensitive Stat5a/b serine kinase. *J. Biol. Chem.*, 272: 15459-15465, 1997.
50. Shual, K., Ziemiecki, A., Wilks, A. F., Harpur, A. G., Sadowski, H. B., Gilman, M. Z., and Darnell, J. E. Polypeptide signalling to the nucleus through tyrosine phosphorylation of Jak and Stat proteins. *Nature (Lond.)*, 366: 580-583, 1993.
51. Petersen, H., and Haldosen, L. A. EGF modulates expression of STAT5 in mammary epithelial cells. *Exp. Cell Res.*, 243: 347-358, 1998.
52. Kohlhuber, F., Rogers, N. C., Watling, D., Feng, J., Guschin, D., Briscoe, J., Witthuhn, B. A., Kutenko, S. V., Pestka, S., Stark, G. R., Ihle, J. N., and Kerr, I. M. A JAK1/JAK2 chimera can sustain α and γ interferon responses. *Mol. Cell. Biol.*, 17: 695-706, 1997.
53. Kutenko, S. V., Izotova, L. S., Pollack, B. P., Muthukumar, G., Pauku, K., Silvennoinen, O., Ihle, J. N., and Pestka, S. Other kinases can substitute for Jak2 in signal transduction by interferon- γ . *J. Biol. Chem.*, 271: 17174-17182, 1996.
54. Lupu, R., Cardillo, M., Cho, C., Harris, L., Hijazi, M., Perez, C., Rosenberg, K., Yang, D., and Tang, C. The significance of heregulin in breast cancer tumor progression and drug resistance. *Breast Cancer Res. Treat.*, 38: 57-66, 1996.
55. Sebastian, J., Richards, R. G., Walker, M. P., Wiesen, J. F., Werb, Z., Derynck, R., Hom, Y. K., Cunha, G. R., and DiAugustine, R. P. Activation and function of the epidermal growth factor receptor and erbB-2 during mammary gland morphogenesis. *Cell Growth Differ.*, 9: 777-785, 1998.
56. Wiesen, J. F., Young, P., Werb, Z., and Cunha, G. R. Signaling through the stromal epidermal growth factor receptor is necessary for mammary ductal development. *Development (Camb.)*, 126: 335-344, 1998.
57. Muller, W. J., Arteaga, C. L., Muthuswamy, S. K., Siegel, P. M., Webster, M. A., Cardiff, R. D., Meise, K. S., Li, F., Halter, S. A., and Coffey, R. J. Synergistic interaction of the Neu proto-oncogene product and transforming growth factor α in the mammary epithelium of transgenic mice. *Mol. Cell. Biol.*, 16: 5726-5736, 1996.
58. Kenney, N. J., Smith, G. H., Maroulakou, I. G., Green, J. H., Muller, W. J., Callahan, R., Salomon, D. S., and Dickson, R. B. Detection of amphiregulin and Cripto-1 in mammary tumors from transgenic mice. *Mol. Carcinog.*, 15: 44-56, 1996.
59. Huang, S., Trujillo, J. M., and Chakrabarty, S. Proliferation of human colon cancer cells: role of epidermal growth factor and transforming growth factor α . *Int. J. Cancer*, 52: 978-986, 1992.
60. Liu, X., Gallego, M. I., Smith, G. H., Robinson, G. W., and Hennighausen, L. Functional release of Stat5a-null mammary tissue through the activation of compensating signals including Stat5b. *Cell Growth Differ.*, 9: 795-803, 1998.
61. Danial, N. N., Losman, J. A., Lu, T., Yip, N., Krishnan, K., Krolewski, J., Goff, S. P., Wang, J. Y., and Rothman, P. B. Direct interaction of Jak1 and v-Ab1 is required for v-Ab1-induced activation of STATs and proliferation. *Mol. Cell. Biol.*, 18: 6795-6804, 1998.
62. Frank, D. A., and Varticovski, L. BCR/ab1 leads to the constitutive activation of Stat proteins and shares an epitope with tyrosine phosphorylated Stats. *Leukemia (Baltimore)*, 10: 1724-1730, 1996.
63. Liu, X., Robinson, G. W., Gouilleux, F., Groner, B., and Hennighausen, L. Cloning and expression of Stat5 and an additional homologue (Stat5b) involved in prolactin signal transduction in mouse mammary tissue. *Proc. Natl. Acad. Sci. USA*, 92: 8831-8835, 1995.
64. Humphreys, R. C., Krajewska, M., Krnacik, S., Jaeger, R., Weiher, H., Krajewski, S., Reed, J. C., and Rosen, J. M. Apoptosis in the terminal endbud of the murine mammary gland: a mechanism of ductal morphogenesis. *Development (Camb.)*, 122: 4013-4022, 1996.

Abstract to poster presentation at the Breast Cancer think Tank Symposium Held At Chantilly Virginia

Signal Transducer and Activator of Transcription 5a (Stat5a) Regulates Tumorigenesis and Epithelial Cell Survival in the Mouse Mammary Gland

Robin C. Humphreys and Lothar Hennighausen

Laboratory of Genetics and Physiology National Institute of Digestive, Diabetes and Kidney Disease
National Institutes of Health, Bethesda, MD 20892

The mammary gland undergoes extensive tissue remodeling and cell death at the end of lactation in a process known as involution. We present evidence that the prolactin-activated transcription factor, Stat5a has a crucial role in the regulation of cell death during mammary gland involution and can affect the progression of epidermal growth factor (EGF)-dependent mammary tumorigenesis. In a transforming growth factor- α (TGF α) transgenic mouse model, that exhibited delayed mammary gland involution, the absence of Stat5a facilitated involution-associated changes in morphology of the gland and the extent and timing of programmed cell death (PCD). These Stat5a-dependent changes also altered EGF receptor-initiated mammary gland tumorigenesis. Overexpression of the TGF α transgene in the mammary epithelium reproducibly generated mammary hyperplasia and tumors. In the presence of TGF α overexpression, complete genetic deletion of Stat5a reduced the rate of tumor formation and delayed initial mammary tumor development by 6 weeks. These observations demonstrate that Stat5a is a survival factor and its presence is required for the epithelium of the mammary gland to resist regression and involution-mediated apoptosis. We suggest that Stat5a is one of the antecedent, locally-acting, molecules that initiate the process of epithelial regression and reorganization during involution. In addition, these studies suggest that disruption of the pathway which leads to Stat5a activation can affect the progression of tumorigenesis in the mammary gland. Supported by USMRMC DAMD 17-99-1-9328.

Abstract for Platform Presentation at Keystone Meeting "Advances in Breast and Prostate Cancer" in March, 2000.

Stat5a Influences Mammary Epithelial Cell Survival and Tumorigenesis

Robin C. Humphreys and Lothar G. Hennighausen. Laboratory of Genetics and Physiology, NIDDK, National Institutes of Health, Bethesda, MD 20892

The *in vivo* relationship between the epidermal growth factor (EGF) and prolactin/Jak/Stat signaling pathways in mammary gland development and tumorigenesis was explored in transgenic mice which overexpress the TGF alpha gene (TGF α TG) in the mammary epithelium and lack the prolactin-activated transcription factor, Stat5a (Stat5aKO). We present evidence that Stat5a has a crucial role in the regulation of cell death during mammary gland involution and the development of EGF-dependent tumorigenesis. TGF α TG mice exhibit delayed mammary gland involution and the absence of Stat5a in the Stat5aKO/TGF α TG gland facilitated involution-associated changes in morphology and the extent and timing of programmed cell death in the mammary gland. Stat5a remained phosphorylated through involution in the TGF α TG mammary glands suggesting that Stat5a is constitutively phosphorylated and "activated" by the EGF signaling pathway. These results suggested that the TGF α epithelium is prevented from progressing into the normal pattern of involution and apoptosis by the "activated" form of Stat5a. Alternatively, the presence of activated Stat5a is required for the epithelium of the TGF α TG mammary gland to resist regression and involution-mediated apoptosis.

These Stat5a-dependent changes also affected EGF-receptor initiated (EGFR) mammary gland tumorigenesis. Overexpression of the TGF α transgene in the mammary epithelium reproducibly generates mammary hyperplasia and tumors. In the presence of the activated EGFR, deletion of Stat5a delayed initial hyperplasia and mammary tumor development by 6 weeks. These observations demonstrate that Stat5a is a survival factor and its presence is required for the epithelium of the mammary gland to resist regression and involution-mediated apoptosis. This effect on involution can influence the development of mammary gland tumorigenesis. We are currently utilizing specific kinase inhibitors to block the phosphorylation of Stat5a by the EGFR signaling pathway in order to determine the mechanism of Stat5a activation. This work is supported by a grant from the DoD, USMRMC to R.C.H. [DAMD17-99-1-9328]



Transforming growth factor alpha and mouse models of human breast cancer

Robin C Humphreys^{*,1} and Lothar Hennighausen¹

¹National Institutes of Health, National Institute of Digestive, Diabetes and Kidney Disease, Laboratory of Genetics and Physiology, Building 8, Room 111, Bethesda, Maryland, MD 20892, USA

Transforming growth factor alpha (TGF α) is a principal molecule in the normal and neoplastic development of the mammary gland. Binding of TGF α to the epidermal growth factor receptor (EGFR), activates the EGFRs' endogenous tyrosine kinase activity and stimulates growth of the epithelium in the virgin and pregnant mouse mammary gland. TGF α expression can be detected in breast cancer cells *in vivo* and *in vitro* and overexpression can elicit partial transformation or immortalized human and rodent mammary epithelial cells. Despite evidence implicating TGF α in the development of mammary neoplasia, the actual mechanism of TGF α -induced transformation is unclear. Transgenic mouse models targeting heterologous TGF α to the mammary gland have established TGF α overexpression can induce hyperproliferation, hyperplasia and occasional carcinoma. These transgenic studies demonstrated a facilitating, proliferative role for TGF α in the development of neoplasia and implicated several oncogenes that can cooperate with TGF α to transform the mammary epithelium. From studies of EGFR signaling pathways, inhibitory and modulating agents such as anti-EGFR antibodies and specific kinases inhibitors have been used to block the action of this pathway and prevent the development of TGF α -induced neoplasia and tumor formation. Studies in Stat5a knockout mice have established that the JAK2/Stat5a pathway can facilitate the survival of the mammary epithelium and can impact the progression of TGF α -mandated mammary tumorigenesis. Together these experiments indicate that TGF α and the EGFR signaling pathway are potentially amenable to therapies for treatment of human breast disease. *Oncogene* (2000) 19, 1085–1091.

Keywords: TGF alpha; mammary gland; transformation; cancer; mouse models; transgenic

Introduction

Development and progression of breast cancer, like many other types of human cancer, is dependent on the progressive corruption and alteration of normal signaling pathways. Signaling mechanisms that transmit growth signals are a predominant target for carcinogenic and oncogenic alterations as they often confer a growth advantage to the pre-neoplastic cell. One of the primary growth factor receptors

involved in normal and neoplastic development of the mammary gland is the EGFR and its extended family of related receptors and peptide ligands (Table 1). EGFR and the ligand TGF α , have expression levels altered in the development of cancer in numerous issues. The related family of ligands that bind to these receptors; EGF, heregulin, amphiregulin, and cripto also display alterations in their expression and activity within the pre-neoplastic and transformed cell (for a review see (Dickson and Lippman, 1995; Gullick *et al.*, 1999; Schroeder and Lee, 1997)). Despite the well characterized role of the EGFR and TGF α in normal mammary growth signaling, in transformation of mammary epithelial cells *in vitro* and *in vivo* and their elevated expression in some breast cancers, the specific alterations of the EGFR signaling mechanism that lead to TGF α -initiated neoplasia are poorly understood. In an attempt to elucidate these alterations, several transgenic mouse models have been generated to explore the role of TGF α and EGFR in mammary neoplasia. This review will focus on TGF α and its role in transformation of the mammary gland. In addition, it will examine those studies that have attempted to identify the mechanisms of TGF α action in the mammary gland and those molecules that are capable of cooperatively interacting with TGF α to promote tumorigenesis.

TGF α and EGFR expression coincides with normal proliferation in vivo

TGF α is structurally and functionally similar to EGF; the peptides share a 42% identity and can elicit the same biological effects in cultured mammary epithelial cells and explants (Daniel and Silberstein, 1985; Salomon *et al.*, 1987; Vonderhaar, 1987). TGF α is often co-expressed with the EGFR and binding to the receptor activates the EGFRs' endogenous tyrosine kinase activity. Mammary epithelial cell lines can be stimulated to proliferate by TGF α (Smith *et al.*, 1989; Zajchowski and Sager, 1991) and it can act as an autocrine growth factor in normal and immortalized human mammary epithelial cells. TGF α autocrine and proliferative activity can be blocked in these cells by an anti-EGFR antibody (Bates *et al.*, 1990; Kenney *et al.*, 1993).

In normal mammary gland development, TGF α and EGF transcripts can be detected in the ductal and lobuloalveolar stages. The virgin mouse mammary gland expresses TGF α in the proliferative cap cells and the stromal fibroblasts around the neck of the terminal endbud, whereas EGF expression is localized

*Correspondence: RC Humphreys

Table 1

Ligand	Receptor
<i>EGF-Like</i>	
EGF	EGFR
TGF α	EGFR
Amphiregulin	EGFR
Epiregulin	EGFR ErbB-4
Betacellulin	EGFR ErbB-3 ErbB-4
Heparin binding-EGF	EGFR ErbB-3 ErbB-4
<i>Heregulin/neuregulins</i>	
Heregulin-1 (α and β)	ErbB-3 ErbB-4
Heregulin-2 (α and β)	ErbB-3 ErbB-4
Heregulin-3	?
Heregulin-4	ErbB-4
Glial growth factor	?
<i>Cripto</i>	
Cripto-1	? unique receptor and erbB4
Cripto-3	?

to the luminal ductal epithelium (Snedeker *et al.*, 1991). In the absence of ovarian steroids, exogenous TGF α and EGF can stimulate ductal growth of the mouse mammary epithelium suggesting that they can act independently of secondary signals (Snedeker *et al.*, 1991). EGF and TGF α mRNA is present in pregnant and lactating rat and human mammary glands and increases 2–3-fold over virgin levels at pregnancy (Liscia *et al.*, 1990). The EGFR is required for ductal development (Wiesen *et al.*, 1999) and can be detected in both the stromal and epithelial components of the virgin mammary gland (Coleman *et al.*, 1988). In the pregnant gland, a rapid increase is observed during midpregnancy precisely at the time of extensive cellular proliferation (Edery *et al.*, 1985). After pregnancy, the levels of EGFR decrease significantly. The coincident expression of TGF α and its receptor with the proliferative phases of mammary epithelial growth and TGF α 's direct mitogenic effect on epithelium *in vitro* and *in vivo* confirms the functional role for this signaling dyad in the development and proliferation of the mammary epithelium.

TGF α expression is associated with human breast cancer

TGF α is expressed and mitotically active in numerous breast cancer cell lines and has been directly implicated as a modulator of transformation *in vivo* (Borellini and Oka, 1989; Daniel and Silberstein, 1985; de Jong *et al.*, 1998a; Salomon *et al.*, 1984; Valverius *et al.*, 1989). In fact, the discovery of TGF α was based on its ability to transform retrovirally-infected cultured fibroblasts (Todaro *et al.*, 1980; de Larco and Todaro, 1978). Expression of TGF α has been identified in pleural effusions from normal mammary gland (Arteaga *et al.*, 1988), in invasive ductal carcinoma (Pilichowska *et al.*, 1997) and correlates with increased neo-angiogenesis (de Jong *et al.*, 1998b) in breast tumors. Carcinomas of the breast that have higher level expression of TGF α also express high levels of EGFR, implicating a functional role for the TGF α /EGFR autocrine loop in tumors (Umekita *et al.*, 1992). The correlation between levels of EGFR expression and neoplastic transformation *in vivo* is controversial (Gullick and Srinivasan, 1998). Robert-

son and colleagues demonstrated that 40–60% of neoplasias examined in one study displayed normal levels of EGFR expression (Robertson *et al.*, 1996). Only a small proportion of breast cancers has elevated levels of EGFR (Slamon *et al.*, 1987), or amplification of EGFR (Peters and Wolff, 1983). This data implies that alterations in EGFR ligands may be more influential on the activity of the EGFR than changes in the levels of receptor expression itself. Alternatively, the formation of heterodimers between EGFR and the other erbB family members may play a role in effecting differential activities in tumorigenesis of the breast.

TGF α can modulate cellular transformation in breast cancer cell lines. In the immortalized human breast cancer cell line, MCF-7, TGF α transfected cells that expressed EGFR were stimulated to grow in a colony forming assay (Ciardiello *et al.*, 1990). In a parallel study with fully transformed MCF-7 cells that lacked EGFR, TGF α transfection did not effect a positive growth advantage (Clarke *et al.*, 1989). Overexpression of TGF α in the immortalized mouse mammary cell line, NOG-8, generated anchorage independent growth but colonies failed to form tumors in nude mice (Shankar *et al.*, 1989). Therefore, expression of TGF α is associated with the ability to stimulate growth, while overexpression is allied with partial transformation *in vitro* and correlated with appearance of carcinomas *in vivo*.

TGF α transgenic mouse models of breast cancer

Mouse models have been developed to study the role of TGF α in the transformation and development of the mammary gland. Several promoters have been utilized to target TGF α both non-specifically and specifically to the mammary gland. These mouse models have established the importance of TGF α in the early stages of neoplastic development of the mammary gland.

Initial studies demonstrating the neoplasia-promoting activity of TGF α were described in mice expressing human TGF α under the control of the zinc-inducible metallothionein (MT) promoter (Jhappan *et al.*, 1990). Despite the fact that expression of the transgene was low in the mammary gland and was generally not inducible, these mice displayed increased cellular proliferation and delayed epithelial penetration of the stromal fatpad during ductal development. In a second parallel study, using the MT promoter directing the expression of rat TGF α , Sandgren and colleagues noticed that mice that had passed through multiple pregnancies developed hyperplastic nodules and dysplasia of the mammary epithelium (Sandgren *et al.*, 1990). Only one mouse developed a secretory adenocarcinoma. These studies demonstrated that there was an association between the *in vivo* expression of TGF α and the development of mammary hyperplasia.

A study directing TGF α expression specifically to the mammary gland with the mouse mammary tumor virus (MMTV) promoter demonstrated unequivocally the growth and neoplasia promoting activity of TGF α (Matsui *et al.*, 1990). Precocious alveolar development, hyperproliferation and hyperplasias were apparent in the mature virgin mouse. The alveolar hyperplasia was present in glands from mature virgin mice, but not in

immature virgin mice, suggesting a requirement for secondary inputs from systemic hormones for any TGF α -dependent hyperplasia to occur. Hyperplastic alveolar nodules and cysts were prominent in the multiparous animal and increased in hyperplastic and dysplastic character with the number of pregnancies. One multiparous mouse developed adenocarcinomas although none were metastatic. A second study was performed with the MMTV-TGF α mice that studied in greater detail the generation of hyperplasia and tumors (Halter *et al.*, 1992). At one year, 65% of multiparous and 45% of virgin mice displayed hyperplasia. By 16 months, 40% of the multiparous and 30% of the virgin mice and mice had generated tumors. These studies demonstrated that TGF α could stimulate the proliferation of the mammary epithelium and generate proliferative late pregnancy mammary gland phenotypes in early pregnant animals. In addition, the hyperplasia, dysplasia and frank carcinoma observed with this mouse model suggested that there was a multistage nature to the progression of TGF α -initiated transformation.

A more dramatic mammary phenotype was achieved by targeting rat TGF α exclusively to the mammary gland with the WAP-TGF α transgenic mouse (Sandgren *et al.*, 1995). This transgenic model achieved high level TGF α expression during pregnancy and lactation and displayed similar proliferative mammary gland phenotypes as described in the MT and MMTV transgenic models but with an increased incidence

and decreased latency of tumors. In addition, the process of involution, in which the gland absorbs and reorganizes a significant portion of its epithelial structure, was delayed in these mice as it was in the original MT mice (Sandgren *et al.*, 1990). Latency of tumor appearance was decreased in comparison to the MMTV-TGF α model, but still required several rounds of pregnancy for initial tumor development. A majority of the tumors were well-differentiated glandular adenomas or carcinomas (53%) and fibroadenomas (35%) and retained the glandular characteristics of late pregnant mammary epithelium. The appearance of multiple types of tumor and the retention of normal glandular morphologies suggested that these different types were stages of TGF α -dependent tumor progression. Interestingly, the levels of cyclinD1 were elevated in the WAP-TGF α transgenic glands. The targeted overexpression of this cell cycle regulatory molecule is known to effect mammary transformation (Wang *et al.*, 1994) and is in a chromosomal region frequently rearranged in human breast cancer (Dickson *et al.*, 1995; Gillett *et al.*, 1994, 1996, 1999). This indicates that cyclinD1 may be a cooperative factor in the development of TGF α induced mammary carcinogenesis. The delay in involution was suggested as a mechanism for promoting transformation of the mammary gland as it provided an expanded population of proliferative epithelial cells that could be predisposed to transformation. This seminal paper demonstrated TGF α could transform the mammary




Figure 1 Involution is enhanced in the absence of Stat5a. Hemotoxylin and eosin staining of inguinal mammary glands from Stat5a null non-transgenic (NTGKO), wildtype (WT), Stat5a null TGF α transgenic (KOTG) and TGF α transgenic (TG) mice at 18 days of pregnancy (p18/a-d), days 1 (i1/e-h), 3 (i3/i-l) and 7 (i7/m-p) of involution. Samples were collected from mice in or immediately after their first pregnancy. Compare epithelial condensation at day 3 and 7 of involution in the wildtype (i-j) versus the KOTG (i-k) and TG (i-l). Magnification is defined by the bar in a = 200 μ m. Reproduced with permission (Humphreys and Hennighausen, 1999)

epithelium and suggested a possible mechanism for neoplastic development. The observation that TGF α could cause a delay in involution was supported by results from a separate study with MT-TGF α mice (Smith *et al.*, 1995). An increase in DNA synthesis in lactation was accompanied by a significant decrease in apoptotic cells after 2 days of involution. Involution was also affected in a mouse model utilizing WAP-TGF α and the Stat5a knockout mouse (Humphreys and Hennighausen, 1999). In the absence of Stat5a, apoptosis levels rose in the mammary gland during pregnancy and involution. This increase in apoptosis enhanced involution of the WAP-TGF α gland (Figure 1) and increased the latency of WAP-TGF α -induced tumors. This result demonstrated the WAP-TGF α -induced delay in involution and tumor formation, could be abrogated by a downstream signaling molecule that regulates cell death in the mammary epithelium.

Subsequent studies examined the role that oncogenes may play in acting cooperatively with TGF α to promote mammary transformation. The oncogene *c-myc* is overexpressed in 25–30% of breast cancer (Bonilla *et al.*, 1988; Callahan and Campbell, 1989; Mariani-Costantini *et al.*, 1988, 1989; Morse *et al.*,

1988) and in rodent mammary epithelial cell lines transfected with *myc*, TGF α could cooperate to support a transformed phenotype (Telang *et al.*, 1990). Mice that overexpressed both *c-myc* and TGF α in the mammary gland had an increased tumor incidence and decreased tumor latency when compared to *c-myc* transgenics (Amundadottir *et al.*, 1995). A second paper utilizing the same bitransgenic mice demonstrated that the synergism between these two proteins was due to a cooperative growth stimulus and inhibition of *c-myc*-induced apoptosis by TGF α (Amundadottir *et al.*, 1996). Apoptotic tumors were present exclusively in mice that expressed only *c-myc*. In cell lines derived from these tumors, exogenous TGF α could inhibit apoptosis. These data again implicated TGF α as a survival factor in the mammary epithelium. Interestingly, tumor cells from the mammary glands of these bitransgenic mice could only become apoptotic when exposed to a specific inhibitor of the EGFR kinase pathway. This result suggested that an intact TGF α /EGFR autocrine loop was required to mediate the survival effects of TGF α .

Interaction between TGF α and the proto-oncogene *neu* was examined in bitransgenic mice expressing both




Figure 2 Model of the TGF α /EGF signal transduction pathway and EGFR signaling inhibitors that affect EGFR-mediated transformation. EGF and TGF α binding to the EGFR stimulate the activation of the endogenous receptor tyrosine kinase. The membrane-bound EGFR kinase activates one of several intracellular signal transduction pathways including the protein kinase A, Ras/Raf/MAPK, *c-src* and Jak/Stat pathways. Direct or indirect phosphorylation and activation of one or more of the Stat proteins, 1, 3, 5a, and 5b can be achieved through several of these mechanisms. The activating kinase, attributes of the targeted Stat and phosphorylated residue can elicit distinct functional consequences for the cell. Multiple intercellular pathways can lead to activation of the Stats and other nuclear factors like *myc* and cyclinD1. Each of these pathways can be blocked by specific inhibitors (shown in red italics). The proliferative stimulus provided by TGF α and EGF in cooperation with transforming oncogenes can lead to cellular transformation. Consequently, a clear understanding of the mechanisms involved and the use of multifocal inhibitors of signaling intermediaries is critical for effectual inhibition of EGFR signal transduction. Some alternative and intermediary intercellular signaling molecules have been omitted for clarity

genes under the control of the MMTV promoter (Muller *et al.*, 1996). The *neu* proto-oncogene is a member of the EGFR family and although it does not bind to TGF α or EGF, it can form heterodimers with the EGFR and it is often found overexpressed and amplified in human breast cancers (Slamon *et al.*, 1987). Tumor latency was decreased in the bitransgenic mice, compared to the TGF α and *neu*, mono-transgenic lines. At 150 days 95% of the bitransgenic mice had mammary tumors vs 6% and 35% for the TGF α and *neu* mice, respectively. Bitransgenic tumors were multifocal whereas mono-transgenic lines generated focal tumors and contained activated *neu*. This paper suggested a novel mechanism for TGF α cooperativity involving the transactivation of *neu* through the EGFR. Treatment of MMTV-TGF α mice with the tumor promoter 7, 12 dimethyl benzanthrane demonstrated that TGF α could accelerate tumor formation and the authors suggested that TGF α could act as a tumor promoter (Coffey *et al.*, 1994).

These studies on the mechanism of TGF α -mediated tumorigenesis and cooperativity in the mammary gland demonstrate that the overexpression of TGF α gives the preneoplastic mammary epithelium a proliferative advantage but in and of itself is not a transforming event. Secondary events, like the activation of proto-oncogenes, can significantly increase the efficiency with which TGF α can transform the mammary epithelium. Importantly, these studies support the theory that TGF α 's influence on the regulation of apoptosis in the mammary gland is a possible mechanism of promoting survival of the neoplastic cell.

Disruption of TGF α signaling and potential therapies for breast cancer

The EGFR is required for mammary gland ductal development (Wiesen *et al.*, 1999; Xie *et al.*, 1997) and, as described previously, is aberrantly expressed in 40% of human breast cancers and occasionally overexpressed in those tumors with poor prognosis. This receptor can interact with several distinct ligands (Table 1) and activates different intracellular and nuclear signaling pathways (Figure 2). Substrates for this type 1 receptor tyrosine kinase are numerous and demonstrate the influence of EGFR on the regulation of cellular growth. Importantly, it is clear that this receptor plays a central role in the development and progression of human breast cancer. Therefore, insight into EGFR signal transduction and the mechanism of substrate selection and activation are critical to understanding the development of breast cancer. The EGFR and its interaction with TGF α has been and continues to be a target for potential anti-cancer therapies that aim to regulate its activity and signaling mechanisms.

A recent study demonstrated that inhibition of the signal from EGFR to its intracellular signaling molecules is critical in blocking tumor progression. Treatment of MMTV-TGF α transgenic mice with a farnesyl transferase inhibitor, which blocks Ras function, demonstrated a significant regression of mammary tumors (Norgaard *et al.*, 1999). This inhibitory effect was ineffectual after tumor accelera-

tion from multipregnancy or after treatment with the carcinogen DMBA supporting the theory that that EGFR signaling is important in the initial stage of neoplastic development. The EGFR can activate the intracellular tyrosine kinases Jak1 and the transcription factors Stat1, 3 and 5b in response to growth signaling from EGF (Leonard and O'Shea, 1998). Recent data has demonstrated that growth hormone activated Ras/Raf/MAPK pathway can directly activate the normally cytokine-activated pathway of Stat5a (Pircher *et al.*, 1999) and EGF can phosphorylate Stat5a through *c-src* (Olayioye *et al.*, 1999). Stat5a, a prolactin-activated transcription factor, has an established role in the development and differentiation of the lactation-competent mammary gland (Liu *et al.*, 1987) but is also associated with signaling in tumor cells (Hayakawa *et al.*, 1998; Richer *et al.*, 1998; Yu *et al.*, 1997; Zhang *et al.*, 1996). Stat5a-null mice interbred with WAP-TGF α transgenic mice revealed inhibition of TGF α -dependent MAPK activation and TGF α -dependent apoptosis inhibition after deletion of Stat5a. The increase in apoptosis permitted a more complete epithelial regression to occur at involution (Figure 1). A more complete regression deleted a significant number of potentially neoplastic cells and this impacted WAP-TGF α driven tumorigenesis. Stat5a null/WAP TGF α transgenic mice had an increase in tumor latency when compared to the WAP-TGF α transgenic. These mice displayed an increase in apoptosis before and during involution. These data suggested Stat5a was acting as a survival factor for the mammary epithelium by blocking the onset of apoptosis. Theoretically, the absence of Stat5a permitted apoptosis to occur and thereby diminished the pool of potential neoplastic cells that could become transformed (Humphreys and Hennighausen, 1999).

A novel mechanism of inhibition of TGF α activity was demonstrated recently in human mammary epithelial cells with an anti-metalloproteinase. Metalloproteinases are extracellular enzymes that cleave components of the extracellular matrix including the EGFR ligands; TGF α and EGF. This cleavage event releases them from the cell surface rendering the growth factor into an active form. These metalloproteinase inhibitors prevented TGF α release from the cell surface, blocked cell migration and decreased proliferation. Additionally, the metalloproteinase inhibitors reduced the growth of EGF-dependent tumor cell lines and could synergize with anti-EGFR antibodies (Dong *et al.*, 1999).

Therapies for inhibiting the action of EGFR and indirectly the action of TGF α , to block tumorigenesis are being explored. Current research into the action of molecules that may have a role in regulating the TGF α signaling pathway like EGFR-specific monoclonal antibodies (Ciardiello *et al.*, 1999), protein kinase A inhibitors, a combination of antibodies and chemotherapeutic agents (Bianco *et al.*, 1997; Ciardiello *et al.*, 1996; Ciardiello and Tortora, 1998), anti-estrogens like taxol, raloxifen, alone or in combination with antibodies like the humanized monoclonal antibody to *neu*; Herceptin (Brenner and Adams, 1999; Hanna *et al.*, 1999; Robertson, 1998; Ross and Fletcher, 1998), provide some promising results for future treatment of breast cancer.

Conclusions

Transgenic mouse model studies have given us insight into possible mechanism of TGF α -initiated breast cancer. Overexpression of TGF α may be an early prognostic factor in the initiation of this disease and in cooperation with secondary transforming events can lead to carcinoma. Importantly, the role of TGF α in the inhibition of apoptosis during normal development could lead to a critical understanding of the role this growth factor plays in normal and neoplastic development in the mammary gland. This growth factor pathway appears to be able to promote growth in the initial phase of neoplastic transformation although it

lacks the dominating transformation ability observed with some of the other EGFR-related mouse models of mammary gland disease like MMTV/erb-2 (Guy *et al.*, 1992; Siegel *et al.*, 1999). In combination with an understanding of TGF α signaling mechanisms, inhibitory agents, which act on the receptor-ligand interaction or on the intracellular kinases stimulated by the EGFR, could lead to practical and efficacious treatment regimens for this disease.

Acknowledgements

This work is supported by a grant from the Breast Cancer Research Program from USMRMC DAMD 17-99-1-9328.

References

- Amundadottir LT, Johnson MD, Merlino G, Smith GH and Dickson RB. (1995). *Cell Growth Differ.*, **6**, 737–748.
- Amundadottir LT, Nass SJ, Berchem GJ, Johnson MD and Dickson RB. (1996). *Oncogene*, **13**, 757–765.
- Arteaga CL, Hanauske AR, Clark GM, Osborne CK, Hazarika P, Pardue RL, Tio F and Von Hoff DD. (1988). *Cancer Res.*, **48**, 5023–5028.
- Bates SE, Valverius EM, Ennis BW, Bronzert DA, Sheridan JP, Stampfer MR, Mendelsohn J, Lippman ME and Dickson RB. (1990). *Endocrinology*, **126**, 596–607.
- Bianco C, Tortora G, Baldassarre G, Caputo R, Fontanini G, Chine S, Bianco AR and Ciardiello F. (1997). *Clin. Cancer Res.*, **3**, 439–448.
- Bonilla M, Ramirez M, Lopez-Cueto J and Gariglio P. (1988). *J. Natl. Cancer Inst.*, **80**, 665–671.
- Borellini F and Oka T. (1989). *Environ. Health Perspect.* **80**, 85–99.
- Brenner TL and Adams VR. (1999). *J. Am. Pharm. Assoc. (Wash.)*, **39**, 236–238.
- Callahan R and Campbell G. (1989). *J. Natl. Cancer Inst.*, **81**, 1780–1786.
- Ciardiello F, Bianco R, Damiano V, De Lorenzo S, Pepe S, De Placido S, Fan Z, Mendelsohn J, Bianco AR and Tortora G. (1999). *Clin. Cancer Res.*, **5**, 909–916.
- Ciardiello F, Damiano V, Bianco R, Bianco C, Fontanini G, De Laurentiis M, De Placido S, Mendelsohn J, Bianco AR and Tortora G. (1996). *J. Natl. Cancer Inst.*, **88**, 1770–1776.
- Ciardiello F, McGeady ML, Kim N, Basolo F, Hynes N, Langton BC, Yokozaki H, Saeki T, Elliott JW, Masui H and *et al.* (1990). *Cell Growth Differ.*, **1**, 407–420.
- Ciardiello F and Tortora G. (1998). *Clin. Cancer Res.*, **4**, 821–828.
- Clarke R, Brunner N, Katz D, Glanz P, Dickson RB, Lippman ME and Kern FG. (1989). *Mol. Endocrinol.*, **3**, 372–380.
- Coffey Jr, RJ, Meise KS, Matsui Y, Hogan BL, Dempsey PJ and Halter SA. (1994). *Cancer Res.*, **54**, 1678–1683.
- Coleman S, Silberstein GB and Daniel CW. (1988). *Dev. Biol.*, **127**, 304–315.
- Daniel CW and Silberstein GB. (1985). *The Mammary Gland: Development, Regulation and Function*, Vol 1. Neville MC and Daniel CW. (eds). Plenum Publishing: New York, pp. 3–36.
- de Jong JS, van Diest PJ, van der Valk P and Baak JP. (1998a). *J. Pathol.*, **184**, 44–52.
- de Jong JS, van Diest PJ, van der Valk P and Baak JP. (1998b). *J. Pathol.*, **184**, 53–57.
- de Larco JE and Todaro GJ. (1978). *Proc. Natl. Acad. Sci. USA*, **75**, 4001–4005.
- Dickson C, Fantl V, Gillett C, Brookes S, Bartek J, Smith R, Fisher C, Barnes D and Peters G. (1995). *Cancer Lett.*, **90**, 43–50.
- Dickson RB and Lippman ME. (1995). *Endocr. Rev.*, **16**, 559–589.
- Dong J, Opresko LK, Dempsey PJ, Lauffenburger DA, Coffey RJ and Wiley HS. (1999). *Proc. Natl. Acad. Sci. USA*, **96**, 6235–6240.
- Edery M, Pang K, Larson L, Colosi T and Nandi S. (1985). *Endocrinology*, **117**, 405–411.
- Gillett C, Fantl V, Smith R, Fisher C, Bartek J, Dickson C, Barnes D and Peters G. (1994). *Cancer Res.*, **54**, 1812–1817.
- Gillett C, Smith P, Gregory W, Richards M, Millis R, Peters G and Barnes D. (1996). *Int. J. Cancer*, **69**, 92–99.
- Gillett CE, Smith P, Peters G, Lu X and Barnes DM. (1999). *J. Pathol.*, **187**, 200–206.
- Gullick WJ, Bianco C, Normanno N, Martinez-Lacaci I, De Santis M, Ebert AD and Salomon DS. (1999). *Women Cancer*, **1**, 29–57.
- Gullick WJ and Srinivasan R. (1998). *Breast Cancer Res. Treat.*, **52**, 43–53.
- Guy CT, Webster MA, Schaller M, Parsons TJ, Cardiff RD and Muller WJ. (1992). *Proc. Natl. Acad. Sci. USA*, **89**, 10578–10582.
- Halter SA, Dempsey P, Matsui Y, Stokes MK, Graves-Deal R, Hogan BL and Coffey RJ. (1992). *Am. J. Pathol.*, **140**, 1131–1146.
- Hanna W, Kahn HJ and Trudeau M. (1999). *Mod. Pathol.*, **12**, 827–834.
- Hayakawa F, Towatari M, Iida H, Wakao H, Kiyoi H, Naoe T and Saito H. (1998). *Br. J. Haematol.*, **101**, 521–528.
- Humphreys RC and Hennighausen L. (1999). *Cell Growth Differ.*, **10**, 685–694.
- Jhappan C, Stahle C, Harkins RN, Fausto N, Smith GH and Merlino GT. (1990). *Cell*, **61**, 1137–1146.
- Kenney NJ, Saeki T, Gottardis M, Kim N, Garcia-Morales P, Martin MB, Normanno N, Ciardiello F, Day A, Cutler ML and *et al.* (1993). *J. Cell Physiol.*, **156**, 497–514.
- Leonard WJ and O'Shea JJ. (1998). *Annu. Rev. Immunol.*, **16**, 293–322.
- Liscia DS, Merlo G, Ciardiello F, Kim N, Smith GH, Callahan R and Salomon DS. (1990). *Dev. Biol.*, **140**, 123–131.
- Liu SC, Sanfilippo B, Perroteau I, Derynck R, Salomon DS and Kidwell WR. (1987). *Mol. Endocrinol.*, **1**, 683–692.
- Mariani-Costantini R, Escot C, Theillet C, Gentile A, Merlo G, Lidereau R and Callahan R. (1988). *Cancer Res.*, **48**, 199–205.
- Mariani-Costantini R, Merlo G and Frati L. (1989). *Tumori*, **75**, 311–320.
- Matsui Y, Halter SA, Holt JT, Hogan BL and Coffey RJ. (1990). *Cell*, **61**, 1147–1155.
- Morse B, Rotherg PG, South VJ, Spandorfer JM and Astrin SM. (1988). *Nature*, **333**, 87–90.

- Muller WJ, Arteaga CL, Muthuswamy SK, Siegel PM, Webster MA, Cardiff RD, Meise KS, Li F, Halter SA and Coffey RJ. (1996). *Mol. Cell. Biol.*, **16**, 5726–5736.
- Norgaard P, Law B, Joseph H, Page DL, Shyr Y, Mays D, Pietenpol JA, Kohl NE, Oliff A, Coffey Jr, RJ., Poulsen HS and Moses HL. (1999). *Clin. Cancer Res.*, **5**, 35–42.
- Olayioye MA, Beuvink I, Horsch K, Daly JM and Hynes NE. (1999). *J. Biol. Chem.*, **274**, 17209–17218.
- Peters GN and Wolff M. (1983). *Cancer*, **52**, 680–686.
- Pilichowska M, Kimura N, Fujiwara H and Nagura H. (1997). *Mod. Pathol.*, **10**, 969–975.
- Pircher TJ, Petersen H, Gustafsson JA and Haldosen LA. (1999). *Mol. Endocrinol.*, **13**, 555–565.
- Richer JK, Lange CA, Manning NG, Owen G, Powell R and Horwitz KB. (1998). *J. Biol. Chem.*, **273**, 31317–31326.
- Robertson D. (1998). *Nat. Biotechnol.*, **16**, 615.
- Robertson KW, Reeves JR, Smith G, Keith WN, Ozanne BW, Cooke TG and Stanton PD. (1996). *Cancer Res.*, **56**, 3823–3830.
- Ross JS and Fletcher JA. (1998). *Oncologist*, **3**, 237–252.
- Salomon DS, Perroteau I, Kidwell WR, Tam J and Derynck R. (1987). *J. Cell Physiol.*, **130**, 397–409.
- Salomon DS, Zwiebel JA, Bano M, Losonczy I, Fehnel P and Kidwell WR. (1984). *Cancer Res.*, **44**, 4069–4077.
- Sandgren EP, Luetke NC, Palmiter RD, Brinster RL and Lee DC. (1990). *Cell*, **61**, 1121–1135.
- Sandgren EP, Schroeder JA, Qui TH, Palmiter RD, Brinster RL and Lee DC. (1995). *Cancer Res.*, **55**, 3915–3927.
- Schroeder JA and Lee DC. (1997). *J. Mammary Gland Biol., Neoplasia*, **2**, 119–130.
- Shankar V, Ciardiello F, Kim N, Derynck R, Liscia DS, Merlo G, Langton BC, Sheer D, Callahan R, Bassin RH *et al.* (1989). *Mol. Carcinog.*, **2**, 1–11.
- Siegel PM, Ryan ED, Cardiff RD and Muller WJ. (1999). *EMBO J.*, **18**, 2149–2164.
- Slamon DJ, Clark GM, Wong SG, Levin WJ, Ullrich A and McGuire WL. (1987). *Science*, **235**, 177–182.
- Smith GH, Sharp R, Kordon EC, Jhappan C and Merlino G. (1995). *Am. J. Pathol.*, **147**, 1081–1096.
- Smith JA, Barraclough R, Fernig DG and Rudland PS. (1989). *J. Cell Physiol.*, **141**, 362–370.
- Snedeker SM, Brown CF and DiAugustine RP. (1991). *Proc. Natl. Acad. Sci. USA*, **88**, 276–280.
- Telang NT, Osborne MP, Sweterlitsch LA and Narayanan R. (1990). *Cell Regul.*, **1**, 863–872.
- Todaro GJ, Callahan R, Rapp UR and De Larco JE. (1980). *Proc. R. Soc. Lond. B. Biol. Sci.*, **210**, 367–385.
- Umekita Y, Enokizono N, Sagara Y, Kuriwaki K, Takasaki T, Yoshida A and Yoshida H. (1992). *Virchows Arch. A. Pathol. Anat. Histopathol.*, **420**, 345–351.
- Valverius EM, Bates SE, Stampfer MR, Clark R, McCormick F, Salomon DS, Lippman ME and Dickson RB. (1989). *Mol. Endocrinol.*, **3**, 203–214.
- Vonderhaar BK. (1987). *J. Cell. Physiol.*, **132**, 581–584.
- Wang TC, Cardiff RD, Zukerberg L, Lees E, Arnold A and Schmidt EV. (1994). *Nature*, **369**, 669–671.
- Wiesen JF, Young P, Werb Z and Cunha GR. (1999). *Development*, **126**, 335–344.
- Xie W, Paterson AJ, Chin E, Nabell LM and Kudlow JE. (1997). *Mol. Endocrinol.*, **11**, 1766–1781.
- Yu CL, Jove R and Burakoff SJ. (1997). *J. Immunol.*, **159**, 5206–5210.
- Zaichowski DA and Sager R. (1991). *Mol. Endocrinol.*, **5**, 1613–1623.
- Zhang Q, Nowak I, Vonderheid EC, Rook AH, Kadin ME, Nowell PC, Shaw LM and Wasik MA. (1996). *Proc. Natl. Acad. Sci. USA*, **93**, 9148–9153.