



Long Term Resource Monitoring Program

# Program Report 2001-P001

---

## 1998 Annual Status Report

*Submersed and Floating-leaf Vegetation  
in Pools 4, 8, 13, and 26 and La Grange Pool  
of the Upper Mississippi River System*



May 2001

20010718 069

---

**1998 Annual Status Report**  
*Submersed and Rooted Floating-leaf Vegetation  
in Pools 4, 8, 13, and 26 and La Grange Pool  
of the Upper Mississippi River System*

by

Yao Yin, Heidi Langrehr, Theresa Blackburn, Megan Moore,  
Jenny Winkelman, Robert Cosgriff, and Thad Cook

May 2001

U.S. Geological Survey  
Upper Midwest Environmental Sciences Center  
2630 Fanta Reed Road  
La Crosse, Wisconsin 54603

## Contents

	<i>Page</i>
Preface .....	v
Abstract .....	1
Introduction .....	2
Study Areas .....	2
Methods .....	4
Stratification .....	4
Site Selection .....	5
Equipment and Definitions .....	6
Site and Subsampling Areas .....	7
Navigation to Sampling Sites .....	7
Sampling and Data Recording .....	7
Computation of Summary Indexes .....	8
Frequency .....	8
Abundance Index .....	8
Percent Cover (Rooted floating-leaf life form) .....	9
References .....	9
Appendix .....	A-1
Chapter 1. Results in Pool 4, Upper Mississippi River .....	1.1
Chapter 2. Results in Pool 8, Upper Mississippi River .....	2.1
Chapter 3. Results in Pool 13, Upper Mississippi River .....	3.1
Chapter 4. Results in Pool 26, Upper Mississippi River .....	4.1
Chapter 5. Results in La Grange Pool, Illinois River .....	5.1

## Preface

The Long Term Resource Monitoring Program (LTRMP) was authorized under the Water Resources Development Act of 1986 (Public Law 99-662) as an element of the U.S. Army Corps of Engineers' Environmental Management Program. The LTRMP is being implemented by the Upper Midwest Environmental Sciences Center, a U.S. Geological Survey science center, in cooperation with the five Upper Mississippi River System (UMRS) States of Illinois, Iowa, Minnesota, Missouri, and Wisconsin. The U.S. Army Corps of Engineers provides guidance and has overall Program responsibility. The mode of operation and respective roles of the agencies are outlined in a 1988 Memorandum of Agreement.

The UMRS encompasses the commercially navigable reaches of the Upper Mississippi River, as well as the Illinois River and navigable portions of the Kaskaskia, Black, St. Croix, and Minnesota Rivers. Congress has declared the UMRS to be both a nationally significant ecosystem and a nationally significant commercial navigation system. The mission of the LTRMP is to provide decision makers with information for maintaining the UMRS as a sustainable large river ecosystem given its multiple-use character. The long-term goals of the Program are to understand the system, determine resource trends and effects, develop management alternatives, manage information, and develop useful products.

This report presents the results of aquatic vegetation stratified random sampling surveys conducted in 1998 by field station personnel under the direction of the UMESC. Pools 4, 8, 13, and 26 of the Upper Mississippi River and La Grange Pool of the Illinois River were surveyed. This report satisfies, for 1998, Task 2.2.4.6, *Evaluate and Summarize Annual Present-day Results* under Goal 2, *Monitor Resource Change* of the Operating Plan (U.S. Fish and Wildlife Service 1993). The purpose of this report is to provide a summary of data regarding the distribution and abundance of submersed and floating-leaf vegetation collected from the field stations for 1998. This report was developed with funding provided by the Long Term Resource Monitoring Program.



# United States Department of the Interior

U.S. GEOLOGICAL SURVEY

Upper Midwest Environmental Sciences Center  
2630 Fanta Reed Road  
La Crosse, Wisconsin 54603

July 9, 2001

Author

Upper Mississippi River Basin Association  
Environmental Management Program Coordinating Committee  
Long Term Resource Monitoring Program (LTRMP) Analysis Team  
Mississippi River Advisory Group  
Science Review Committee  
U.S. Geological Survey Science Centers and Regional Offices  
Minnesota/Wisconsin Boundary Area Commission  
U.S. Army Corps of Engineers

Dear LTRMP Participant:

I am enclosing "1998 Annual Status Report: Submersed and floating-leaf vegetation in Pools 4, 8, 13, and 26 and La Grange Pool of the Upper Mississippi River System" by Yao Yin, Heidi Langrehr, Theresa Blackburn, Megan Moore, Jenny Winkelman, Robert Cosgriff, and Thad Cook. This report may also be found on our web site at

[http://www.umesc.usgs.gov/reports\\_publications/ltrmp\\_rep\\_list.html](http://www.umesc.usgs.gov/reports_publications/ltrmp_rep_list.html)

in the form of a PDF file. We recommend using Adobe Acrobat Reader 3.0 to view the document.

Please contact me at 608-781-6221, if you have questions regarding this report.

Sincerely,

Leslie E. Holland-Bartels  
Center Director

Enclosure:

LTRMP 2001-P001

**1998 Annual Status Report**  
*Submersed and Rooted Floating-Leaf Vegetation  
in Pools 4, 8, 13, and 26 and La Grange Pool  
of the Upper Mississippi River System*

by

Yao Yin

*U.S. Geological Survey, Upper Midwest Environmental Sciences Center  
2630 Fanta Reed Road, La Crosse, Wisconsin 54603*

Heidi Langrehr

*Wisconsin Department of Natural Resources, Onalaska Field Station  
575 Lester Avenue, Onalaska, Wisconsin 54650*

Theresa Blackburn

*Iowa Department of Natural Resources, Mississippi River Monitoring Station  
206 Rose Street, Bellevue, Iowa 52031*

Megan Moore and Jenny Winkelman

*Minnesota Department of Natural Resources, Lake City Field Station  
1801 South Oak Street, Lake City, Minnesota 55041*

Robert Cosgriff

*Illinois Natural History Survey, Alton Field Station  
8450 Montclair Avenue, Brighton, Illinois 62012*

Thad Cook

*Illinois Natural History Survey, Havana Field Station  
704 North Schrader Avenue, Havana, Illinois 62644*

**Abstract:** Aquatic vegetation was investigated in five navigation pools in the Upper Mississippi River System using a new protocol named "stratified random sampling" or SRS protocol for the first time in 1998. The five pools were Pools 4, 8, 13, and 26 of the Upper Mississippi River and La Grange Pool of the Illinois River. The results on submersed aquatic vegetation and rooted floating-leaf aquatic vegetation were summarized in this report. The percent frequencies of submersed aquatic vegetation in shallow water areas ( $\leq 3$  m deep at flat-pool condition) in the five pools were 36.6%, 47.6%, 42%, 6.1%, and 0%, respectively. The aquatic area strata that were directly influenced by the flow in the main channel, such as the main channel borders and secondary channels, had lower percent frequencies of submersed aquatic vegetation than the aquatic area strata that were less directly influenced by the flow in the main channel, such as the contiguous and isolated backwaters. The percent covers of rooted floating-leaf vegetation were 4.1%, 7.5%, 6.5%, 0.9%, and 0%. The majority of aquatic vegetation that was recorded in Pool 26 was from one isolated backwater area. Aquatic vegetation was not recorded at any of the sampling sites in La Grange Pool.

**Key words:** Annual report, aquatic, floating-leaf, Illinois River, La Grange, LTRMP, Mississippi River, submersed vegetation.

---




Figure 1. Sampling sites and aquatic area strata in Pool 4, Upper Mississippi River System, 1998.




Figure 2. Sampling sites and aquatic area strata in Pool 8, Upper Mississippi River System, 1998.

of aquatic habitat. Similar to pools upstream, Pool 13 contains many high bank islands adjacent to the main channel in the upper section, braided backwater channels and sloughs in the middle section, and a large open lake-like area in the lower section of the pool. Major tributaries include the Apple and Plum Rivers on the Illinois side and Maquoketa and Elk Rivers on the Iowa side. The shallow water areas are divided into five strata (Table 1; Figure 3) for sampling and analysis.

The Pool 26 study area includes water bodies along the Upper Mississippi River from Lock and Dam 25 (Winfield, Missouri) to Lock and Dam 26 (Alton, Illinois) and the lower Illinois River from its confluence with the Mississippi River north to Illinois River (river mile 12). This reach of the two rivers is bordered by high bluffs on the Illinois side and low elevation floodplain on the Missouri side. The reach encompasses 9,500 ha (23,700 acres) of aquatic habitat. Presently, most of the backwaters of the lower Illinois River are isolated from the river by low levees so as to decrease sedimentation and allow management for waterfowl. Likewise, many of the secondary channels of the Mississippi River are isolated from the river on the up-stream side to create backwaters




Figure 4. Sampling sites and aquatic area strata in Pool 26, Upper Mississippi River System, 1998.

support aquatic vegetation. Shallow water areas were divided into general habitat types (strata), including main channel borders, secondary channels, contiguous backwaters, isolated backwaters, and impounded areas. Sampling efforts were generally proportional to acreage and perceived habitat heterogeneity (Table 1) of each stratum, except for the isolated backwater areas whose sampling sizes were kept small to ensure a timely completion of the investigation. Some areas were excluded from the

sampling sites because of safety concerns and accessibility difficulties.

### Site Selection

Sites to be investigated were selected from a computer program using a random number generator. A 50- × 50-m grid was generated and overlaid onto the stratified aquatic areas. Nodes of the grid were geo-spatially registered (Universal Transverse

about 3 m long, with a rope extension, and is scaled at 10-cm increments. Aquatic vegetation or aquatic species refer to the following plant types/life forms: submersed and rooted floating-leaf.

### Site and Subsampling Areas

Most sites were investigated with the field crew standing in a 5-m (16-ft) boat or an airboat. Each site was represented by a 2-m wide buffer, about 44 m<sup>2</sup> in area, extending off the perimeter of the boat. Six subsampling areas were clustered at each site; each of them was an imaginary (no marked boundary) rectangle of 1.5 m long and 0.36 m wide (the width of the rake head). Four of the subsampling areas were located off the corners of the boat and the other two were located off the left and right sides (Figure 7).

### Navigation to Sampling Sites

The field crew navigated in a boat to the general area of a site using an enlarged hard-copy map and then switched to global positioning system (GPS) equipment with differentially corrected signals as the boat approached the targeted location. The boat was anchored at bow and stern when both the easting and northing coordinates displayed on the GPS unit were within 10 m (- or +) of their respective target readings. The actual GPS coordinates were read and recorded twice at each site, once immediately after the boat had been anchored and again before the boat was released for departure.

### Sampling and Data Recording

Individual species and different life forms of aquatic vegetation were recorded as either present or absent at each subsampling area based on visual examination and a rake sample. When present, submersed species and the filamentous algae were given a density rating based on their thickness on the rake teeth (Table 2). When present, rooted floating-leaf and emergent species were given a percent cover rating (Table 3). Species that had not been recorded in the six subsampling areas but were observed at the site were recorded and marked as "additional species." Fassett (1957), Voss (1972, 1985), and Gleason and Cronquist (1991) were the primary references used for plant identification. Scientific nomenclature and common names (Appendix) are based on those found in the U.S. Department of Agriculture's




Figure 7. Placement of the six subsampling areas around the boat.

where  $m$  is the total number of strata,  $A$  is the abundance index of the species in stratum  $j$ , and  $S_j$  is the acreage of stratum  $j$ .

### Percent Cover (Rooted floating-leaf life form)

The percent cover of rooted floating-leaf life forms in a stratum is computed using the following formula:

$$C = \frac{\sum_{j=1}^m L_j \cdot A}{m}$$

where  $m$  is the total number of sites in the stratum,  $L$  is the cover rating at individual sites, and  $A$  is the mid-point of the corresponding percent cover (Table 3). Percent cover in a pool is computed as the average of all shallow water strata, weighted by acreage:

$$C = \frac{\sum_{j=1}^m C_j \cdot S_j}{\sum_{j=1}^m S_j}$$

where  $m$  is the total number of strata,  $C$  is percent cover in stratum  $j$ , and  $S_j$  is the acreage of stratum  $j$ .

### References

- Deppe, E. R., and R. C. Lathrop. 1992. A comparison of two rake sampling techniques for sampling aquatic macrophytes. Wisconsin Department of Natural Resources Research Management Findings. PUBL-RS-732. 4 pp.
- Fassett, N. C. 1957. A manual of aquatic plants. University of Wisconsin Press, Madison. 405 pp.
- Gleason, H. A., and A. Cronquist. 1991. A manual of vascular plants of northeastern United States and adjacent Canada. 2nd edition. The New York Botanical Garden, Bronx. 910 pp.
- Jessen, R., and R. Lound. 1962. An evaluation of survey techniques for submerged aquatic plants. Minnesota Department of Conservation. Game Investigational Report 6, St. Paul. 10 pp.
- U.S. Fish and Wildlife Service. 1993. Operating Plan for the Upper Mississippi River System Long Term Resource Monitoring Program. Environmental Management Technical Center, Onalaska, Wisconsin, Revised September 1993. EMTC 91-P002R. 179 pp. (NTIS #PB94-160199)
- Voss, E. G. 1972. Michigan flora, Part I, gymnosperms and monocots. Cranbrook Institute of Science, Bloomfield Hills, Michigan. 488 pp.
- Voss, E. G. 1985. Michigan flora, Part II, dicots. Regents of the University of Michigan, Ann Arbor. 724 pp.
- Yin, Y., J. S. Winkelman, and H. A. Langrehr. 2000. Long Term Resource Monitoring Program procedures: Aquatic vegetation monitoring. U.S. Geological Survey, Upper Midwest Environmental Sciences Center, La Crosse, Wisconsin, April 2000. LTRMP 95-P002-7. 8 pp. + Appendixes A-C

## Appendix

Aquatic vegetation species found during stratified random sampling in Pools 4, 8, 13, and 26 of the Mississippi River and La Grange Pool of the Illinois River, 1998<sup>a</sup>

Common name	Scientific name	Species Code	Family	Life form <sup>b</sup>
Coontail, coon's tail	<i>Ceratophyllum demersum</i> L.	CEDE4	Ceratophyllaceae	S
Chara	<i>Chara</i> spp.	CH?AR	Characeae	S
Watermilfoil	<i>Myriophyllum</i> spp.	MY?RI	Haloragaceae	S
Northern watermilfoil, shortspike watermilfoil	<i>M. sibiricum</i> Komarov	MYSI	Haloragaceae	S
Eurasian watermilfoil, spike watermilfoil	<i>M. spicatum</i> L.	MYSP2	Haloragaceae	S
Canadian waterweed	<i>Elodea canadensis</i> Michx.	ELCA7	Hydrocharitaceae	S
Wild celery, American eelgrass	<i>Vallisneria spiralis</i> L. <i>Utricularia macrorhiza</i> Le Conte synonymy	VAAM3	Hydrocharitaceae	S
Common bladderwort	<i>U. vulgaris</i> L.	UTMA	Lentibulariaceae	S
Bushy pondweed, slender naiad, nodding water nymph	<i>Najas flexilis</i> (Willd.) Rostk. and Schmidt	NAFL	Najadaceae	S
Southern water nymph	<i>N. guadalupensis</i> (Spreng.) Magnus	NAGU	Najadaceae	S
American lotus	<i>Nelumbo lutea</i> Willd.	NELU	Nelumbonaceae	F
Yellow pond-lily	<i>Nuphar variegata</i> Durand. <sup>c</sup>	NULU	Nymphaeaceae	F
White waterlily	<i>Nymphaea odorata</i> Ait. synonymy <i>N. tuberosa</i> Paine	NYTU	Nymphaeaceae	F
Water stargrass, grassleaf mudplantain	<i>Heteranthera dubia</i> (Jacq.) MacM. synonymy <i>Zosterella dubia</i> Jacq.	ZODU	Pontederiaceae	S
Leafy/small pondweed	<i>Potamogeton foliosus</i> Raf./ <i>P. pusillus</i> L.	NLPW	Potamogetonaceae	S
Curly pondweed, curlyleaf pondweed	<i>P. crispus</i> L.	POCR3	Potamogetonaceae	S
Leafy pondweed	<i>P. foliosus</i> Raf.	POFO3	Potamogetonaceae	S
Longleaf pondweed, American pondweed	<i>P. nodosus</i> Poir	PONO2	Potamogetonaceae	S
Sago pondweed	<i>P. pectinatus</i> L.	POPE6	Potamogetonaceae	S
Small pondweed, slender pondweed	<i>P. pusillus</i> L.	POPU7	Potamogetonaceae	S
Richardson's pondweed	<i>P. richardsonii</i> (Benn.) Rydb.	PORI2	Potamogetonaceae	S
Flatstem pondweed	<i>P. zosteriformis</i> Fern.	POZO	Potamogetonaceae	S
Longbeak buttercup	<i>Ranunculus longirostris</i> Godr. <sup>d</sup>	RALO2	Ranunculaceae	S
Horned pondweed	<i>Zannichellia palustris</i> L.	ZAPA	Zannichelliaceae	S

<sup>a</sup>Scientific nomenclature and common names follow the U.S. Department of Agriculture's Internet PLANTS Database (1996). Common names used by Upper Mississippi River managers are also included.

<sup>b</sup>F = rooted floating-leaf      S = submersed

<sup>c</sup>Scientific nomenclature follows Gleason and Cronquist (1991). *Nuphar lutea* spp. *variegata* in PLANTS database.

<sup>d</sup>*Ranunculus longirostris* and *R. trichophyllus* were combined (Voss 1985).

## Chapter 1. Results in Pool 4, Upper Mississippi River

### Sampling Efforts

Sampling began June 22 and ended August 11, 1998. Of the 550 sites targeted for sampling at the beginning of the season, 545 were actually sampled. Five sites were not sampled because dense emergent vegetation and low water impeded access.

### Submersed Vegetation

The status of submersed aquatic vegetation (SAV) varied between different sections of Pool 4 and the different strata sampled (Table 1.1; Figure 1.1). The SAV was scarce in and above Lake Pepin, but a much greater abundance (measured by both percent frequency of occurrence and the abundance index) was found in lower Pool 4, below Lake Pepin. When comparisons were made between strata, the greatest abundance was found in the isolated backwaters, followed by the contiguous backwaters. The distribution of SAV along the main and secondary channels was limited. Poolwide, about 36.6% of the shallow water areas supported SAV. Considering the acreage of SAV, contiguous backwaters below Lake Pepin were the primary habitat for SAV in Pool 4.

Species richness of submersed plants seemed to be closely correlated with abundance. That is, where there was greater abundance, there was generally more species. A total of 17 submersed plant species/taxon were recorded in the entire pool. The isolated

backwaters and the contiguous backwaters below Lake Pepin were rich in species.

Where SAV was scarce, such as in and above Lake Pepin and along the main and secondary channels, sago pondweed (*Potamogeton pectinatus*) was generally the most abundant species. Coontail (*Ceratophyllum demersum*) and sago pondweed were the most abundant species in the isolated backwaters, although Canadian waterweed (*Elodea canadensis*) also thrived in that stratum. Coontail and Canadian waterweed were also the most abundant species in the contiguous backwaters below Lake Pepin, closely followed by wild celery (*Vallisneria spiralis*). Lower Pool 4, especially the contiguous backwaters, was the most important habitat for wild celery—a species that is known to produce high quality food (tubers) for migrating waterfowl in the fall.

### Rooted Floating-leaf Vegetation

White waterlily (*Nymphaea odorata*), American lotus (*Nelumbo lutea*), and yellow pond-lily (*Nuphar variegata*) were the three rooted floating-leaf species recorded (Table 1.2; Figure 1.1). White waterlily was recorded above and below Lake Pepin but not in Lake Pepin, American lotus was recorded below Lake Pepin, and yellow pond-lily was recorded at two contiguous backwater sites below Lake Pepin. Poolwide, rooted floating-leaf vegetation covered about 4.1% of the shallow water area.




Figure 1.1. Sampling sites where species were recorded within Pool 4, Upper Mississippi River System, 1998.




Figure 1.1. Continued.



## Chapter 2. Results in Pool 8, Upper Mississippi River

### *Sampling Efforts*

Sampling began June 23 and ended July 21, 1998. Of the 550 sites targeted for sampling at the beginning of the season, 545 were actually sampled. Four sites could not be accessed because dense emergent vegetation and low water inhibited access. One site could not be reached because a barge was docked on the site on the day of sampling.


### *Submersed Vegetation*

The status of submersed aquatic vegetation (SAV) in Pool 8 varied among the strata sampled. Isolated backwaters had the highest abundance of SAV (Table 2.1; Figure 2.1). Contiguous backwaters, secondary channels, impounded areas, and main channel border areas followed in decreasing order. Sizable beds were found throughout most of the shallow water areas, except in the main channel border areas and the lower impounded areas of the pool where sizable beds were generally absent. The SAV covered about 47.6% of the shallow water areas poolwide.


Except for the main channel border areas, each of the other four strata sampled harbored a rich assemblage of submersed plant species. A total of 16 species were recorded in the entire pool. Coontail (*Ceratophyllum demersum*) and Canadian waterweed (*Elodea canadensis*) were the two most abundant species in the contiguous backwaters while coontail and leafy/small pondweed (*Potamogeton foliosus/P. pusillus*) were most abundant in isolated backwaters. No species was significantly more abundant in the impounded areas and main and secondary channels.

### *Rooted Floating-leaf Vegetation*

White waterlily (*Nymphaea odorata*), American lotus (*Nelumbo lutea*), and yellow pond-lily (*Nuphar variegata*) were the three rooted floating-leaf species recorded (Table 2.2; Figure 2.1). The percent cover of rooted floating-leaf species was the highest in the isolated backwaters. American lotus and white waterlily were scattered in much of the isolated and contiguous backwaters, while yellow pond-lily had a much more limited distribution in contiguous and isolated backwaters. The three species together covered about 7.5% of the shallow water areas.



**Figure 2.1.** Sampling sites where species were recorded within Pool 8, Upper Mississippi River System, 1998.



**Figure 2.1.** Continued.

## Chapter 3. Results in Pool 13, Upper Mississippi River

### *Sampling Efforts*

Sampling began June 24 and ended July 29, 1998. Of the 550 sites targeted for investigation at the beginning of the season, 549 were actually sampled. One isolated backwater site was not accessible.

### *Submersed Aquatic Vegetation*

The submersed aquatic vegetation (SAV) was recorded throughout most of the shallow water areas, but much less frequently in the northern portion of the pool (Table 3.1; Figure 3.1). Isolated backwaters had the highest abundance of SAV, measured by both percent frequency of occurrence and the abundance index. Contiguous backwaters, impounded areas, secondary channels, and main channel border areas followed in decreasing order. However, most of the SAV were recorded in impounded areas and contiguous backwaters. About 42% of the shallow water areas in the pool supported SAV.

A total of 14 submersed aquatic plant species were recorded. Isolated backwaters, contiguous

backwaters, and impounded areas all supported a diverse assemblage of species; main channel border areas and secondary channels supported fewer species. Coontail (*Ceratophyllum demersum*) was the most abundant species in the isolated and contiguous backwaters, wild celery (*Vallisneria americana*) and water stargrass (*Heteranthera dubia*) were the most abundant species in the impounded areas, and sago pondweed (*Potamogeton pectinatus*) was the most abundant species along the main channel border areas and in the secondary channels.

### *Rooted Floating-leaf Vegetation*

American lotus (*Nelumbo lutea*) and white waterlily (*Nymphaea odorata*) were the two rooted floating-leaf species recorded (Table 3.2; Figure 3.1). The two species had a similar distribution in the pool. Together they covered about 6.5% of the shallow water areas, mostly in contiguous backwaters. However, the percent cover of rooted floating-leaf vegetation was higher in isolated backwaters than in the contiguous backwaters. Rooted floating-leaf vegetation was not recorded in main channel border areas.




Figure 3.1. Sampling sites where species were recorded within Pool 13, Upper Mississippi River System, 1998.




Figure 3.1. Continued.

## Chapter 4. Results in Pool 26, Upper Mississippi River

### *Sampling Efforts*

Sampling began June 15 and ended August 10, 1998. All of the 550 sites targeted for sampling at the beginning of the season were sampled.

### *Submersed Vegetation*

Only one sizable bed of submersed aquatic vegetation (SAV) was found in Pool 26. It was located in an isolated backwater of the Illinois River locally known as the Stump Lake (Table 4.1; Figure 4.1). The existence of SAV outside the Stump Lake was negligible—coontail (*Ceratophyllum demersum*) was recorded at two sites while sago pondweed (*Potamogeton pectinatus*) was recorded at one site. No SAV was recorded along the Mississippi River and its backwater areas above the confluence with

the Illinois River. The SAV in Stump Lake consisted of six species, among which sago pondweed and coontail were most abundant. Although localized in distribution, SAV was estimated to cover about 6.1% of the shallow water areas in Pool 26.

### *Rooted Floating–Leaf Vegetation*

American lotus (*Nelumbo lutea*) and floating primrose-willow (*Ludwigia peploides*) were the two rooted floating–leaf species recorded in Pool 26 (Table 4.2; Figure 4.1). No rooted floating–leaf vegetation was recorded along the Mississippi River and its backwater areas above the confluence with the Illinois River. Although very localized in distribution, the two species together covered about 0.9% of the shallow water areas.




Figure 4.1. Sampling sites where species were recorded within Pool 26, Upper Mississippi River System, 1998.




Figure 4.1. Continued.

## Chapter 5. Results in La Grange Pool, Illinois River

### Sampling Efforts

Sampling began June 7 and ended August 4, 1998. Of the 550 sites targeted for sampling at the beginning of the season, 526 were actually sampled. Twenty-four sites were not sampled because of unforeseen inaccessibility, lack of permission from private landowners, equipment failure, and oversight in tracking sampling sites.


### Submersed Vegetation

No submersed aquatic vegetation (SAV) was recorded at the sampling sites. Some small beds of

sago pondweed (*Potamogeton pectinatus*) were observed along the main channel border and secondary channel border areas during informal surveys. However, the amount of SAV was negligible poolwide.

### Rooted Floating-leaf Vegetation

One species (primrose-willow, [*Ludwigia peploides*]) was recorded at one isolated backwater site (Figure 5.1), which indicates that the amount of rooted floating-leaf vegetation was negligible poolwide.



**Figure 5.1.** Sampling sites where species were recorded in La Grange Pool, Upper Mississippi River System, 1998.

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, D.C. 20503			
1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE May 2001	3. REPORT TYPE AND DATES COVERED	
4. TITLE AND SUBTITLE 1998 Annual Status Report: Submersed and rooted floating-leaf vegetation in Pools 4, 8, 13, and 26 and La Grange Pool of the Upper Mississippi River System		5. FUNDING NUMBERS	
6. AUTHOR(S) Yao Yin, <sup>1</sup> Heidi Langrehr, <sup>2</sup> Theresa Blackburn, <sup>3</sup> Megan Moore, <sup>4</sup> Jenny Winkleman, <sup>4</sup> Robert Cosgriff, <sup>5</sup> and Thad Cook <sup>6</sup>			
7. PERFORMING ORGANIZATION NAME AND ADDRESS <sup>1</sup> U.S. Geological Survey, Upper Midwest Environmental Sciences, 2630 Fanta Reed Road, La Crosse, Wisconsin 54603; <sup>2</sup> Wisconsin Department of Natural Resources, Onalaska Field Station, 575 Lester Avenue, Onalaska, Wisconsin 54650; <sup>3</sup> Iowa Department of Natural Resources, Mississippi River Monitoring Station, 206 Rose Street, Bellevue, Iowa 52031; <sup>4</sup> Minnesota Department of Natural Resources, Lake City Field Station, 1801 South Oak Street, Lake City, Minnesota 55041; <sup>5</sup> Illinois Natural History Survey, Alton Field Station, 8450 Montclair Avenue, Brighton, Illinois 62012; and <sup>6</sup> Illinois Natural History Survey, Havana Field Station, 704 North Schrader Avenue, Havana, Illinois 62644.		8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Geological Survey, Upper Midwest Environmental Sciences Center, 2630 Fanta Reed Road, La Crosse, Wisconsin 54603		10. SPONSORING/MONITORING AGENCY REPORT NUMBER 2001-P001	
11. SUPPLEMENTARY NOTES			
12a. DISTRIBUTION/AVAILABILITY STATEMENT Release unlimited. Available from National Technical Information Service, 5285 Port Royal Road, Springfield, VA 22161 (1-800-553-6847 or 703-487-4650. Available to registered users from the Defense Technical Information Center, Attn: Help Desk, 8725 Kingman Road, Suite 0944, Fort Belvoir, VA 22060-6218 (1-800-225-3842 or 703-767-9050).		12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) Aquatic vegetation was investigated in five navigation pools in the Upper Mississippi River System using a new protocol named "stratified random sampling" or SRS protocol for the first time in 1998. The five pools were Pools 4, 8, 13, and 26 of the Upper Mississippi River and La Grange Pool of the Illinois River. The results on submersed aquatic vegetation and rooted floating-leaf aquatic vegetation were summarized in this report. The percent frequencies of submersed aquatic vegetation in shallow water areas ( $\leq 3$ m deep at flat-pool condition) in the five pools were 36.6%, 47.6%, 42%, 6.1%, and 0%, respectively. The aquatic area strata that were directly influenced by the flow in the main channel, such as the main channel borders and secondary channels, had lower percent frequencies of submersed aquatic vegetation than the aquatic area strata that were less directly influenced by the flow in the main channel, such as the contiguous and isolated backwaters. The percent covers of rooted floating-leaf vegetation were 4.1%, 7.5%, 6.5%, 0.9%, and 0%. The majority of aquatic vegetation that was recorded in Pool 26 was from one isolated backwater area. Aquatic vegetation was not recorded at any of the sampling sites in La Grange Pool.			
14. SUBJECT TERMS Annual report, aquatic, floating-leaf, Illinois River, La Grange, LTRMP, Mississippi River, submersed vegetation.		15. NUMBER OF PAGES 9 pp. + Appendix + Chapters 1-5	
		16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT