

UCRL-Trans-1003(L)

# D T I E MICROCARD ISSUANCE DATE

17 / 13

1964

20011011 065

DEC 3 1963

UCRL-Trans-1003(L)

## SHOCK WAVES IN A PLASTIC COMPACTING MEDIUM

MAS: [unclear]

A. S. Kompaneets

Translated from  
Akademiia Nauk SSSR, Doklady  
109, 49-52 (1956)

DISTRIBUTION STATEMENT A  
Approved for Public Release  
Distribution Unlimited

University of California  
Lawrence Radiation Laboratory  
Livermore, California  
Translated by: Sergey Shewchuck  
October 1963

**LEGAL NOTICE**  
This report was prepared as an account of Government sponsored work. Neither the United States nor the Government makes any warranty, representation, or guarantee, expressed or implied, with respect to the accuracy or completeness of the information contained in this report, or that the use of any apparatus, apparatus, method, or process described in this report may not infringe upon any patent rights.  
The Government is authorized to reproduce and distribute reprints for Government purposes not withstanding any copyright notation that may appear hereon.  
Persons acting in behalf of the "Commission" include the work employees or employees of the Commission, or employees of any contractor performing work under contract with the Commission, or any subcontractor with such contractor.

Photostat Price \$ 1.10  
Microfilm Price \$ .80  
Available from the  
Office of Technical Services  
Department of Commerce  
Washington 25, D. C.

COLLEGE TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
LIVERMORE LIBRARY

The problem of slow propagation of a plastic deformation was solved by S. L. Sobolev.<sup>1</sup> We shall consider a shock propagation of a plastic deformation in a medium which is characterized by the following properties. In the initial state its density is  $\rho_0$ , at this density it exerts a negligibly small resistance to compression. When brought to a density  $\rho_1$ , the medium is incompressible and in this state is plastic, whereupon it is assumed that the absolute value of the most "tangential" tension is linearly dependent on the average normal tension (the plasticity condition of Prandtl)<sup>2</sup>. This kind of plasticity, for example, can be possessed by sand.

A shock wave emerges (is formed) under the effect of an explosion in some very small spherical volume with radius  $R_0$ . The main axes of the stress tensor coincides with the coordinate lines of the circle system, whose center is superposed (congruent) with the center of explosion. The main normal tensions we shall name as  $\sigma_r$  and  $\sigma_\phi = \sigma_\theta$ . The condition of plasticity, according to the assumption, must be written thus:

$$\sigma_r - \sigma_\phi = k + m(\sigma_r + 2\sigma_\phi) \quad (1)$$

The equations of motion of the compacted medium have the form

$$\rho_1 \left( \frac{\partial u}{\partial t} + u \frac{\partial u}{\partial r} \right) = \frac{\partial \sigma_r}{\partial r} + \frac{2(\sigma_\phi - \sigma_r)}{r} \quad (2)$$

$$\frac{1}{r^2} \frac{\partial}{\partial r} (r^2 \dot{\epsilon}) = 0 \quad (3)$$

From Eq. (3) it follows that

$$u = \frac{\dot{\epsilon}(t)}{r} \quad (4)$$

Substituting this in (2) and eliminating  $\sigma_\phi$  with the help of (1), we derive after integration:

$$\sigma_r = -\frac{k}{3m} + \frac{1}{(\alpha-1)r} \left[ \frac{\dot{\epsilon}}{\alpha-1} - 2\dot{\epsilon}_0 \frac{k^2}{(\alpha-1)^2} + C(t) r^{-\alpha} \right] \quad (5)$$

where

$$\alpha = \frac{m}{1+2m} \quad (6)$$

The boundary conditions on the front of the shock wave have the form

$$\rho_0 \dot{R} = \rho_1 (\dot{R} - u(R)), \quad (7)$$

$$\rho_0 \dot{R}^2 = \rho_1 (\dot{R} - u(R))^2 - \sigma_r(R). \quad (8)$$

Here  $R$  is the radius of the shock wave front.

At the boundary of the expanding cavity one must equate the pressure to the normal tension, taken with the opposite sign:

$$p = -\sigma_r(R_0). \quad (9)$$

The radius of the cavity is easily determined by starting with conservation of mass. If  $R_0$  is the initial radius of the cavity, we derive

$$R_0 = \left[ \left( 1 - \frac{\rho_0}{\rho_1} \right) R^3 + \frac{2\rho_0}{\rho_1} R^2 \right]^{1/3} \cong R \left( 1 - \frac{\rho_0}{\rho_1} \right)^{1/3}. \quad (10)$$

One can disregard the quantity  $R_0^3$  here, even if  $1 - \rho_0/\rho_1$  is of the order of several percent.

Assuming that the substance in the cavity expands like an ideal gas with an isentrope index  $\gamma$ , we find the expression for  $p$ :

$$p = \rho_0 \left( \frac{R_0}{R} \right)^{3\gamma} \left( 1 - \frac{\rho_0}{\rho_1} \right)^{-\gamma}. \quad (11)$$

Excluding  $C(t)$  and  $\lambda(t)$  from (7), (8) and (9), we come to the differential equation:

$$R \ddot{R} + \dot{R}^2 \left[ 2 + \frac{\rho_0}{\rho_1} \frac{\alpha-1}{1 - \left( 1 - \frac{\rho_0}{\rho_1} \right)^{\frac{1-\alpha}{\alpha}}} - 2 \frac{\left( \alpha-1 \right) \left[ \left( 1 - \frac{\rho_0}{\rho_1} \right)^{\frac{1-\alpha}{\alpha}} - 1 \right]}{\left( \alpha-1 \right) \left[ \left( 1 - \frac{\rho_0}{\rho_1} \right)^{\frac{1-\alpha}{\alpha}} - 1 \right]} \right] = \left\{ \frac{\rho_1}{\alpha-1} \left( 1 - \frac{\rho_0}{\rho_1} \right)^{1-\alpha} \left[ \left( 1 - \frac{\rho_0}{\rho_1} \right)^{\frac{1-\alpha}{\alpha}} - 1 \right] \right\}^{-1} \left\{ p + \frac{k}{3m} \left[ \left( 1 - \frac{\rho_0}{\rho_1} \right)^{\frac{1-\alpha}{\alpha}} - 1 \right] \right\}. \quad (12)$$

It is integrated in quadratures. Before writing down the quadrature, it is convenient to study the expression in the right hand side. The equation must, in any case, permit a solution which answers to the equilibrium,  $\dot{R} = 0$ ,  $R = R_0$ . In the right hand side of (12) there will stand 0, if

$$\left( \frac{\dot{R}_0}{R_0} \right)^{3\gamma} = -\frac{k}{\rho_0} \frac{1}{3m} \left[ \left( 1 - \frac{\rho_0}{\rho_1} \right)^{\frac{1-\alpha}{\alpha}} - 1 \right] \left( 1 - \frac{\rho_0}{\rho_1} \right)^{\gamma}. \quad (13)$$

Equilibrium must be achieved also then when  $m = 0$ , just as in the plasticity theory of Sen-Venana, because  $k$  and  $m$  are not connected with each other. But in this case it is necessary for equilibrium that  $k$  be less than zero. The ratio  $|k|/\rho_0$  is a very small number, of the order of  $10^{-4}$  or less. This justifies the approximation, made in (10), applicable to the condition of equilibrium (13) for not very large values of  $\sigma$ .

The condition of equilibrium (13) for  $k < 0$  is satisfied for all values  $m \geq -1/2$ . But if  $-1/2 \leq m < 0$ , then the equilibrium radius is greater than for  $m = 0$ . This is not very probable, because  $m$  describes the additional (auxiliary) friction in the condition of Prandtl. Therefore, one must consider that  $m < 0$ . For very large values of  $m$  the equilibrium radius again becomes greater than for  $m = 0$ . Consequently, these values of  $m$  lie beyond the limits of applicability of the linear formula (1) in the problem under consideration.

Let us now introduce the following dimensionless quantities:

$$\frac{R}{R_0} = x; \quad R \sqrt{\frac{2\mu}{\rho_0}} = y; \quad 1 - \frac{2\mu}{\rho_0} \xi = \zeta; \quad (14)$$

$$1 - \frac{2\mu}{\rho_0} \xi = \zeta; \quad 1 - \frac{2\mu}{\rho_0} \xi = \zeta; \quad 1 - \frac{2\mu}{\rho_0} \xi = \zeta; \quad (15)$$

$$A = 2 \frac{(\alpha - 1)(1 - \xi)}{\xi^2 (\xi^2 - 1)}; \quad (16)$$

In these variables the Eq. (12) is rewritten thus:

$$x \frac{d(y^2)}{dx} + y^2 = A \left( \frac{x^{-2\alpha}}{\xi^2} - \frac{|k|}{\rho_0} \left| \xi^2 - \frac{2}{3m} \right| \right); \quad (17)$$

Its first integral is

$$y^2 = \frac{1}{\xi} x^{-\alpha} + Ax^{-\alpha} \int x^{\alpha-1} \left( \frac{x^{-2\alpha}}{\xi^2} - \frac{|k|}{\rho_0} \left| \xi^2 - \frac{2}{3m} \right| \right) dx; \quad (18)$$

The approximation (10), and thus all those following, is justified only in the case if in the integral (18) the region close to 1 does not give a material

contribution, i. e., if  $\mu - 3\gamma > 0$ . This condition does not seem particularly restrictive, as evident from Table 1 of the values  $\mu$  for different  $\alpha$  and porosity  $\xi = 1 - \rho_0/\rho_1$ .

Table 1

$\xi$	$\alpha$	0	1	2	3
0.01		3.26	4.44	6.00	7.84
0.1		3.78	4.78	5.93	7.40
0.2		4.12	5.00	5.96	7.12

Considering that condition  $\mu - 3\gamma > 0$  is satisfied, one can disregard also the first term in Eq. (18), of which one is easily convinced by comparing the coefficient  $A/\xi^2$  with the quantity  $(1 - \xi)/\xi$ . Then a simple expression is derived for  $y^2$ :

$$y^2 = A \left( \frac{x^{-2\alpha}}{(\mu - 3\gamma)\xi^2} - \frac{|k|}{\rho_0} \left| \xi^2 - \frac{2}{3m} \right| \right); \quad (19)$$

from which there is obtained the value of the maximum radius of wave expansion

$$R_{m=0} = R_0 \left[ \frac{3m\mu}{\mu - 3\gamma} \frac{1}{\left( \xi^2 - \frac{2}{3m} \right) \xi^2} - \frac{1}{|k|} \right]^{\frac{1}{2\alpha}}; \quad (20)$$

The substance remains compressed even after load release, since the compression is irreversible.  $R_m$  is larger than the equilibrium radius in the ratio  $[\mu/(\mu - 3\gamma)]^{1/3\alpha}$ .

The time of full expansion is equal to

$$t_m = R_0 \sqrt{\frac{2\mu}{\rho_0} \frac{(\mu - 3\gamma)^{2\alpha}}{A} \left( \frac{R_m}{R_0} \right)^{2\alpha} \frac{1}{\Gamma\left(\frac{1}{3\alpha} + 1\right)}}; \quad (21)$$

Note that there did not go into the problem any relationship between the tensions and the deformations or the deformation velocities. Here we find an analogy with the statically determinable flat problems of the plasticity theory.

The Institute of Chemical Physics  
of the Academy of Sciences, USSR  
Submitted Nov. 1, 1956

Cited Literature

1. S. L. Sobolev. Trudy of seismological instr. 49 (1935).
2. L. S. Leibenzon. Elements of mathematical theory of plasticity, 1943, page 56.

END