

11761

U.S. Army Center for Health Promotion
and Preventive Medicine

**TRAINING MUNITIONS HEALTH RISK
ASSESSMENT
NO. 39-EJ-1485-00
RESIDENTIAL EXPOSURE FROM INHALATION OF
AIR EMISSIONS FROM THE
M17 .50 CALIBER TRACER CARTRIDGE
DEPARTMENT OF DEFENSE IDENTIFICATION CODE: A571**

Prepared by:

Environmental Health Risk Assessment Program

Prepared for:

U.S. Army Environmental Center

Published date:

15 June 2001

Approved for public release; distribution unlimited



U
S
C
H
P
P
M

20011218 130

Readiness Thru Health

U.S. Army Center for Health Promotion and Preventive Medicine

The lineage of the U.S. Army Center for Health Promotion and Preventive Medicine (USACHPPM) can be traced back over 50 years. This organization began as the U.S. Army Industrial Hygiene Laboratory, established during the industrial buildup for World War II, under the direct supervision of the Army Surgeon General. Its original location was at the Johns Hopkins School of Hygiene and Public Health. Its mission was to conduct occupational health surveys and investigations within the Department of Defense's (DOD's) industrial production base. It was staffed with three personnel and had a limited annual operating budget of three thousand dollars.

Most recently, it became internationally known as the U.S. Army Environmental Hygiene Agency (AEHA). Its mission expanded to support worldwide preventive medicine programs of the Army, DOD, and other Federal agencies as directed by the Army Medical Command or the Office of The Surgeon General, through consultations, support services, investigations, on-site visits, and training.

On 1 August 1994, AEHA was redesignated the U.S. Army Center for Health Promotion and Preventive Medicine with a provisional status and a commanding general officer. On 1 October 1995, the nonprovisional status was approved with a mission of providing preventive medicine and health promotion leadership, direction, and services for America's Army.

The organization's quest has always been one of excellence and the provision of quality service. Today, its goal is to be an established world-class center of excellence for achieving and maintaining a fit, healthy, and ready force. To achieve that end, the CHPPM holds firmly to its values which are steeped in rich military heritage:

★ *Integrity is the foundation*

★ *Excellence is the standard*

★ *Customer satisfaction is the focus*

★ *Its people are the most valued resource*

★ *Continuous quality improvement is the pathway*

This organization stands on the threshold of even greater challenges and responsibilities. It has been reorganized and reengineered to support the Army of the future. The CHPPM now has three direct support activities located in Fort Meade, Maryland; Fort McPherson, Georgia; and Fitzsimons Army Medical Center, Aurora, Colorado; to provide responsive regional health promotion and preventive medicine support across the U.S. There are also two CHPPM overseas commands in Landstuhl, Germany and Camp Zama, Japan who contribute to the success of CHPPM's increasing global mission. As CHPPM moves into the 21st Century, new programs relating to fitness, health promotion, wellness, and disease surveillance are being added. As always, CHPPM stands firm in its commitment to Army readiness. It is an organization proud of its fine history, yet equally excited about its challenging future.

REPORT DOCUMENTATION PAGE					Form Approved OMB No. 0704-0188	
The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.						
1. REPORT DATE (DD-MM-YYYY) 06/15/2001		2. REPORT TYPE Technical Report		3. DATES COVERED (From - To) March 1999-August 2001		
4. TITLE AND SUBTITLE Training Munitions Health Risk Assessment No.39-EJ-1485-00 Residential Exposure from Inhalation of the Air Emissions from the M17 .50 Caliber Tracer Cartridge , Department of Defesnse Identification Code: A571				5a. CONTRACT NUMBER		
				5b. GRANT NUMBER		
				5c. PROGRAM ELEMENT NUMBER		
				5d. PROJECT NUMBER		
6. AUTHOR(S) Laura Peters, Stafford D.F.R.Coakley, Joleen Mobley				5e. TASK NUMBER		
				5f. WORK UNIT NUMBER		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) U.S. Army Center for Health Promotion and Preventive Medicine 5158 Blackhawk Road Aberdeen Proving Ground, Maryland 21010-5422				8. PERFORMING ORGANIZATION REPORT NUMBER Risk Assessment # 39-EJ-1485-00		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Environmental Center ATTN: SFIM-AEC-PC Aberdeen Proving Ground, MD 21010-5401				10. SPONSOR/MONITOR'S ACRONYM(S) USAEC		
				11. SPONSOR/MONITOR'S REPORT NUMBER(S) SFIM-AEC-PC-CR-200141		
12. DISTRIBUTION/AVAILABILITY STATEMENT Distribution Unlimited						
13. SUPPLEMENTARY NOTES Point of Contact: Tamera Rush 410-436-6849						
14. ABSTRACT This assessment evaluated the potential for human health effects to offsite residents breathing air emissions following use of the M17 .50 Caliber Tracer Cartridge. This document present the evaluation of the potential for adverse human health effects to the offsite residents breathing air emissions following the use of military firing ranges during training exercises. Study results showed no protential for health risks to the hypothetical resident from inhalation of air emissions from the .50 Caliber Cartridge. To conduct this study, air emissions from the M17 .50 Caliber Tracer Cartridge were collected in a test chamber (at Aberdeen Test Center, Aberdeen, MD). This information was then used in an air dispersion model to determine ambient air concentrations at a location downwind from the site where the item was activated. Modeled air concentrations were combined with exposure information to estimate the amount of substances the hypothetical resident breathes. This intake was combined with the substance's health information, to determine if there is a potential for health risks from inhjalation of these substances. The health risk included both long-term and short term exposures to the modeled substance concentrations. Study results showed no potential for helath risks from inhalation of air emissions from the M17 .50 Caliber Tracer Cartridge						
15. SUBJECT TERMS emissions, aberdeen test center, characterization, health risk, munitions, firing point,						
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UU	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON Tamera Rush	
a. REPORT U	b. ABSTRACT U	c. THIS PAGE U			19b. TELEPHONE NUMBER (Include area code) 410-436-6849	



REPLY TO
ATTENTION OF

DEPARTMENT OF THE ARMY
U.S. ARMY CENTER FOR HEALTH PROMOTION AND PREVENTIVE MEDICINE
5158 BLACKHAWK ROAD
ABERDEEN PROVING GROUND, MARYLAND 21010-5403

MCHB-TS-EHR

TRAINING MUNITIONS HEALTH RISK ASSESSMENT NO. 39-EJ-1485-00
RESIDENTIAL EXPOSURE FROM INHALATION OF AIR EMISSIONS
FROM THE M17 .50 CALIBER TRACER CARTRIDGE

EXECUTIVE SUMMARY

This assessment evaluated the potential for human health effects to offsite residents breathing air emissions following use of the M17 .50 Caliber Tracer Cartridge (M17) on firing ranges during training exercises.

To conduct this assessment, air emissions from the M17 were collected in a test chamber at the U.S. Army Aberdeen Test Center, Maryland. The data collected from the Firing Point Emission Study provided the amount and types of substances released from the M17. This information was then used in an air dispersion model to determine ambient air concentrations at locations downwind from the M17 firing location. Since the training facility in this assessment is hypothetical, the air model used assumptions that provided conservative estimates of air concentrations.

Modeled air concentrations were combined with exposure information (e.g., number of cartridges used per year) to estimate the amount of each substance the hypothetical offsite resident breathes. This estimate was then compared with the substance's health-based screening level, which was obtained from agencies such as the U.S. Environmental Protection Agency, to determine if there is a potential for health effects from inhalation.

The health risk assessment included both long-term (30 years) and short-term (15-minute or 1-hour) exposures to modeled substance concentrations. Assessment results, generated using conservative methods, showed that the hypothetical offsite resident breathing air as close as 200 meters (656 feet) from the M17 firing location is safe from these emissions. At locations where offsite residents are located less than 200 meter from the M17 firing locations, a more site-specific evaluation is necessary. It should be noted that at most training installations, training areas are over 1,000 meters (over half a mile) away from populated areas.

TABLE OF CONTENTS

1. PURPOSE	1
2. AUTHORITY	1
3. REFERENCES	1
4. BACKGROUND	1
4.1 CARTRIDGES AND THEIR USE	1
4.2 WHAT IS THE M17?	1
4.3 USE OF THE M17	1
4.4 ASSESSMENT SUMMARY	2
5. DATA COLLECTION AND AIR MODELING	3
5.1 EMISSION FACTORS	3
5.2 BACKGROUND AND DESCRIPTION	3
5.3 MODEL ASSUMPTIONS	3
5.4 GENERAL METHODOLOGY	5
5.5 USE OF MODEL OUTPUT	5
5.6 DETERMINATION OF SUBSTANCE-SPECIFIC EMISSION RATES	6
6. RISK ASSESSMENT	7
6.1 EXPOSURE ASSUMPTIONS	7
6.2 TIME-AVERAGING	7
6.3 TOXICITY ASSESSMENT	10
7. RISK CHARACTERIZATION	14
7.1 CHRONIC HEALTH RISK	14
7.2 ACUTE HEALTH RISK	15
7.3 FACT SHEET	16
8. UNCERTAINTY DISCUSSION	16
9. CONCLUSION	18
10. RECOMMENDATIONS	18
11. POINT OF CONTACT	19

LIST OF APPENDICES

REFERENCES	APPENDIX A
AIR DISPERSION MODELING OUTPUT DATA	APPENDIX B
HEALTH-BASED SCREENING LEVELS AND ACUTE TOXICITY VALUES	APPENDIX C
RISK ASSESSMENT DATA	APPENDIX D
FACT SHEET SUBMITTED TO THE U.S. ARMY ENVIRONMENTAL CENTER.....	APPENDIX E

LIST OF TABLES

TABLE 1 – SOURCE PARAMETERS	4
TABLE 2 – WORST-CASE METEOROLOGICAL PARAMETERS.....	5
TABLE 3 – AIR MODEL INPUT PARAMETERS	5
TABLE 4 – FREQUENCY OF USE FOR THE M17.....	7
TABLE 5 – EXPOSURE PARAMETERS USED TO DETERMINE TIME-AVERAGED CHRONIC AIR CONCENTRATIONS	8
TABLE 6 – SUMMARY OF RfCs USED FOR PETROLEUM HYDROCARBONS	12
TABLE 7– TYPES OF UNCERTAINTY	17

LIST OF ACRONYMS

AEC	U.S. Army Environmental Center
AEGL	Acute Exposure Guideline Levels
AIHA	American Industrial Hygiene Association
Al	Aluminum
ATC	U.S. Army Aberdeen Test Center
ATV	Acute Toxicity Value
CO ₂	Carbon Dioxide
DODIC	Department of Defense Identification Code
DOE	U.S. Department of Energy
EPA	U.S. Environmental Protection Agency
ERPG	Emergency Response Planning Guidelines
HBSL	Health-Based Screening Level
INPUFF	Integrated PUFF Model
NAAQS	National Ambient Air Quality Standards
NEW	Net Explosive Weight
OEL	Occupational Exposure Limit
PM _{2.5}	Particulate Matter under 2.5 microns in size
PM ₁₀	Particulate Matter under 10 microns in size
PRG	Preliminary Remediation Goals
RBC	Risk-Based Concentration
RfC	Reference Concentration
TEEL	Temporary Emergency Exposure Limits
TPH	Total Petroleum Hydrocarbons
TSP	Total Suspended Particulates
USACHPPM	U.S. Army Center for Health Promotion and Preventive Medicine

TRAINING MUNITIONS HEALTH RISK ASSESSMENT NO. 39-EJ-1485-00
RESIDENTIAL EXPOSURE FROM INHALATION OF AIR EMISSIONS FROM THE
M17 .50 CALIBER TRACER CARTRIDGE

1. PURPOSE

This document presents the assessment of the potential for human health effects to offsite residents breathing air emissions following use of the M17 .50 Caliber Tracer Cartridge (M17) on firing ranges during training exercises.

2. AUTHORITY

Statement of Work, 30 November 2000, Training Munitions Inhalation Risk Evaluations.

3. REFERENCES

See Appendix A for a list of references.

4. BACKGROUND

4.1 CARTRIDGES AND THEIR USE

Cartridges are cases that contain a primer, propelling charge, and projectile. The primer is needed to activate the propelling charge, which provides the force to send the projectile to a target. Examples of projectiles include bullets, rockets, and missiles. Cartridges are also referred to as "rounds" and are fired from weapons such as pistols or rifles.

4.2 WHAT IS THE M17?

The M17 is a tracer cartridge, which is used to track the path of the bullet. When fired at night, the tracer leaves a visible trail to show the direction in which the bullet is traveling. Each tracer cartridge is about as long as the length of a soda can and can be identified by its brown tip (Reference 1).

The M17 has a brass cartridge case and a bullet consisting of a brass jacket and steel core. It also contains a propelling charge that is made up mostly of nitrocellulose. Nitrocellulose is commonly used in the production of lacquers and artificial leathers. It is a primary ingredient in smokeless propellant for both military and commercial use.

4.3 USE OF THE M17

The M17 is fired from .50 caliber machine guns. Each weapon operates differently and is used for different functions. Training with the M17 is very important as

it teaches our soldiers to operate these weapons safely and effectively. This will prepare them for combat situations.

4.4 ASSESSMENT SUMMARY

The general assessment approach consisted of two main parts: air dispersion modeling and exposure assessment, which are briefly discussed in the paragraphs below. Sections 5 through 7 present a discussion of the methodology used for this assessment.

Emissions data used in the air dispersion modeling were obtained from the Firing Point Emission Study, conducted by the U.S. Army Aberdeen Test Center (ATC), at Aberdeen Proving Ground, Maryland (Reference 2). This assessment was funded by the U.S. Army Environmental Center (AEC) with the purpose of identifying and quantifying emissions from weapons firing. Data from this study were generated by firing munitions in a test chamber with weapons that are representative of those used by the U.S. Army during training operations. Emissions data for the M17 were generated by firing it from the M2 machine gun.

The emissions data for the M17 were used with an atmospheric dispersion model to estimate the average concentrations that may be experienced by an offsite resident. Since this assessment is designed to provide results that would be applicable to most Army training facilities, the training area used in this assessment was a hypothetical one. While most training areas are at least 1,000 meters away from populated areas, as a conservative distance, it was initially assumed that a person could reside 100 meters downwind from the firing point (location where the machine gun is positioned). In addition, air-modeling parameters were selected to mimic worst-case conditions.

The exposure assessment included calculations of time-averaged concentrations for both long-term (chronic) and short-term (acute) exposures. For the purpose of this assessment, air concentrations were averaged over 30 years for chronic exposures and 1-hour or 15 minutes for acute exposures. Using a screening approach, a substance's estimated time-averaged air concentration was then compared to chronic health-based screening levels (HBSLs) established by the U.S. Environmental Protection Agency (EPA) or acute toxicity values (ATVs) established by selected agencies depending on the exposure duration (i.e., 30 years versus 1-hour or 15 minutes). The comparison was made using the ratio of the HBSL or ATV to estimated air concentration for each of the substances evaluated. If this ratio was less than one, no further evaluation was needed. This approach is conservative because the exposure assumptions used by the agencies, to establish HBSLs and ATVs, are likely to overestimate the exposures experienced by offsite residents living near firing ranges. If the chronic or acute averaged concentrations (C_{chronic} and C_{acute}) were greater than these screening levels, producing a ratio greater than one, further analysis would be warranted to determine the potential for health effects. Note that concentrations greater than the screening levels do not indicate an onset of health effects, but rather the potential for such.

5. DATA COLLECTION AND AIR MODELING

5.1 EMISSION FACTORS

Emission factors, used to derive the air modeling emission rates used in this assessment, were generated from the Firing Point Emission Study conducted by the ATC (Reference 2). The data provided by the ATC included the net explosive weight (NEW), the substances sampled, and substance-specific emission factors. Emissions data from the Firing Point Emission Study are included in the first four columns of the table located in Appendix B.

5.2 BACKGROUND AND DESCRIPTION

Air dispersion models are available to mathematically simulate plume behavior and to estimate downwind concentrations of substances emitted from various sources. However, specific models are not available to determine the dispersion of emissions from munitions used during training. Estimating the magnitude and location of these concentrations depends on many factors including the amount and type of emissions, the behavior of the source, and meteorological conditions. Since a specific model is not available for modeling the use of munitions during training, the U.S. Army Center for Health Promotion and Preventive Medicine (USACHPPM) evaluated numerous air models to determine which would be suitable for use with munitions used during training. The USACHPPM recommended using the Integrated PUFF (INPUFF) model to estimate the dispersion of emissions from various munitions sources (Reference 3).

The INPUFF Model (Reference 4) was developed to simulate dispersion from instantaneous or semi-continuous point sources. This Gaussian-integrated puff model is capable of addressing a cloud type release over short periods of time, and computations can be performed for a single point source for multiple receptors. The algorithms used to calculate concentrations assume a vertically uniform wind direction (with no chemical reaction) to compute the contribution of each cloud at a receptor for each time step/interval.

5.3 MODEL ASSUMPTIONS

Some assumptions were made to best represent the firing of the M17 cartridges. These assumptions were as follows:

- Typically, with conventional point sources (such as incinerators), the cloud rise and formation are determined by characterizing flue gas exit velocity, temperature, and stack diameter. However, the M17 cartridges are used in conjunction with machine guns. For unconventional sources with no real physical stack dimensions, such as machine guns, the stack height and diameter were assumed to be equal to the height of the barrel and the bore diameter. No exit velocity was used with this source because the emission rates generated from the test data were obtained from sampling a stabilized

cloud with no exit velocity. Table 1 includes the source parameters used to model the M17 cartridges.

TABLE 1: SOURCE PARAMETERS

Parameter	Model Input
Source/Stack Diameter	0.01 meters
Source/Stack Height	1 meter
Source Exit Temperature	298.15 degrees Kelvin ($^{\circ}$ K) (or 77 $^{\circ}$ F)
Exit Velocity	0 meters/second
Initial horizontal dispersion coefficient (σ_y)	0.87 meters
Initial vertical dispersion coefficient (σ_z)	1.07 meters

- Initial cloud dimensions are preferred to model the air emissions from these types of releases. The dimensions are used to define the initial horizontal and vertical dispersion values (σ_y and σ_z) of the released cloud. This information was not measured during the studies at the ATC; therefore, the cloud dimensions were based on the test chamber dimensions and the volume of air sampled. By assuming an elliptical cloud with the prevailing wind direction being perpendicular to the muzzle when fired, the test chamber's radius would be equal to the initial vertical dispersion (σ_z), and the initial horizontal dispersion (σ_y), would be equal to one half the length of the test chamber. The cloud exit temperature was assumed to be equal to the test chamber temperature.
- For the purposes of this assessment, a hypothetical offsite resident was assumed to be located first at 100 meters, then at 200 meters directly downwind from the source. The meander of the cloud is a major factor when estimating concentrations at given locations downwind from the source. Assuming that the resident is directly downwind from the source is the same as assuming that there is no cloud meander and that the center of the cloud migrates directly over the hypothetical offsite resident. This assumption provides the most conservative modeled concentrations.
- Since this assessment does not look at a specific training site, generic, worst-case meteorological data were used. To determine the worst-case meteorological conditions that would result in the highest air emission concentrations, the modeling was performed using the EPA Risk Management Program Guidance (Reference 5). This guidance includes tables for estimating the footprint of chemical releases and is intended to inform emergency responders of potential accidental releases. The EPA has defined most default conditions for meteorological modeling parameters. Table 2 lists the meteorological parameters that were used in the air model.

TABLE 2: WORST-CASE METEOROLOGICAL PARAMETERS

Parameter	Input Value
Wind Speed	1 meter/second
Atmospheric Stability	Category F
Wind Direction	270°
Ambient Temperature	293 degrees Kelvin (°K) (or 68 °F)

5.4 GENERAL METHODOLOGY

The model was run for a total calculation time of 200 seconds for the 100-meter location and 300 seconds for the 200-meter location. This was done to simulate a single round being fired and to ensure that the total mass of the cloud had passed the hypothetical resident locations. Concentrations were calculated every 2 or 3 seconds, depending on the location being modeled. The model results indicated that the initial cloud reached the hypothetical offsite resident at 200 meters within 154 seconds and dissipated below the lowest concentration the model calculated ($1 \times 10^{-12} \text{ g/m}^3$) within 275 seconds. Table 3 contains the air model input parameters used in this assessment.

TABLE 3: AIR MODEL INPUT PARAMETERS

Parameter	Input Value	
	100 meters	200 meters
Number of meteorological periods (NTIME)	1	1
Duration of each meteorological period (ITIME)	200 seconds	300 seconds
Number of updates to the source (NSRCDS)	100	100
Duration/time step between each source update (ISUPDT)	2 seconds	3 seconds
Total time modeled/Simulation Period (NTIME) (ITIME)= (NSRCDS) (ISUPDT)	200 seconds	300 seconds

5.5 USE OF MODEL OUTPUT

The concentrations provided by the INPUFF model were based on a unit emission rate (ER_{unit}) of 1 gram/second from an emission source, and did not represent any substance-specific concentrations from the use of any weapons system. This unit emission rate is typically used for ease of modeling purposes. The relationship between the emission rate and predicted concentration is linear. Therefore, the ratio of the predicted concentration to the unit emission rate was multiplied by each substance-specific emission rate to provide substance-specific concentrations.

5.6 DETERMINATION OF SUBSTANCE-SPECIFIC EMISSION RATES

The actual substance emission rate for one item (ER_1) for each substance was calculated using Equation 1. Example 1 contains a sample calculation using this equation.

$$ER_1 = \frac{EF \cdot CV}{t} \quad \text{Equation 1}$$

Where:

ER_1 = emission rate for one item ((g/item)/sec)

EF = average adjusted emission factor (lb/item)

CV = conversion factor (453.59 g/lb)

t = release duration obtained from the INPUFF model (sec)

Example 1
Sample Calculation Using Equation 1:

$$ER_1 = \frac{(5.69 \text{ E} - 03)(453.59)}{(3)}$$

$$= 8.610 \text{ E} - 01 \text{ g/sec}$$

Calculation provided for Carbon Dioxide (CO_2) at the 200-meter location.
Appendix B provides the average adjusted emission factor of CO_2 in lb/item.

Substance-specific ambient concentrations for one item (CONC) were calculated using Equation 2. A sample calculation using this equation is provided in Example 2. Appendix B contains the estimated air concentrations for both the 100 and 200-meter locations.

$$\text{CONC} = ER_1 \cdot \frac{UC}{ER_{unit}} \quad \text{Equation 2}$$

Where:

CONC = substance concentration based on one item (g/m³)

ER₁ = emission rate for one item (g/sec)

ER_{unit} = unit emission rate as used in the model (g/sec)

UC = concentration based on the unit emission rate (g/m³)

Example 2

Sample Calculation Using Equation 2:

$$CONC = (8.610E-01) \frac{(6.870E-05)}{(1)}$$

$$= 5.915E-05 \text{ g/m}^3$$

Calculation provided for CO₂ at the 200-meter location.

6. RISK ASSESSMENT

6.1 EXPOSURE ASSUMPTIONS

Exposure assumptions were selected using a typical use scenario for the M17 during training exercises. The typical use scenario was provided by the AEC and is based on consultation with their senior training advisor (References 6, 7). The frequency of use for the M17 was required to determine how much substance an offsite resident would be exposed to in the time period of interest (i.e., acute or chronic exposure). Table 4 summarizes the general use scenario for the M17.

TABLE 4: FREQUENCY OF USE FOR THE M17

Parameter	Value Used
Number of cartridges used per year	323,952
Maximum number of cartridges used in 1- hour	3,000

6.2 TIME-AVERAGING

For the chronic assessment, time-averaged concentrations were calculated by assuming that the hypothetical offsite resident would be exposed for 30 years. This is consistent with the exposure duration used by the EPA, which assumes that the resident spends 30 years at the same residence. By using the same exposure duration,

the estimated time-averaged concentrations can be compared with their respective HBSLs, which are derived using standard EPA default assumptions.

Using the default residence time established by the EPA, the assumption was made that someone could be exposed to air emissions from 323,952 cartridges per year for 30 years. Table 5 lists the exposure parameters used to estimate concentrations for the chronic assessment. These parameters are based on the typical use scenario provided by the AEC (Table 4) and the assumptions used in the air model run.

TABLE 5: EXPOSURE PARAMETERS USED TO DETERMINE TIME-AVERAGED CHRONIC AIR CONCENTRATIONS

Exposure Parameter	Value Used	
	100 meters	200 meters
Exposure Time (ET _{ctg})	3.33 min/cartridge ¹	5 min/cartridge ¹
Exposure Frequency (EF _{ctg})	323,952 cartridges/year	
Exposure Duration (ED)	30 years ²	

¹Based on the total model time of 200 seconds (3.33 minutes) or 300 seconds (5 minutes) used in the air model run.

²EPA default value.

Chronic averaged concentrations were calculated using Equation 3. Example 3 shows how this calculation was performed using the total suspended particulates (TSP) concentration at 200 meters as an example. Since TSP is classified as a noncarcinogen, the averaging time (AT) is the same as the exposure duration.

$$C_{chronic} = \frac{CONC \cdot 10^6 \cdot ET_{ctg} \cdot EF_{ctg} \cdot ED}{525,600 \cdot AT} \quad \text{Equation 3}$$

Where:

- $C_{chronic}$ = average chronic concentration ($\mu\text{g}/\text{m}^3$)
- CONC = average modeled concentration for one cartridge (g/m^3)
- 10^6 = unit conversion ($\mu\text{g}/\text{g}$)
- ET_{ctg} = exposure time per cartridge (minutes/cartridge)
- EF_{ctg} = exposure frequency (cartridges/year)
- ED = exposure duration (years)
- 525,600 = unit conversion (minutes/year)
- AT = averaging time (years)
(carcinogenic endpoint: AT = 70 years
noncarcinogenic endpoint: AT = ED)

Example 3

Sample Calculation Using Equation 3:

$$C_{chronic(TSP)} = \frac{(4.277E-06)(10^6)(5)(323,952)(30)}{(525,600)(30)}$$

$$= 1.32E+01 \mu\text{g}/\text{m}^3$$

Appendix B provides the average modeled concentration for one cartridge (CONC). Table 5 includes the exposure parameters.

Unlike the chronic assessment, only limited guidance for evaluating acute exposures is currently available. Since many cartridges may be fired in a short period of time, however, acute exposures cannot be overlooked. For the purpose of this assessment, acute exposure is defined as a 1-hour or 15-minute exposure. The 1-hour or 15-minute acute exposure averaging times allow for comparison with guidelines developed specifically for emergency planning purposes (see discussion on acute toxicity below).

The exposure frequency is based on the number of cartridges used per 1-hour or 15 minutes depending on the guideline used for comparison. This information is based on the use scenario provided in Table 4. To estimate air concentrations for potential acute health effects, it was conservatively assumed that 3,000 M17s are fired in 1-hour. The average acute concentrations were computed using Equation 4. Example 4 contains a sample calculation at 200 meters using this equation. Since TSP does not have an ATV, aluminum (Al) is used as the example substance.

$$C_{acute} = \frac{CONC \cdot 10^6 \cdot ET_{ctg} \cdot EF_{ctg}}{60} \quad \text{Equation 4}$$

Where:

- C_{acute} = average acute concentration ($\mu\text{g}/\text{m}^3$)
- CONC = average modeled concentration for one cartridge (g/m^3)
- 10^6 = unit conversion ($\mu\text{g}/\text{g}$)
- ET_{ctg} = exposure time per cartridge (minutes/cartridge)
- EF_{ctg} = exposure frequency (cartridges/hour)*
- 60 = unit conversion (minutes/hour)

* Based on 1-hour or 15 minute (0.25 hour) ATV

Example 4
Sample Calculation Using Equation 4:

$$C_{acute(AI)} = \frac{(9.060E-09)(10^6)(5)(3,000/0.25)}{60}$$
$$= 9.06E+00 \mu\text{g}/\text{m}^3$$

Appendix B provides the average modeled concentration for one cartridge (CONC) for AI.

6.3 TOXICITY ASSESSMENT

The potential for health effects was determined by comparing time-averaged air concentrations to HBSLs and ATVs, which are developed from a substance's known toxicity. These toxicity values typically include different levels of safety factors depending on the level of confidence of the critical study. Appendix C contains a table of screening toxicity values used for the chronic and acute assessments.

6.3.1 CHRONIC ASSESSMENT

The chronic assessment was conducted using a screening approach. Using this method, a substance's estimated time-averaged air concentration was compared to its HBSL. If this ratio was less than one, no further analysis was required. This approach is conservative because the exposure assumptions used by the EPA, to establish HBSLs, assume that the resident is continuously exposed for 350 days per year (assuming 2 weeks vacation per year). In contrast, exposure to air emissions from actual training activities at a firing range is intermittent and is not likely to occur on a daily basis year round.

A hierarchy of sources was developed for selection of the HBSLs to quantitatively evaluate as many of the identified substances as possible. The hierarchy of sources used was as follows:

- Clean Air Act, EPA National Ambient Air Quality Standards (NAAQS) (Reference 11)
- EPA Region 9 Preliminary Remediation Goals (PRGs) (Reference 10)
- EPA Region 3 Risk-Based Concentrations (RBCs) (Reference 9)

Some substances have neither PRGs nor RBCs because they have their own set of regulatory standards. Under the Clean Air Act, the EPA is required to establish NAAQS for several substances considered harmful to public health and the

environment. Currently, NAAQS are available for seven substances. The NAAQS for the longer averaging time were used for the chronic assessment. Depending on the substance, this can range from an 8-hour average to an annual average. In addition, since the majority of the measured TSP was PM₁₀ (particulate matter under 10 microns in size) (Reference 2), the NAAQS for PM₁₀ was used to evaluate the potential for health effects from exposure to TSP.

Next on the hierarchy, after the NAAQS, are the EPA Region 9 PRGs and the EPA Region 3 RBCs. Since the methodology used by EPA Region 9 to develop the PRGs generally results in lower values than the EPA Region 3 RBCs, the PRGs were first on the hierarchy of sources. RBCs were used when a PRG was not available. To ensure that the most recent information was used, the Internet sites of both EPA Regions were checked. The HBSLs used for this assessment are presented in Appendix C.

Although the general approach used by both EPA Region 3 and Region 9 is the same, the exposure assumptions differ enough so that final recommended values can vary to a certain degree. In both methods, a substance's screening concentration was selected using the toxicity endpoint that derives a lower concentration. For example, if a substance has a known systemic toxicity and is a carcinogen, the screening concentration was calculated using both toxicity values. To maintain a conservative approach, EPA then selected the lower screening concentration as the recommended PRG or RBC.

Example 5 shows a sample calculation of how a substance's estimated chronic concentration was compared to its HBSL using the TSP concentration at 200 meters.

Example 5
Sample Calculation Comparing a Substance's Estimated Chronic Concentration to Its HBSL:

$$\frac{C_{\text{chronic(TSP)}}}{\text{HBSL}} = \frac{1.32\text{E} + 01}{5.00\text{E} + 01}$$
$$= 2.64\text{E}-01 < 1$$

In this case, the resulting ratio is less than one, indicating further evaluation is not necessary.

Many petroleum hydrocarbons were detected but do not have specific screening levels. Therefore, the approach recommended by the Total Petroleum Hydrocarbon Criteria Working Group (Reference 12) was adopted to evaluate petroleum hydrocarbon mixtures. Based on the working group's assessment of various hydrocarbons, it was recommended that mixtures be separated according to a

substance's number of carbons and its chemical class (i.e., aliphatic or aromatic¹). Generally, as a substance's carbon number increases, its molecular weight increases, and it is, therefore, not a substance of concern via inhalation. The working group also concluded that aromatic hydrocarbons tend to be more toxic than aliphatic hydrocarbons (Reference 12). Table 6 tabulates the inhalation toxicity values used to evaluate exposure to petroleum mixtures. To be consistent with the methodology used in this assessment, the reference concentrations (RfCs) were converted to PRGs using Region 9 exposure assumptions. The resulting PRGs were used as the HBSLs for the petroleum hydrocarbons in this assessment. These values are presented in Appendix D.

TABLE 6: SUMMARY OF RfCs USED FOR PETROLEUM HYDROCARBONS¹

Carbon Range	Aromatic Inhalation RfC (mg/m ³)	Aliphatic Inhalation RfC (mg/m ³)
C ₅ – C ₆ C _{>6} – C ₈		18.4
C _{>7} – C ₈	0.4	
C _{>8} – C ₁₀ C _{>10} – C ₁₂ C _{>12} – C ₁₆	0.2	1.0
C _{>16} – C ₂₁ C _{>21} – C ₃₅	NA	NA
¹ Reference 13 NA = not applicable for high molecular weight TPHs (Total Petroleum Hydrocarbons) (C _{>16}) because substances in this carbon range are not volatile and therefore, inhalation is not a pathway of concern.		

6.3.2 ACUTE ASSESSMENT

An established method for assessing acute health effects is not currently available. In 1995 the EPA recognized the need for acute exposure guidelines for emergency response purposes and created the National Advisory Committee for Acute Exposure Guideline Levels (AEGLs) for Hazardous Substances. Currently, AEGLs are available for only a few substances.

To overcome the absence of acute toxicity data, several state regulatory agencies have suggested that guidelines developed for emergency purposes be used in the interim. Although suggestions have been made to use occupational exposure limits (OELs) by applying additional safety factors (References 14, 15), OELs were not used in this assessment because they introduce even more uncertainty than the use of

¹ Aliphatic hydrocarbons are hydrocarbons in which the carbon atoms are joined by single covalent bonds consisting of two shared electrons (e.g., butane). Aromatic hydrocarbons have ring structures (e.g., benzene) (Reference 13).

emergency guidelines. The OELs are designed to protect the workplace environment, and assume 8 hours a day, 5 days a week exposures. By definition, these exposures are more chronic than acute.

In comparison, emergency planning guidelines are more appropriate because they are typically developed for exposures of 1-hour or less. In addition, safety factors are included as part of the guideline development, so that the values would be protective of the general population.

Emergency Response Planning Guidelines (ERPGs) published by the American Industrial Hygiene Association (AIHA) (Reference 16) and the Temporary Emergency Exposure Limits (TEELs) developed by the U.S. Department of Energy (DOE) (Reference 17) were used for this assessment, specifically the ERPG-1s and the TEEL-1s. Since TEEL-1s are intended for exposures up to 15-minutes, air concentrations compared to TEELs were averaged over a 15-minute period. Air concentrations compared to ERPGs and AEGLs were averaged over 1-hour, as these values are intended for 1-hour exposures.

For this assessment, the hierarchy of sources for ATV selection was as follows with each ATV defined below:

- EPA AEGL-1. "AEGL-1 is the airborne concentration of a substance above which it is predicted that the general population, including susceptible individuals, could experience notable discomfort, irritation, or certain asymptomatic, nonsensory effects. However, the effects are not disabling and are transient and reversible upon cessation of exposure."
- AIHA ERPG-1. "The maximum concentration in air below which it is believed nearly all individuals could be exposed for up to 1- hour without experiencing other than mild transient adverse health effects or perceiving a clearly defined objectionable odor."
- DOE TEEL-1. "The maximum concentration in air below which it is believed nearly all individuals could be exposed without experiencing other than mild transient adverse health effects or perceiving a clearly defined objectionable odor."

AEGLs were used first when available since they are developed specifically for the purpose of acute exposure assessments. The ERPGs were selected next, prior to a substance's TEEL, because they are vigorously reviewed before they are published whereas the TEELs are not.

Example 6 shows a sample calculation of how a substance's estimated acute concentration was compared to its ATV using the aluminum concentration at 200 meters.

Example 6

Sample Calculation Comparing a Substance's Estimated Acute Concentration to Its ATV:

$$\frac{C_{acute(AI)}}{ATV} = \frac{9.06E+00}{3.00E+04}$$
$$= 3.02E-04 < 1$$

In this example with AI, the ratio is less than one, indicating that further analysis is not necessary.

7. RISK CHARACTERIZATION

As previously described, the exposure assessment included calculations of time-averaged concentrations for both long-term (chronic) and short-term (acute) exposures. Using a screening approach, a substance's estimated time-averaged air concentration was then compared to chronic HBSLs or ATVs. The comparison was made using the ratio of the HBSL or ATV to the estimated concentration. This approach is conservative because the exposure assumptions used by the EPA, to establish HBSLs and ATVs, are likely to overestimate the exposures experienced by offsite residents living near firing ranges.

If this ratio was less than one, no further evaluation was needed. If the chronic or acute averaged concentrations ($C_{chronic}$ and C_{acute}) were greater than the screening levels, resulting in a ratio greater than one, further evaluation would be warranted to determine the potential for health effects. Note that concentrations greater than the screening levels do not indicate an onset of health effects, but rather, the potential for such.

The chronic and acute assessments were conducted as outlined in Section 6.3. Appendix D presents results from the M17 risk characterization.

7.1 CHRONIC HEALTH RISK

The chronic assessment, at the 100-meter downwind hypothetical offsite resident location, indicated that estimated concentrations of hydrogen cyanide, $PM_{2.5}$, barium, and 1,3-butadiene from the M17 emissions were greater than the HBSLs. The ratios of all other substances to their HBSLs were below one. Estimated concentrations of M17 air emissions were remodeled to a distance 200 meters downwind from the firing location. The results showed that the estimated concentrations of the four substances, which were greater than the HBSLs at 100 meters, all decreased to safe levels. The estimated concentrations for all other substances were further reduced with all ratios below one.

The ratio of the estimated hydrogen cyanide concentration to its HBSL was 2.14 at the 100-meter location. Hydrogen cyanide is a colorless gas with a faint, bitter, almond-like odor. It is naturally produced by some microorganisms and can be found in a number of foods and plants. Examples of industrial uses of hydrogen cyanide include chemical production, photographic development, and some mining processes. There are no reports that hydrogen cyanide causes cancer. Long-term inhalation of hydrogen cyanide may result in breathing difficulties, heart pains, vomiting, blood changes, headaches, and enlargement of the thyroid gland (Reference 17).

The ratio of the estimated $PM_{2.5}$ concentration to its HBSL was 1.12 at the 100-meter location. $PM_{2.5}$ refers to particulate matter that is 2.5 microns or less in size. The term particulate matter is used to refer to a mixture of solid particles and liquid droplets found in the air. $PM_{2.5}$ is a product of fuel combustion from sources such as automobiles, power plants, industrial processes, wood burning, and diesel-powered vehicles (Reference 18). $PM_{2.5}$ can penetrate deep into the lungs due to its small size. Inhalation of $PM_{2.5}$ has been linked to a series of health effects including asthma, bronchitis, and other respiratory system effects. Those most at risk are the elderly, children, and people with existing health conditions such as respiratory and cardiopulmonary diseases (References 19).

The ratio of the estimated barium concentration to its HBSL was 1.57 at the 100-meter location. Barium is a silvery-white metal that is naturally found in the environment. Barium compounds are also produced and used by industry and in medicine. The respiratory system is thought to be the primary system affected from barium inhalation. Studies have suggested that inhalation of barium may also be associated with systemic effects such as hypertension (Reference 20).

The ratio of the estimated 1,3-Butadiene concentration to its HBSL was 1.41 at the 100-meter location. 1,3-Butadiene is a colorless gas with a mild gasoline-like odor and is made from the processing of petroleum. 1,3-Butadiene is used to make rubber and plastics. 1,3-Butadiene is classified as a known human carcinogen based on occupational exposures and animal studies. Chronic inhalation has been associated with increased risk for lymph, blood, and digestive system cancers, diabetes mellitus, and arteriosclerosis (Reference 21).

7.2 ACUTE HEALTH RISK

The acute assessment, at the 100-meter downwind hypothetical offsite resident location, indicated that estimated copper and lead concentrations from the M17 emissions were greater than their ATVs. The ratios of all other substances were less than one. Estimated concentrations were remodeled to a distance 200 meters downwind from the firing location. The results showed that the estimated concentrations of copper and lead both decreased to safe levels. The estimated concentrations for all other substances were further reduced with all ratios below one.

The ratio of estimated copper concentration to the ATV was 1.10 at the 100-meter location. Copper is a naturally occurring reddish metal found in the earth's crust. Copper is widely used because it can be easily molded or shaped and may be mixed with other metals. The acute health effects from inhalation exposure to copper are not well known. However, some studies have indicated that short-term inhalation of copper may result in nose and respiratory irritation (Reference 22).

The ratio of estimated lead concentration to the ATV was 2.29. Lead is a naturally occurring bluish-gray metal found in the earth's crust in small amounts. It is commonly used in lead-acid batteries for automotive and industrial applications. Exposure to lead in the air primarily results from emissions from industrial processes. The main target for lead toxicity is the nervous system. Information is limited on the health effects from short-term inhalation exposure to lead (Reference 23).

Again, it should be noted that an estimated concentration, which is greater than the HBSL or ATV, does not indicate an onset of health effects, but rather the potential for such.

7.3 FACT SHEET

Appendix E includes a copy of the fact sheet submitted to the AEC. The fact sheet used results from this assessment to address health concerns related to inhalation of M17 air emissions.

8. UNCERTAINTY DISCUSSION

The limitations inherent in modeling and the added conservatism of the assessment contribute to the uncertainty of the assessment results. The risk assessment methodology typically includes safety factors that are embedded in the toxicity data to ensure adequate protection of the general population, particularly, susceptible individuals such as the sick, elderly, and children. Table 7 identifies areas of uncertainty associated with this assessment.

TABLE 7: TYPES OF UNCERTAINTY

Issue	Uncertainty	Direction of Effect
Emissions Modeling		
Modeled versus real-time sampling	The air concentrations in this assessment were modeled. Actual air concentrations taken from the field may be higher or lower.	Varies
Frequency of use for the M17	Actual frequency of use for these munitions during training exercises may be different from those stated in this report.	Varies
Hypothetical offsite resident assumed to be located directly downwind	Unless the area around the training facility is populated, the chances that a person living directly downwind is low.	Overestimates
Use of worst-case meteorological conditions	To ensure that this assessment is applicable to most training areas, worst-case meteorological conditions were used in the air model.	Overestimates
Exposure Assessment		
Estimating time-averaged concentrations	Actual exposure from the M17 is intermittent. If one were to plot a person's exposure profile, the plot would consist of a series of spikes. Since current risk assessment methodology does not allow the assessment of the potential for health effects as a function of time, a single concentration, averaged over the exposure duration was used. In this assessment, the exposure durations used were 30 years and 1-hour or 15 minutes.	Varies
Comparing estimated concentration to established screening levels	The Region 3 and Region 9 HBSLs were developed assuming that the resident is exposed 350 days per year. It is unlikely for training with the M17 to occur for 350 days per year at a particular firing range.	Overestimates
Comparing estimated concentrations to established screening levels	Comparison to screening levels does not account for possible cumulative effects of exposure to more than one substance.	Underestimates

TABLE 7: TYPES OF UNCERTAINTY

Issue	Uncertainty	Direction of Effect
Screening assessment versus calculating an average daily intake	Calculating an average daily intake allows the use of scenario-specific assumptions. However, unless the ratio of concentration to screening level approaches one, a screening assessment is useful as a first-cut evaluation.	Varies
Exposure to other munitions	Other munitions are typically used during the same training exercise. These items may contain similar or different substances from those detected in the M17 emissions.	Underestimates
Toxicity Assessment		
Lack of toxicity data	Some substances were not quantitatively evaluated because they have no known toxicity data.	Underestimates
Modifying and uncertainty factors for toxicity data	Modifying factors and uncertainty factors of varying degree are typically applied to toxicological values. These factors are used to conservatively account for extrapolating from animal studies for human health evaluation, and to conservatively account for variation in human populations.	Overestimates

9. CONCLUSION

Using conservative assumptions, the assessment indicated that residents who live as close as 200 meters downwind from the firing location are safe from breathing air emissions from the M17. It is believed that the assumptions contained in this analysis are conservative enough to be protective of all the population including the sick, elderly, and children.

10. RECOMMENDATIONS

At installations where offsite residents are located less than 200-meters from the M17 firing location, a more site-specific evaluation is recommended. However, it should be noted that most training areas are located at least 1,000 meters (over half a mile) away from populated areas.

The results from this assessment are intended for a hypothetical training facility, and actual results can vary depending on site-specific conditions. This assessment used


conservative assumptions (e.g., worst-case meteorological conditions, receptor located directly downwind, etc.) and it is believed that most site-specific analyses would result in even lower concentrations. Therefore, the results from this assessment should be applicable to most training facilities unless site-specific conditions vary significantly.

11. POINT OF CONTACT


Questions about this report may be directed to Ms. Joleen Mobley at (800) 222-9698 (ext 2953) or (410) 436-2953.

PREPARED BY:

APPROVED BY:


for LAURA L. PETERS
Environmental Engineer
Environmental Health Risk Assessment
Program


DAVID L. DAUGHDRILL
Program Manager
Environmental Health Risk Assessment


JOLEEN MOBLEY
Environmental Scientist
Environmental Health Risk Assessment
Program


STAFFORD D.F.R. COAKLEY
Environmental Engineer
Environmental Health Risk Assessment
Program

APPENDIX A

REFERENCES

1. U.S. Army (1994). *Technical Manual, Army Ammunition Data Sheets for Small Caliber Ammunition*. TM-43-0001-27.
2. U.S. Army. Email communication between Ms. Tamera Clark-Rush, AEC, and Ms. Hsieng-Ye Chang, USACHPPM. Subject: Electronic copy of Firing Point Emission Study Series 3 Emission Factors, 16 August 2000.
3. USACHPPM (Aug. 2000). *Ambient Air Quality Consultation NO. 43-EL-1485-00 Air Dispersion Modeling Evaluation For Military Munitions*, Aberdeen Proving Ground, MD.
4. Bowman Environmental, Inc. (1999). *INPUFF2, Multiple Source Integrated Puff Model*, Version 4.1.
5. Title 40, Code of Federal Regulations, Part 68 (40 CFR 68), Chemical Accident Prevention Provisions, 1 July 1998.
6. U.S. Army. Email communication between Ms. Tamera Clark-Rush, AEC, and Ms. Hsieng-Ye Chang, USACHPPM. Subject: Scenarios (Small Caliber Ammunition), 12 September 2000.
7. Army Training Evaluation Protocol (ARTEP) 7-20-MTP, *Mission Training Plan for the Infantry Battalion*.
8. EPA (April 2000). *Region 3 Risk Based Concentration (RBC) Tables*. Available online at www.epa.gov/reghwmd/risk/riskmenu.htm
9. EPA (Nov. 2000). *Region 9 Preliminary Remediation Goals (PRG)*. Available online at www.epa.gov/region09/waste/sfund/prg/index.html
10. EPA. *National Ambient Air Quality Standards*. Available online at <http://www.epa.gov/airprogm/airs/criteria.html>
11. Total Petroleum Hydrocarbon Criteria Working Group (1997). *Development of Fraction Specific Reference Doses (RfDs) and Reference Concentrations (RfCs) for Total Petroleum Hydrocarbons (TPH)*, Volume 4. Amherst Scientific Publishers. Amherst, MA.
12. Manahan, Stanley (1994). *Environmental Chemistry*. Sixth edition. CRC Press, Inc. Boca Raton, FL.
13. U.S. Army (1996). *Final Screening Risk Assessment for the Anniston Chemical Agent Disposal Facility at the Anniston Army Depot, Alabama*. Revision No. 5. Prepared by the U.S. Army Center for Health Promotion and Preventive Medicine for the Program Manager for Chemical Demilitarization. Aberdeen Proving Ground, Maryland.

14. U.S. Army (1997). *Final Screening Risk Assessment for the Pine Bluff Chemical Agent Disposal Facility at the Pine Bluff Arsenal, Arkansas*. Revision No. 1. Prepared by the U.S. Army Center for Health Promotion and Preventive Medicine for the Program Manager for Chemical Demilitarization. Aberdeen Proving Ground, Maryland.
15. American Industrial Hygiene Association (AIHA) (1999). *Emergency Response Planning Guidelines*. AIHA Press, Fairfax, VA.
16. Department of Energy (1998). *Temporary Emergency Exposure Limits*, Revision 15. <http://www.scapa.bnl.gov>.
17. Agency for Toxic Substances and Disease Registry (1995). *Toxicological Profile for Cyanide*. Prepared by Clement International Corporation and Syracuse Research Corporation for ATSDR.
18. U.S. Environmental Protection Agency (2001). *PM_{2.5} Objectives and History*. Region 4. Available online at www.epa.gov/region4/sesd/pm25/p2.htm.
19. U.S. Environmental Protection Agency (1999). *Air Criteria Document for Particulate Matter, External Draft Review*. EPA 600/P-99/002a-c. Office of Research and Development, Washington, DC 20460.
20. Agency for Toxic Substances and Disease Registry (1992). *Toxicological Profile for Barium*. Prepared by Clement International Corporation for ATSDR.
21. Agency for Toxic Substances and Disease Registry (1992). *Toxicological Profile for 1,3-Butadiene*. Prepared by Clement International Corporation and Syracuse Research Corporation for ATSDR.
22. Agency for Toxic Substances and Disease Registry (1990). *Toxicological Profile for Copper*. Prepared by Syracuse Research Corporation for ATSDR.
23. Agency for Toxic Substances and Disease Registry (1999). *Toxicological Profile for Lead*.

APPENDIX B

AIR DISPERSION MODELING OUTPUT DATA

Table B-1: Air Modeling Output Data for the Cartridge, 0.50 Caliber Tracer, M17 (M2) - 100 meter location

Cartridge, 0.50 caliber, Tracer, M17 (M2)									
DODIC: A571									
Number of items tested = 3									
Net Explosive Weight (lbs) = 3.21E-02									
Compound	ATC Firing Test Results				Total Mass of Substance Emitted (grams/item) M	Average Modeled Concentration for One Item (grams/m³) CONC	Pollutant Emission Rate for One Item (g/sec) ER ₁		
	Measured Actual Concentration (mg/m³)	Measured Background Concentration (mg/m³)	Average Adjusted Emission Factor EF	Average Adjusted Emission Factor (lb/lb NEW)					
	Permanent Gases								
	Ammonia (NH ₃)	2.03E+01	NA	2.20E-04	6.83E-03	9.96E-02	8.147E-06	4.98E-02	
	Carbon Dioxide (CO ₂)	5.27E+02	NA	5.69E-03	1.77E-01	2.58E+00	2.113E-04	1.29E+00	
	Carbon Monoxide (CO)	1.01E+03	NA	1.09E-02	3.40E-01	4.96E+00	4.059E-04	2.48E+00	
	Oxides of Nitrogen (NOx)	4.45E+01	NA	4.81E-04	1.50E-02	2.18E-01	1.784E-05	1.09E-01	
	Sulfur Dioxide (SO ₂)	2.62E-01	NA	2.83E-06	8.81E-05	1.29E-03	1.051E-07	6.43E-04	
	Acid Gases								
	Hydrogen Fluoride	2.20E-01	2.20E-01	ND	ND	ND	ND	ND	
Hydrogen Chloride	2.20E-01	2.10E-01	ND	ND	ND	ND	ND		
Hydrogen Bromide	2.10E-01	2.10E-01	ND	ND	ND	ND	ND		
Nitric Acid	2.10E-01	2.10E-01	ND	ND	ND	ND	ND		
Phosphoric Acid	2.10E-01	2.10E-01	ND	ND	ND	ND	ND		
Sulfuric Acid	2.55E-01	2.10E-01	4.66E-06	1.45E-04	2.11E-03	1.728E-07	1.06E-03		
Cyanide									
Particulate Cyanide	5.80E-01	1.20E-02	7.07E-06	2.20E-04	3.20E-03	2.622E-07	1.60E-03		
Hydrogen Cyanide	7.22E+00	1.30E-02	8.78E-05	2.73E-03	3.98E-02	3.257E-06	1.99E-02		
Particulates									
Total Suspended Particulate	3.39E+01	NA	4.12E-04	1.28E-02	1.87E-01	1.528E-05	9.34E-02		
Particulate Matter <10 microns	3.11E+01	NA	3.78E-04	1.18E-02	1.71E-01	1.403E-05	8.57E-02		
Particulate Matter <2.5 microns	1.81E+01	NA	2.20E-04	6.86E-03	1.00E-01	8.178E-06	5.00E-02		
Metals									
Aluminum	7.19E-02	4.30E-02	8.72E-07	2.71E-05	3.96E-04	3.236E-08	1.98E-04		
Antimony	5.65E-01	4.30E-02	6.88E-06	2.14E-04	3.12E-03	2.551E-07	1.56E-03		
Arsenic	1.05E-02	1.08E-02	ND	ND	ND	ND	ND		
Barium	8.81E-01	4.30E-02	1.07E-05	3.34E-04	4.87E-03	3.985E-07	2.44E-03		
Beryllium	4.21E-02	4.30E-02	ND	ND	ND	ND	ND		
Cadmium	4.21E-02	4.30E-02	ND	ND	ND	ND	ND		
Calcium	4.62E-01	1.96E-01	3.50E-06	1.09E-04	1.59E-03	1.297E-07	7.93E-04		

Table B-1: Air Modeling Output Data for the Cartridge, 0.50 Caliber Tracer, M17 (M2) - 100 meter location

Cartridge, 0.50 caliber, Tracer, M17 (M2)					Number of Rounds (I):		1 round	
DODIC: A571					Release duration (t):		2 seconds	
Number of items tested = 3					Unit Concentration (UC):		1.636E-04 (g/m³)/(g/s)	
Net Explosive Weight (lbs) = 3.21E-02								
Compound	ATC Firing Test Results							
	Measured Actual Concentration (mg/m³)	Measured Background Concentration (mg/m³)	Average Adjusted Emission Factor (lb/item) EF	Average Adjusted Emission Factor (lb/lb NEW)	Total Mass of Substance Emitted (grams/item) M	Average Modeled Concentration for One Item (grams/m³) CONC	Pollutant Emission Rate for One Item (g/sec) ER ₁	
	Chromium	4.21E-02	4.30E-02	ND	ND	ND	ND	ND
	Cobalt	4.21E-02	4.30E-02	ND	ND	ND	ND	ND
	Copper	1.11E+01	9.55E-02	1.33E-04	4.15E-03	6.06E-02	4.953E-06	3.03E-02
	Lead	1.15E+00	4.30E-02	1.39E-05	4.33E-04	6.31E-03	5.162E-07	3.16E-03
	Magnesium	2.26E-01	4.30E-02	2.76E-06	8.58E-05	1.25E-03	1.023E-07	6.26E-04
	Manganese	4.21E-02	4.30E-02	ND	ND	ND	ND	ND
	Nickel	4.21E-02	4.30E-02	ND	ND	ND	ND	ND
	Selenium	1.05E-02	1.08E-02	ND	ND	ND	ND	ND
	Silver	4.21E-02	4.30E-02	ND	ND	ND	ND	ND
	Thallium	4.21E-02	4.30E-02	ND	ND	ND	ND	ND
	Vanadium	4.21E-02	4.30E-02	ND	ND	ND	ND	ND
	Zinc	1.81E+00	4.30E-02	2.20E-05	6.84E-04	9.97E-03	8.159E-07	4.99E-03
	TO-11 Carbonyls							
	Formaldehyde	2.46E-02	1.23E-01	2.98E-07	9.27E-06	1.35E-04	1.106E-08	6.76E-05
	Acetaldehyde	1.80E-01	1.80E-01	ND	ND	ND	ND	ND
	Acetone	1.19E+00	1.19E+00	ND	ND	ND	ND	ND
	Acrolein	2.29E-01	2.29E-01	ND	ND	ND	ND	ND
	Propionaldehyde	2.37E-01	2.37E-01	ND	ND	ND	ND	ND
Crotonaldehyde	2.87E-01	2.87E-01	ND	ND	ND	ND	ND	
Butyraldehyde	2.95E-01	2.95E-01	ND	ND	ND	ND	ND	
Benzaldehyde	4.34E-01	4.34E-01	ND	ND	ND	ND	ND	
Isovaleraldehyde	3.52E-01	3.52E-01	ND	ND	ND	ND	ND	
Valeraldehyde	3.52E-01	3.52E-01	ND	ND	ND	ND	ND	
o,m,p-Tolualdehyde	4.91E-01	4.91E-01	ND	ND	ND	ND	ND	
Hexaldehyde	4.10E-01	4.10E-01	ND	ND	ND	ND	ND	
2,5-Dimethylbenzaldehyde	4.10E-01	4.10E-01	ND	ND	ND	ND	ND	
VOCs								
Propene	5.94E-02	3.44E-04	7.15E-07	2.23E-05	3.25E-04	2.655E-08	1.62E-04	
Dichlorodifluoromethane	2.97E-03	3.46E-03	ND	ND	ND	ND	ND	

Table B-1: Air Modeling Output Data for the Cartridge, 0.50 Caliber Tracer, M17 (M2) - 100 meter location

Compound	Cartridge, 0.50 caliber, Tracer, M17 (M2)				Number of Rounds (I):		1 round
	DODIC: A571				Release duration (I):		2 seconds
	Number of items tested = 3				Unit Concentration (UC):		1.636E-04 (g/m ³)/(g/s)
	Net Explosive Weight (lbs) = 3.21E-02						
	ATC Firing Test Results						
	Measured Actual Concentration (mg/m ³)	Measured Background Concentration (mg/m ³)	Average Adjusted Emission Factor (lb/item) EF	Average Adjusted Emission Factor (lb/lb NEW)	Total Mass of Substance Emitted (grams/item) M	Average Modeled Concentration for One Item (grams/m ³) CONC	Pollutant Emission Rate for One Item (g/sec) ER ₁
Chlorodifluoromethane	3.54E-03	3.54E-03	ND	ND	ND	ND	ND
Freon 114	6.99E-03	6.99E-03	ND	ND	ND	ND	ND
Chloromethane	1.65E-03	1.45E-03	4.41E-09	1.37E-07	2.00E-06	1.636E-10	1.00E-06
Vinyl Chloride	2.56E-03	2.56E-03	ND	ND	ND	ND	ND
1,3-Butadiene	1.33E-02	2.21E-03	1.61E-07	5.00E-06	7.29E-05	5.966E-09	3.65E-05
Bromomethane	3.88E-03	3.88E-03	ND	ND	ND	ND	ND
Chloroethane	2.64E-03	2.64E-03	ND	ND	ND	ND	ND
Dichlorofluoromethane	4.21E-03	4.21E-03	ND	ND	ND	ND	ND
Trichlorofluoromethane	1.69E-03	1.69E-03	2.22E-09	6.91E-08	1.01E-06	8.239E-11	5.04E-07
Pentane	8.85E-04	2.95E-03	1.07E-08	3.34E-07	4.87E-06	3.986E-10	2.44E-06
Acrolein	2.29E-03	2.29E-03	ND	ND	ND	ND	ND
1,1-Dichloroethene	4.05E-03	4.05E-03	ND	ND	ND	ND	ND
Freon 113	7.68E-03	7.68E-03	ND	ND	ND	ND	ND
Acetone	1.84E-01	1.07E-01	1.10E-06	3.41E-05	4.98E-04	4.072E-08	2.49E-04
Methyl Iodide	5.81E-03	5.81E-03	ND	ND	ND	ND	ND
Carbon Disulfide	3.11E-03	3.11E-03	ND	ND	ND	ND	ND
Acetonitrile	9.82E-02	8.39E-03	1.10E-06	3.41E-05	4.97E-04	4.067E-08	2.49E-04
3-Chloropropene	3.13E-03	3.13E-03	ND	ND	ND	ND	ND
Methylene Chloride	1.39E-01	4.52E-02	1.20E-06	3.74E-05	5.45E-04	4.458E-08	2.72E-04
tert-Butyl Alcohol	3.03E-03	3.03E-03	ND	ND	ND	ND	ND
Acrylonitrile	1.95E-02	2.17E-03	2.36E-07	7.35E-06	1.07E-04	8.765E-09	5.36E-05
trans-1,2-Dichloroethene	3.96E-03	3.96E-03	ND	ND	ND	ND	ND
Methyl t-Butyl Ether	3.61E-03	3.61E-03	ND	ND	ND	ND	ND
Hexane	9.52E-02	1.37E-01	ND	ND	ND	ND	ND
1,1-Dichloroethane	3.97E-03	3.97E-03	ND	ND	ND	ND	ND
Vinyl Acetate	3.52E-03	3.52E-03	ND	ND	ND	ND	ND
cis-1,2-Dichloroethene	3.96E-03	3.96E-03	ND	ND	ND	ND	ND
2-Butanone	2.95E-03	2.95E-03	ND	ND	ND	ND	ND
Ethyl Acetate	1.08E-02	3.60E-03	1.30E-07	4.04E-06	5.89E-05	4.821E-09	2.95E-05

Table B-1: Air Modeling Output Data for the Cartridge, 0.50 Caliber Tracer, M17 (M2) - 100 meter location

Cartridge, 0.50 caliber, Tracer, M17 (M2)					Number of Rounds (I):		1 round	
					Release duration (t):		2 seconds	
DODIC: A571					Unit Concentration (UC):		1.636E-04 (g/m³)/(g/s)	
Number of items tested = 3								
Net Explosive Weight (lbs) = 3.21E-02								
Air Sampling Test Results:								
Compound	Measured Actual Concentration (mg/m³)	Measured Background Concentration (mg/m³)	Average Adjusted Emission Factor (lb/item) EF	Average Adjusted Emission Factor (lb/lb NEW)	Total Mass of Substance Emitted (grams/item) M	Average Modeled Concentration for One Item (grams/m³) CONC	Pollutant Emission Rate for One Item (g/sec) ER₁	
Methyl Acrylate	3.52E-03	3.52E-03	ND	ND	ND	ND	ND	ND
Chloroform	4.88E-03	4.88E-03	ND	ND	ND	ND	ND	ND
1,1,1-Trichloroethane	3.82E-03	3.82E-03	4.91E-09	1.53E-07	2.23E-06	1.820E-10	1.11E-06	1.11E-06
Carbon Tetrachloride	6.29E-03	6.29E-03	ND	ND	ND	ND	ND	ND
1,2-Dichloroethane	6.07E-03	4.05E-03	7.34E-08	2.28E-06	3.33E-05	2.725E-09	1.67E-05	1.67E-05
Benzene	3.12E-01	9.59E-04	3.76E-06	1.17E-04	1.71E-03	1.396E-07	8.53E-04	8.53E-04
Isooctane	4.67E-03	4.67E-03	ND	ND	ND	ND	ND	ND
Heptane	4.10E-03	2.46E-03	ND	ND	ND	ND	ND	ND
Trichloroethane	4.88E-03	4.88E-03	ND	ND	ND	ND	ND	ND
Ethyl Acrylate	4.09E-03	4.09E-03	ND	ND	ND	ND	ND	ND
1,2-Dichloropropane	4.62E-03	4.62E-03	ND	ND	ND	ND	ND	ND
Methyl Methacrylate	4.09E-03	4.09E-03	ND	ND	ND	ND	ND	ND
Dibromomethane	7.11E-03	7.11E-03	ND	ND	ND	ND	ND	ND
1,4-Dioxane	3.60E-03	3.60E-03	ND	ND	ND	ND	ND	ND
Bromodichloromethane	6.70E-03	6.70E-03	ND	ND	ND	ND	ND	ND
4-Methyl-2-Pentanone	4.10E-03	4.10E-03	ND	ND	ND	ND	ND	ND
Toluene	3.58E-02	7.54E-04	4.26E-07	1.33E-05	1.93E-04	1.581E-08	9.66E-05	9.66E-05
Octane	4.67E-03	4.67E-03	ND	ND	ND	ND	ND	ND
trans-1,3-Dichloropropene	4.54E-03	4.54E-03	ND	ND	ND	ND	ND	ND
Ethyl Methacrylate	4.67E-03	4.67E-03	ND	ND	ND	ND	ND	ND
1,1,2-Trichloroethane	5.46E-03	5.46E-03	ND	ND	ND	ND	ND	ND
Tetrachloroethene	6.78E-03	6.78E-03	ND	ND	ND	ND	ND	ND
2-Hexanone	4.10E-03	4.10E-03	ND	ND	ND	ND	ND	ND
Dibromochloromethane	8.52E-03	8.52E-03	ND	ND	ND	ND	ND	ND
1,2-Dibromoethane	7.68E-03	7.68E-03	ND	ND	ND	ND	ND	ND
Chlorobenzene	4.60E-03	4.60E-03	ND	ND	ND	ND	ND	ND
1,1,1,2-Tetrachloroethane	6.87E-03	6.87E-03	ND	ND	ND	ND	ND	ND
Ethylbenzene	2.82E-03	4.34E-03	3.42E-08	1.06E-06	1.55E-05	1.268E-09	7.75E-06	7.75E-06
m/p-Xylene	6.51E-03	1.30E-03	6.47E-08	2.01E-06	2.93E-05	2.400E-09	1.47E-05	1.47E-05

Table B-1: Air Modeling Output Data for the Cartridge, 0.50 Caliber Tracer, M17 (M2) - 100 meter location

Cartridge, 0.50 caliber, Tracer, M17 (M2)					Number of Rounds (I):		1 round	
DODIC: A571					Release duration (I):		2 seconds	
Number of items tested = 3					Unit Concentration (UC):		1.636E-04 (g/m ³)/(g/s)	
Net Explosive Weight (lbs) = 3.21E-02								
Compound	At-Cartridge Test Results							
	Measured Actual Concentration (mg/m ³)	Measured Background Concentration (mg/m ³)	Average Adjusted Emission Factor (lb/item) EF	Average Adjusted Emission Factor (lb/lb NEW)	Total Mass of Substance Emitted (grams/item) M	Average Modeled Concentration for One Item (grams/m ³) CONC	Pollutant Emission Rate for One Item (g/sec) ER _i	
	o-Xylene	4.34E-03	4.34E-03	5.27E-08	1.64E-06	2.39E-05	1.955E-09	1.19E-05
	Styrene	1.07E-02	4.26E-03	1.29E-07	4.01E-06	5.85E-05	4.786E-09	2.93E-05
	Bromoform	1.03E-02	1.03E-02	ND	ND	ND	ND	ND
	Cumene	4.92E-03	4.92E-03	ND	ND	ND	ND	ND
	1,1,2,2-Tetrachlorethane	6.87E-03	6.87E-03	ND	ND	ND	ND	ND
	1,2,3-Trichloropropane	6.03E-03	6.03E-03	ND	ND	ND	ND	ND
	Bromobenzene	6.42E-03	6.42E-03	ND	ND	ND	ND	ND
	4-Ethyltoluene	4.92E-03	4.92E-03	ND	ND	ND	ND	ND
	1,3,5-Trimethylbenzene	4.92E-03	4.92E-03	ND	ND	ND	ND	ND
	Alpha Methyl Styrene	4.83E-03	4.83E-03	ND	ND	ND	ND	ND
	1,2,4-Trimethylbenzene	9.83E-04	4.92E-03	1.18E-08	3.68E-07	5.36E-06	4.384E-10	2.68E-06
	1,3-Dichlorobenzene	6.01E-03	6.01E-03	ND	ND	ND	ND	ND
	1,4-Dichlorobenzene	6.01E-03	6.01E-03	ND	ND	ND	ND	ND
	Benzyl Chloride	5.18E-03	5.18E-03	ND	ND	ND	ND	ND
	1,2-Dichlorobenzene	6.01E-03	6.01E-03	ND	ND	ND	ND	ND
Hexachlorethane	9.68E-03	9.68E-03	ND	ND	ND	ND	ND	
1,2,4-Trichlorobenzene	7.42E-03	7.42E-03	ND	ND	ND	ND	ND	
Hexachlorobutadiene	1.07E-02	1.07E-02	ND	ND	ND	ND	ND	
VOC Tentatively Identified Compounds (TICs)								
Hydrocarbons								
Methane	4.10E+00	1.10E+00	3.79E-05	1.18E-03	1.72E-02	1.405E-06	8.59E-03	
Ethylene	2.00E-01	2.29E-02	2.43E-06	7.57E-05	1.10E-03	9.024E-08	5.52E-04	
Acetylene	2.74E-02	2.13E-02	3.32E-07	1.03E-05	1.51E-04	1.234E-08	7.54E-05	
Ethane	1.13E-01	2.46E-02	1.37E-06	4.27E-05	6.23E-04	5.099E-08	3.12E-04	
Propylene	5.42E-02	3.44E-02	6.59E-07	2.05E-05	2.99E-04	2.444E-08	1.49E-04	
Propane	3.61E-02	3.61E-02	ND	ND	ND	ND	ND	
Propyne	3.20E-02	3.20E-02	ND	ND	ND	ND	ND	
Isobutane	4.75E-02	4.75E-02	ND	ND	ND	ND	ND	
1-Butene/Isobutylene	9.18E-02	9.18E-02	ND	ND	ND	ND	ND	

Table B-1: Air Modeling Output Data for the Cartridge, 0.50 Caliber Tracer, M17 (M2) - 100 meter location

Compound	Cartridge, 0.50 caliber, Tracer, M17 (M2)				Number of Rounds (I):		Pollutant Emission Rate for One Item (g/sec) ER ₁
	DODIC: A571				Release duration (t):		
	Number of items tested = 3				Unit Concentration (UC):		
	Net Explosive Weight (lbs) = 3.21E-02				1.636E-04 (g/m ³)/(g/s)		
	Net Explosive Weight (lbs) = 3.21E-02				1.636E-04 (g/m ³)/(g/s)		
Compound	ATC Firing Test Results				Total Mass of Substance Emitted (grams/item) M	Average Modeled Concentration for One Item (grams/m ³) CONC	Pollutant Emission Rate for One Item (g/sec) ER ₁
	Measured Actual Concentration (mg/m ³)	Measured Background Concentration (mg/m ³)	Average Adjusted Emission Factor (lb/item) EF	Average Adjusted Emission Factor (lb/lb NEW)			
	1,3-Butadiene/butane	4.59E-02	4.59E-02	ND	ND	ND	ND
	cis-butene	4.59E-02	4.59E-02	ND	ND	ND	ND
	1-Butyne	4.59E-02	4.59E-02	ND	ND	ND	ND
	trans-Butene	4.59E-02	4.59E-02	ND	ND	ND	ND
	2-Butyne	4.42E-02	4.42E-02	ND	ND	ND	ND
	n-Pentane	5.90E-02	5.90E-02	ND	ND	ND	ND
	n-Hexane	1.99E-01	7.05E-02	2.41E-06	7.50E-05	1.09E-03	5.47E-04
	SVOCs						
	N-nitrosodimethylamine	1.81E-02	1.83E-02	ND	ND	ND	ND
	Bis(2-chloroethyl)ether	1.81E-02	1.83E-02	ND	ND	ND	ND
	Phenol	1.81E-02	1.83E-02	ND	ND	ND	ND
	2-chlorophenol	1.81E-02	1.83E-02	ND	ND	ND	ND
	1,3-dichlorobenzene	1.81E-02	1.83E-02	ND	ND	ND	ND
	1,4-dichlorobenzene	1.81E-02	1.83E-02	ND	ND	ND	ND
	1,2-dichlorobenzene	1.81E-02	1.83E-02	ND	ND	ND	ND
	Benzyl alcohol	1.81E-02	1.83E-02	ND	ND	ND	ND
	Bis(2-chloroisopropyl)ether	1.81E-02	1.83E-02	ND	ND	ND	ND
	2-methylphenol	1.81E-02	1.83E-02	ND	ND	ND	ND
Hexachloroethane	1.81E-02	1.83E-02	ND	ND	ND	ND	
N-nitroso-di-n-propylamine	1.81E-02	1.83E-02	ND	ND	ND	ND	
4-methylphenol	1.81E-02	1.83E-02	ND	ND	ND	ND	
Nitrobenzene	1.81E-02	1.83E-02	ND	ND	ND	ND	
Isophorone	1.81E-02	1.83E-02	ND	ND	ND	ND	
2-nitrophenol	1.81E-02	1.83E-02	ND	ND	ND	ND	
2,4-dimethylphenol	1.81E-02	1.83E-02	ND	ND	ND	ND	
Bis(2-chloroethoxy)methane	1.81E-02	1.83E-02	ND	ND	ND	ND	
2,4-dichlorophenol	1.81E-02	1.83E-02	ND	ND	ND	ND	
1,2,4-trichlorobenzene	1.81E-02	1.83E-02	ND	ND	ND	ND	
Naphthalene	1.79E-02	1.83E-02	2.17E-07	6.76E-06	9.86E-05	4.93E-05	

Table B-1: Air Modeling Output Data for the Cartridge, 0.50 Caliber Tracer, M17 (M2) - 100 meter location

Cartridge, 0.50 caliber, Tracer, M17 (M2)				Number of Rounds (l):		1 round
DODIC: A571				Release duration (t):		2 seconds
Number of items tested = 3				Unit Concentration (UC):		1.636E-04 (g/m ³)/(g/s)
Net Explosive Weight (lbs) = 3.21E-02						
Compound	ATC Modeling Results					Pollutant Emission Rate for One Item (g/sec) ER ₁
	Measured Actual Concentration (mg/m ³)	Measured Background Concentration (mg/m ³)	Average Adjusted Emission Factor (lb/item) EF	Average Adjusted Emission Factor (lb/lb NEW)	Average Modeled Concentration for One Item (grams/m ³) CONC	
4-chloroaniline	1.81E-02	1.83E-02	ND	ND	ND	ND
Hexachlorobutadiene	1.81E-02	1.83E-02	ND	ND	ND	ND
4-chloro-3-methylphenol	1.81E-02	1.83E-02	ND	ND	ND	ND
2-methylnaphthalene	1.81E-02	1.83E-02	ND	ND	ND	ND
Hexachlorocyclopentadiene	1.81E-02	1.83E-02	ND	ND	ND	ND
2,4,6-trichlorophenol	1.81E-02	1.83E-02	ND	ND	ND	ND
2,4,5-trichlorophenol	1.81E-02	1.83E-02	ND	ND	ND	ND
2-chloronaphthalene	1.81E-02	1.83E-02	ND	ND	ND	ND
2-nitroaniline	1.81E-02	1.83E-02	ND	ND	ND	ND
Acenaphthylene	1.81E-02	1.83E-02	ND	ND	ND	ND
Dimethylphthalate	1.81E-02	1.83E-02	ND	ND	ND	ND
2,6-dinitrotoluene	1.81E-02	1.83E-02	ND	ND	ND	ND
Acenaphthene	3.62E-02	3.67E-02	ND	ND	ND	ND
3-nitroaniline	3.62E-02	3.67E-02	ND	ND	ND	ND
2,4-dinitrophenol	1.81E-02	1.83E-02	ND	ND	ND	ND
Dibenzofuran	1.81E-02	1.83E-02	ND	ND	ND	ND
2,4-dinitrotoluene	3.62E-02	3.67E-02	ND	ND	ND	ND
4-nitrophenol	1.81E-02	1.83E-02	ND	ND	ND	ND
Fluorene	1.81E-02	1.83E-02	ND	ND	ND	ND
4-chlorophenyl-phenylether	1.81E-02	1.83E-02	ND	ND	ND	ND
Diethylphthalate	1.81E-02	1.83E-02	ND	ND	ND	ND
4-nitroaniline	3.62E-02	3.67E-02	ND	ND	ND	ND
4,6-dinitro-2-methylphenol	3.62E-02	3.67E-02	ND	ND	ND	ND
N-nitrosodiphenylamine(1)	1.81E-02	1.83E-02	ND	ND	ND	ND
4-bromophenyl-phenylether	1.81E-02	1.83E-02	ND	ND	ND	ND
Hexachlorobenzene	3.62E-02	3.67E-02	ND	ND	ND	ND
Pentachlorophenol	1.81E-02	1.83E-02	ND	ND	ND	ND
Phenanthrene	1.81E-02	1.83E-02	ND	ND	ND	ND
Anthracene	1.81E-02	1.83E-02	ND	ND	ND	ND

Table B-1: Air Modeling Output Data for the Cartridge, 0.50 Caliber Tracer, M17 (M2) - 100 meter location

Cartridge, 0.50 caliber, Tracer, M17 (M2)					Number of Rounds (I):		1 round	
DODIC: A571					Release duration (I):		2 seconds	
Number of items tested = 3					Unit Concentration (UC):		1.636E-04 (g/m³)/(g/s)	
Net Explosive Weight (lbs) = 3.21E-02								
Compound	ATC Filtration Test Results				Total Mass of Substance Emitted (grams/item) M	Average Modeled Concentration for One Item (grams/m³) CONC	Pollutant Emission Rate for One Item (g/sec) ER ₁	
	Measured Actual Concentration (mg/m³)	Measured Background Concentration (mg/m³)	Average Adjusted Emission Factor (lb/item) EF	Average Adjusted Emission Factor (lb/lb NEW)				
Di-n-butylphthalate	1.81E-02	1.83E-02	ND	ND	ND	ND	ND	
Fluoranthene	1.81E-02	1.83E-02	ND	ND	ND	ND	ND	
Pyrene	1.81E-02	1.83E-02	ND	ND	ND	ND	ND	
Butylbenzylphthalate	1.81E-02	1.83E-02	ND	ND	ND	ND	ND	
Benzo(a)anthracene	1.81E-02	1.83E-02	ND	ND	ND	ND	ND	
Chrysene	1.81E-02	1.83E-02	ND	ND	ND	ND	ND	
3,3-dichlorobenzidine	1.81E-02	1.83E-02	ND	ND	ND	ND	ND	
Bis(2-ethylhexyl)phthalate	9.11E-01	2.02E-01	8.94E-06	2.78E-04	4.06E-03	3.318E-07	2.03E-03	
Di-n-octylphthalate	1.81E-02	1.83E-02	ND	ND	ND	ND	ND	
Benzo(b)fluoranthene	1.81E-02	1.83E-02	ND	ND	ND	ND	ND	
Benzo(k)fluoranthene	1.81E-02	1.83E-02	ND	ND	ND	ND	ND	
Benzo(a)pyrene	1.81E-02	1.83E-02	ND	ND	ND	ND	ND	
Indeno(1,2,3-cd)pyrene	1.81E-02	1.83E-02	ND	ND	ND	ND	ND	
Dibenz(a,h)anthracene	1.81E-02	1.83E-02	ND	ND	ND	ND	ND	
Benzo(g,h,i)perylene	1.81E-02	1.83E-02	ND	ND	ND	ND	ND	
SVOC Tentatively Identified Compounds (TICs)								
TO-13 (PAHs)								
Naphthalene	1.61E-02	5.50E-03	1.36E-07	4.23E-06	6.17E-05	5.044E-09	3.08E-05	
Acenaphthylene	7.87E-04	1.83E-05	9.55E-09	2.97E-07	4.33E-06	3.545E-10	2.17E-06	
Acenaphthene	1.35E-04	4.77E-05	1.12E-09	3.48E-08	5.08E-07	4.157E-11	2.54E-07	
Fluorene	3.62E-04	3.85E-05	3.98E-09	1.24E-07	1.80E-06	1.475E-10	9.02E-07	
Phenanthrene	3.35E-04	7.88E-05	3.21E-09	9.98E-08	1.46E-06	1.191E-10	7.28E-07	
Anthracene	4.43E-05	1.83E-05	5.38E-10	1.67E-08	2.44E-07	1.995E-11	1.22E-07	
Fluoranthene	3.35E-04	1.83E-05	4.05E-09	1.26E-07	1.84E-06	1.504E-10	9.19E-07	
Pyrene	4.80E-04	1.83E-05	5.81E-09	1.81E-07	2.64E-06	2.156E-10	1.32E-06	
Benzo(a)anthracene	2.47E-04	1.83E-05	2.98E-09	9.27E-08	1.35E-06	1.105E-10	6.76E-07	
Chrysene	2.54E-04	1.83E-05	3.07E-09	9.54E-08	1.39E-06	1.138E-10	6.96E-07	
Benzo(b)fluoranthene	3.98E-04	1.83E-05	4.82E-09	1.50E-07	2.19E-06	1.790E-10	1.09E-06	
Benzo(k)fluoranthene	2.33E-04	1.83E-05	2.82E-09	8.76E-08	1.28E-06	1.045E-10	6.39E-07	

Table B-1: Air Modeling Output Data for the Cartridge, 0.50 Caliber Tracer, M17 (M2) - 100 meter location

Cartridge, 0.50 caliber, Tracer, M17 (M2)									
Number of Rounds (I):									
1 round									
Release duration (t):									
2 seconds									
Unit Concentration (UC):									
1.636E-04 (g/m ³)/(g/s)									
DODIC: A571									
Number of Items tested = 3									
Net Explosive Weight (lbs) = 3.21E-02									
ATC Firing Test Results									
Compound	Measured Actual Concentration (mg/m ³)	Measured Background Concentration (mg/m ³)	Average Adjusted Emission Factor (lb/item) EF	Average Adjusted Emission Factor (lb/lb NEW)	Total Mass of Substance Emitted (grams/item) M	Average Modeled Concentration for One Item (grams/m ³) CONC	Pollutant		
							Emission Rate for One Item (g/sec) ER ₁		
Benzo(e)pyrene	4.80E-04	1.83E-05	5.82E-09	1.81E-07	2.64E-06	2.158E-10	1.32E-06		
Benzo(a)pyrene	3.44E-04	1.83E-05	4.17E-09	1.30E-07	1.89E-06	1.547E-10	9.46E-07		
Indeno(1,2,3-cd)pyrene	3.71E-04	1.83E-05	4.50E-09	1.40E-07	2.04E-06	1.669E-10	1.02E-06		
Dibenz(a,h)anthracene	4.17E-05	1.83E-05	5.04E-10	1.57E-08	2.29E-07	1.871E-11	1.14E-07		
Benzo(g,h,i)perylene	7.05E-04	1.83E-05	8.56E-09	2.66E-07	3.88E-06	3.177E-10	1.94E-06		
Dioxins and Furans									
2378-Tetrachlorodibenzo-p-dioxin	3.75E-09	3.61E-09	ND	ND	ND	ND	ND		
12378-Pentachlorodibenzo-p-dioxin	3.59E-09	3.40E-09	ND	ND	ND	ND	ND		
123478-Hexachlorodibenzo-p-dioxin	4.38E-09	4.20E-09	ND	ND	ND	ND	ND		
123678-Hexachlorodibenzo-p-dioxin	4.53E-09	4.25E-09	ND	ND	ND	ND	ND		
123789-Hexachlorodibenzo-p-dioxin	7.24E-09	6.88E-09	ND	ND	ND	ND	ND		
1234678-Heptachlorodibenzo-p-dioxin	1.40E-08	5.35E-09	1.12E-13	3.49E-12	5.09E-11	4.166E-15	2.55E-11		
OCDD	2.14E-07	6.58E-08	1.89E-12	5.88E-11	8.58E-10	7.016E-14	4.29E-10		
2378-Tetrachlorodibenzo-p-furan	3.84E-09	3.61E-09	ND	ND	ND	ND	ND		
12378-Pentachlorodibenzo-p-furan	3.80E-09	3.65E-09	ND	ND	ND	ND	ND		
23478-Pentachlorodibenzo-o-furan	2.33E-09	2.16E-09	ND	ND	ND	ND	ND		
123478-Hexachlorodibenzo-p-furan	3.17E-09	2.97E-09	ND	ND	ND	ND	ND		
123678-Hexachlorodibenzo-p-furan	3.31E-09	3.02E-09	ND	ND	ND	ND	ND		
123789-Hexachlorodibenzo-p-furan	2.87E-09	2.70E-09	ND	ND	ND	ND	ND		
234678-Hexachlorodibenzo-p-furan	2.27E-09	2.13E-09	ND	ND	ND	ND	ND		
1234678-Heptachlorodibenzo-p-furan	2.79E-09	1.94E-09	3.38E-14	1.05E-12	1.53E-11	1.253E-15	7.66E-12		
1234789-Heptachlorodibenzo-p-furan	5.78E-09	5.62E-09	ND	ND	ND	ND	ND		
OCDF	9.51E-09	5.10E-09	1.16E-13	3.60E-12	5.25E-11	4.293E-15	2.62E-11		
Energetics									
Nitrobenzene	3.51E-03	NA	ND	ND	ND	ND	ND		
2-Nitrotoluene	3.51E-03	NA	ND	ND	ND	ND	ND		
3-Nitrotoluene	3.51E-03	NA	ND	ND	ND	ND	ND		
4-Nitrotoluene	3.51E-03	NA	ND	ND	ND	ND	ND		
Nitroglycerine	3.51E-03	NA	ND	ND	ND	ND	ND		

Table B-1: Air Modeling Output Data for the Cartridge, 0.50 Caliber Tracer, M17 (M2) - 100 meter location

Cartridge, 0.50 caliber, Tracer, M17 (M2)					Number of Rounds (I):		1 round	
DODIC: A571					Release duration (t):		2 seconds	
Number of items tested =					3			
Net Explosive Weight (lbs) =					3.21E-02		1.636E-04 (g/m³)/(g/s)	
Compound	ATC Firing Test Results						Pollutant Emission Rate for One Item (g/sec) ER ₁	
	Measured Actual Concentration (mg/m³)	Measured Background Concentration (mg/m³)	Average Adjusted Emission Factor (lb/item) EF	Average Adjusted Emission Factor (lb/lb NEW)	Total Mass of Substance Emitted (grams/item) M	Average Modeled Concentration for One Item (grams/m³) CONC		
	1,3-Dinitrobenzene	3.51E-03	NA	ND	ND	ND		ND
	2,6-Dinitrotoluene	3.51E-03	NA	ND	ND	ND		ND
	2,4-Dinitrotoluene	3.51E-03	NA	ND	ND	ND		ND
	1,3,5-Trinitrobenzene	3.51E-03	NA	ND	ND	ND		ND
	2,4,6-Trinitrotoluene	3.51E-03	NA	ND	ND	ND		ND
	RDX	3.51E-03	NA	ND	ND	ND		ND
	4-Amino-2,6-Dinitrotoluene	3.51E-03	NA	ND	ND	ND		ND
	2-Amino-4,6-Dinitrotoluene	3.51E-03	NA	ND	ND	ND		ND
	Tetryl	3.51E-03	NA	ND	ND	ND		ND
	HMX	7.01E-03	NA	ND	ND	ND		ND
	Pentaerythritol tetranitrate	7.01E-03	NA	ND	ND	ND		ND
	Dibutyl phthalate	1.75E-01	NA	ND	ND	ND		ND
	Diethyl phthalate	1.75E-01	NA	ND	ND	ND		ND
	Diphenylamine	8.77E-02	NA	ND	ND	ND		ND

Footnotes:

¹ATC = Aberdeen Test Center (for additional information on the data, refer to the Firing Point Emission Study)

NA = Not Applicable

ND = Not Detected

Footnotes:

¹ATC = Aberdeen Test Center (for additional information on the data, refer to the Firing Point Emission Study)

NA = Not Applicable

ND = Not Detected

Table B-2: Air Modeling Output Data for the Cartridge, 0.50 Caliber Tracer, M17 (M2) - 200 meter location

Compound	Cartridge, 0.50 caliber, Tracer, M17 (M2)				Number of Rounds (I):		1 round
	DODIC: A571				Release duration (t):		3 seconds
	Number of items tested = 3				Unit Concentration (UC):		6.870E-05 (g/m³)/(g/s)
	Net Explosive Weight (lbs) = 3.21E-02						
	ATC-Field Test Results						
	Measured Actual Concentration (mg/m³)	Measured Background Concentration (mg/m³)	Average Adjusted Emission Factor (lb/item) EF	Average Adjusted Emission Factor (lb/lb NEW)	Total Mass of Substance Emitted (grams/item) M	Average Modeled Concentration for One Item (grams/m³) CONC	Pollutant Emission Rate for One Item (g/sec) ER ₁
Permanent Gases							
	2.03E+01	NA	2.20E-04	6.83E-03	9.96E-02	2.281E-06	3.32E-02
	5.27E+02	NA	5.69E-03	1.77E-01	2.58E+00	5.915E-05	8.61E-01
	1.01E+03	NA	1.09E-02	3.40E-01	4.96E+00	1.136E-04	1.65E+00
	4.45E+01	NA	4.81E-04	1.50E-02	2.18E-01	4.996E-06	7.27E-02
	2.62E-01	NA	2.83E-06	8.81E-05	1.29E-03	2.944E-08	4.28E-04
Acid Gases							
	2.20E-01	2.20E-01	ND	ND	ND	ND	ND
	2.20E-01	2.10E-01	ND	ND	ND	ND	ND
	2.10E-01	2.10E-01	ND	ND	ND	ND	ND
	2.10E-01	2.10E-01	ND	ND	ND	ND	ND
	2.10E-01	2.10E-01	ND	ND	ND	ND	ND
Phosphoric Acid							
	2.55E-01	2.10E-01	4.66E-06	1.45E-04	2.11E-03	4.836E-08	7.04E-04
Sulfuric Acid							
Cyanide							
	5.80E-01	1.20E-02	7.07E-06	2.20E-04	3.20E-03	7.339E-08	1.07E-03
Particulate Cyanide		1.30E-02	8.78E-05	2.73E-03	3.98E-02	9.119E-07	1.33E-02
Hydrogen Cyanide							
Particulates							
	3.39E+01	NA	4.12E-04	1.28E-02	1.87E-01	4.277E-06	6.23E-02
	3.11E+01	NA	3.78E-04	1.18E-02	1.71E-01	3.927E-06	5.72E-02
	1.81E+01	NA	2.20E-04	6.86E-03	1.00E-01	2.289E-06	3.33E-02
Metals							
	7.19E-02	4.30E-02	8.72E-07	2.71E-05	3.96E-04	9.060E-09	1.32E-04
	5.65E-01	4.30E-02	6.88E-06	2.14E-04	3.12E-03	7.142E-08	1.04E-03
						ND	ND
	1.05E-02	1.08E-02	ND	ND	ND	ND	ND
	8.81E-01	4.30E-02	1.07E-05	3.34E-04	4.87E-03	1.116E-07	1.62E-03
	4.21E-02	4.30E-02	ND	ND	ND	ND	ND
	4.21E-02	4.30E-02	ND	ND	ND	ND	ND

Table B-2: Air Modeling Output Data for the Cartridge, 0.50 Caliber Tracer, M17 (M2) - 200 meter location

Compound	Cartridge, 0.50 caliber, Tracer, M17 (M2)					Number of Rounds (l):		1 round
	DODIC: A571					Release duration (t):		3 seconds
	Number of items tested = 3					Unit Concentration (UC):		6.870E-05 (g/m³)/(g/s)
	Net Explosive Weight (lbs) = 3.21E-02							
	Auto Firing Test Results							
	Measured Actual Concentration (mg/m³)	Measured Background Concentration (mg/m³)	Average Adjusted Emission Factor (lb/item) EF	Average Adjusted Emission Factor (lb/lb NEW)	Total Mass of Substance Emitted (grams/item) M	Average Modeled Concentration for One Item (grams/m³) CONC	Pollutant Emission Rate for One Item (g/sec) ER₁	
Chromium	4.21E-02	4.30E-02	ND	ND	ND	ND	ND	
Cobalt	4.21E-02	4.30E-02	ND	ND	ND	ND	ND	
Copper	1.11E+01	9.55E-02	1.33E-04	4.15E-03	6.06E-02	1.387E-06	2.02E-02	
Lead	1.15E+00	4.30E-02	1.39E-05	4.33E-04	6.31E-03	1.445E-07	2.10E-03	
Magnesium	2.26E-01	4.30E-02	2.76E-06	8.58E-05	1.25E-03	2.865E-08	4.17E-04	
Manganese	4.21E-02	4.30E-02	ND	ND	ND	ND	ND	
Nickel	4.21E-02	4.30E-02	ND	ND	ND	ND	ND	
Selenium	1.05E-02	1.08E-02	ND	ND	ND	ND	ND	
Silver	4.21E-02	4.30E-02	ND	ND	ND	ND	ND	
Thallium	4.21E-02	4.30E-02	ND	ND	ND	ND	ND	
Vanadium	4.21E-02	4.30E-02	ND	ND	ND	ND	ND	
Zinc	1.81E+00	4.30E-02	2.20E-05	6.84E-04	9.97E-03	2.284E-07	3.32E-03	
TO-11 Carbonyls								
Formaldehyde	2.46E-02	1.23E-01	2.98E-07	9.27E-06	1.35E-04	3.096E-09	4.51E-05	
Acetaldehyde	1.80E-01	1.80E-01	ND	ND	ND	ND	ND	
Acetone	1.19E+00	1.19E+00	ND	ND	ND	ND	ND	
Acrolein	2.29E-01	2.29E-01	ND	ND	ND	ND	ND	
Propionaldehyde	2.37E-01	2.37E-01	ND	ND	ND	ND	ND	
Crotonaldehyde	2.87E-01	2.87E-01	ND	ND	ND	ND	ND	
Butyraldehyde	2.95E-01	2.95E-01	ND	ND	ND	ND	ND	
Benzaldehyde	4.34E-01	4.34E-01	ND	ND	ND	ND	ND	
Isovaleraldehyde	3.52E-01	3.52E-01	ND	ND	ND	ND	ND	
Valeraldehyde	3.52E-01	3.52E-01	ND	ND	ND	ND	ND	
o,m,p-Tolualdehyde	4.91E-01	4.91E-01	ND	ND	ND	ND	ND	
Hexaldehyde	4.10E-01	4.10E-01	ND	ND	ND	ND	ND	
2,5-Dimethylbenzaldehyde	4.10E-01	4.10E-01	ND	ND	ND	ND	ND	
VOCs								
Propene	5.94E-02	3.44E-04	7.15E-07	2.23E-05	3.25E-04	7.431E-09	1.08E-04	
Dichlorodifluoromethane	2.97E-03	3.46E-03	ND	ND	ND	ND	ND	

Table B-2: Air Modeling Output Data for the Cartridge, 0.50 Caliber Tracer, M17 (M2) - 200 meter location

Cartridge, 0.50 caliber, Tracer, M17 (M2)				Number of Rounds (I):		1 round	
DODIC: A571				Release duration (t):		3 seconds	
Number of items tested =				Unit Concentration (UC):		6.870E-05 (g/m³)/(g/s)	
Net Explosive Weight (lbs) =				3.21E-02			
Compound	ATC Firing Test Results						
	Measured Actual Concentration (mg/m³)	Measured Background Concentration (mg/m³)	Average Adjusted Emission Factor (lb/item) EF	Average Adjusted Emission Factor (lb/lb NEW)	Total Mass of Substance Emitted (grams/item) M	Average Modeled Concentration for One Item (grams/m³) CONC	Pollutant Emission Rate for One Item (g/sec) ER₁
Chlorodifluoromethane	3.54E-03	3.54E-03	ND	ND	ND	ND	ND
Freon 114	6.99E-03	6.99E-03	ND	ND	ND	ND	ND
Chloromethane	1.65E-03	1.45E-03	4.41E-09	1.37E-07	2.00E-06	4.581E-11	6.67E-07
Vinyl Chloride	2.56E-03	2.56E-03	ND	ND	ND	ND	ND
1,3-Butadiene	1.33E-02	2.21E-03	1.61E-07	5.00E-06	7.29E-05	1.670E-09	2.43E-05
Bromomethane	3.88E-03	3.88E-03	ND	ND	ND	ND	ND
Chloroethane	2.64E-03	2.64E-03	ND	ND	ND	ND	ND
Dichlorofluoromethane	4.21E-03	4.21E-03	ND	ND	ND	ND	ND
Trichlorofluoromethane	1.69E-03	1.69E-03	2.22E-09	6.91E-08	1.01E-06	2.307E-11	3.36E-07
Pentane	8.85E-04	2.95E-03	1.07E-08	3.34E-07	4.87E-06	1.116E-10	1.62E-06
Acrolein	2.29E-03	2.29E-03	ND	ND	ND	ND	ND
1,1-Dichloroethene	4.05E-03	4.05E-03	ND	ND	ND	ND	ND
Freon 113	7.68E-03	7.68E-03	ND	ND	ND	ND	ND
Acetone	1.84E-01	1.07E-01	1.10E-06	3.41E-05	4.98E-04	1.140E-08	1.66E-04
Methyl Iodide	5.81E-03	5.81E-03	ND	ND	ND	ND	ND
Carbon Disulfide	3.11E-03	3.11E-03	ND	ND	ND	ND	ND
Acetonitrile	9.82E-02	8.39E-03	1.10E-06	3.41E-05	4.97E-04	1.138E-08	1.66E-04
3-Chloropropene	3.13E-03	3.13E-03	ND	ND	ND	ND	ND
Methylene Chloride	1.39E-01	4.52E-02	1.20E-06	3.74E-05	5.45E-04	1.248E-08	1.82E-04
tert-Butyl Alcohol	3.03E-03	3.03E-03	ND	ND	ND	ND	ND
Acrylonitrile	1.95E-02	2.17E-03	2.36E-07	7.35E-06	1.07E-04	2.454E-09	3.57E-05
trans-1,2-Dichloroethene	3.96E-03	3.96E-03	ND	ND	ND	ND	ND
Methyl t-Butyl Ether	3.61E-03	3.61E-03	ND	ND	ND	ND	ND
Hexane	9.52E-02	1.37E-01	ND	ND	ND	ND	ND
1,1-Dichloroethane	3.97E-03	3.97E-03	ND	ND	ND	ND	ND
Vinyl Acetate	3.52E-03	3.52E-03	ND	ND	ND	ND	ND
cis-1,2-Dichloroethene	3.96E-03	3.96E-03	ND	ND	ND	ND	ND
2-Butanone	2.95E-03	2.95E-03	ND	ND	ND	ND	ND
Ethyl Acetate	1.08E-02	3.60E-03	1.30E-07	4.04E-06	5.89E-05	1.350E-09	1.96E-05

Table B-2: Air Modeling Output Data for the Cartridge, 0.50 Caliber Tracer, M17 (M2) - 200 meter location

Cartridge, 0.50 caliber, Tracer, M17 (M2)									
DODIC: A571									
Number of items tested = 3									
Net Explosive Weight (lbs) = 3.21E-02									
AIC Firing Test Results									
Compound	Measured Actual Concentration (mg/m³)	Measured Background Concentration (mg/m³)	Average Adjusted Emission Factor (lb/item) EF	Average Adjusted Emission Factor (lb/lb NEW)	Total Mass of Substance Emitted (grams/item) M	Average Modeled Concentration for One Item (grams/m³) CONC	Pollutant Emission Rate for One Item (g/sec) ER ₁	Number of Rounds (I):	
								Release duration (t):	Unit Concentration (UC):
								1 round	3 seconds
								6.870E-05 (g/m³)/(g/s)	
Methyl Acrylate	3.52E-03	3.52E-03	ND	ND	ND	ND	ND	ND	ND
Chloroform	4.88E-03	4.88E-03	ND	ND	ND	ND	ND	ND	ND
1,1,1-Trichloroethane	3.82E-03	3.82E-03	4.91E-09	1.53E-07	2.23E-06	5.096E-11	7.42E-07		
Carbon Tetrachloride	6.29E-03	6.29E-03	ND	ND	ND	ND	ND	ND	ND
1,2-Dichloroethane	6.07E-03	4.05E-03	7.34E-08	2.28E-06	3.33E-05	7.628E-10	1.11E-05		
Benzene	3.12E-01	9.59E-04	3.76E-06	1.17E-04	1.71E-03	3.907E-08	5.69E-04		
Isooctane	4.67E-03	4.67E-03	ND	ND	ND	ND	ND	ND	ND
Heptane	4.10E-03	2.46E-03	ND	ND	ND	ND	ND	ND	ND
Trichloroethane	4.88E-03	4.88E-03	ND	ND	ND	ND	ND	ND	ND
Ethyl Acrylate	4.09E-03	4.09E-03	ND	ND	ND	ND	ND	ND	ND
1,2-Dichloropropane	4.62E-03	4.62E-03	ND	ND	ND	ND	ND	ND	ND
Methyl Methacrylate	4.09E-03	4.09E-03	ND	ND	ND	ND	ND	ND	ND
Dibromomethane	7.11E-03	7.11E-03	ND	ND	ND	ND	ND	ND	ND
1,4-Dioxane	3.60E-03	3.60E-03	ND	ND	ND	ND	ND	ND	ND
Bromodichloromethane	6.70E-03	6.70E-03	ND	ND	ND	ND	ND	ND	ND
4-Methyl-2-Pentanone	4.10E-03	4.10E-03	ND	ND	ND	ND	ND	ND	ND
Toluene	3.58E-02	7.54E-04	4.26E-07	1.33E-05	1.93E-04	4.426E-09	6.44E-05		
Octane	4.67E-03	4.67E-03	ND	ND	ND	ND	ND	ND	ND
trans-1,3-Dichloropropene	4.54E-03	4.54E-03	ND	ND	ND	ND	ND	ND	ND
Ethyl Methacrylate	4.67E-03	4.67E-03	ND	ND	ND	ND	ND	ND	ND
1,1,2-Trichloroethane	5.46E-03	5.46E-03	ND	ND	ND	ND	ND	ND	ND
Tetrachloroethene	6.78E-03	6.78E-03	ND	ND	ND	ND	ND	ND	ND
2-Hexanone	4.10E-03	4.10E-03	ND	ND	ND	ND	ND	ND	ND
Dibromochloromethane	8.52E-03	8.52E-03	ND	ND	ND	ND	ND	ND	ND
1,2-Dibromoethane	7.68E-03	7.68E-03	ND	ND	ND	ND	ND	ND	ND
Chlorobenzene	4.60E-03	4.60E-03	ND	ND	ND	ND	ND	ND	ND
1,1,1,2-Tetrachloroethane	6.87E-03	6.87E-03	ND	ND	ND	ND	ND	ND	ND
Ethylbenzene	2.82E-03	4.34E-03	3.42E-08	1.06E-06	1.55E-05	3.550E-10	5.17E-06		
m/p-Xylene	6.51E-03	1.30E-03	6.47E-08	2.01E-06	2.93E-05	6.720E-10	9.78E-06		

Table B-2: Air Modeling Output Data for the Cartridge, 0.50 Caliber Tracer, M17 (M2) - 200 meter location

Cartridge, 0.50 caliber, Tracer, M17 (M2)							
DODIC: A571							
Number of items tested = 3							
Net Explosive Weight (lbs) = 3.21E-02							
Compound	ATC Emissions Results						
	Measured Actual Concentration (mg/m ³)	Measured Background Concentration (mg/m ³)	Average Adjusted Emission Factor (lb/item) EF	Average Adjusted Emission Factor (lb/lb NEW)	Total Mass of Substance Emitted (grams/item) M	Average Modeled Concentration for One Item (grams/m ³) CONC	Pollutant Emission Rate for One Item (g/sec) ER ₁
o-Xylene	4.34E-03	4.34E-03	5.27E-08	1.64E-06	2.39E-05	5.473E-10	7.97E-06
Styrene	1.07E-02	4.26E-03	1.29E-07	4.01E-06	5.85E-05	1.340E-09	1.95E-05
Bromoform	1.03E-02	1.03E-02	ND	ND	ND	ND	ND
Cumene	4.92E-03	4.92E-03	ND	ND	ND	ND	ND
1,1,2,2-Tetrachlorethane	6.87E-03	6.87E-03	ND	ND	ND	ND	ND
1,2,3-Trichloropropane	6.03E-03	6.03E-03	ND	ND	ND	ND	ND
Bromobenzene	6.42E-03	6.42E-03	ND	ND	ND	ND	ND
4-Ethyltoluene	4.92E-03	4.92E-03	ND	ND	ND	ND	ND
1,3,5-Trimethylbenzene	4.92E-03	4.92E-03	ND	ND	ND	ND	ND
Alpha Methyl Styrene	4.83E-03	4.83E-03	ND	ND	ND	ND	ND
1,2,4-Trimethylbenzene	9.83E-04	4.92E-03	1.18E-08	3.68E-07	5.36E-06	1.227E-10	1.79E-06
1,3-Dichlorobenzene	6.01E-03	6.01E-03	ND	ND	ND	ND	ND
1,4-Dichlorobenzene	6.01E-03	6.01E-03	ND	ND	ND	ND	ND
Benzyl Chloride	5.18E-03	5.18E-03	ND	ND	ND	ND	ND
1,2-Dichlorobenzene	6.01E-03	6.01E-03	ND	ND	ND	ND	ND
Hexachlorethane	9.68E-03	9.68E-03	ND	ND	ND	ND	ND
1,2,4-Trichlorobenzene	7.42E-03	7.42E-03	ND	ND	ND	ND	ND
Hexachlorobutadiene	1.07E-02	1.07E-02	ND	ND	ND	ND	ND
VOC Tentatively Identified Compounds (TICs)							
Hydrocarbons							
Methane	4.10E+00	1.10E+00	3.79E-05	1.18E-03	1.72E-02	3.934E-07	5.73E-03
Ethylene	2.00E-01	2.29E-02	2.43E-06	7.57E-05	1.10E-03	2.526E-08	3.68E-04
Acetylene	2.74E-02	2.13E-02	3.32E-07	1.03E-05	1.51E-04	3.453E-09	5.03E-05
Ethane	1.13E-01	2.46E-02	1.37E-06	4.27E-05	6.23E-04	1.427E-08	2.08E-04
Propylene	5.42E-02	3.44E-02	6.59E-07	2.05E-05	2.99E-04	6.842E-09	9.96E-05
Propane	3.61E-02	3.61E-02	ND	ND	ND	ND	ND
Propyne	3.20E-02	3.20E-02	ND	ND	ND	ND	ND
Isobutane	4.75E-02	4.75E-02	ND	ND	ND	ND	ND
1-Butene/Isobutylene	9.18E-02	9.18E-02	ND	ND	ND	ND	ND

Table B-2: Air Modeling Output Data for the Cartridge, 0.50 Caliber Tracer, M17 (M2) - 200 meter location

Cartridge, 0.50 caliber, Tracer, M17 (M2)				Number of Rounds (I):		1 round	
DODIC: A571				Release duration (t):		3 seconds	
Number of items tested = 3				Unit Concentration (UC):		6.870E-05 (g/m ³)/(g/s)	
Net Explosive Weight (lbs) = 3.21E-02							
Compound	ATC Firing Test Results				Total Mass of Substance Emitted (grams/item) M	Average Modeled Concentration for One Item (grams/m ³) CONC	Pollutant Emission Rate for One Item (g/sec) ER ₁
	Measured Actual Concentration (mg/m ³)	Measured Background Concentration (mg/m ³)	Average Adjusted Emission Factor (lb/item) EF	Average Adjusted Emission Factor (lb/lb NEW)			
1,3-Butadiene/butane	4.59E-02	4.59E-02	ND	ND	ND	ND	ND
cis-butene	4.59E-02	4.59E-02	ND	ND	ND	ND	ND
1-Butyne	4.59E-02	4.59E-02	ND	ND	ND	ND	ND
trans-Butene	4.59E-02	4.59E-02	ND	ND	ND	ND	ND
2-Butyne	4.42E-02	4.42E-02	ND	ND	ND	ND	ND
n-Pentane	5.90E-02	5.90E-02	ND	ND	ND	ND	ND
n-Hexane	1.99E-01	7.05E-02	2.41E-06	7.50E-05	1.09E-03	2.505E-08	3.65E-04
SVOCs							
N-nitrosodimethylamine	1.81E-02	1.83E-02	ND	ND	ND	ND	ND
Bis(2-chloroethyl)ether	1.81E-02	1.83E-02	ND	ND	ND	ND	ND
Phenol	1.81E-02	1.83E-02	ND	ND	ND	ND	ND
2-chlorophenol	1.81E-02	1.83E-02	ND	ND	ND	ND	ND
1,3-dichlorobenzene	1.81E-02	1.83E-02	ND	ND	ND	ND	ND
1,4-dichlorobenzene	1.81E-02	1.83E-02	ND	ND	ND	ND	ND
1,2-dichlorobenzene	1.81E-02	1.83E-02	ND	ND	ND	ND	ND
Benzyl alcohol	1.81E-02	1.83E-02	ND	ND	ND	ND	ND
Bis(2-chloroisopropyl)ether	1.81E-02	1.83E-02	ND	ND	ND	ND	ND
2-methylphenol	1.81E-02	1.83E-02	ND	ND	ND	ND	ND
Hexachloroethane	1.81E-02	1.83E-02	ND	ND	ND	ND	ND
N-nitroso-di-n-propylamine	1.81E-02	1.83E-02	ND	ND	ND	ND	ND
4-methylphenol	1.81E-02	1.83E-02	ND	ND	ND	ND	ND
Nitrobenzene	1.81E-02	1.83E-02	ND	ND	ND	ND	ND
Isophorone	1.81E-02	1.83E-02	ND	ND	ND	ND	ND
2-nitrophenol	1.81E-02	1.83E-02	ND	ND	ND	ND	ND
2,4-dimethylphenol	1.81E-02	1.83E-02	ND	ND	ND	ND	ND
Bis(2-chloroethoxy)methane	1.81E-02	1.83E-02	ND	ND	ND	ND	ND
2,4-dichlorophenol	1.81E-02	1.83E-02	ND	ND	ND	ND	ND
1,2,4-trichlorobenzene	1.81E-02	1.83E-02	ND	ND	ND	ND	ND
Naphthalene	1.79E-02	1.83E-02	2.17E-07	6.76E-06	9.86E-05	2.259E-09	3.29E-05

Table B-2: Air Modeling Output Data for the Cartridge, 0.50 Caliber Tracer, M17 (M2) - 200 meter location

Cartridge, 0.50 caliber, Tracer, M17 (M2)						Number of Rounds (l):		1 round	
DODIC: A571						Release duration (t):		3 seconds	
Number of items tested =						3		6.870E-05 (g/m³)/(g/s)	
Net Explosive Weight (lbs) =						3.21E-02			
Compound	ATC Firing Test Results					Total Mass of Substance Emitted (grams/item) M	Average Modeled Concentration for One Item (grams/m³) CONC	Pollutant Emission Rate for One Item (g/sec) ER ₁	
	Measured Actual Concentration (mg/m³)	Measured Background Concentration (mg/m³)	Average Adjusted Emission Factor (lb/item) EF	Average Adjusted Emission Factor (lb/lb NEW)					
4-chloroaniline	1.81E-02	1.83E-02	ND	ND	ND	ND	ND	ND	
Hexachlorobutadiene	1.81E-02	1.83E-02	ND	ND	ND	ND	ND	ND	
4-chloro-3-methylphenol	1.81E-02	1.83E-02	ND	ND	ND	ND	ND	ND	
2-methylnaphthalene	1.81E-02	1.83E-02	ND	ND	ND	ND	ND	ND	
Hexachlorocyclopentadiene	1.81E-02	1.83E-02	ND	ND	ND	ND	ND	ND	
2,4,6-trichlorophenol	1.81E-02	1.83E-02	ND	ND	ND	ND	ND	ND	
2,4,5-trichlorophenol	1.81E-02	1.83E-02	ND	ND	ND	ND	ND	ND	
2-chloronaphthalene	1.81E-02	1.83E-02	ND	ND	ND	ND	ND	ND	
2-nitroaniline	1.81E-02	1.83E-02	ND	ND	ND	ND	ND	ND	
Acenaphthylene	1.81E-02	1.83E-02	ND	ND	ND	ND	ND	ND	
Dimethylphthalate	1.81E-02	1.83E-02	ND	ND	ND	ND	ND	ND	
2,6-dinitrotoluene	1.81E-02	1.83E-02	ND	ND	ND	ND	ND	ND	
Acenaphthene	3.62E-02	3.67E-02	ND	ND	ND	ND	ND	ND	
3-nitroaniline	3.62E-02	3.67E-02	ND	ND	ND	ND	ND	ND	
2,4-dinitrophenol	1.81E-02	1.83E-02	ND	ND	ND	ND	ND	ND	
Dibenzofuran	1.81E-02	1.83E-02	ND	ND	ND	ND	ND	ND	
2,4-dinitrotoluene	3.62E-02	3.67E-02	ND	ND	ND	ND	ND	ND	
4-nitrophenol	1.81E-02	1.83E-02	ND	ND	ND	ND	ND	ND	
Fluorene	1.81E-02	1.83E-02	ND	ND	ND	ND	ND	ND	
4-chlorophenyl-phenylether	1.81E-02	1.83E-02	ND	ND	ND	ND	ND	ND	
Diethylphthalate	1.81E-02	1.83E-02	ND	ND	ND	ND	ND	ND	
4-nitroaniline	3.62E-02	3.67E-02	ND	ND	ND	ND	ND	ND	
4,6-dinitro-2-methylphenol	3.62E-02	3.67E-02	ND	ND	ND	ND	ND	ND	
N-nitrosodiphenylamine(1)	1.81E-02	1.83E-02	ND	ND	ND	ND	ND	ND	
4-bromophenyl-phenylether	1.81E-02	1.83E-02	ND	ND	ND	ND	ND	ND	
Hexachlorobenzene	1.81E-02	1.83E-02	ND	ND	ND	ND	ND	ND	
Pentachlorophenol	3.62E-02	3.67E-02	ND	ND	ND	ND	ND	ND	
Phenanthrene	1.81E-02	1.83E-02	ND	ND	ND	ND	ND	ND	
Anthracene	1.81E-02	1.83E-02	ND	ND	ND	ND	ND	ND	

Table B-2: Air Modeling Output Data for the Cartridge, 0.50 Caliber Tracer, M17 (M2) - 200 meter location

Cartridge, 0.50 caliber, Tracer, M17 (M2)									
DODIC: A571									
Number of items tested = 3									
Net Explosive Weight (lbs) = 3.21E-02									
Compound	ATC Firing Test Results					Total Mass of Substance Emitted (grams/item) M	Average Modeled Concentration for One Item (grams/m³) CONC	Pollutant Emission Rate for One Item (g/sec) ER ₁	
	Measured Actual Concentration (mg/m³)	Measured Background Concentration (mg/m³)	Average Adjusted Emission Factor (lb/item) EF	Average Adjusted Emission Factor (lb/lb NEW)					
Di-n-butylphthalate	1.81E-02	1.83E-02	ND	ND	ND	ND	ND	ND	ND
Fluoranthene	1.81E-02	1.83E-02	ND	ND	ND	ND	ND	ND	ND
Pyrene	1.81E-02	1.83E-02	ND	ND	ND	ND	ND	ND	ND
Butylbenzylphthalate	1.81E-02	1.83E-02	ND	ND	ND	ND	ND	ND	ND
Benzo(a)anthracene	1.81E-02	1.83E-02	ND	ND	ND	ND	ND	ND	ND
Chrysene	1.81E-02	1.83E-02	ND	ND	ND	ND	ND	ND	ND
3,3-dichlorobenzidine	1.81E-02	1.83E-02	ND	ND	ND	ND	ND	ND	ND
Bis(2-ethylhexyl)phthalate	9.11E-01	2.02E-01	8.94E-06	2.78E-04	4.06E-03	9.288E-08	1.35E-03		
Di-n-octylphthalate	1.81E-02	1.83E-02	ND	ND	ND	ND	ND	ND	ND
Benzo(b)fluoranthene	1.81E-02	1.83E-02	ND	ND	ND	ND	ND	ND	ND
Benzo(k)fluoranthene	1.81E-02	1.83E-02	ND	ND	ND	ND	ND	ND	ND
Benzo(a)pyrene	1.81E-02	1.83E-02	ND	ND	ND	ND	ND	ND	ND
Indeno(1,2,3-cd)pyrene	1.81E-02	1.83E-02	ND	ND	ND	ND	ND	ND	ND
Dibenz(a,h)anthracene	1.81E-02	1.83E-02	ND	ND	ND	ND	ND	ND	ND
Benzo(g,h,i)perylene	1.81E-02	1.83E-02	ND	ND	ND	ND	ND	ND	ND
SVOC Tentatively Identified Compounds (TICs)									
TO-13 (PAHs)									
Naphthalene	1.61E-02	5.50E-03	1.36E-07	4.23E-06	6.17E-05	1.412E-09	2.06E-05		
Acenaphthylene	7.87E-04	1.83E-05	9.55E-09	2.97E-07	4.33E-06	9.924E-11	1.44E-06		
Acenaphthene	1.35E-04	4.77E-05	1.12E-09	3.48E-08	5.08E-07	1.164E-11	1.69E-07		
Fluorene	3.62E-04	3.85E-05	3.98E-09	1.24E-07	1.80E-06	4.130E-11	6.01E-07		
Phenanthrene	3.35E-04	7.88E-05	3.21E-09	9.98E-08	1.46E-06	3.333E-11	4.85E-07		
Anthracene	4.43E-05	1.83E-05	5.38E-10	1.67E-08	2.44E-07	5.585E-12	8.13E-08		
Fluoranthene	3.35E-04	1.83E-05	4.05E-09	1.26E-07	1.84E-06	4.210E-11	6.13E-07		
Pyrene	4.80E-04	1.83E-05	5.81E-09	1.81E-07	2.64E-06	6.036E-11	8.79E-07		
Benzo(a)anthracene	2.47E-04	1.83E-05	2.98E-09	9.27E-08	1.35E-06	3.094E-11	4.50E-07		
Chrysene	2.54E-04	1.83E-05	3.07E-09	9.54E-08	1.39E-06	3.187E-11	4.64E-07		
Benzo(b)fluoranthene	3.98E-04	1.83E-05	4.82E-09	1.50E-07	2.19E-06	5.011E-11	7.29E-07		
Benzo(k)fluoranthene	2.33E-04	1.83E-05	2.82E-09	8.76E-08	1.28E-06	2.926E-11	4.26E-07		

Table B-2: Air Modeling Output Data for the Cartridge, 0.50 Caliber Tracer, M17 (M2) - 200 meter location

Compound	Cartridge, 0.50 caliber, Tracer, M17 (M2)				Number of Rounds (I):		1 round	
	DODIC: A571				Release duration (t):		3 seconds	
	Number of items tested =				Unit Concentration (UC):		6.870E-05 (g/m ³)/(g/s)	
	Net Explosive Weight (lbs) =							
	3.21E-02							
Compound	ATC Firing Test Results				Total Mass of Substance Emitted (grams/item)	Average Modeled Concentration for One Item (grams/m ³)	Pollutant Emission Rate for One Item (g/sec)	
	Measured Actual Concentration (mg/m ³)	Measured Background Concentration (mg/m ³)	Average Adjusted Emission Factor (lb/item) EF	Average Adjusted Emission Factor (lb/lb NEW)				M
	Benzo(e)pyrene	4.80E-04	1.83E-05	5.82E-09	1.81E-07	2.64E-06	6.042E-11	8.80E-07
	Benzo(a)pyrene	3.44E-04	1.83E-05	4.17E-09	1.30E-07	1.89E-06	4.331E-11	6.30E-07
	Indeno(1,2,3-cd)pyrene	3.71E-04	1.83E-05	4.50E-09	1.40E-07	2.04E-06	4.673E-11	6.80E-07
	Dibenz(a,h)anthracene	4.17E-05	1.83E-05	5.04E-10	1.57E-08	2.29E-07	5.238E-12	7.62E-08
	Benzo(g,h,i)perylene	7.05E-04	1.83E-05	8.56E-09	2.66E-07	3.88E-06	8.895E-11	1.29E-06
	Dioxins and Furans							
	2378-Tetrachlorodibenzo-p-dioxin	3.75E-09	3.61E-09	ND	ND	ND	ND	ND
	12378-Pentachlorodibenzo-p-dioxin	3.59E-09	3.40E-09	ND	ND	ND	ND	ND
	123478-Hexachlorodibenzo-p-dioxin	4.38E-09	4.20E-09	ND	ND	ND	ND	ND
	123678-Hexachlorodibenzo-p-dioxin	4.53E-09	4.25E-09	ND	ND	ND	ND	ND
	123789-Hexachlorodibenzo-p-dioxin	7.24E-09	6.88E-09	ND	ND	ND	ND	ND
	1234678-Heptachlorodibenzo-p-dioxin	1.40E-08	5.35E-09	1.12E-13	3.49E-12	5.09E-11	1.166E-15	1.70E-11
	OCDD	2.14E-07	6.58E-08	1.89E-12	5.88E-11	8.58E-10	1.964E-14	2.86E-10
	2378-Tetrachlorodibenzo-p-furan	3.84E-09	3.61E-09	ND	ND	ND	ND	ND
	12378-Pentachlorodibenzo-p-furan	3.80E-09	3.65E-09	ND	ND	ND	ND	ND
	23478-Pentachlorodibenzo-o-furan	2.33E-09	2.16E-09	ND	ND	ND	ND	ND
	123478-Hexachlorodibenzo-p-furan	3.17E-09	2.97E-09	ND	ND	ND	ND	ND
	123678-Hexachlorodibenzo-p-furan	3.31E-09	3.02E-09	ND	ND	ND	ND	ND
123789-Hexachlorodibenzo-p-furan	2.87E-09	2.70E-09	ND	ND	ND	ND	ND	
234678-Hexachlorodibenzo-p-furan	2.27E-09	2.13E-09	ND	ND	ND	ND	ND	
1234678-Heptachlorodibenzo-p-furan	2.79E-09	1.94E-09	3.38E-14	1.05E-12	1.53E-11	3.508E-16	5.11E-12	
1234789-Heptachlorodibenzo-p-furan	5.78E-09	5.62E-09	ND	ND	ND	ND	ND	
OCDF	9.51E-09	5.10E-09	1.16E-13	3.60E-12	5.25E-11	1.202E-15	1.75E-11	
Energetics								
Nitrobenzene	3.51E-03	NA	ND	ND	ND	ND	ND	
2-Nitrotoluene	3.51E-03	NA	ND	ND	ND	ND	ND	
3-Nitrotoluene	3.51E-03	NA	ND	ND	ND	ND	ND	
4-Nitrotoluene	3.51E-03	NA	ND	ND	ND	ND	ND	
Nitroglycerine	3.51E-03	NA	ND	ND	ND	ND	ND	

Table B-2: Air Modeling Output Data for the Cartridge, 0.50 Caliber Tracer, M17 (M2) - 200 meter location

Cartridge, 0.50 caliber, Tracer, M17 (M2)					Number of Rounds (I):		1 round
DODIC: A571					Release duration (t):		3 seconds
Number of Items tested =					3		
Net Explosive Weight (lbs) =					3.21E-02		6.870E-05 (g/m ³)/(g/s)
Compound	ATC Firing Test Results				Total Mass of Substance Emitted (grams/item) M	Average Modeled Concentration for One Item (grams/m ³) CONC	Pollutant Emission Rate for One Item (g/sec) ER ₁
	Measured Actual Concentration (mg/m ³)	Measured Background Concentration (mg/m ³)	Average Adjusted Emission Factor EF	Average Adjusted Emission Factor (lb/lb NEW)			
	1,3-Dinitrobenzene	3.51E-03	NA	ND	ND	ND	ND
	2,6-Dinitrotoluene	3.51E-03	NA	ND	ND	ND	ND
	2,4-Dinitrotoluene	3.51E-03	NA	ND	ND	ND	ND
	1,3,5-Trinitrobenzene	3.51E-03	NA	ND	ND	ND	ND
	2,4,6-Trinitrotoluene	3.51E-03	NA	ND	ND	ND	ND
	RDX	3.51E-03	NA	ND	ND	ND	ND
	4-Amino-2,6-Dinitrotoluene	3.51E-03	NA	ND	ND	ND	ND
	2-Amino-4,6-Dinitrotoluene	3.51E-03	NA	ND	ND	ND	ND
	Tetryl	3.51E-03	NA	ND	ND	ND	ND
	HMX	7.01E-03	NA	ND	ND	ND	ND
	Pentaerythritol tetranitrate	7.01E-03	NA	ND	ND	ND	ND
	Dibutyl phthalate	1.75E-01	NA	ND	ND	ND	ND
	Diethyl phthalate	1.75E-01	NA	ND	ND	ND	ND
	Diphenylamine	8.77E-02	NA	ND	ND	ND	ND

Footnotes:
'ATC' = Aberdeen Test Center (for additional information on the data, refer to the Firing Point Emission Study)
NA = Not Applicable
ND = Not Detected

Footnotes:

'ATC = Aberdeen Test Center (for additional information on the data, refer to the Firing Point Emission Study)

NA = Not Applicable

ND = Not Detected

APPENDIX C

HEALTH-BASED SCREENING LEVELS AND ACUTE TOXICITY VALUES

Appendix C: Health-Based Screening Levels and Acute Toxicity Values

Compound	CAS #	Region 9 PRG (µg/m ³)	Toxicity Endpoint (c or nc)	Region 9 RBC (µg/m ³)	Toxicity Endpoint (c or nc)	HBSL (µg/m ³)	ERRG (µg/m ³)	TBEL (µg/m ³)	AEGI (µg/m ³)	Source (T or E)	ATV (µg/m ³)
Permanent Gases											
Ammonia (NH ₃)	7664-41-7	1.04E+02	nc	104.39		1.04E+02	1.75E+04	1.75E+04	NA	E	1.75E+04
Carbon Dioxide (CO ₂)	124-38-9	NA		NA		NA	NA	5.40E+07	NA	T	5.40E+07
Carbon Monoxide (CO)	630-08-0	1.00E+04	nc	NA		1.00E+04	2.30E+05	2.28E+05	NA	E	2.30E+05
Oxides of Nitrogen (as NO)	10102-43-9	1.00E+02	nc	NA		1.00E+02	NA	3.08E+04	NA	T	3.08E+04
Sulfur Dioxide (SO ₂)	7446-09-5	8.00E+01	nc	NA		8.00E+01	7.89E+02	7.86E+02	NA	E	7.89E+02
Acid Gases											
Hydrogen fluoride	7664-39-3	NA		NA		NA	1.60E+03	1.64E+03	NA	E	1.60E+03
Hydrogen chloride	7647-01-0	2.08E+01	nc	2.08E+01		2.08E+01	4.50E+03	4.47E+03	NA	E	4.50E+03
Hydrogen bromide	10035-10-6	NA		NA		NA	NA	9.93E+03	NA	T	9.93E+03
Nitric Acid	7697-37-2	NA		NA		NA	NA	2.58E+03	1.30E+03	A	1.30E+03
Phosphoric acid	7664-38-2	1.04E+01	nc	1.06E+01		1.04E+01	NA	3.00E+03	NA	T	3.00E+03
Sulfuric Acid	7664-93-9	NA		NA		NA	2.00E+03	2.00E+03	NA	E	2.00E+03
Cyanide											
Particulate Cyanide	57-12-5	NA		7.30E+01		7.30E+01	NA	5.00E+03	NA	T	5.00E+03
Hydrogen Cyanide	74-90-8	3.13E+00	nc	3.14E+00		3.13E+00	NA	5.17E+03	NA	T	5.17E+03
Particulates											
Total Suspended Particulate	12789-66-1	5.00E+01	nc	NA		5.00E+01	NA	NA	NA		NA
PM ₁₀		5.00E+01	nc	NA		5.00E+01	NA	NA	NA		NA
PM _{2.5}		1.50E+01	nc	NA		1.50E+01	NA	NA	NA		NA
Metals											
Aluminum	7429-90-5	5.11E+00	nc	3.65E+00		5.11E+00	NA	3.00E+04	NA	T	3.00E+04
Antimony	7440-36-0	NA		1.46E+00		1.46E+00	NA	1.50E+03	NA	T	1.50E+03
Arsenic	7440-38-2	4.47E-04	c	4.15E-04		4.47E-04	NA	3.00E+01	NA	T	3.00E+01
Barium	7440-39-3	5.21E-01	nc	5.11E-01		5.21E-01	NA	1.50E+03	NA	T	1.50E+03
Beryllium	7440-41-7	8.00E-04	c	7.45E-04		8.00E-04	NA	5.00E+00	NA	T	5.00E+00
Cadmium	7440-43-9	1.07E-03	c	9.94E-04		1.07E-03	NA	3.00E+01	NA	T	3.00E+01
Calcium	7440-70-2	NA		NA		NA	NA	3.00E+04	NA	T	3.00E+04
Chromium	7440-47-3		c	1.53E-04		1.53E-04	NA	1.50E+03	NA	T	1.50E+03
Cobalt	7440-48-4	NA		2.20E+02		2.20E+02	NA	6.00E+01	NA	T	6.00E+01
Copper	7440-50-8	NA		1.46E+02		1.46E+02	NA	3.00E+03	NA	T	3.00E+03
Lead	7439-92-1	1.50E+00	nc	NA		1.50E+00	NA	1.50E+02	NA	T	1.50E+02
Magnesium	7439-95-4	NA		NA		NA	NA	3.00E+04	NA	T	3.00E+04
Manganese	7439-96-5	5.11E-02	nc	5.22E-02		5.11E-02	NA	3.00E+03	NA	T	3.00E+03
Nickel	7440-02-0	NA		7.30E+01		7.30E+01	NA	3.00E+03	NA	T	3.00E+03

Appendix C: Health-Based Screening Levels and Acute Toxicity Values

Compound	CAS #	Region 9 PRG (µg/m³)	Toxicity Endpoint (c or nc)	Region 3 RBC (µg/m³)	Toxicity Endpoint (c or nc)	HBSL (µg/m³)	ERPG (µg/m³)	TEEL (µg/m³)	SAEGL (µg/m³)	Source (T or E)	ATV (µg/m³)
Selenium	7782-49-2	NA		1.83E+01		1.83E+01	NA	6.00E+02	NA	T	6.00E+02
Silver	7740-22-4	NA		1.83E+01		1.83E+01	NA	3.00E+02	NA	T	3.00E+02
Thallium	7440-28-0	NA		2.56E-01		2.56E-01	NA	3.00E+02	NA	T	3.00E+02
Vanadium	7440-62-2	NA		2.56E+01		2.56E+01	NA	1.50E+02	NA	T	1.50E+02
Zinc	7440-66-6	NA		1.10E+03		1.10E+03	NA	3.00E+04	NA	T	3.00E+04
TO-11 Carbonyls											
Formaldehyde	50-00-0	1.48E-01	c	1.39E-01		1.48E-01	1.23E+03	1.23E+03	NA	E	1.23E+03
Acetaldehyde	75-07-0	8.73E-01	c	8.13E-01		8.73E-01	1.80E+04	1.80E+04	NA	E	1.80E+04
Acetone	67-64-1	3.65E+02	nc	3.65E+02		3.65E+02	NA	2.37E+06	NA	T	2.37E+06
Acrolein	107-02-8	2.09E-02	nc	2.08E-02		2.09E-02	2.30E+02	2.29E+02	NA	E	2.30E+02
Propionaldehyde	123-38-6	NA		NA		NA	NA	7.50E+04	NA	T	7.50E+04
Crotonaldehyde	4170-30-3	3.54E-03	c	3.30E-03		3.54E-03	5.72E+03	5.72E+03	NA	E	5.72E+03
Butyraldehyde	123-72-8	NA		NA		NA	NA	7.38E+04	NA	T	7.38E+04
Benzaldehyde	100-52-7	3.65E+02	nc	3.65E+02		3.65E+02	NA	1.50E+04	NA	T	1.50E+04
Isovaleraldehyde	590-86-3	NA		NA		NA	NA	NA	NA		NA
Valeraldehyde	110-62-3	NA		NA		NA	NA	NA	NA		NA
o,m,p-Tolualdehyde	1334-78-7	NA		NA		NA	NA	NA	NA		NA
Hexaldehyde	66-25-1	NA		NA		NA	NA	NA	NA		NA
2,5-Dimethylbenzaldehyde	5779-94-2	NA		NA		NA	NA	NA	NA		NA
VOCs											
Propene	115-07-1	NA		NA		NA	NA	NA	NA		NA
Dichlorodifluoromethane	75-71-8	2.09E+02	nc	1.83E+02		2.09E+02	NA	1.48E+07		T	1.48E+07
Chlorodifluoromethane	75-45-6	5.11E+04	nc	5.11E+04		5.11E+04	NA	4.41E+06		T	4.41E+06
Freon 114	76-14-2	NA		NA		NA	NA	2.10E+07		T	2.10E+07
Chloromethane	74-87-3	1.07E+00	c	1.79E+00		1.07E+00	NA	2.06E+05		T	2.06E+05
Vinyl Chloride	75-01-4	2.20E-02	c	2.10E-02		2.20E-02	NA	1.28E+04		T	1.28E+04
1,3-Butadiene	106-99-0	3.74E-03	c	3.48E-03		3.74E-03	2.20E+04	2.21E+04		E	2.20E+04
Bromomethane	74-83-9	5.21E+00	nc	5.11E+00		5.21E+00	NA	5.82E+04		T	5.82E+04
Chloroethane	75-00-3	2.32E+00	nc	NA		2.32E+00	NA	2.64E+06		T	2.64E+06
Dichlorofluoromethane	75-71-8	2.09E+02	nc	1.83E+02		2.09E+02	NA	1.48E+07		T	1.48E+07
Trichlorofluoromethane	75-69-4	7.30E+02	nc	7.30E+02		7.30E+02	NA	2.81E+06		T	2.81E+06
Pentane	109-66-0	NA		NA		NA	NA	1.80E+06		T	1.80E+06
Acrolein	107-02-8	2.09E-02	nc	2.08E-02		2.09E-02	2.30E+02	2.29E+02		E	2.30E+02
1,1-Dichloroethene	75-35-4	5.21E+02	nc	5.11E+02		5.21E+02	NA	7.92E+04		T	7.92E+04
Freon 113	76-13-1	3.13E+04	nc	3.14E+04		3.13E+04	NA	9.58E+06		T	9.58E+06
Acetone	67-64-1	3.65E+02	nc	3.65E+02		3.65E+02	NA	2.37E+06		T	2.37E+06

Appendix C: Health-Based Screening Levels and Acute Toxicity Values

Compound	GAS #	Region 9 PRG ($\mu\text{g}/\text{m}^3$)	Toxicity Endpoint ($\text{g}/\text{org}/\text{nc}$)	Region 3 RBC ($\mu\text{g}/\text{m}^3$)	Toxicity Endpoint ($\text{g}/\text{org}/\text{nc}$)	HBSL ($\mu\text{g}/\text{m}^3$)	ERPG ($\mu\text{g}/\text{m}^3$)	TEEL ($\mu\text{g}/\text{m}^3$)	AEGL ($\mu\text{g}/\text{m}^3$)	P-Source ($\mu\text{g}/\text{m}^3$)	ATV ($\mu\text{g}/\text{m}^3$)
Methyl Iodide	74-88-4	NA		NA		NA	145000	1.45E+05		E	1.45E+05
Carbon Disulfide	75-15-0	7.30E+02	nc	7.30E+02	nc	7.30E+02	NA	3.11E+04		T	3.11E+04
Acetonitrile	75-05-8	6.20E+01	nc	6.21E+01	nc	6.20E+01	NA	1.01E+05		T	1.01E+05
3-Chloropropene	107-05-1	1.04E+00	nc	NA		1.04E+00	9.39E+03	9.39E+03		E	9.39E+03
Methylene Chloride	75-09-2	4.09E+00	c	3.79E+00	c	4.09E+00	696000	6.94E+05		E	6.96E+05
tert-Butyl Alcohol	75-65-0	NA		NA		NA	NA	4.55E+05		T	4.55E+05
Acrylonitrile	107-13-1	2.83E-02	c	2.61E-02	c	2.83E-02	21700	2.17E+04		E	2.17E+04
trans-1,2-Dichloroethene	156-60-5	7.30E+01	nc	7.30E+01	nc	7.30E+01	NA	4.95E+04		T	4.95E+04
Methyl t-Butyl Ether	1634-04-4	3.13E+03	nc	3.13E+03	nc	3.13E+03	NA	4.32E+05		T	4.32E+05
Hexane	110-54-3	2.09E+02	nc	2.08E+02	nc	2.09E+02	NA	5.28E+05		T	5.28E+05
1,1-Dichloroethane	75-34-3	5.21E+02	nc	5.11E+02	nc	5.21E+02	NA	1.21E+06		T	1.21E+06
Vinyl Acetate	108-05-4	2.09E+02	nc	2.08E+02	nc	2.09E+02	19150	1.76E+04		E	1.92E+04
cis-1,2-Dichloroethene	156-59-2	3.65E+01	nc	3.65E+01	nc	3.65E+01	NA	7.92E+05		T	7.92E+05
2-Butanone	78-93-3	1.04E+03	nc	1.04E+03	nc	1.04E+03	NA	8.85E+05		T	8.85E+05
Ethyl Acetate	141-78-6	3.29E+03	nc	3.29E+03	nc	3.29E+03	NA	1.44E+06		T	1.44E+06
Methyl Acrylate	96-33-3	1.10E+02	nc	1.10E+02	nc	1.10E+02	NA	NA			NA
Chloroform	67-66-3	8.35E-02	c	7.73E-02	c	8.35E-02	NA	9.76E+03		T	9.76E+03
1,1,1-Trichloroethane	71-55-6	1.04E+03	nc	2.30E+03	nc	1.04E+03	1.94E+06	1.91E+06		E	1.94E+06
Carbon Tetrachloride	56-23-5	1.28E-01	c	1.18E-01	c	1.28E-01	1.28E+05	1.26E+05		E	1.28E+05
1,2-Dichloroethane	107-06-2	7.39E-02	c	6.88E-02	c	7.39E-02	NA	8.08E+03		T	8.08E+03
Benzene	71-43-2	2.49E-01	c	2.16E-01	c	2.49E-01	1.56E+05	1.60E+05		E	1.56E+05
Isooctane (2,2,4-trimethylpentane)	540-84-1	NA		NA		NA	NA	3.50E+05		T	3.50E+05
Heptane	142-82-5	NA		NA		NA	NA	1.80E+06		T	1.80E+06
Trichloroethane	71-55-6	1.04E+03	nc	2.30E+03	nc	1.04E+03	1.94E+06	1.91E+06		E	1.94E+06
Ethyl Acrylate	140-88-5	1.40E-01	c	NA		1.40E-01	NA	6.14E+04		T	6.14E+04
1,2-Dichloropropane	78-87-5	9.89E-02	c	9.21E-02	c	9.89E-02	NA	5.08E+05		T	5.08E+05
Methyl Methacrylate	80-62-6	7.30E+02	nc	7.30E+02	nc	7.30E+02	NA	4.09E+05		T	4.09E+05
Dibromomethane	74-95-3	3.65E+01	nc	3.65E+01	nc	3.65E+01	NA	2.50E+05		T	2.50E+05
1,4-Dioxane	123-91-1	6.11E-01	c	5.69E-01	c	6.11E-01	NA	9.00E+04		T	9.00E+04
Bromodichloromethane	75-27-4	1.08E-01	c	1.01E-01	c	1.08E-01	NA	4.00E+03		T	4.00E+03
4-Methyl-2-Pentanone	108-10-1	8.34E+01	nc	7.30E+01	nc	8.34E+01	NA	3.07E+05		T	3.07E+05
Toluene	108-88-3	4.02E+02	nc	4.16E+02	nc	4.02E+02	1.88E+05	1.89E+05		E	1.88E+05
Octane	111-65-9	NA		NA		NA	NA	NA			NA
trans-1,3-Dichloropropene	10061-02-6	5.17E-02	c	4.82E-02	c	5.17E-02	NA	NA			NA
Ethyl Methacrylate	97-63-2	3.29E+02	nc	3.29E+02	nc	3.29E+02	NA	NA			NA
1,1,2-Trichloroethane	79-00-5	1.20E-01	c	1.12E-01	c	1.20E-01	NA	1.64E+05		T	1.64E+05

Appendix C: Health-Based Screening Levels and Acute Toxicity Values

Compound	CAS#	Region 9 PRG (µg/m ³)	Toxicity Endpoint (organo)	Region 3 RBC (µg/m ³)	Toxicity Endpoint (organo)	HBSL (µg/m ³)	ERPG (µg/m ³)	TEEL (µg/m ³)	AEGL (µg/m ³)	Source (T or E)	ATV (µg/m ³)
Tetrachloroethene	127-18-4	3.31E+00	c	3.13E+00	c	3.31E+00	NA	6.78E+05		T	6.78E+05
2-Hexanone	591-78-6	NA		5.11E+00		5.11E+00	NA	4.09E+04		T	4.09E+04
Dibromochloromethane	124-48-1	8.00E-02	c	7.45E-02	c	8.00E-02	NA	6.00E+03		T	6.00E+03
1,2-Dibromoethane	106-93-4	8.73E-03	c	8.24E-03	c	8.73E-03	NA	1.54E+05		T	1.54E+05
Chlorobenzene	108-90-7	6.21E+01	nc	6.21E+01	nc	6.21E+01	NA	1.38E+05		T	1.38E+05
1,1,1,2-Tetrachloroethane	630-20-6	2.60E-01	c	2.41E-01	c	2.60E-01	NA	5.15E+04		T	5.15E+04
Ethylbenzene	100-41-4	1.06E+03	nc	1.06E+03	nc	1.06E+03	NA	5.43E+05		T	5.43E+05
m&p-Xylene	108-38-3 106-42-3	7.30E+02	nc	7.30E+03	nc	7.30E+02	NA	6.51E+05		T	6.51E+05
o-Xylene	95-47-6	7.30E+02	nc	7.30E+03	nc	7.30E+02	NA	6.51E+05		T	6.51E+05
Styrene	100-42-5	1.06E+03	nc	1.04E+03	nc	1.06E+03	2.13E+05	2.13E+05		E	2.13E+05
Bromoform	75-25-2	1.75E+00	c	1.61E+00	c	1.75E+00	NA	6.20E+03		T	6.20E+03
Cumene	98-82-8	4.02E+02	nc	4.02E+02	nc	4.02E+02	NA	2.46E+05		T	2.46E+05
1,1,2,2-Tetrachloroethane	79-34-5	3.31E-02	c	3.13E-02	c	3.31E-02	NA	2.06E+04		T	2.06E+04
1,2,3-Trichloropropane	96-18-4	9.61E-04	c	3.13E-03	c	9.61E-04	NA	6.03E+04		T	6.03E+04
Bromobenzene	108-86-1	1.04E+01	nc	NA	nc	1.04E+01	NA	4.82E+04		T	4.82E+04
4-Ethyltoluene	622-96-8	NA		NA		NA	NA	1.25E+05		T	1.25E+05
1,3,5-Trimethylbenzene	108-67-8	6.21E+00	nc	6.21E+00	nc	6.21E+00	NA	3.68E+05		T	3.68E+05
Alpha Methyl Styrene	98-83-9	2.56E+02	nc	2.56E+02	nc	2.56E+02	NA	NA			NA
1,2,4-Trimethylbenzene	95-63-6	6.21E+00	nc	6.21E+00	nc	6.21E+00	NA	1.80E+05		T	1.80E+05
1,3-Dichlorobenzene	541-73-1	3.29E+00	nc	3.29E+00	nc	3.29E+00	NA	3.61E+04		T	3.61E+04
1,4-Dichlorobenzene	106-46-7	3.06E-01	c	2.85E-01	c	3.06E-01		6.61E+05		T	6.61E+05
Benzyl Chloride	100-44-7	3.96E-02	c	3.68E-02	c	3.96E-02	5.20E+03	5.17E+03		E	5.20E+03
1,2-Dichlorobenzene	95-50-1	2.09E+02	nc	3.29E+01	nc	2.09E+02	NA	3.01E+05		T	3.01E+05
Hexachlorethane	67-72-1	4.80E-01	c	4.47E-01	c	4.80E-01	NA	2.90E+04		T	2.90E+04
1,2,4-Trichlorobenzene	120-82-1	2.08E+02	nc	2.08E+02	nc	2.08E+02	NA	3.71E+04		T	3.71E+04
Hexachlorobutadiene	87-68-3	8.73E-02	c	8.03E-02	c	8.73E-02	3.21E+04	3.20E+04		E	3.21E+04
Hydrocarbons											
Methane	74-82-8	NA		NA		NA	NA	3.30E+06		T	3.30E+06
Ethylene	74-85-1	NA		NA		NA	NA	4.60E+05		T	4.60E+05
Acetylene	74-86-2	NA		NA		NA	NA	NA			NA
Ethane	74-84-0	NA		NA		NA	NA	NA			NA
Propylene	115-07-1	NA		NA		NA	NA	NA			NA
Propane	74-98-6	NA		NA		NA	NA	3.78E+06		T	3.78E+06
Propyne (methyl acetylene)	74-99-7	NA		NA		NA	NA	2.79E+06		T	2.79E+06

Appendix C: Health-Based Screening Levels and Acute Toxicity Values

Compound	CAS	Region 9 PRG (µg/m³)	Toxicity Endpoint (c or nc)	Region 3 RBC (µg/m³)	Toxicity Endpoint (c or nc)	HBSL (µg/m³)	ERPG (µg/m³)	TEEL (µg/m³)	AEGL (µg/m³)	Source (T or E)	ATV (µg/m³)
Isobutane	75-28-5	NA		NA		NA	NA	9.52E+05		T	9.52E+05
1-Butene/isobutylene (115-11-7)	106-98-9	NA		NA		NA	NA	6.87E+06		T	6.87E+06
1,3-Butadiene/butane	106-99-0	3.74E-03	c	3.48E-03		3.74E-03	2.20E+04	2.21E+04		E	2.20E+04
cis-butene	25167-67-3	NA		NA		NA	NA	1.72E+04	NA	T	1.72E+04
1-Butyne	107-00-6	NA		NA		NA	NA	NA			NA
trans-Butene	25167-67-3	NA		NA		NA	NA	1.72E+04	NA	T	1.72E+04
2-Butyne (crotonylene)	503-17-3	NA		NA		NA	NA	NA			NA
n-Pentane	109-66-0	NA		NA		NA	NA	1.80E+06		T	1.80E+06
n-Hexane	110-54-3	2.10E+02	nc	2.08E+02	nc	2.10E+02	NA	5.28E+05		T	5.28E+05
SVOCs											
n-nitrosodimethylamine	62-75-9	1.37E-04	c	1.23E-04	c	1.37E-04	NA	2.50E+03		T	2.50E+03
bis(2-chloroethyl)ether	111-44-4	5.82E-03	c	5.69E-03	c	5.82E-03	NA	5.85E+04		T	5.85E+04
phenol	108-95-2	2.19E+03	nc	2.19E+03	nc	2.19E+03	NA	3.85E+04		T	3.85E+04
2-chlorophenol	95-57-8	1.83E+01	nc	1.83E+01	nc	1.83E+01	NA	5.25E+03		T	5.25E+03
1,3-Dichlorobenzene	541-73-1	3.29E+00	nc	3.29E+00	nc	3.29E+00	NA	3.61E+04		T	3.61E+04
1,4-dichlorobenzene	106-46-7	3.06E-01	c	2.85E-01	c	3.06E-01	NA	6.61E+05		T	6.61E+05
1,2-dichlorobenzene	95-50-1	2.09E+02	nc	3.29E+01	nc	2.09E+02	NA	3.01E+05		T	3.01E+05
benzyl alcohol	100-51-6	1.10E+03	nc	1.10E+03	nc	1.10E+03	NA	5.53E+04		T	5.53E+04
bis(2-chloroisopropyl)ether	108-60-1	1.92E-01	c	1.79E-01	c	1.92E-01	NA	6.99E+04		T	6.99E+04
2-methylphenol	95-48-7	1.83E+02	nc	1.83E+02	nc	1.83E+02	NA	NA			NA
hexachloroethane	67-72-1	4.80E-01	c	4.47E-01	c	4.80E-01	NA	2.90E+04		T	2.90E+04
n-nitroso-di-n-propylamine	621-64-7	9.61E-04	c	8.94E-04	c	9.61E-04	NA	2.00E+02		T	2.00E+02
4-methylphenol	106-44-5	1.83E+02	nc	1.83E+02	nc	1.83E+02	NA	NA			NA
nitrobenzene	98-95-3	2.09E+00	nc	2.19E+00	nc	2.09E+00	NA	1.51E+04		T	1.51E+04
isophorone	78-59-1	7.08E+00	c	6.59E+00	c	7.08E+00	NA	2.83E+04		T	2.83E+04
2-nitrophenol	88-75-5	NA		NA		NA	NA	NA			NA
2,4-dimethylphenol	105-67-9	7.30E+01	nc	7.30E+01	nc	7.30E+01	NA	NA			NA
bis(2-chloroethoxy)methane	111-91-1	NA		NA		NA	NA	NA			NA
2,4-dichlorophenol	120-83-2	1.10E+01	nc	1.10E+01	nc	1.10E+01	NA	3.00E+04		T	3.00E+04
1,2,4-trichlorobenzene	120-82-1	2.08E+02	nc	2.08E+02	nc	2.08E+02	NA	3.71E+04		T	3.71E+04
naphthalene	91-20-3	3.13E+00	nc	3.29E+00	nc	3.13E+00	NA	7.86E+04		T	7.86E+04
4-chloroaniline	106-47-8	1.46E+01	nc	1.46E+01	nc	1.46E+01	NA	3.00E+04		T	3.00E+04
hexachlorobutadiene	87-68-3	8.62E-02	c	8.03E-02	c	8.62E-02	3.21E+04	3.20E+04		E	3.21E+04
4-chloro-3-methylphenol	59-50-7	NA		NA		NA	NA	2.00E+04		T	2.00E+04
2-methylnaphthalene	91-57-6	NA		7.30E+01	nc	7.30E+01	NA	2.00E+04		T	2.00E+04
hexachlorocyclopentadiene	77-47-4	7.30E-02	nc	7.30E-02	nc	7.30E-02	NA	2.23E+02		T	2.23E+02

Appendix C: Health-Based Screening Levels and Acute Toxicity Values

Compound	CAS #	Region 9 PRG (µg/m ³)	Toxicity Endpoint (c or nc)	Region 3 RBC (µg/m ³)	Toxicity Endpoint (c or nc)	HBSL (µg/m ³)	ERPG (µg/m ³)	HEEL (µg/m ³)	AEGL (µg/m ³)	Source (T or E)	ATV (µg/m ³)
2,4,6-trichlorophenol	88-06-2	1.10E+02	nc	1.10E+02	nc	1.10E+02	NA	3.00E+04		T	3.00E+04
2,4,5-trichlorophenol	95-95-4	3.65E+02	nc	3.65E+02	nc	3.65E+02	NA	3.00E+04		T	3.00E+04
2-chloronaphthalene	91-58-7	2.92E+02	nc	2.92E+02	nc	2.92E+02	NA	6.00E+02		T	6.00E+02
2-nitroaniline	88-74-4	2.09E-01	nc	2.08E-01	nc	2.09E-01	NA	NA			NA
Acenaphthylene	208-96-8	NA		NA		NA	NA	2.00E+02		T	2.00E+02
dimethylphthalate	131-11-3	3.65E+04	nc	3.65E+04	nc	3.65E+04	NA	1.50E+04		T	1.50E+04
2,6-dinitrotoluene	606-20-2	3.65E+00	nc	3.65E+00	nc	3.65E+00	NA	6.00E+02		T	6.00E+02
acenaphthene	83-32-9	2.19E+02	nc	2.19E+02	nc	2.19E+02	NA	1.25E+03		T	1.25E+03
3-nitroaniline	99-09-2	NA		NA		NA	NA	NA			NA
2,4-dinitrophenol	51-28-5	7.30E+00	nc	7.30E+00	nc	7.30E+00	NA	7.50E+03		T	7.50E+03
dibenzofuran	132-64-9	1.46E+01	nc	1.46E+01	nc	1.46E+01	NA	NA			NA
2,4-dinitrotoluene	121-14-2	7.30E+00	nc	7.30E+00	nc	7.30E+00	NA	6.00E+02		T	6.00E+02
4-nitrophenol	100-02-7	2.92E+01	nc	2.92E+01	nc	2.92E+01	NA	3.00E+04		T	3.00E+04
Fluorene	86-73-7	1.46E+02	nc	1.46E+02	nc	1.46E+02	NA	7.50E+04		T	7.50E+04
4-chlorophenyl-phenylether	7005-72-3	NA		NA		NA	NA	NA			NA
diethylphthalate	84-66-2	2.92E+03	nc	2.92E+03	nc	2.92E+03	NA	1.50E+04		T	1.50E+04
4-nitroaniline	100-01-6	NA		NA		NA	NA	9.00E+03		T	9.00E+03
4,6-dinitro-2-methylphenol	534-52-1	NA		3.65E-01	nc	3.65E-01	NA	5.00E+02		T	5.00E+02
n-nitrosodiphenylamine(1)	86-30-6	1.37E+00	c	1.28E+00	c	1.37E+00	NA	NA			NA
4-bromophenyl-phenylether	101-55-3	NA		NA		NA	NA	NA			NA
hexachlorobenzene	118-74-1	4.18E-03	c	3.91E-03	c	4.18E-03	NA	7.50E+01		T	7.50E+01
pentachlorophenol	87-86-5	5.60E-02	c	5.22E-02	c	5.60E-02	NA	1.50E+03		T	1.50E+03
phenanthrene	85-01-8	NA		NA		NA	NA	2.00E+03		T	2.00E+03
anthracene	120-12-7	1.10E+03	nc	1.10E+03	nc	1.10E+03	NA	6.00E+03		T	6.00E+03
di-n-butylphthalate	84-74-2	3.65E+02	nc	3.65E+02	nc	3.65E+02	NA	1.50E+04		T	1.50E+04
fluoranthene	206-44-0	1.46E+02	nc	1.46E+02	nc	1.46E+02	NA	3.00E+01		T	3.00E+01
pyrene	129-00-0	1.10E+02	nc	1.10E+02	nc	1.10E+02	NA	1.50E+04		T	1.50E+04
butylbenzylphthalate	85-68-7	7.30E+02	nc	7.30E+02	nc	7.30E+02	NA	5.00E+05		T	5.00E+05
benzo(a)anthracene	56-55-3	2.17E-02	c	8.58E-03	c	2.17E-02	NA	6.00E+02		T	6.00E+02
chrysene	218-01-9	2.17E+00	c	8.58E-01	c	2.17E+00	NA	2.00E+02		T	2.00E+02
3,3-dichlorobenzidine	91-94-1	1.50E-02	c	1.39E-02	c	1.50E-02	NA	6.21E+03		T	6.21E+03
bis(2-ethylhexyl)phthalate	117-81-7	4.80E-01	c	4.47E-01	c	4.80E-01	NA	1.00E+04		T	1.00E+04
di-n-octylphthalate	117-84-0	7.30E+01	nc	7.30E+01	nc	7.30E+01	NA	1.50E+05		T	1.50E+05
benzo(b)fluoranthene	205-99-2	2.17E-02	c	8.58E-03	c	2.17E-02	NA	NA			NA
benzo(k)fluoranthene	207-08-9	2.17E-01	c	8.58E-02	c	2.17E-01	NA	NA			NA
benzo(a)pyrene	50-32-8	2.17E-03	c	2.02E-03	c	2.17E-03	NA	7.50E+03		T	7.50E+03

Appendix C: Health-Based Screening Levels and Acute Toxicity Values

Compound	GAS #	Region 9 PRG (µg/m³)	Toxicity Endpoint (c or nc)	Region 3 RBC (µg/m³)	Toxicity Endpoint (c or nc)	HBSL (µg/m³)	ERPG (µg/m³)	TEEL (µg/m³)	AEGL (µg/m³)	Source (T or E)	ATV (µg/m³)
indeno(1,2,3-cd)pyrene	193-39-5	2.17E-02	c	8.58E-03		2.17E-02	NA	NA			NA
dibenz(a,h)anthracene	53-70-3	2.17E-03	c	8.58E-04		2.17E-03	NA	3.00E+04		T	3.00E+04
benzo(g,h,i)perylene	191-24-2	NA		NA		NA	NA	3.00E+04		T	3.00E+04
TO-13 (PAHs)											
naphthalene	91-20-3	3.13E+00	nc	3.29E+00		3.13E+00	NA	7.86E+04		T	7.86E+04
acenaphthylene	208-96-8	NA		NA		NA	NA	2.00E+02		T	2.00E+02
Acenaphthene	83-32-9	2.19E+02	nc	2.19E+02		2.19E+02	NA	1.25E+03		T	1.25E+03
fluorene	86-73-7	1.46E+02	nc	1.46E+02		1.46E+02	NA	7.50E+04		T	7.50E+04
phenanthrene	85-01-8	NA		NA		NA	NA	2.00E+03		T	2.00E+03
anthracene	120-12-7	1.10E+03	nc	1.10E+03		1.10E+03	NA	6.00E+03		T	6.00E+03
fluoranthene	206-44-0	1.46E+02	nc	1.46E+02		1.46E+02	NA	3.00E+01		T	3.00E+01
pyrene	129-00-0	1.10E+02	nc	1.10E+02		1.10E+02	NA	1.50E+04		T	1.50E+04
benzo(a)anthracene	56-55-3	2.17E-02	c	8.58E-03		2.17E-02	NA	6.00E+02		T	6.00E+02
chrysene	218-01-9	2.17E+00	c	8.58E-01		2.17E+00	NA	2.00E+02		T	2.00E+02
benzo(b)fluoranthene	205-99-2	2.17E-02	c	8.58E-03		2.17E-02	NA	NA			NA
benzo(k)fluoranthene	207-08-9	2.17E-01	c	8.58E-02		2.17E-01	NA	NA			NA
Benzo(e)pyrene	192-97-2	NA		NA		NA	NA	NA	NA		NA
benzo(a)pyrene	50-32-8	2.17E-03	c	2.02E-03		2.17E-03	NA	7.50E+03		T	7.50E+03
indeno(1,2,3-cd)pyrene	193-39-5	2.17E-02	c	8.58E-03		2.17E-02	NA	NA			NA
dibenz(a,h)anthracene	53-70-3	2.17E-03	c	8.58E-04		2.17E-03	NA	3.00E+04		T	3.00E+04
benzo(g,h,i)perylene	191-24-2	NA		NA		NA	NA	3.00E+04		T	3.00E+04
Dioxins and Furans											
2378-Tetrachlorodibenzo-p-dioxin	1746-01-6	4.48E-08	c	4.17E-08		4.48E-08	NA	3.50E+00		T	3.50E+00
12378-Pentachlorodibenzo-p-dioxin	40321-76-4	NA		NA		NA	NA	2.50E+00		T	2.50E+00
123478-Hexachlorodibenzo-p-dioxin	39227-28-6	NA		NA		NA	NA	NA			NA
123678-Hexachlorodibenzo-p-dioxin	57653-85-7	NA		NA		NA	NA	1.50E+01		T	1.50E+01
123789-Hexachlorodibenzo-p-dioxin	19408-74-3	1.48E-06	c	1.38E-06		1.48E-06	NA	NA			NA
1234678-Heptachlorodibenzo-p-dioxin	35822-46-9	NA		NA		NA	NA	NA			NA
Octachlorodibenzo(p)dioxin	3268-87-9	NA		NA		NA	NA	1.50E+02		T	1.50E+02
2376-Tetrachlorodibenzo-p-furan	51207-31-9	NA		NA		NA	NA	2.00E+00		T	2.00E+00
12378-Pentachlorodibenzo-p-furan	57117-41-6	NA		NA		NA	NA	NA			NA
23478-Pentachlorodibenzo-o-furan	57117-31-4	NA		NA		NA	NA	7.50E-02		T	7.50E-02
123478-Hexachlorodibenzo-p-furan	70648-26-9	NA		NA		NA	NA	7.50E+00		T	7.50E+00
123678-Hexachlorodibenzo-p-furan	57117-44-9	NA		NA		NA	NA	2.50E+00		T	2.50E+00
123789-Hexachlorodibenzo-p-furan	72918-21-9	NA		NA		NA	NA	NA			NA

Appendix C: Health-Based Screening Levels and Acute Toxicity Values

Compound	CAS #	Region 9 PRG (µg/m³)	Toxicity Endpoint (G or nc)	Region 3 RBC (µg/m³)	Toxicity Endpoint (G or nc)	HBSL (µg/m³)	ERPG (µg/m³)	TEEL (µg/m³)	AEGL (µg/m³)	Source (T or E)	ATV (µg/m³)
234678-Hexachlorodibenzo-p-furan	60851-34-5	NA		NA		NA	NA	1.50E+00		T	1.50E+00
1234678-Heptachlorodibenzo-p-furan	67562-39-4	NA		NA		NA	NA	NA			NA
1234789-Heptachlorodibenzo-p-furan	55673-89-7	NA		NA		NA	NA	NA			NA
Octachlorodibenzofuran	39001-02-0	NA		NA		NA	NA	3.00E+02		T	3.00E+02
Energetics											
Nitrobenzene	98-95-3	2.09E+00	nc	2.19E+00	nc	2.09E+00	NA	1.51E+04		T	1.51E+04
2-Nitrotoluene	88-72-2	3.65E+01	nc	3.65E+01	nc	3.65E+01	NA	NA			NA
3-Nitrotoluene	99-08-1	3.65E+01	nc	7.30E+01	nc	3.65E+01	NA	NA			NA
4-Nitrotoluene	99-99-0	3.65E+01	nc	3.65E+01	nc	3.65E+01	NA	3.37E+04		T	3.37E+04
Nitroglycerine	55-63-0	4.80E-01	c	4.47E-01	c	4.80E-01	NA	NA			NA
1,3-Dinitrobenzene	99-65-0	3.65E-01	nc	3.65E-01	nc	3.65E-01	NA	3.00E+03		T	3.00E+03
2,6-Dinitrotoluene	606-20-2	3.65E+00	nc	3.65E+00	nc	3.65E+00	NA	6.00E+02		T	6.00E+02
2,4-Dinitrotoluene	121-14-2	7.30E+00	nc	7.30E+00	nc	7.30E+00	NA	6.00E+02	NA	T	6.00E+02
1,3,5-Trinitrobenzene	99-35-4	1.10E+02	nc	1.10E+02	nc	1.10E+02	NA	3.00E+04		T	3.00E+04
2,4,6-Trinitrotoluene	118-96-7	2.24E-01	c	2.09E-01	c	2.24E-01	NA	2.50E+04		T	2.50E+04
RDX	121-82-4	6.11E-02	c	5.69E-02	c	6.11E-02	NA	NA			NA
4-Amino-2,6-Dinitrotoluene	19406-51-0	NA		NA		NA	NA	NA			NA
2-Amino-2,6-Dinitrotoluene	35572-78-2	NA		NA		NA	NA	1.50E+04		T	1.50E+04
Tetryl	479-45-8	3.65E+01	nc	3.65E+01	nc	3.65E+01	NA	NA			NA
HMX	2691-41-0	1.83E+02	nc	1.83E+02	nc	1.83E+02	NA	NA			NA
Pentaerythritoltetranitrate	78-11-5	NA		NA		NA	NA	5.00E+01		T	5.00E+01
Dibutyl Phthalate	84-74-2	3.65E+02	nc	3.65E+02	nc	3.65E+02	NA	1.50E+04		T	1.50E+04
Diocetyl Phthalate	117-81-7	4.80E-01	c	4.47E-01	c	4.80E-01	NA	1.00E+04		T	1.00E+04
Diphenylamine	122-39-4	9.13E+01	nc	9.13E+01	nc	9.13E+01	NA	3.00E+04		T	3.00E+04

Footnotes:

PRG = Preliminary Remediation Goals

c = cancer

nc = non-cancer

RBC = Risk-Based Concentration

HBSL = Health-Based Screening Level

(E) ERPG = Emergency Response Planning Guidelines

(T) TEEL = Temporary Emergency Exposure Limits

(A) AEGL = Acute Exposure Guideline Level

ATV = Acute Toxicity Value

NA = Not Available

APPENDIX D

RISK ASSESSMENT DATA

Table D-1: Comparison of Modeled Air Concentrations with Health-Based Values (M17) - 100 meter location

Cartridge, 0.50 caliber, Tracer, M17 (M2) DODIC: A571									
Compound	C _{chronic} (µg/m ³)	Health-Based Screening Level (µg/m ³)	C _{chronic} / HBSL	> 1?	Maximum Concentration (µg/m ³)	C _{acute} (µg/m ³)	Acute Toxicity Value (µg/m ³)	C _{acute} / ATV	> 1?
Permanent Gases									
Ammonia (NH3)	1.67E+01	1.04E+02	1.60E-01	no	1.36E+03	1.36E+03	1.75E+04	7.76E-02	no
Carbon Dioxide (CO2)	4.34E+02	NV		na	1.41E+05	1.41E+05	5.40E+07	2.61E-03	no
Carbon Monoxide (CO)	8.34E+02	1.00E+04	8.34E-02	no	6.77E+04	6.77E+04	2.30E+05	2.94E-01	no
Oxides of Nitrogen (as NO)	3.67E+01	1.00E+02	3.67E-01	no	1.19E+04	1.19E+04	3.08E+04	3.87E-01	no
Sulfur Dioxide (SO2)	2.16E-01	8.00E+01	2.70E-03	no	1.75E+01	1.75E+01	7.89E+02	2.22E-02	no
Acid Gases									
Hydrogen fluoride	NA	NV		na		NA	1.60E+03		na
Hydrogen chloride	NA	2.08E+01		na		NA	4.50E+03		na
Hydrogen bromide	NA	NV		na		NA	9.93E+03		na
Nitric Acid	NA	NV		na		NA	1.30E+03		na
Phosphoric acid	NA	1.04E+01		na		NA	3.00E+03		na
Sulfuric Acid	3.55E-01	NV		na	2.88E+01	2.88E+01	2.00E+03	1.44E-02	no
Cyanide									
Particulate Cyanide	5.39E-01	7.30E+01	7.38E-03	no	1.75E+02	1.75E+02	5.00E+03	3.50E-02	no
Hydrogen Cyanide	6.69E+00	3.13E+00	2.14E+00	yes	2.17E+03	2.17E+03	5.17E+03	4.20E-01	no
Particulates									
Total Suspended Particulate									
PM10	3.14E+01	5.00E+01	6.28E-01	no	2.55E+03	2.55E+03	NA		na
PM2.5	2.88E+01	5.00E+01	5.76E-01	no	2.34E+03	2.34E+03	NA		na
	1.68E+01	1.50E+01	1.12E+00	yes	1.36E+03	1.36E+03	NA		na
Metals									
Aluminum	6.65E-02	5.11E+00	1.30E-02	no	2.16E+01	2.16E+01	3.00E+04	7.19E-04	no
Antimony	5.24E-01	1.46E+00	3.59E-01	no	1.70E+02	1.70E+02	1.50E+03	1.13E-01	no
Arsenic	NA	4.47E-04		na		NA	3.00E+01		na
Barium	8.19E-01	5.21E-01	1.57E+00	yes	2.66E+02	2.66E+02	1.50E+03	1.77E-01	no
Beryllium	NA	8.00E-04		na		NA	5.00E+00		na
Cadmium	NA	1.07E-03		na		NA	3.00E+01		na
Calcium	2.66E-01	NV		na	8.65E+01	8.65E+01	3.00E+04	2.88E-03	no
Chromium	NA	1.53E-04		na		NA	1.50E+03		na
Cobalt	NA	2.20E+02		na		NA	6.00E+01		na
Copper	1.02E+01	1.46E+02	6.97E-02	no	3.30E+03	3.30E+03	3.00E+03	1.10E+00	yes
Lead	1.06E+00	1.50E+00	7.07E-01	no	3.44E+02	3.44E+02	1.50E+02	2.29E+00	yes
Magnesium	2.10E-01	NV		na	6.82E+01	6.82E+01	3.00E+04	2.27E-03	no
Manganese	NA	5.11E-02		na		NA	3.00E+03		na
Nickel	NA	7.30E+01		na		NA	3.00E+03		na
Selenium	NA	1.83E+01		na		NA	6.00E+02		na

Table D-1: Comparison of Modeled Air Concentrations with Health-Based Values (M17) - 100 meter location

Cartridge 0.50 caliber, Tracer, M17 (M2) DODIC: A571									
Compound	C _{chronic} (µg/m ³)	Health-Based Screening Level (µg/m ³)	C _{chronic} / HBSL	> 1?	Maximum Concentration (µg/m ³)	C _{acute} (µg/m ³)	Acute Toxicity Value (µg/m ³)	C _{acute} / ATV	> 1?
Silver	NA	1.83E+01		na		NA	3.00E+02		na
Thallium	NA	2.56E-01		na		NA	3.00E+02		na
Vanadium	NA	2.56E+01		na		NA	1.50E+02		na
Zinc	1.68E+00	1.10E+03	1.53E-03	no	5.44E+02	5.44E+02	3.00E+04	1.81E-02	no
TO-11 Carbonyls									
Formaldehyde	9.74E-03	1.48E-01	6.59E-02	no	1.84E+00	1.84E+00	1.23E+03	1.50E-03	no
Acetaldehyde	NA	8.73E-01		na		NA	1.80E+04		na
Acetone	NA	3.65E+02		na		NA	2.37E+06		na
Acrolein	NA	2.09E-02		na		NA	2.30E+02		na
Propionaldehyde	NA	NV		na		NA	7.50E+04		na
Crotonaldehyde	NA	3.54E-03		na		NA	5.72E+03		na
Butyraldehyde	NA	NV		na		NA	7.38E+04		na
Benzaldehyde	NA	3.65E+02		na		NA	1.50E+04		na
Isovaleraldehyde	NA	NV		na		NA	NA		na
Valeraldehyde	NA	NV		na		NA	NA		na
o,m,p-Tolualdehyde	NA	NV		na		NA	NA		na
Hexaldehyde	NA	NV		na		NA	NA		na
2,5-Dimethylbenzaldehyde	NA	NV		na		NA	NA		na
VOCs									
Propene	5.45E-02	NV		na	4.42E+00	4.42E+00	NA		na
Dichlorodifluoromethane	NA	2.09E+02		na		NA	1.48E+07		na
Chlorodifluoromethane	NA	5.11E+04		na		NA	4.41E+06		na
Freon 114	NA	NV		na		NA	2.10E+07		na
Chloromethane	1.44E-04	1.07E+00	1.35E-04	no	1.09E-01	1.09E-01	2.06E+05	5.30E-07	no
Vinyl Chloride	NA	2.20E-02		na		NA	1.28E+04		na
1,3-Butadiene	5.25E-03	3.74E-03	1.41E+00	yes	9.94E-01	9.94E-01	2.20E+04	4.52E-05	no
Bromomethane	NA	5.21E+00		na		NA	5.82E+04		na
Chloroethane	NA	2.32E+00		na		NA	2.64E+06		na
Dichlorofluoromethane	NA	2.09E+02		na		NA	1.48E+07		na
Trichlorofluoromethane	1.69E-04	7.30E+02	2.32E-07	no	5.49E-02	5.49E-02	2.81E+06	1.96E-08	no
Pentane	8.19E-04	NV		na	2.66E-01	2.66E-01	1.80E+06	1.48E-07	no
Acrolein	NA	2.09E-02		na		NA	2.30E+02		na
1,1-Dichloroethene	NA	5.21E+02		na		NA	7.92E+04		na
Freon 113	NA	3.13E+04		na		NA	9.58E+06		na
Acetone	8.37E-02	3.65E+02	2.29E-04	no	2.71E+01	2.71E+01	2.37E+06	1.15E-05	no
Methyl Iodide	NA	NV		na		NA	1.45E+05		na

Table D-1: Comparison of Modeled Air Concentrations with Health-Based Values (M17) - 100 meter location

Cartridge, 0.50 caliber, Tracer, M17 (M2) DODIC: A571									
Compound	C _{chronic} (µg/m ³)	Health-Based Screening Level (µg/m ³)	C _{chronic} / HBSL	> 1?	Maximum Concentration (µg/m ³)	C _{acute} (µg/m ³)	Acute Toxicity Value (µg/m ³)	C _{acute} / ATV	> 1?
Carbon Disulfide	NA	7.30E+02		na		NA	3.11E+04		na
Acetonitrile	8.35E-02	6.20E+01	1.35E-03	no	2.71E+01	2.71E+01	1.01E+05	2.69E-04	no
3-Chloropropene	NA	1.04E+00		na		NA	9.39E+03		na
Methylene Chloride	3.93E-02	4.09E+00	9.60E-03	no	7.43E+00	7.43E+00	6.96E+05	1.07E-05	no
tert-Butyl Alcohol	NA	NV		na		NA	4.55E+05		na
Acrylonitrile	7.72E-03	2.83E-02	2.73E-01	no	1.46E+00	1.46E+00	2.17E+04	6.73E-05	no
trans-1,2-Dichloroethene	NA	7.30E+01		na		NA	4.95E+04		na
Methyl t-Butyl Ether	NA	3.13E+03		na		NA	4.32E+05		na
Hexane	NA	2.09E+02		na		NA	5.28E+05		na
1,1-Dichloroethane	NA	5.21E+02		na		NA	1.21E+06		na
Vinyl Acetate	NA	2.09E+02		na		NA	1.92E+04		na
cis-1,2-Dichloroethene	NA	3.65E+01		na		NA	7.92E+05		na
2-Butanone	NA	1.04E+03		na		NA	8.85E+05		na
Ethyl Acetate	9.90E-03	3.29E+03	3.02E-06	no	3.21E+00	3.21E+00	1.44E+06	2.23E-06	no
Methyl Acrylate	NA	1.10E+02		na		NA	NA		na
Chloroform	NA	8.35E-02		na		NA	9.76E+03		na
1,1,1-Trichloroethane	3.74E-04	1.04E+03	3.59E-07	no	3.03E-02	3.03E-02	1.94E+06	1.56E-08	no
Carbon Tetrachloride	NA	1.28E-01		na		NA	1.28E+05		na
1,2-Dichloroethane	2.40E-03	7.39E-02	3.25E-02	no	1.82E+00	1.82E+00	8.08E+03	2.25E-04	no
Benzene	1.23E-01	2.49E-01	4.93E-01	no	2.33E+01	2.33E+01	1.56E+05	1.49E-04	no
Isooctane (2,2,4-trimethylpentane)	NA	NV		na		NA	3.50E+05		na
Heptane	NA	NV		na		NA	1.80E+06		na
Trichloroethane	NA	1.04E+03		na		NA	1.94E+06		na
Ethyl Acrylate	NA	1.40E-01		na		NA	6.14E+04		na
1,2-Dichloropropane	NA	9.89E-02		na		NA	5.08E+05		na
Methyl Methacrylate	NA	7.30E+02		na		NA	4.09E+05		na
Dibromomethane	NA	3.65E+01		na		NA	2.50E+05		na
1,4-Dioxane	NA	6.11E-01		na		NA	9.00E+04		na
Bromodichloromethane	NA	1.08E-01		na		NA	4.00E+03		na
4-Methyl-2-Pentanone	NA	8.34E+01		na		NA	3.07E+05		na
Toluene	3.25E-02	4.02E+02	8.09E-05	no	2.63E+00	2.63E+00	1.88E+05	1.41E-05	no
Octane	NA	NV		na		NA	NA		na
trans-1,3-Dichloropropene	NA	5.17E-02		na		NA	NA		na
Ethyl Methacrylate	NA	3.29E+02		na		NA	NA		na
1,1,2-Trichloroethane	NA	1.20E-01		na		NA	1.64E+05		na
Tetrachloroethene	NA	3.31E+00		na		NA	6.78E+05		na

Table D-1: Comparison of Modeled Air Concentrations with Health-Based Values (M17) - 100 meter location

Cartridge, 0.50 caliber, Tracer, M17 (M2) DODIC: A571									
Compound	C _{chronic} (µg/m ³)	Health-Based Screening Level (µg/m ³)	C _{chronic} / HBSL	> 1?	Maximum Concentration (µg/m ³)	C _{acute} (µg/m ³)	Acute Toxicity Value (µg/m ³)	C _{acute} / ATV	> 1?
2-Hexanone	NA	5.11E+00		na		NA	4.09E+04		na
Dibromochloromethane	NA	8.00E-02		na		NA	6.00E+03		na
1,2-Dibromoethane	NA	8.73E-03		na		NA	1.54E+05		na
Chlorobenzene	NA	6.21E+01		na		NA	1.38E+05		na
1,1,1,2-Tetrachloroethane	NA	2.60E-01		na		NA	5.15E+04		na
Ethylbenzene	2.60E-03	1.06E+03	2.46E-06	no	8.45E-01	8.45E-01	5.43E+05	1.56E-06	no
m&p-Xylene	4.93E-03	7.30E+02	6.76E-06	no	1.60E+00	1.60E+00	6.51E+05	2.46E-06	no
o-Xylene	4.02E-03	7.30E+02	5.50E-06	no	1.30E+00	1.30E+00	6.51E+05	2.00E-06	no
Styrene	9.83E-03	1.06E+03	9.29E-06	no	7.98E-01	7.98E-01	2.13E+05	3.74E-06	no
Bromoform	NA	1.75E+00		na		NA	6.20E+03		na
Cumene	NA	4.02E+02		na		NA	2.46E+05		na
1,1,2,2-Tetrachloroethane	NA	3.31E-02		na		NA	2.06E+04		na
1,2,3-Trichloropropane	NA	9.61E-04		na		NA	6.03E+04		na
Bromobenzene	NA	1.04E+01		na		NA	4.82E+04		na
4-Ethyltoluene	NA	NV		na		NA	1.25E+05		na
1,3,5-Trimethylbenzene	NA	6.21E+00		na		NA	3.68E+05		na
Alpha Methyl Styrene	NA	2.56E+02		na		NA	NA		na
1,2,4-Trimethylbenzene	9.01E-04	6.21E+00	1.45E-04	no	2.92E-01	2.92E-01	1.80E+05	1.62E-06	no
1,3-Dichlorobenzene	NA	3.29E+00		na		NA	3.61E+04		na
1,4-Dichlorobenzene	NA	3.06E-01		na		NA	6.61E+05		na
Benzyl Chloride	NA	3.96E-02		na		NA	5.20E+03		na
1,2-Dichlorobenzene	NA	2.09E+02		na		NA	3.01E+05		na
Hexachloroethane	NA	4.80E-01		na		NA	2.90E+04		na
1,2,4-Trichlorobenzene	NA	2.08E+02		na		NA	3.71E+04		na
Hexachlorobutadiene	NA	8.73E-02		na		NA	3.21E+04		na
Hydrocarbons									
Methane	2.89E+00	NV		na	9.37E+02	9.37E+02	3.30E+06	2.84E-04	no
Ethylene	1.85E-01	NV		na	6.02E+01	6.02E+01	4.60E+05	1.31E-04	no
Acetylene	2.53E-02	NV		na	2.06E+00	2.06E+00	NA		na
Ethane	1.05E-01	NV		na	8.50E+00	8.50E+00	NA		na
Propylene	5.02E-02	NV		na	4.07E+00	4.07E+00	NA		na
Propane	NA	NV		na		NA	3.78E+06		na
Propyne (methyl acetylene)	NA	NV		na		NA	2.79E+06		na
Isobutane	NA	NV		na		NA	9.52E+05		na
1-Butene/Isobutylene (115-11-7)	NA	NV		na		NA	6.87E+06		na

Table D-1: Comparison of Modeled Air Concentrations with Health-Based Values (M17) - 100 meter location

Cartridge, 0.50-caliber, Tracer, M17 (M2) DODIG: A571									
Compound	C _{chronic} (µg/m ³)	Health-Based Screening Level (µg/m ³)	C _{chronic} / HBSL	> 1?	Maximum Concentration (µg/m ³)	C _{acute} (µg/m ³)	Acute Toxicity Value (µg/m ³)	C _{acute} / ATV	> 1?
1,3-Butadiene/butane	NA	3.74E-03		na		NA	2.20E+04		na
cis-butene	NA	NV		na		NA	1.72E+04		na
1-Butyne	NA	NV		na		NA	NA		na
trans-Butene	NA	NV		na		NA	1.72E+04		na
2-Butyne (crotonylene)	NA	NV		na		NA	NA		na
n-Pentane	NA	NV		na		NA	1.80E+06		na
n-Hexane	1.84E-01	2.10E+02	8.75E-04	no	5.96E+01	5.96E+01	5.28E+05	1.13E-04	no
SVOCs									
n-nitrosodimethylamine	NA	1.37E-04		na		NA	2.50E+03		na
bis(2-chloroethyl)ether	NA	5.82E-03		na		NA	5.85E+04		na
phenol	NA	2.19E+03		na		NA	3.85E+04		na
2-chlorophenol	NA	1.83E+01		na		NA	5.25E+03		na
1,3-Dichlorobenzene	NA	3.29E+00		na		NA	3.61E+04		na
1,4-dichlorobenzene	NA	3.06E-01		na		NA	6.61E+05		na
1,2-dichlorobenzene	NA	2.09E+02		na		NA	3.01E+05		na
benzyl alcohol	NA	1.10E+03		na		NA	5.53E+04		na
bis(2-chloroisopropyl)ether	NA	1.92E-01		na		NA	6.99E+04		na
2-methylphenol	NA	1.83E+02		na		NA	NA		na
hexachloroethane	NA	4.80E-01		na		NA	2.90E+04		na
n-nitroso-di-n-propylamine	NA	9.61E-04		na		NA	2.00E+02		na
4-methylphenol	NA	1.83E+02		na		NA	NA		na
nitrobenzene	NA	2.09E+00		na		NA	1.51E+04		na
isophorone	NA	7.08E+00		na		NA	2.83E+04		na
2-nitrophenol	NA	NV		na		NA	NA		na
2,4-dimethylphenol	NA	7.30E+01		na		NA	NA		na
bis(2-chloroethoxy)methane	NA	NV		na		NA	NA		na
2,4-dichlorophenol	NA	1.10E+01		na		NA	3.00E+04		na
1,2,4-trichlorobenzene	NA	2.08E+02		na		NA	3.71E+04		na
naphthalene	1.66E-02	3.13E+00	5.30E-03	no	5.38E+00	5.38E+00	7.86E+04	6.84E-05	no
4-chloroaniline	NA	1.46E+01		na		NA	3.00E+04		na
hexachlorobutadiene	NA	8.62E-02		na		NA	3.21E+04		na
4-chloro-3-methylphenol	NA	NV		na		NA	2.00E+04		na
2-methylnaphthalene	NA	7.30E+01		na		NA	2.00E+04		na
hexachlorocyclopentadiene	NA	7.30E-02		na		NA	2.23E+02		na
2,4,6-trichlorophenol	NA	1.10E+02		na		NA	3.00E+04		na
2,4,5-trichlorophenol	NA	3.65E+02		na		NA	3.00E+04		na

Table D-1: Comparison of Modeled Air Concentrations with Health-Based Values (M17) - 100 meter location

Cartridge, 0.50-caliber, Tracer, M17 (M2) DODIC: A571									
Compound	C _{chronic} (µg/m ³)	Health-Based Screening Level (µg/m ³)	C _{chronic} / HBSL	> 1?	Maximum Concentration (µg/m ³)	C _{acute} (µg/m ³)	Acute Toxicity Value (µg/m ³)	C _{acute} / ATV	> 1?
2-chloronaphthalene	NA	2.92E+02		na		NA	6.00E+02		na
2-nitroaniline	NA	2.09E-01		na		NA	NA		na
Acenaphthylene	NA	NV		na		NA	2.00E+02		na
dimethylphthalate	NA	3.65E+04		na		NA	1.50E+04		na
2,6-dinitrotoluene	NA	3.65E+00		na		NA	6.00E+02		na
acenaphthene	NA	2.19E+02		na		NA	1.25E+03		na
3-nitroaniline	NA	NV		na		NA	NA		na
2,4-dinitrophenol	NA	7.30E+00		na		NA	7.50E+03		na
dibenzofuran	NA	1.46E+01		na		NA	NA		na
2,4-dinitrotoluene	NA	7.30E+00		na		NA	6.00E+02		na
4-nitrophenol	NA	2.92E+01		na		NA	3.00E+04		na
Fluorene	NA	1.46E+02		na		NA	7.50E+04		na
4-chlorophenyl-phenylether	NA	NV		na		NA	NA		na
diethylphthalate	NA	2.92E+03		na		NA	1.50E+04		na
4-nitroaniline	NA	NV		na		NA	9.00E+03		na
4,6-dinitro-2-methylphenol	NA	3.65E-01		na		NA	5.00E+02		na
n-nitrosodiphenylamine(1)	NA	1.37E+00		na		NA	NA		na
4-bromophenyl-phenylether	NA	NV		na		NA	NA		na
hexachlorobenzene	NA	4.18E-03		na		NA	7.50E+01		na
pentachlorophenol	NA	5.60E-02		na		NA	1.50E+03		na
phenanthrene	NA	NV		na		NA	2.00E+03		na
anthracene	NA	1.10E+03		na		NA	6.00E+03		na
di-n-butylphthalate	NA	3.65E+02		na		NA	1.50E+04		na
fluoranthene	NA	1.46E+02		na		NA	3.00E+01		na
pyrene	NA	1.10E+02		na		NA	1.50E+04		na
butylbenzylphthalate	NA	7.30E+02		na		NA	5.00E+05		na
benzo(a)anthracene	NA	2.17E-02		na		NA	6.00E+02		na
chrysene	NA	2.17E+00		na		NA	2.00E+02		na
3,3-dichlorobenzidine	NA	1.50E-02		na		NA	6.21E+03		na
bis(2-ethylhexyl)phthalate	2.92E-01	4.80E-01	6.08E-01	no	2.21E+02	2.21E+02	1.00E+04	2.21E-02	no
di-n-octylphthalate	NA	7.30E+01		na		NA	1.50E+05		na
benzo(b)fluoranthene	NA	2.17E-02		na		NA	NA		na
benzo(k)fluoranthene	NA	2.17E-01		na		NA	NA		na
benzo(a)pyrene	NA	2.17E-03		na		NA	7.50E+03		na
indeno(1,2,3-cd)pyrene	NA	2.17E-02		na		NA	NA		na
dibenz(a,h)anthracene	NA	2.17E-03		na		NA	3.00E+04		na

Table D-1: Comparison of Modeled Air Concentrations with Health-Based Values (M17) - 100 meter location

Cartridge, 0.50 caliber, Tracer, M17 (M2) DODIC: A571									
Compound	C _{chronic} (µg/m ³)	Health-Based Screening Level (µg/m ³)	C _{chronic} / HBSL	> 1?	Maximum Concentration (µg/m ³)	C _{acute} (µg/m ³)	Acute Toxicity Value (µg/m ³)	C _{acute} / ATV	> 1?
benzo(g,h,i)perylene	NA	NV		na		NA	3.00E+04		na
TO-13 (PAHs)									
naphthalene	1.04E-02	3.13E+00	3.31E-03	no	3.36E+00	3.36E+00	7.86E+04	4.28E-05	no
acenaphthylene	7.28E-04	NV		na	2.36E-01	2.36E-01	2.00E+02	1.18E-03	no
Acenaphthene	8.54E-05	2.19E+02	3.90E-07	no	2.77E-02	2.77E-02	1.25E+03	2.22E-05	no
fluorene	3.03E-04	1.46E+02	2.08E-06	no	9.84E-02	9.84E-02	7.50E+04	1.31E-06	no
phenanthrene	2.45E-04	NV		na	7.94E-02	7.94E-02	2.00E+03	3.97E-05	no
anthracene	4.10E-05	1.10E+03	3.74E-08	no	1.33E-02	1.33E-02	6.00E+03	2.22E-06	no
fluoranthene	3.09E-04	1.46E+02	2.12E-06	no	1.00E-01	1.00E-01	3.00E+01	3.34E-03	no
pyrene	4.43E-04	1.10E+02	4.05E-06	no	1.44E-01	1.44E-01	1.50E+04	9.58E-06	no
benzo(a)anthracene	9.73E-05	2.17E-02	4.49E-03	no	7.37E-02	7.37E-02	6.00E+02	1.23E-04	no
chrysene	1.00E-04	2.17E+00	4.62E-05	no	7.59E-02	7.59E-02	2.00E+02	3.79E-04	no
benzo(b)fluoranthene	1.58E-04	2.17E-02	7.27E-03	no	2.98E-02	2.98E-02	NA		na
benzo(k)fluoranthene	9.20E-05	2.17E-01	4.24E-04	no	1.74E-02	1.74E-02	NA		na
Benzo(e)pyrene	4.43E-04	NV		na	3.60E-02	3.60E-02	NA		na
benzo(a)pyrene	1.36E-04	2.17E-03	6.28E-02	no	1.03E-01	1.03E-01	7.50E+03	1.38E-05	no
indeno(1,2,3-cd)pyrene	1.47E-04	2.17E-02	6.78E-03	no	2.78E-02	2.78E-02	NA		na
dibenz(a,h)anthracene	1.65E-05	2.17E-03	7.60E-03	no	1.25E-02	1.25E-02	3.00E+04	4.16E-07	no
benzo(g,h,i)perylene	6.53E-04	NV		na	2.12E-01	2.12E-01	3.00E+04	7.06E-06	no
Dioxins and Furans									
2378-Tetrachlorodibenzo-p-dioxin	NA	4.48E-08		na		NA	3.50E+00		na
12378-Pentachlorodibenzo-p-dioxin	NA	NV		na		NA	2.50E+00		na
123478-Hexachlorodibenzo-p-dioxin	NA	NV		na		NA	NA		na
123678-Hexachlorodibenzo-p-dioxin	NA	NV		na		NA	1.50E+01		na
123789-Hexachlorodibenzo-p-dioxin	NA	1.48E-06		na		NA	NA		na
1234678-Heptachlorodibenzo-p-dioxin	8.56E-09	NV		na	6.94E-07	6.94E-07	NA		na
OCDD	1.44E-07	NV		na	4.68E-05	4.68E-05	1.50E+02	3.12E-07	no
2378-Tetrachlorodibenzo-p-furan	NA	NV		na		NA	2.00E+00		na
12378-Pentachlorodibenzo-p-furan	NA	NV		na		NA	NA		na
23478-Pentachlorodibenzo-o-furan	NA	NV		na		NA	7.50E-02		na
123478-Hexachlorodibenzo-p-furan	NA	NV		na		NA	7.50E+00		na
123678-Hexachlorodibenzo-p-furan	NA	NV		na		NA	2.50E+00		na
123789-Hexachlorodibenzo-p-furan	NA	NV		na		NA	NA		na
234678-Hexachlorodibenzo-p-furan	NA	NV		na		NA	1.50E+00		na
1234678-Heptachlorodibenzo-p-furan	2.57E-09	NV		na	2.09E-07	2.09E-07	NA		na

Table D-1: Comparison of Modeled Air Concentrations with Health-Based Values (M17) - 100 meter location

Cartridge, 0.50 caliber, Tracer, M17 (M2) DODIC: A571									
Compound	C _{chronic} (µg/m ³)	Health-Based Screening Level (µg/m ³)	C _{chronic} / HBSL	> 1?	Maximum Concentration (µg/m ³)	C _{acute} (µg/m ³)	Acute Toxicity Value (µg/m ³)	C _{acute} / ATV	> 1?
1234789-Heptachlorodibenzo-p-furan	NA	NV		na		NA	NA		na
OCDF	8.82E-09	NV		na	2.86E-06	2.86E-06	3.00E+02	9.54E-09	no
Energetics									
Nitrobenzene	NA	2.09E+00		na		NA	1.51E+04		na
2-Nitrotoluene	NA	3.65E+01		na		NA	NA		na
3-Nitrotoluene	NA	3.65E+01		na		NA	NA		na
4-Nitrotoluene	NA	3.65E+01		na		NA	3.37E+04		na
Nitroglycerine	NA	4.80E-01		na		NA	NA		na
1,3-Dinitrobenzene	NA	3.65E-01		na		NA	3.00E+03		na
2,6-Dinitrotoluene	NA	3.65E+00		na		NA	6.00E+02		na
2,4-Dinitrotoluene	NA	7.30E+00		na		NA	6.00E+02		na
1,3,5-Trinitrobenzene	NA	1.10E+02		na		NA	3.00E+04		na
2,4,6-Trinitrotoluene	NA	2.24E-01		na		NA	2.50E+04		na
RDX	NA	6.11E-02		na		NA	NA		na
4-Amino-2,6-Dinitrotoluene	NA	NV		na		NA	NA		na
2-Amino-2,6-Dinitrotoluene	NA	NV		na		NA	1.50E+04		na
Tetryl	NA	3.65E+01		na		NA	NA		na
HMX	NA	1.83E+02		na		NA	NA		na
Pentaerythritoltrinitrate	NA	NV		na		NA	5.00E+01		na
Dibutyl Phthalate	NA	3.65E+02		na		NA	1.50E+04		na
Diocyl Phthalate	NA	4.80E-01		na		NA	1.00E+04		na
Diphenylamine	NA	9.13E+01		na		NA	3.00E+04		na

Footnotes:

NA: Not applicable because compound was not detected

na: Not available because health-based screening value is not available or not applicable if compound was not detected

NV: No value available

C_{chronic}: Chronic time-averaged concentration

HBSL: Chronic health-based screening level

C_{acute}: Acute concentration

ATV: Acute toxicity value

Footnotes:

NA: Not applicable because compound was not detected

na: Not available because health-based screening value is not available or not applicable if compound was not detected

NV: No value available

C_{chronic}: Chronic time-averaged concentration

HBSL: Chronic health-based screening level

C_{acute}: Acute concentration

ATV: Acute toxicity value

Cartridge 0.50 caliber, Tracer M17 (M2)
DODIC A571

Footnotes:
>1? = Is the ratio greater than one?
NA = Not Applicable because compound was not detected
C_{chronic} = chronic averaged air Concentration
HBSL = Health-Based Screening Level

Table D-3: Comparison of Modeled Air Concentrations with Health-Based Values (M17) - 200 meter location

Cartridge 0.50 caliber Tracer, M17 (M2) DODIC A571								
Compound	C _{chronic} (µg/m ³)	Health-Based Screening Level (µg/m ³)	C _{chronic} / HBSL	> 1?	C _{acute} (µg/m ³)	Acute Toxicity Value (µg/m ³)	C _{acute} /ATV	> 1?
Permanent Gases								
Ammonia (NH3)	7.03E+00	1.04E+02	6.74E-02	no	5.70E+02	1.75E+04	3.26E-02	no
Carbon Dioxide (CO2)	1.82E+02	NV		na	5.92E+04	5.40E+07	1.10E-03	no
Carbon Monoxide (CO)	3.50E+02	1.00E+04	3.50E-02	no	2.84E+04	2.30E+05	1.24E-01	no
Oxides of Nitrogen (as NO)	1.54E+01	1.00E+02	1.54E-01	no	5.00E+03	3.08E+04	1.62E-01	no
Sulfur Dioxide (SO2)	9.07E-02	8.00E+01	1.13E-03	no	7.36E+00	7.89E+02	9.32E-03	no
Acid Gases								
Hydrogen fluoride	NA	NV		na	NA	1.60E+03		na
Hydrogen chloride	NA	2.08E+01		na	NA	4.50E+03		na
Hydrogen bromide	NA	NV		na	NA	9.93E+03		na
Nitric Acid	NA	NV		na	NA	1.30E+03		na
Phosphoric acid	NA	1.04E+01		na	NA	3.00E+03		na
Sulfuric Acid	1.49E-01	NV		na	1.21E+01	2.00E+03	6.05E-03	no
Cyanide								
Particulate Cyanide	2.26E-01	7.30E+01	3.10E-03	no	7.34E+01	5.00E+03	1.47E-02	no
Hydrogen Cyanide	2.81E+00	3.13E+00	8.98E-01	no	9.12E+02	5.17E+03	1.76E-01	no
Particulates								
Total Suspended Particulate	1.32E+01	5.00E+01	2.64E-01	no	1.07E+03	NA		na
PM10	1.21E+01	5.00E+01	2.42E-01	no	9.82E+02	NA		na
PM2.5	7.06E+00	1.50E+01	4.70E-01	no	5.72E+02	NA		na
Metals								
Aluminum	2.79E-02	5.11E+00	5.46E-03	no	9.06E+00	3.00E+04	3.02E-04	no
Antimony	2.20E-01	1.46E+00	1.51E-01	no	7.14E+01	1.50E+03	4.76E-02	no
Arsenic	NA	4.47E-04		na	NA	3.00E+01		na
Barium	3.44E-01	5.21E-01	6.59E-01	no	1.12E+02	1.50E+03	7.44E-02	no
Beryllium	NA	8.00E-04		na	NA	5.00E+00		na
Cadmium	NA	1.07E-03		na	NA	3.00E+01		na
Calcium	1.12E-01	NV		na	3.63E+01	3.00E+04	1.21E-03	no
Chromium	NA	1.53E-04		na	NA	1.50E+03		na
Cobalt	NA	2.20E+02		na	NA	6.00E+01		na
Copper	4.27E+00	1.46E+02	2.93E-02	no	1.39E+03	3.00E+03	4.62E-01	no
Lead	4.45E-01	1.50E+00	2.97E-01	no	1.45E+02	1.50E+02	9.63E-01	no
Magnesium	8.83E-02	NV		na	2.86E+01	3.00E+04	9.55E-04	no
Manganese	NA	5.11E-02		na	NA	3.00E+03		na

Table D-3: Comparison of Modeled Air Concentrations with Health-Based Values (M17) - 200 meter location

Cartridge, 0.50 caliber, Tracer, M17 (M2) DODIC: A571								
Compound	C _{chronic} (µg/m ³)	Health-Based Screening Level (µg/m ³)	C _{chronic} / HBSL	> 1?	C _{acute} (µg/m ³)	Acute Toxicity Value (µg/m ³)	C _{acute} /ATV	> 1?
Nickel	NA	7.30E+01		na	NA	3.00E+03		na
Selenium	NA	1.83E+01		na	NA	6.00E+02		na
Silver	NA	1.83E+01		na	NA	3.00E+02		na
Thallium	NA	2.56E-01		na	NA	3.00E+02		na
Vanadium	NA	2.56E+01		na	NA	1.50E+02		na
Zinc	7.04E-01	1.10E+03	6.43E-04	no	2.28E+02	3.00E+04	7.61E-03	no
TO-11 Carbonyls								
Formaldehyde	4.09E-03	1.48E-01	2.77E-02	no	7.74E-01	1.23E+03	6.29E-04	no
Acetaldehyde	NA	8.73E-01		na	NA	1.80E+04		na
Acetone	NA	3.65E+02		na	NA	2.37E+06		na
Acrolein	NA	2.09E-02		na	NA	2.30E+02		na
Propionaldehyde	NA	NV		na	NA	7.50E+04		na
Crotonaldehyde	NA	3.54E-03		na	NA	5.72E+03		na
Butyraldehyde	NA	NV		na	NA	7.38E+04		na
Benzaldehyde	NA	3.65E+02		na	NA	1.50E+04		na
Isovaleraldehyde	NA	NV		na	NA	NA		na
Valeraldehyde	NA	NV		na	NA	NA		na
o,m,p-Tolualdehyde	NA	NV		na	NA	NA		na
Hexaldehyde	NA	NV		na	NA	NA		na
2,5-Dimethylbenzaldehyde	NA	NV		na	NA	NA		na
VOCs								
Propene	2.29E-02	NV		na	1.86E+00	NA		na
Dichlorodifluoromethane	NA	2.09E+02		na	NA	1.48E+07		na
Chlorodifluoromethane	NA	5.11E+04		na	NA	4.41E+06		na
Freon 114	NA	NV		na	NA	2.10E+07		na
Chloromethane	6.05E-05	1.07E+00	5.67E-05	no	4.58E-02	2.06E+05	2.22E-07	no
Vinyl Chloride	NA	2.20E-02		na	NA	1.28E+04		na
1,3-Butadiene	2.21E-03	3.74E-03	5.91E-01	no	4.18E-01	2.20E+04	1.90E-05	no
Bromomethane	NA	5.21E+00		na	NA	5.82E+04		na
Chloroethane	NA	2.32E+00		na	NA	2.64E+06		na
Dichlorofluoromethane	NA	2.09E+02		na	NA	1.48E+07		na
Trichlorofluoromethane	7.11E-05	7.30E+02	9.74E-08	no	2.31E-02	2.81E+06	8.22E-09	no
Pentane	3.44E-04	NV		na	1.12E-01	1.80E+06	6.20E-08	no
Acrolein	NA	2.09E-02		na	NA	2.30E+02		na

Table D-3: Comparison of Modeled Air Concentrations with Health-Based Values (M17) - 200 meter location

Cartridge, 0.50 caliber, Tracer, M17 (M2) DODIC: A571								
Compound	C _{chronic} (µg/m ³)	Health-Based Screening Level (µg/m ³)	C _{chronic} / HBSL	> 1?	C _{acute} (µg/m ³)	Acute Toxicity Value (µg/m ³)	C _{acute} /ATV	> 1?
1,1-Dichloroethene	NA	5.21E+02		na	NA	7.92E+04		na
Freon 113	NA	3.13E+04		na	NA	9.58E+06		na
Acetone	3.51E-02	3.65E+02	9.63E-05	no	1.14E+01	2.37E+06	4.81E-06	no
Methyl Iodide	NA	NV		na	NA	1.45E+05		na
Carbon Disulfide	NA	7.30E+02		na	NA	3.11E+04		na
Acetonitrile	3.51E-02	6.20E+01	5.66E-04	no	1.14E+01	1.01E+05	1.13E-04	no
3-Chloropropene	NA	1.04E+00		na	NA	9.39E+03		na
Methylene Chloride	1.65E-02	4.09E+00	4.03E-03	no	3.12E+00	6.96E+05	4.48E-06	no
tert-Butyl Alcohol	NA	NV		na	NA	4.55E+05		na
Acrylonitrile	3.24E-03	2.83E-02	1.15E-01	no	6.13E-01	2.17E+04	2.83E-05	no
trans-1,2-Dichloroethene	NA	7.30E+01		na	NA	4.95E+04		na
Methyl t-Butyl Ether	NA	3.13E+03		na	NA	4.32E+05		na
Hexane	NA	2.09E+02		na	NA	5.28E+05		na
1,1-Dichloroethane	NA	5.21E+02		na	NA	1.21E+06		na
Vinyl Acetate	NA	2.09E+02		na	NA	1.92E+04		na
cis-1,2-Dichloroethene	NA	3.65E+01		na	NA	7.92E+05		na
2-Butanone	NA	1.04E+03		na	NA	8.85E+05		na
Ethyl Acetate	4.16E-03	3.29E+03	1.27E-06	no	1.35E+00	1.44E+06	9.37E-07	no
Methyl Acrylate	NA	1.10E+02		na	NA	NA		na
Chloroform	NA	8.35E-02		na	NA	9.76E+03		na
1,1,1-Trichloroethane	1.57E-04	1.04E+03	1.51E-07	no	1.27E-02	1.94E+06	6.56E-09	no
Carbon Tetrachloride	NA	1.28E-01		na	NA	1.28E+05		na
1,2-Dichloroethane	1.01E-03	7.39E-02	1.36E-02	no	7.63E-01	8.08E+03	9.44E-05	no
Benzene	5.16E-02	2.49E-01	2.07E-01	no	9.77E+00	1.56E+05	6.26E-05	no
Isooctane (2,2,4-trimethylpentane)	NA	NV		na	NA	3.50E+05		na
Heptane	NA	NV		na	NA	1.80E+06		na
Trichloroethane	NA	1.04E+03		na	NA	1.94E+06		na
Ethyl Acrylate	NA	1.40E-01		na	NA	6.14E+04		na
1,2-Dichloropropane	NA	9.89E-02		na	NA	5.08E+05		na
Methyl Methacrylate	NA	7.30E+02		na	NA	4.09E+05		na
Dibromomethane	NA	3.65E+01		na	NA	2.50E+05		na
1,4-Dioxane	NA	6.11E-01		na	NA	9.00E+04		na
Bromodichloromethane	NA	1.08E-01		na	NA	4.00E+03		na
4-Methyl-2-Pentanone	NA	8.34E+01		na	NA	3.07E+05		na

Table D-3: Comparison of Modeled Air Concentrations with Health-Based Values (M17) - 200 meter location

Cartridge, 0.50 caliber, Tracer, M17 (M2) DODIC: A571								
Compound	C _{chronic} (µg/m ³)	Health-Based Screening Level (µg/m ³)	C _{chronic} / HBSL	> 1?	C _{acute} (µg/m ³)	Acute Toxicity Value (µg/m ³)	C _{acute} /ATV	> 1?
Toluene	1.36E-02	4.02E+02	3.40E-05	no	1.11E+00	1.88E+05	5.90E-06	no
Octane	NA	NV		na	NA	NA		na
trans-1,3-Dichloropropene	NA	5.17E-02		na	NA	NA		na
Ethyl Methacrylate	NA	3.29E+02		na	NA	NA		na
1,1,2-Trichloroethane	NA	1.20E-01		na	NA	1.64E+05		na
Tetrachloroethene	NA	3.31E+00		na	NA	6.78E+05		na
2-Hexanone	NA	5.11E+00		na	NA	4.09E+04		na
Dibromochloromethane	NA	8.00E-02		na	NA	6.00E+03		na
1,2-Dibromoethane	NA	8.73E-03		na	NA	1.54E+05		na
Chlorobenzene	NA	6.21E+01		na	NA	1.38E+05		na
1,1,1,2-Tetrachloroethane	NA	2.60E-01		na	NA	5.15E+04		na
Ethylbenzene	1.09E-03	1.06E+03	1.03E-06	no	3.55E-01	5.43E+05	6.54E-07	no
m&p-Xylene	2.07E-03	7.30E+02	2.84E-06	no	6.72E-01	6.51E+05	1.03E-06	no
o-Xylene	1.69E-03	7.30E+02	2.31E-06	no	5.47E-01	6.51E+05	8.41E-07	no
Styrene	4.13E-03	1.06E+03	3.90E-06	no	3.35E-01	2.13E+05	1.57E-06	no
Bromoform	NA	1.75E+00		na	NA	6.20E+03		na
Cumene	NA	4.02E+02		na	NA	2.46E+05		na
1,1,2,2-Tetrachloroethane	NA	3.31E-02		na	NA	2.06E+04		na
1,2,3-Trichloropropane	NA	9.61E-04		na	NA	6.03E+04		na
Bromobenzene	NA	1.04E+01		na	NA	4.82E+04		na
4-Ethyltoluene	NA	NV		na	NA	1.25E+05		na
1,3,5-Trimethylbenzene	NA	6.21E+00		na	NA	3.68E+05		na
Alpha Methyl Styrene	NA	2.56E+02		na	NA	NA		na
1,2,4-Trimethylbenzene	3.78E-04	6.21E+00	6.10E-05	no	1.23E-01	1.80E+05	6.82E-07	no
1,3-Dichlorobenzene	NA	3.29E+00		na	NA	3.61E+04		na
1,4-Dichlorobenzene	NA	3.06E-01		na	NA	6.61E+05		na
Benzyl Chloride	NA	3.96E-02		na	NA	5.20E+03		na
1,2-Dichlorobenzene	NA	2.09E+02		na	NA	3.01E+05		na
Hexachlorethane	NA	4.80E-01		na	NA	2.90E+04		na
1,2,4-Trichlorobenzene	NA	2.08E+02		na	NA	3.71E+04		na
Hexachlorobutadiene	NA	8.73E-02		na	NA	3.21E+04		na
Hydrocarbons Methane	1.21E+00	NV		na	3.93E+02	3.30E+06	1.19E-04	no

Table D-3: Comparison of Modeled Air Concentrations with Health-Based Values (M17) - 200 meter location

Cartridge, 0.50 caliber, Tracer, M17 (M2) DODIC: A571								
Compound	C _{chronic} (µg/m ³)	Health-Based Screening Level (µg/m ³)	C _{chronic} / HBSL	> 1?	C _{acute} (µg/m ³)	Acute Toxicity Value (µg/m ³)	C _{acute} / ATV	> 1?
Ethylene	7.79E-02	NV		na	2.53E+01	4.60E+05	5.49E-05	no
Acetylene	1.06E-02	NV		na	8.63E-01	NA		na
Ethane	4.40E-02	NV		na	3.57E+00	NA		na
Propylene	2.11E-02	NV		na	1.71E+00	NA		na
Propane	NA	NV		na	NA	3.78E+06		na
Propyne (methyl acetylene)	NA	NV		na	NA	2.79E+06		na
Isobutane	NA	NV		na	NA	9.52E+05		na
1-Butene/Isobutylene (115-11-7)	NA	NV		na	NA	6.87E+06		na
1,3-Butadiene/butane	NA	3.74E-03		na	NA	2.20E+04		na
cis-butene	NA	NV		na	NA	1.72E+04		na
1-Butyne	NA	NV		na	NA	NA		na
trans-Butene	NA	NV		na	NA	1.72E+04		na
2-Butyne (crotonylene)	NA	NV		na	NA	NA		na
n-Pentane	NA	NV		na	NA	1.80E+06		na
n-Hexane	7.72E-02	2.10E+02	3.68E-04	no	2.50E+01	5.28E+05	4.74E-05	no
SVOCs								
n-nitrosodimethylamine	NA	1.37E-04		na	NA	2.50E+03		na
bis(2-chloroethyl)ether	NA	5.82E-03		na	NA	5.85E+04		na
phenol	NA	2.19E+03		na	NA	3.85E+04		na
2-chlorophenol	NA	1.83E+01		na	NA	5.25E+03		na
1,3-Dichlorobenzene	NA	3.29E+00		na	NA	3.61E+04		na
1,4-dichlorobenzene	NA	3.06E-01		na	NA	6.61E+05		na
1,2-dichlorobenzene	NA	2.09E+02		na	NA	3.01E+05		na
benzyl alcohol	NA	1.10E+03		na	NA	5.53E+04		na
bis(2-chloroisopropyl)ether	NA	1.92E-01		na	NA	6.99E+04		na
2-methylphenol	NA	1.83E+02		na	NA	NA		na
hexachloroethane	NA	4.80E-01		na	NA	2.90E+04		na
n-nitroso-di-n-propylamine	NA	9.61E-04		na	NA	2.00E+02		na
4-methylphenol	NA	1.83E+02		na	NA	NA		na
nitrobenzene	NA	2.09E+00		na	NA	1.51E+04		na
isophorone	NA	7.08E+00		na	NA	2.83E+04		na
2-nitrophenol	NA	NV		na	NA	NA		na
2,4-dimethylphenol	NA	7.30E+01		na	NA	NA		na
bis(2-chloroethoxy)methane	NA	NV		na	NA	NA		na

Table D-3: Comparison of Modeled Air Concentrations with Health-Based Values (M17) - 200 meter location

Cartridge, 0.50 caliber, Tracer, M17 (M2) DODIC: A571								
Compound	C _{chronic} (µg/m ³)	Health-Based Screening Level (µg/m ³)	C _{chronic} / HBSL	> 1?	C _{acute} (µg/m ³)	Acute Toxicity Value (µg/m ³)	C _{acute} / ATV	> 1?
2,4-dichlorophenol	NA	1.10E+01		na	NA	3.00E+04		na
1,2,4-trichlorobenzene	NA	2.08E+02		na	NA	3.71E+04		na
naphthalene	6.96E-03	3.13E+00	2.23E-03	no	2.26E+00	7.86E+04	2.87E-05	no
4-chloroaniline	NA	1.46E+01		na	NA	3.00E+04		na
hexachlorobutadiene	NA	8.62E-02		na	NA	3.21E+04		na
4-chloro-3-methylphenol	NA	NV		na	NA	2.00E+04		na
2-methylnaphthalene	NA	7.30E+01		na	NA	2.00E+04		na
hexachlorocyclopentadiene	NA	7.30E-02		na	NA	2.23E+02		na
2,4,6-trichlorophenol	NA	1.10E+02		na	NA	3.00E+04		na
2,4,5-trichlorophenol	NA	3.65E+02		na	NA	3.00E+04		na
2-chloronaphthalene	NA	2.92E+02		na	NA	6.00E+02		na
2-nitroaniline	NA	2.09E-01		na	NA	NA		na
Acenaphthylene	NA	NV		na	NA	2.00E+02		na
dimethylphthalate	NA	3.65E+04		na	NA	1.50E+04		na
2,6-dinitrotoluene	NA	3.65E+00		na	NA	6.00E+02		na
acenaphthene	NA	2.19E+02		na	NA	1.25E+03		na
3-nitroaniline	NA	NV		na	NA	NA		na
2,4-dinitrophenol	NA	7.30E+00		na	NA	7.50E+03		na
dibenzofuran	NA	1.46E+01		na	NA	NA		na
2,4-dinitrotoluene	NA	7.30E+00		na	NA	6.00E+02		na
4-nitrophenol	NA	2.92E+01		na	NA	3.00E+04		na
Fluorene	NA	1.46E+02		na	NA	7.50E+04		na
4-chlorophenyl-phenylether	NA	NV		na	NA	NA		na
diethylphthalate	NA	2.92E+03		na	NA	1.50E+04		na
4-nitroaniline	NA	NV		na	NA	9.00E+03		na
4,6-dinitro-2-methylphenol	NA	3.65E-01		na	NA	5.00E+02		na
n-nitrosodiphenylamine(1)	NA	1.37E+00		na	NA	NA		na
4-bromophenyl-phenylether	NA	NV		na	NA	NA		na
hexachlorobenzene	NA	4.18E-03		na	NA	7.50E+01		na
pentachlorophenol	NA	5.60E-02		na	NA	1.50E+03		na
phenanthrene	NA	NV		na	NA	2.00E+03		na
anthracene	NA	1.10E+03		na	NA	6.00E+03		na
di-n-butylphthalate	NA	3.65E+02		na	NA	1.50E+04		na
fluoranthene	NA	1.46E+02		na	NA	3.00E+01		na

Table D-3: Comparison of Modeled Air Concentrations with Health-Based Values (M17) - 200 meter location

Cartridge, 0.50 caliber, Tracer, M17 (M2) DODIC: A571							
Compound	C _{chronic} (µg/m ³)	Health-Based Screening Level (µg/m ³)	C _{chronic} / HBSL	> 1?	C _{acute} (µg/m ³)	Acute Toxicity Value (µg/m ³)	C _{acute} /ATV
pyrene	NA	1.10E+02		na	NA	1.50E+04	na
butylbenzylphthalate	NA	7.30E+02		na	NA	5.00E+05	na
benzo(a)anthracene	NA	2.17E-02		na	NA	6.00E+02	na
chrysene	NA	2.17E+00		na	NA	2.00E+02	na
3,3-dichlorobenzidine	NA	1.50E-02		na	NA	6.21E+03	na
bis(2-ethylhexyl)phthalate	1.23E-01	4.80E-01	2.55E-01	no	9.29E+01	1.00E+04	9.29E-03
di-n-octylphthalate	NA	7.30E+01		na	NA	1.50E+05	na
benzo(b)fluoranthene	NA	2.17E-02		na	NA	NA	na
benzo(k)fluoranthene	NA	2.17E-01		na	NA	NA	na
benzo(a)pyrene	NA	2.17E-03		na	NA	7.50E+03	na
indeno(1,2,3-cd)pyrene	NA	2.17E-02		na	NA	NA	na
dibenz(a,h)anthracene	NA	2.17E-03		na	NA	3.00E+04	na
benzo(g,h,i)perylene	NA	NV		na	NA	3.00E+04	na
TO-13 (PAHs)							
naphthalene	4.35E-03	3.13E+00	1.39E-03	no	1.41E+00	7.86E+04	1.80E-05
acenaphthylene	3.06E-04	NV		na	9.92E-02	2.00E+02	4.96E-04
Acenaphthene	3.59E-05	2.19E+02	1.64E-07	no	1.16E-02	1.25E+03	9.31E-06
fluorene	1.27E-04	1.46E+02	8.72E-07	no	4.13E-02	7.50E+04	5.51E-07
phenanthrene	1.03E-04	NV		na	3.33E-02	2.00E+03	1.67E-05
anthracene	1.72E-05	1.10E+03	1.57E-08	no	5.59E-03	6.00E+03	9.31E-07
fluoranthene	1.30E-04	1.46E+02	8.89E-07	no	4.21E-02	3.00E+01	1.40E-03
pyrene	1.86E-04	1.10E+02	1.70E-06	no	6.04E-02	1.50E+04	4.02E-06
benzo(a)anthracene	4.09E-05	2.17E-02	1.88E-03	no	3.09E-02	6.00E+02	5.16E-05
chrysene	4.21E-05	2.17E+00	1.94E-05	no	3.19E-02	2.00E+02	1.59E-04
benzo(b)fluoranthene	6.62E-05	2.17E-02	3.05E-03	no	1.25E-02	NA	na
benzo(k)fluoranthene	3.86E-05	2.17E-01	1.78E-04	no	7.31E-03	NA	na
Benzo(e)pyrene	1.86E-04	NV		na	1.51E-02	NA	na
benzo(a)pyrene	5.72E-05	2.17E-03	2.64E-02	no	4.33E-02	7.50E+03	5.78E-06
indeno(1,2,3-cd)pyrene	6.17E-05	2.17E-02	2.85E-03	no	1.17E-02	NA	na
dibenz(a,h)anthracene	6.92E-06	2.17E-03	3.19E-03	no	5.24E-03	3.00E+04	1.75E-07
benzo(g,h,i)perylene	2.74E-04	NV		na	8.90E-02	3.00E+04	2.97E-06
Dioxins and Furans							
2378-Tetrachlorodibenzo-p-dioxin	NA	4.48E-08		na	NA	3.50E+00	na

Table D-3: Comparison of Modeled Air Concentrations with Health-Based Values (M17) - 200 meter location

Cartridge, 0.50 caliber, Tracer, M17 (M2) DODIC: A571								
Compound	C _{chronic} (µg/m ³)	Health-Based Screening Level (µg/m ³)	C _{chronic} / HBSL	> 1?	C _{acute} (µg/m ³)	Acute Toxicity Value (µg/m ³)	C _{acute} / ATV	> 1?
12378-Pentachlorodibenzo-p-dioxin	NA	NV		na	NA	2.50E+00		na
123478-Hexachlorodibenzo-p-dioxin	NA	NV		na	NA	NA		na
123678-Hexachlorodibenzo-p-dioxin	NA	NV		na	NA	1.50E+01		na
123789-Hexachlorodibenzo-p-dioxin	NA	1.48E-06		na	NA	NA		na
1234678-Heptachlorodibenzo-p-dioxin	3.59E-09	NV		na	2.92E-07	NA		na
OCDD	6.05E-08	NV		na	1.96E-05	1.50E+02	1.31E-07	no
2378-Tetrachlorodibenzo-p-furan	NA	NV		na	NA	2.00E+00		na
12378-Pentachlorodibenzo-p-furan	NA	NV		na	NA	NA		na
23478-Pentachlorodibenzo-o-furan	NA	NV		na	NA	7.50E-02		na
123478-Hexachlorodibenzo-p-furan	NA	NV		na	NA	7.50E+00		na
123678-Hexachlorodibenzo-p-furan	NA	NV		na	NA	2.50E+00		na
123789-Hexachlorodibenzo-p-furan	NA	NV		na	NA	NA		na
234678-Hexachlorodibenzo-p-furan	NA	NV		na	NA	1.50E+00		na
1234678-Heptachlorodibenzo-p-furan	1.08E-09	NV		na	8.77E-08	NA		na
1234789-Heptachlorodibenzo-p-furan	NA	NV		na	NA	NA		na
OCDF	3.70E-09	NV		na	1.20E-06	3.00E+02	4.01E-09	no
Energetics								
Nitrobenzene	NA	2.09E+00		na	NA	1.51E+04		na
2-Nitrotoluene	NA	3.65E+01		na	NA	NA		na
3-Nitrotoluene	NA	3.65E+01		na	NA	NA		na
4-Nitrotoluene	NA	3.65E+01		na	NA	3.37E+04		na
Nitroglycerine	NA	4.80E-01		na	NA	NA		na
1,3-Dinitrobenzene	NA	3.65E-01		na	NA	3.00E+03		na
2,6-Dinitrotoluene	NA	3.65E+00		na	NA	6.00E+02		na
2,4-Dinitrotoluene	NA	7.30E+00		na	NA	6.00E+02		na
1,3,5-Trinitrobenzene	NA	1.10E+02		na	NA	3.00E+04		na
2,4,6-Trinitrotoluene	NA	2.24E-01		na	NA	2.50E+04		na
RDX	NA	6.11E-02		na	NA	NA		na
4-Amino-2,6-Dinitrotoluene	NA	NV		na	NA	NA		na
2-Amino-2,6-Dinitrotoluene	NA	NV		na	NA	1.50E+04		na
Tetryl	NA	3.65E+01		na	NA	NA		na
HMX	NA	1.83E+02		na	NA	NA		na
Pentaerythritoltetranitrate	NA	NV		na	NA	5.00E+01		na
Dibutyl Phthalate	NA	3.65E+02		na	NA	1.50E+04		na

Table D-3: Comparison of Modeled Air Concentrations with Health-Based Values (M17) - 200 meter location

Cartridge, 0.50 caliber, Tracer, M17 (M2) DODIC: A571								
Compound	C _{chronic} (µg/m ³)	Health-Based Screening Level (µg/m ³)	C _{chronic} / HBSL	> 1?	C _{acute} (µg/m ³)	Acute Toxicity Value (µg/m ³)	C _{acute} /ATV	> 1?
Diethyl Phthalate	NA	4.80E-01		na	NA	1.00E+04		na
Diphenylamine	NA	9.13E+01		na	NA	3.00E+04		na
Footnotes: NA: Not applicable because compound was not detected na: Not available because health-based screening value is not available or not applicable if compound was not detected NV: No value available C _{chronic} : Chronic time-averaged concentration HBSL: Chronic health-based screening level C _{acute} : Acute concentration ATV: Acute toxicity value								

Table D-4: Comparison of Modeled Air Concentrations with Health-Based Values (M17): Total Petroleum Hydrocarbons - 200 meter location

Cartridge 0.60 caliber Tracer M17 (M12) DODIC A574				
Compound	C _{chronic} (µg/m ³)	C _{chronic} (µg/m ³)	C _{chronic} (µg/m ³)	C _{chronic} (µg/m ³)
	Aliphatic: C≤8	Aliphatic: C>8	Aromatic: C≤8	Aromatic: C>8
Permanent Gases				
Pentane	3.44E-04	NA	NA	NA
Benzene	NA	NA	1.20E-01	NA
Toluene	NA	NA	1.36E-02	NA
Ethylbenzene	NA	NA	1.09E-03	NA
m&p-Xylene	NA	NA	2.07E-03	NA
o-Xylene	NA	NA	1.69E-03	NA
Styrene	NA	NA	NA	4.13E-03
1,2,4-Trimethylbenzene	NA	NA	NA	3.78E-04
Propylene	2.11E-02	NA	NA	NA
n-Hexane	7.72E-02	NA	NA	NA
naphthalene	NA	NA	NA	6.96E-03
naphthalene	NA	NA	NA	4.35E-03
acenaphthylene	NA	NA	NA	3.06E-04
Acenaphthene	NA	NA	NA	3.59E-05
fluorene	NA	NA	NA	1.27E-04
phenanthrene	NA	NA	NA	1.03E-04
anthracene	NA	NA	NA	1.72E-05
fluoranthene	NA	NA	NA	1.30E-04
Total (µg/m ³)	9.86E-02	0.00E+00	1.39E-01	1.65E-02
Derived Health-Based Screening Level	1.92E+04	1.04E+03	4.17E+02	2.09E+02
C _{chronic} /HBSL	5.14E-06	0.00E+00	3.33E-04	7.93E-05
>1?	no	no	no	no
Footnotes:				
>1? = Is the ratio greater than one?				
NA = Not Applicable because compound was not detected				
C _{chronic} = chronic averaged air Concentration				
HBSL = Health-Based Screening Level				

APPENDIX E

**FACT SHEET SUBMITTED TO THE
U.S. ARMY ENVIRONMENTAL CENTER**


U.S. Army Environmental Center

Training Munitions Fact Sheet

M17 .50 Caliber Tracer Cartridge

Department of Defense Identification Code: A571

Breathing air emissions from the M17 .50 caliber tracer cartridge will not impact the health of residents who live as close as 200 meters (656 feet) from the firing location.



To be fully prepared to protect our country, U.S. soldiers must train with many different weapons and munitions, including the M17 .50 caliber tracer cartridge. This training is important because it helps prepare our soldiers for a variety of combat situations. While the Army recognizes the value of such comprehensive training on our installations, we also work hard to ensure the safety and health of surrounding communities.

WILL BREATHING AIR EMISSIONS FROM THE M17 .50 CALIBER TRACER CARTRIDGE AFFECT MY HEALTH?

To answer this question, the U.S. Army tested the air emissions that are released when the M17 is fired. The information gathered during these tests was then analyzed to determine if there would be a potential for health effects from inhalation to residents who live near training areas. Study results, generated using conservative methods, showed that offsite residents breathing air as close as 200 meters (656 feet or about the length of two football fields) from the firing location are safe from these emissions. If offsite residents are located less than 200 meters from the firing locations, a more site-specific evaluation would be necessary. It should be noted that at most locations, training areas are at least 1,000 meters (over half a mile) away from populated areas and the distance to firing locations may be even farther.

HOW WAS THE STUDY CONDUCTED?

To gather data for this study, the M17 was fired from the M2 machine gun in a test chamber. The air in the chamber was then tested to identify the types and amounts of substances released. About 300 different substances were looked for during this part of the study.

This information was then used in an U.S. Environmental Protection Agency (USEPA) approved air model (a computer program that allows estimation of air concentrations) to determine the amount of each substance to which someone

living near a training site might be exposed. Downwind concentrations were estimated based on a typical use scenario for the M17 during training exercises. Since this study did not look at any one specific training area, the assumptions used in the model would, in most cases, predict higher downwind air concentrations than those expected at an actual training site.

These estimated air concentrations were then compared to screening levels established by the U.S. Environmental Protection Agency and other federal agencies. If the air concentrations are less than these screening levels, they are considered safe for the general population, including sensitive people such as the sick, elderly, and children.

WHAT ARE THE STUDY LIMITATIONS?

Many steps were taken to ensure that the results of this study are protective of residents who live near training facilities. However, as with any study, this study has limitations. For example, the study does not consider exposure to other types of munitions that could also be used during the same training event. Due to these limitations, conservative model conditions were used to ensure the protection of public health from breathing M17 air emissions.

WHAT EXACTLY IS THE M17 .50 CALIBER TRACER CARTRIDGE?

The M17 is a tracer cartridge used to track the path of a bullet. When fired at night, the tracer leaves a visible trail to show the direction in which the bullet is traveling. The M17 has a brass cartridge case and a bullet consisting of a brass jacket and steel core. The cartridge contains a propelling charge that is made up mostly of nitrocellulose. Nitrocellulose is commonly used in furniture lacquers, printing inks, nail polish, and as a primary ingredient in smokeless propellants for military and commercial use. Each M17 cartridge is about the length of a soda can and can be identified by its brown tip.

WHERE CAN I GET MORE INFORMATION?

For more information on the M17 or other military munitions, please call the Army Environmental Hotline at 1-800-USA-3845, visit our Web site at www.aec.army.mil, or e-mail t2hotline@aec.apgea.army.mil.